

**ANNEX I**

**SUMMARY OF PRODUCT CHARACTERISTICS**

## **1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 50 micrograms powder and solvent for solution for injection

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each vial of PegIntron, powder for solution for injection contains 50 micrograms of peginterferon alfa-2b as measured on a protein basis.

Each vial provides 50 micrograms/0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

The active substance is a covalent conjugate of recombinant interferon alfa-2b\* with monomethoxy polyethylene glycol. The potency of this product should not be compared to that of another pegylated or non-pegylated protein of the same therapeutic class. For more information, see section 5.1.

\*produced by rDNA technology in *E.coli* cells harbouring a genetically engineered plasmid hybrid encompassing an interferon alfa-2b gene from human leukocytes

For a full list of excipients, see section 6.1. PegIntron contains 40 mg of sucrose per 0.5 ml.

## **3. PHARMACEUTICAL FORM**

Powder and solvent for solution for injection

White powder.

Clear and colourless solvent.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

PegIntron is indicated for the treatment of adult patients with chronic hepatitis C who have elevated transaminases without liver decompensation and who are positive for serum HCV-RNA or anti-HCV, including naïve patients with clinically stable HIV co-infection (see section 4.4).

The best way to use PegIntron in this indication is in combination with ribavirin.

This combination is indicated in naïve patients including patients with clinically stable HIV co-infection. This combination of PegIntron with ribavirin is also indicated in patients who have previously responded (with normalisation of ALT at the end of treatment) to interferon alpha monotherapy but who have subsequently relapsed.

Interferon monotherapy, including PegIntron, is indicated mainly in case of intolerance or contraindication to ribavirin.

Please refer also to the ribavirin Summary of Product Characteristics (SPC) when PegIntron is to be used in combination with ribavirin.

### **4.2 Posology and method of administration**

Treatment should be initiated and monitored only by a physician experienced in the management of patients with hepatitis C.

#### **Dose to be administered**

PegIntron should be administered as a once weekly subcutaneous injection. The dose administered depends on whether it is used in combination with ribavirin or as monotherapy.

## Combination therapy

PegIntron 1.5 micrograms/kg/week in combination with ribavirin capsules.

The intended dose of 1.5 µg/kg of PegIntron to be used in combination with ribavirin may be delivered in weight categories with the pen/vial strengths according to **Table 1**. Ribavirin capsules are to be administered orally each day in two divided doses with food (morning and evening).

**Table 1 - Dosing for Combination Therapy**

Body Weight (kg)	PegIntron		Ribavirin Capsules	
	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Total Daily Dose (mg)	Number of Capsules (200 mg)
< 40	50	0.5	800	4 <sup>a</sup>
40-50	80	0.4	800	4 <sup>a</sup>
51-64	80	0.5	800	4 <sup>a</sup>
65-75	100	0.5	1,000	5 <sup>b</sup>
76-85	120	0.5	1,000	5 <sup>b</sup>
> 85	150	0.5	1,200	6 <sup>c</sup>

a: 2 morning, 2 evening

b: 2 morning, 3 evening

c: 3 morning, 3 evening

## Duration of treatment

**Predictability of sustained virological response:** Patients infected with virus genotype 1 who fail to achieve virological response at Week 12 are highly unlikely to become sustained virological responders (see also section 5.1).

- **Genotype 1:** For patients who exhibit virological response at week 12, treatment should be continued for another nine month period (i.e., a total of 48 weeks).  
In the subset of patients with genotype 1 infection and low viral load (< 600,000 IU/ml) who become HCV-RNA negative at treatment week 4 and remain HCV-RNA negative at week 24, the treatment could either be stopped after this 24 week treatment course or pursued for an additional 24 weeks (i.e. overall 48 weeks treatment duration). However, an overall 24 weeks treatment duration may be associated with a higher risk of relapse than a 48 weeks treatment duration (see section 5.1).
- **Genotypes 2 or 3:** It is recommended that all patients be treated for 24 weeks, except for HCV/HIV co-infected patients who should receive 48 weeks of treatment.
- **Genotype 4:** In general, patients infected with genotype 4 are considered harder to treat and limited study data (n=66) indicate they are compatible with a duration of treatment as for genotype 1.

## HCV/HIV Co-infection

The recommended duration of dosing for HCV/HIV co-infected patients is 48 weeks, regardless of genotype.

### *Predictability of response and non-response in HCV/HIV Co-infection*

Early virological response by Week 12, defined as a 2 log viral load decrease or undetectable levels of HCV-RNA, has been shown to be predictive for sustained response. The negative predictive value for sustained response in HCV/HIV co-infected patients treated with PegIntron in combination with ribavirin was 99 % (67/68; Study 1) (see section 5.1). A positive predictive value of 50 % (52/104; Study 1) was observed for HCV/HIV co-infected patients receiving combination therapy.

### PegIntron monotherapy

As monotherapy the PegIntron regimen is 0.5 or 1.0 microgram/kg/week. The lowest vial or pen strength available is 50 µg/0.5 ml; therefore for patients prescribed 0.5 µg/kg/week, doses must be adjusted by volume as shown in **Table 2**. For the 1.0 µg/kg dose, similar volume adjustments can be made or alternate vial strengths can be used as shown in **Table 2**. PegIntron monotherapy was not studied in HCV/HIV co-infected patients.

**Table 2- Monotherapy Dosing**

	0.5 µg/kg		1.0 µg/kg	
Body Weight (kg)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)
30-35	50*	0.15	50	0.3
36-45	50*	0.2	50	0.4
46-56	50*	0.25	50	0.5
57-72	50	0.3	80	0.4
73-88	50	0.4	80	0.5
89-106	50	0.5	100	0.5
> 106**	80	0.4	120	0.5
* Must use vial. Minimum delivery for pen is 0.3 ml.				
** For patients > 120 kg, use 80 µg/0.5 ml vial				

### Duration of treatment

For patients who exhibit virological response at Week 12, treatment should be continued for at least another three-month period (i.e., a total of six months). The decision to extend therapy to one year of treatment should be based on prognostic factors (e.g., genotype, age > 40 years, male gender, bridging fibrosis).

### Dose modification for all patients

If severe adverse reactions or laboratory abnormalities develop during treatment with PegIntron monotherapy or PegIntron in combination with ribavirin, modify the dosages of each product as appropriate, until the adverse reactions abate. As adherence might be of importance for outcome of therapy, the dose should be kept as close as possible to the recommended standard dose. Guidelines were developed in clinical trials for dose modification.

### Combination Therapy Dose Reduction Guidelines

<b>Table 2a</b> Dose modification guidelines for combination therapy (with ribavirin)			
Laboratory values	Reduce only ribavirin dose to 600 mg/day* if:	Reduce only PegIntron dose to one-half dose if:	Discontinue combination therapy if:
Haemoglobin	< 10 g/dl	-	< 8.5 g/dl
Haemoglobin in: Patients with history of stable cardiac disease	≥ 2 g/dl decrease in haemoglobin during any four week period during treatment (permanent dose reduction)		< 12 g/dl after four weeks of dose reduction
White blood cells	-	< 1.5 x 10 <sup>9</sup> /l	< 1.0 x 10 <sup>9</sup> /l
Neutrophils	-	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	-	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l
Bilirubin – direct	-	-	2.5 x ULN**

Bilirubin - indirect	> 5 mg/dl	-	> 4 mg/dl (for > 4 weeks)
Creatinine	-	-	> 2.0 mg/dl
ALT/AST	-	-	2 x baseline and > 10 x ULN**

\* Patients whose dose of ribavirin is reduced to 600 mg daily receive one 200 mg capsule in the morning and two 200 mg capsules in the evening.

\*\* Upper limit of normal

Dose reduction of PegIntron may be accomplished by either reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 2b**.

<b>Table 2b – Reduced PegIntron Dosing for Combination Therapy</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5 ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
< 40	25	50*	0.25	25
40-50	32	50	0.3	30
51-64	40	50	0.4	40
65-75	50	50	0.5	50
76-85	60	80	0.4	64
> 85	75	100	0.4	80
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### **PegIntron Monotherapy Dose Reduction Guidelines**

Dose modification guidelines for patients who use PegIntron monotherapy are shown in **Table 3a**.

<b>Table 3a Dose modification guidelines for PegIntron monotherapy</b>		
<b>Laboratory values</b>	<b>Reduce PegIntron to one-half dose if:</b>	<b>Discontinue PegIntron if:</b>
Neutrophils	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l

Dose reduction for patients who use 0.5 µg/kg PegIntron monotherapy must be accomplished by reducing the prescribed volume by one-half. The 50 µg/0.5 ml vial must be used if necessary since the pen can only deliver a minimum volume of 0.3 ml.

For patients who use 1.0 µg/kg PegIntron monotherapy, dose reduction may be accomplished by reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 3b**.

<b>Table 3b – Reduced PegIntron Dose for the 1.0 µg/kg Monotherapy Regimen</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
30-35	15	50*	0.15	15
36-45	20	50*	0.20	20
46-56	25	50*	0.25	25
57-72	32	50	0.3	30
73-89	40	50	0.4	40
90-106	50	50	0.5	50
> 106	60	80	0.4	64
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### Special populations

**Use in renal impairment: *Monotherapy*:** PegIntron should be used with caution in patients with moderate to severe renal impairment. In patients with moderate renal dysfunction (creatinine clearance 30-50 ml/minute), the starting dose of PegIntron should be reduced by 25 %. Patients with severe renal dysfunction (creatinine clearance 15-29 ml/minute) should have the starting dose of PegIntron reduced by 50 %. Data are not available for the use of PegIntron in patients with creatinine clearance < 15 ml/minute (see section 5.2). Patients with severe renal impairment, including those on hemodialysis, should be closely monitored. If renal function decreases during treatment, PegIntron therapy should be discontinued.

***Combination therapy*:** Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see ribavirin SPC). When administered in combination with ribavirin, subjects with impaired renal function should be more carefully monitored with respect to the development of anaemia.

**Use in hepatic impairment:** The safety and efficacy of PegIntron therapy has not been evaluated in patients with severe hepatic dysfunction, therefore PegIntron must not be used for these patients.

**Use in the elderly (≥ 65 years of age):** There are no apparent age-related effects on the pharmacokinetics of PegIntron. Data from elderly patients treated with a single dose of PegIntron suggest no alteration in PegIntron dose is necessary based on age (see section 5.2).

**Use in patients under the age of 18 years:** There is no experience in children (see section 5.2).

### 4.3 Contraindications

- Hypersensitivity to the active substance or to any interferon or to any of the excipients;
- A history of severe pre-existing cardiac disease, including unstable or uncontrolled cardiac disease in the previous six months (see section 4.4);
- Severe, debilitating medical conditions;
- Autoimmune hepatitis or a history of autoimmune disease;
- Severe hepatic dysfunction or decompensated cirrhosis of the liver;
- Pre-existing thyroid disease unless it can be controlled with conventional treatment;
- Epilepsy and/or compromised central nervous system (CNS) function;
- Initiation of PegIntron is contraindicated in HCV/HIV patients with cirrhosis and a Child-Pugh score ≥ 6.

*Combination therapy with ribavirin:* Also see ribavirin Summary of the Product Characteristics (SPC) if PegIntron is to be administered in combination with ribavirin in patients with chronic hepatitis C.

#### 4.4 Special warnings and precautions for use

**Psychiatric and Central Nervous System (CNS):** Severe CNS effects, particularly depression, suicidal ideation and attempted suicide have been observed in some patients during PegIntron therapy, and even after treatment discontinuation mainly during the 6-month follow-up period. Other CNS effects including aggressive behaviour (sometimes directed against others), confusion and alterations of mental status have been observed with alpha interferons. Patients should be closely monitored for any signs or symptoms of psychiatric disorders. If such symptoms appear, the potential seriousness of these undesirable effects must be borne in mind by the prescribing physician and the need for adequate therapeutic management should be considered. If psychiatric symptoms persist or worsen, or suicidal ideation is identified, it is recommended that treatment with PegIntron be discontinued, and the patient followed, with psychiatric intervention as appropriate.

*Patients with existence of, or history of severe psychiatric conditions:* If treatment with peginterferon alfa-2b is judged necessary in patients with existence or history of severe psychiatric conditions, this should only be initiated after having ensured appropriate individualised diagnostic and therapeutic management of the psychiatric condition.

There is no experience with PegIntron in combination with ribavirin in patients who have relapsed after interferon alpha + ribavirin therapy.

All patients in the chronic hepatitis C studies had a liver biopsy before inclusion, but in certain cases (i.e. patients with genotype 2 and 3), treatment may be possible without histological confirmation. Current treatment guidelines should be consulted as to whether a liver biopsy is needed prior to commencing treatment.

**Acute hypersensitivity:** Acute hypersensitivity reactions (e.g., urticaria, angioedema, bronchoconstriction, anaphylaxis) have been observed rarely during interferon alfa-2b therapy. If such a reaction develops during treatment with PegIntron, discontinue treatment and institute appropriate medical therapy immediately. Transient rashes do not necessitate interruption of treatment.

**Cardiovascular system:** As with interferon alfa-2b, patients with a history of congestive heart failure, myocardial infarction and/or previous or current arrhythmic disorders, receiving PegIntron therapy require close monitoring. It is recommended that patients who have pre-existing cardiac abnormalities have electrocardiograms taken prior to and during the course of treatment. Cardiac arrhythmias (primarily supraventricular) usually respond to conventional therapy but may require discontinuation of PegIntron therapy.

More significant obtundation and coma, including cases of encephalopathy, have been observed in some patients, usually elderly, treated at higher doses for oncology indications. While these effects are generally reversible, in a few patients full resolution took up to three weeks. Very rarely, seizures have occurred with high doses of interferon alpha.

**Liver function:** As with all interferons, discontinue treatment with PegIntron in patients who develop prolongation of coagulation markers which might indicate liver decompensation.

**Fever:** While fever may be associated with the flu-like syndrome reported commonly during interferon therapy, other causes of persistent fever must be ruled out.

**Hydration:** Adequate hydration must be maintained in patients undergoing PegIntron therapy since hypotension related to fluid depletion has been seen in some patients treated with alpha interferons. Fluid replacement may be necessary.

**Pulmonary changes:** Pulmonary infiltrates, pneumonitis, and pneumonia, occasionally resulting in fatality, have been observed rarely in interferon alpha treated patients. Any patient developing fever,

cough, dyspnea or other respiratory symptoms must have a chest X-ray taken. If the chest X-ray shows pulmonary infiltrates or there is evidence of pulmonary function impairment, the patient is to be monitored closely, and, if appropriate, discontinue interferon alpha. Prompt discontinuation of interferon alpha administration and treatment with corticosteroids appear to be associated with resolution of pulmonary adverse events.

**Autoimmune disease:** The development of auto-antibodies and autoimmune disorders has been reported during treatment with alpha interferons. Patients predisposed to the development of autoimmune disorders may be at increased risk. Patients with signs or symptoms compatible with autoimmune disorders should be evaluated carefully, and the benefit-risk of continued interferon therapy should be reassessed (see also section 4.4 Thyroid changes and 4.8).

**Ocular changes:** Ophthalmologic disorders, including retinal haemorrhages, cotton wool spots, and retinal artery or vein obstruction have been reported in rare instances after treatment with alpha interferons (see section 4.8). All patients should have a baseline eye examination. Any patient complaining of ocular symptoms, including loss of visual acuity or visual field must have a prompt and complete eye examination. Periodic visual examinations are recommended during PegIntron therapy, particularly in patients with disorders that may be associated with retinopathy, such as diabetes mellitus or hypertension. Discontinuation of PegIntron should be considered in patients who develop new or worsening ophthalmological disorders.

**Thyroid changes:** Infrequently, patients treated for chronic hepatitis C with interferon alpha have developed thyroid abnormalities, either hypothyroidism or hyperthyroidism. Determine thyroid stimulating hormone (TSH) levels if, during the course of therapy, a patient develops symptoms consistent with possible thyroid dysfunction. In the presence of thyroid dysfunction, PegIntron treatment may be continued if TSH levels can be maintained in the normal range by medication.

**Metabolic disturbances:** Hypertriglyceridemia and aggravation of hypertriglyceridemia, sometimes severe, have been observed. Monitoring of lipid levels is, therefore, recommended.

#### **HCV/HIV Co-infection**

Mitochondrial toxicity and lactic acidosis:

Patients co-infected with HIV and receiving Highly Active Anti-Retroviral Therapy (HAART) may be at increased risk of developing lactic acidosis. Caution should be used when adding PegIntron and ribavirin to HAART therapy (see ribavirin SPC).

Moreover, patients treated with PegIntron and ribavirin combination therapy and zidovudine are at increased risk of developing anaemia.

Hepatic decompensation in HCV/HIV co-infected patients with advanced cirrhosis:

Co-infected patients with advanced cirrhosis receiving HAART may be at increased risk of hepatic decompensation and death. Adding treatment with alfa interferons alone or in combination with ribavirin may increase the risk in this patient subset. Other baseline factors in co-infected patients that may be associated with a higher risk of hepatic decompensation include treatment with didanosine and elevated bilirubin serum concentration.

Co-infected patients receiving both antiretroviral (ARV) and anti-hepatitis treatment should be closely monitored, assessing their Child-Pugh score during treatment. Patients progressing to hepatic decompensation should have their anti-hepatitis treatment immediately discontinued and the ARV treatment reassessed.

Haematological abnormalities in HCV/HIV co-infected patients:

HCV/HIV co-infected patients receiving peginterferon alfa-2b/ribavirin treatment and HAART may be at increased risk to develop haematological abnormalities (as neutropenia, thrombocytopenia and anaemia) compared to HCV mono-infected patients. Although, the majority of them could be managed by dose reduction, close monitoring of haematological parameters should be undertaken in this population of patients (see section 4.2 and below “Laboratory tests” and section 4.8).

Patients with low CD4 counts:



In patients co-infected with HCV/HIV, limited efficacy and safety data (N = 25) are available in subjects with CD4 counts less than 200 cells/ $\mu$ l. Caution is therefore warranted in the treatment of patients with low CD4 counts.

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron and ribavirin.

**Dental and periodontal disorders:** Dental and periodontal disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and mucous membranes of the mouth during long-term treatment with the combination of PegIntron and ribavirin. Patients should brush their teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If this reaction occurs, they should be advised to rinse out their mouth thoroughly afterwards.

**Organ transplant recipients:** The safety and efficacy of PegIntron alone or in combination with ribavirin for the treatment of hepatitis C in liver or other organ transplant recipients have not been studied. Preliminary data indicates that interferon alpha therapy may be associated with an increased rate of kidney graft rejection. Liver graft rejection has also been reported.

**Other:** Due to reports of interferon alpha exacerbating pre-existing psoriatic disease and sarcoidosis, use of PegIntron in patients with psoriasis or sarcoidosis is recommended only if the potential benefit justifies the potential risk.

**Laboratory tests:** Standard haematologic tests, blood chemistry and a test of thyroid function must be conducted in all patients prior to initiating therapy. Acceptable baseline values that may be considered as a guideline prior to initiation of PegIntron therapy are:

- |                    |                              |
|--------------------|------------------------------|
| • Platelets        | $\geq 100,000/\text{mm}^3$   |
| • Neutrophil count | $\geq 1,500/\text{mm}^3$     |
| • TSH level        | must be within normal limits |

Laboratory evaluations are to be conducted at weeks 2 and 4 of therapy, and periodically thereafter as clinically appropriate.

**Important information about some of the ingredients of PegIntron:**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

Patients with rare hereditary problems of fructose intolerance, glucose galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Results from a multiple-dose probe study assessing P450 substrates in chronic hepatitis C patients receiving once weekly PegIntron (1.5  $\mu$ g/kg) for 4 weeks demonstrated an increase in activity of CYP2D6 and CYP2C8/9. No change in activity of CYP1A2, CYP3A4, or N-acetyltransferase was observed.

Caution should be used when administering peginterferon alfa-2b with medicines metabolised by CYP2D6 and CYP2C8/9, especially those with narrow therapeutic window, such as warfarin and phenytoin (CYP2C9) and flecainide (CYP2D6).

These findings may partly relate to improved metabolic capacity due to reduced hepatic inflammation in patients undergoing treatment with PegIntron. Caution is therefore advised when PegIntron

treatment is initiated for chronic hepatitis in patients treated with medication with a narrow therapeutic window and sensitive to mild metabolic impairment of the liver.

No pharmacokinetic interactions were noted between PegIntron and ribavirin in a multiple-dose pharmacokinetic study.

**Methadone:** In patients with chronic hepatitis C that were on stable methadone maintenance therapy and naïve to peginterferon alfa-2b, addition of 1.5 microgram/kg/week of PegIntron subcutaneously for 4 weeks increased R-methadone AUC by approximately 15 % (95 % CI for AUC ratio estimate 103 – 128 %). The clinical significance of this finding is unknown; however, patients should be monitored for signs and symptoms of increased sedative effect, as well as respiratory depression. Especially in patients on a high dose of methadone, the risk for QTc prolongation should be considered.

#### *HCV/HIV Co-infection*

Nucleoside analogs: ribavirin has been shown *in vitro* to inhibit phosphorylation of zidovudine and stavudine. The clinical significance of these findings is unknown. However, these *in vitro* findings raise the possibility that concurrent use of ribavirin with either zidovudine or stavudine might lead to increased HIV plasma viraemia. Therefore, it is recommended that plasma HIV RNA levels be closely monitored in patients treated with ribavirin concurrently with either of these two agents. If HIV RNA levels increase, the use of ribavirin concomitantly with reverse transcriptase inhibitors must be reviewed (see ribavirin SPC).

Use of nucleoside analogs, alone or in combination with other nucleosides, has resulted in lactic acidosis. Pharmacologically, ribavirin increases phosphorylated metabolites of purine nucleosides *in vitro*. This activity could potentiate the risk of lactic acidosis induced by purine nucleoside analogs (e.g. didanosine or abacavir). Co-administration of ribavirin and didanosine is not recommended. Reports of mitochondrial toxicity, in particular lactic acidosis and pancreatitis, of which some fatal, have been reported (see ribavirin SPC).

## **4.6 Pregnancy and lactation**

PegIntron is recommended for use in fertile women only when they are using effective contraception during the treatment.

There are no adequate data from the use of interferon alfa-2b in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3). Interferon alfa-2b has been shown to be abortifacient in primates. PegIntron is likely to also cause this effect.

The potential risk in humans is unknown. PegIntron is to be used during pregnancy only if the potential benefit justifies the potential risk to the foetus.

**Lactation:** It is not known whether the components of this medicinal product are excreted in human milk. Because of the potential for adverse reactions in breast-fed infants, breast-feeding should be discontinued prior to initiation of treatment.

#### *Combination therapy with ribavirin:*

Ribavirin causes serious birth defects when administered during pregnancy. Ribavirin therapy is contraindicated in women who are pregnant. Extreme care must be taken to avoid pregnancy in female patients or in partners of male patients taking PegIntron in combination with ribavirin. Females of childbearing potential and their partners must each use an effective contraceptive during treatment and for 4 months after treatment has been concluded. Male patients and their female partners must each use an effective contraceptive during treatment and for 7 months after treatment has been concluded (see Ribavirin SPC).

## **4.7 Effects on ability to drive and use machines**

Patients who develop fatigue, somnolence or confusion during treatment with PegIntron are cautioned to avoid driving or operating machinery.

#### 4.8 Undesirable effects

The safety of PegIntron is evaluated from data from two clinical trials: one with PegIntron monotherapy, one with PegIntron in combination with ribavirin. In both cases, patients were treated for one year.

**Table 4** describes the regimens and patient exposure for one year of treatment in patients with no previous exposure to interferon (interferon-naïve patients). Because of a significant overlap in the pattern of undesirable effects with PegIntron monotherapy, groups of patients have been brought together in **Table 5** to show the pattern of reported effects for all monotherapy groups.

<b>Table 4</b> Regimens and patient exposure		
<b>Treatment</b>	<b>Regimen</b>	<b>Number of patients treated for one year</b>
PegIntron + ribavirin	PegIntron (1.5 micrograms/kg/week) + ribavirin (> 10.6 mg/kg/day)	188
Interferon alfa-2b + ribavirin	Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day)	505
PegIntron monotherapy	PegIntron (0.5 microgram/kg/week)	315
	PegIntron (1.0 microgram/kg/week)	297
	PegIntron (1.5 micrograms/kg/week)	304

<b>Table 5</b> Undesirable effects very commonly reported in clinical trials > 10 % of patients in PegIntron + ribavirin group Very common (> 1/10)			
	<b>PegIntron + ribavirin</b>	<b>Interferon alfa- 2b + ribavirin</b>	<b>PegIntron monotherapy</b>
<b>Infections and infestations</b>			
Infection viral	10 %	5 %	4-5 %
<b>Metabolism and nutrition disorders</b>			
Weight decrease	30 %	19 %	8-18 %
<b>Psychiatric disorders</b>			
Depression	34 %	32 %	26 %
Irritability	32 %	34 %	19 %
Insomnia	37 %	41 %	16-19 %
Anxiety	14 %	14 %	8 %
Concentration impaired	18 %	21 %	9-10 %
Emotional lability	11 %	10 %	5 %
<b>Nervous system disorders</b>			
Headache	58 %	57 %	57-63 %
Mouth dry	10 %	8 %	4-8 %
<b>Respiratory, thoracic and mediastinal disorders</b>			
Pharyngitis	10 %	7 %	3 %
Coughing	14 %	11 %	4 %
Dyspnea	26 %	22 %	5 %
<b>Gastrointestinal disorders</b>			
Nausea	43 %	31 %	20-23 %
Anorexia	35 %	26 %	10-25 %
Diarrhoea	20 %	13 %	14-17 %
Abdominal pain	12 %	9 %	11 %
Vomiting	16 %	10 %	4-7 %
<b>Skin and subcutaneous tissue disorders</b>			
Alopecia	45 %	32 %	20-34 %
Pruritus	27 %	27 %	7-9 %
Skin dry	23 %	21 %	6-9 %
Rash	21 %	21 %	5-7 %
<b>Musculoskeletal and connective tissue disorders</b>			
Myalgia	49 %	49 %	46-60 %
Arthralgia	31 %	26 %	23-28 %
Musculoskeletal pain	15 %	11 %	11-13 %
<b>General disorders and administration site conditions</b>			
Injection site inflammation	20 %	17 %	39-44 %
Injection site reaction	54 %	36 %	7-9 %
Fatigue	56 %	59 %	43 %
Rigors	42 %	40 %	33-43 %
Fever	39 %	32 %	29-43 %
Flu-like symptoms	21 %	23 %	18-25 %
Asthenia	28 %	17 %	12-14 %
Dizziness	17 %	16 %	7-12 %

<b>Table 6</b> Undesirable effects commonly reported in clinical trials in patients treated with PegIntron + ribavirin or PegIntron monotherapy Common (> 1/100, < 1/10)		
<b>Body system</b>	<b>5-10%</b>	<b>1-&lt;5%</b>
Infections and infestations		otitis media, fungal infection, bacterial infection, herpes simplex
Blood and lymphatic system disorders	Anaemia, leukopaenia	Thrombocytopenia, lymphadenopathy
Endocrine disorders	hypothyroidism	hyperthyroidism
Metabolism and nutrition disorders		hyperuricemia, hypocalcemia, thirst
Psychiatric disorders	agitation, nervousness	aggressive behaviour, somnolence, behavior disorder, apathy, appetite increased, sleep disorder, dreaming abnormal, decreased libido
Nervous system disorders	paresthesia, increased sweating	hypoesthesia, hyperaesthesia, hypertonia, confusion, tremor, vertigo, migraine, ataxia, neuralgia
Eye disorders		blurred vision, conjunctivitis, lacrimal gland disorder, eye pain
Cardiac disorders		tachycardia, palpitation
Vascular disorders		hypotension, hypertension, syncope, flushing
Ear and labyrinth disorders		tinnitus, hearing impairment/loss
Respiratory, thoracic and mediastinal disorders		nonproductive cough, rhinitis, sinusitis, bronchitis, respiratory disorder, nasal congestion, rhinorrhea, dysphonia, epistaxis
Gastrointestinal disorders	dyspepsia	constipation, taste perversion, loose stools, stomatitis, ulcerative stomatitis, gingival bleeding, glossitis, flatulence, hemorrhoids, gastroesophageal reflux, gingivitis, dehydration
Hepatobiliary disorders		hepatomegaly, hyperbilirubinemia
Skin and subcutaneous tissue disorders		erythematous rash, eczema, photosensitivity reaction, maculopapular rash, abnormal hair texture, acne, dermatitis, furunculosis, nail disorder, psoriasis, urticaria, erythema, face or peripheral oedema
Musculoskeletal and connective tissue disorders		arthritis
Renal and urinary disorders		micturition frequency, urine abnormal
Reproductive system and breast disorders	menstrual disorder, menorrhagia	ovarian disorder, vaginal disorder, sexual dysfunction (not specified), impotence, breast pain, amenorrhoea, prostatitis
General disorders and administration site conditions	RUQ pain, malaise, chest pain	injection site pain

Most cases of neutropaenia and thrombocytopaenia were mild (WHO grades 1 or 2). There were some cases of more severe neutropenia in patients treated with the recommended doses of PegIntron in combination with ribavirin (WHO grade 3: 39 of 186 [21 %]; and WHO grade 4: 13 of 186 [7 %]).

In a clinical trial, approximately 1.2 % of patients treated with PegIntron or interferon alfa-2b in combination with ribavirin reported life-threatening psychiatric events during treatment. These events

included suicidal ideation and attempted suicide (see section 4.4). Following marketing, psychosis and hallucination have been reported rarely.

Rarely ( $> 1/10,000$ ,  $< 1/1,000$ ) or very rarely ( $< 1/10,000$ ) reported events with interferon alfa-2b, including PegIntron, include:

**Blood and lymphatic system disorders:**

Very rarely interferon alfa-2b or PegIntron used alone or in combination with ribavirin may be associated with aplastic anaemia. Pure red cell aplasia has been reported.

**Immune system disorders:**

very rarely: sarcoidosis or exacerbation of sarcoidosis

**Endocrine disorders:**

rarely: diabetes

**Nervous system disorders:**

rarely: seizure, peripheral neuropathy

very rarely: cerebrovascular ischaemia, cerebrovascular haemorrhage, encephalopathy

**Cardiac disorders:**

rarely: arrhythmia

very rarely: cardiac ischaemia, myocardial infarction

**Respiratory, thoracic and mediastinal disorders:**

very rarely: interstitial lung disease

**Gastrointestinal disorders:**

rarely: pancreatitis

very rarely: ulcerative and ischaemic colitis

**Skin and subcutaneous tissue disorders:**

very rarely: erythema multiforme, Stevens Johnson syndrome, toxic epidermal necrolysis, injection site necrosis

**Musculoskeletal and connective tissue disorders:**

rarely: rhabdomyolysis, myositis

**Renal and urinary disorders:**

rarely: renal insufficiency and renal failure

Cardiovascular (CVS) adverse events, particularly arrhythmia, appeared to be correlated mostly with pre-existing CVS disease and prior therapy with cardiotoxic agents (see section 4.4). Cardiomyopathy, that may be reversible upon discontinuation of interferon alpha, has been reported rarely in patients without prior evidence of cardiac disease.

Ophthalmological disorders that have been reported rarely with alpha interferons include retinopathies (including macular oedema), retinal haemorrhages, retinal artery or vein obstruction, cotton wool spots, loss of visual acuity or visual field, optic neuritis, and papilloedema (see section 4.4).

A wide variety of autoimmune and immune-mediated disorders have been reported with alpha interferons including thyroid disorders, systemic lupus erythematosus, rheumatoid arthritis (new or aggravated), idiopathic and thrombotic thrombocytopenic purpura, vasculitis, neuropathies including mononeuropathies (see also section 4.4, **Autoimmune disorders**).

**HCV/HIV co-infected patients**

For HCV/HIV co-infected patients receiving PegIntron in combination with ribavirin, other undesirable effects (that were not reported in mono-infected patients) which have been reported in the larger studies with a frequency  $> 5\%$  were: oral candidiasis (14 %), lipodystrophy acquired (13 %), CD4 lymphocytes decreased (8 %), appetite decreased (8 %), gamma-glutamyltransferase increased (9 %), back pain (5 %), blood amylase increased (6 %), blood lactic acid increased (5 %), cytolytic hepatitis (6 %), lipase increased (6 %) and pain in limb (6 %).

*Mitochondrial toxicity:*

Mitochondrial toxicity and lactic acidosis have been reported in HIV-positive patients receiving NRTI regimen and associated ribavirin for co-HCV infection (see section 4.4).

#### *Laboratory values for HCV/HIV co-infected patients:*

Although haematological toxicities of neutropenia, thrombocytopenia and anaemia occurred more frequently in HCV/HIV co-infected patients, the majority could be managed by dose modification and rarely required premature discontinuation of treatment (see section 4.4). Haematological abnormalities were more frequently reported in patients receiving PegIntron in combination with ribavirin when compared to patients receiving interferon alfa-2b in combination with ribavirin. In Study 1 (see section 5.1), decrease in absolute neutrophil count levels below 500 cells/mm<sup>3</sup> was observed in 4 % (8/194) of patients and decrease in platelets below 50,000/mm<sup>3</sup> was observed in 4 % (8/194) of patients receiving PegIntron in combination with ribavirin. Anaemia (hemoglobin < 9.4g/dl) was reported in 12% (23/194) of patients treated with PegIntron in combination with ribavirin.

#### *CD4 lymphocytes decrease:*

Treatment with PegIntron in combination with ribavirin was associated with decreases in absolute CD4+ cell counts within the first 4 weeks without a reduction in CD4+ cell percentage. The decrease in CD4+ cell counts was reversible upon dose reduction or cessation of therapy. The use of PegIntron in combination with ribavirin had no observable negative impact on the control of HIV viraemia during therapy or follow-up. Limited safety data (N= 25) are available in co-infected patients with CD4+ cell counts < 200/μl (see section 4.4).

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron in combination with ribavirin.

## **4.9 Overdose**

In clinical trials, cases of accidental overdose, at never more than twice the prescribed dose, were reported. There were no serious reactions. Undesirable effects resolved during continued administration of PegIntron.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Immunostimulants, Cytokines and immunomodulators, Interferons, Peginterferon alfa-2b, ATC code: L03A B10.

Recombinant interferon alfa-2b is covalently conjugated with monomethoxy polyethylene glycol at an average degree of substitution of 1 mole of polymer/mole of protein. The average molecular mass is approximately 31,300 daltons of which the protein moiety constitutes approximately 19,300.

#### **Interferon alfa-2b**

*In vitro* and *in vivo* studies suggest that the biological activity of PegIntron is derived from its interferon alfa-2b moiety.

Interferons exert their cellular activities by binding to specific membrane receptors on the cell surface. Studies with other interferons have demonstrated species specificity. However, certain monkey species, e.g., Rhesus monkeys, are susceptible to pharmacodynamic stimulation upon exposure to human type 1 interferons.

Once bound to the cell membrane, interferon initiates a complex sequence of intracellular events that include the induction of certain enzymes. It is thought that this process, at least in part, is responsible for the various cellular responses to interferon, including inhibition of virus replication in virus-infected cells, suppression of cell proliferation and such immunomodulating activities as enhancement of the phagocytic activity of macrophages and augmentation of the specific cytotoxicity of lymphocytes for target cells. Any or all of these activities may contribute to interferon's therapeutic effects.

Recombinant interferon alfa-2b also inhibits viral replication *in vitro* and *in vivo*. Although the exact antiviral mode of action of recombinant interferon alfa-2b is unknown, it appears to alter the host cell metabolism. This action inhibits viral replication or if replication occurs, the progeny virions are unable to leave the cell.

### **PegIntron**

PegIntron pharmacodynamics were assessed in a rising single-dose trial in healthy subjects by examining changes in oral temperature, concentrations of effector proteins such as serum neopterin and 2'5'-oligoadenylate synthetase (2'5'-OAS), as well as white cell and neutrophil counts. Subjects treated with PegIntron showed mild dose-related elevations in body temperature. Following single doses of PegIntron between 0.25 and 2.0 micrograms/kg/week, serum neopterin concentration was increased in a dose-related manner. Neutrophil and white cell count reductions at the end of week 4 correlated with the dose of PegIntron.

### **PegIntron clinical trials**

Two pivotal trials have been conducted, one (C/I97-010) with PegIntron monotherapy; the other (C/I98-580) with PegIntron in combination with ribavirin. Eligible patients for these trials had chronic hepatitis C confirmed by a positive HCV-RNA polymerase chain reaction (PCR) assay (> 30 IU/ml), a liver biopsy consistent with a histological diagnosis of chronic hepatitis with no other cause for the chronic hepatitis, and abnormal serum ALT.

In the PegIntron monotherapy trial, a total of 916 naïve chronic hepatitis C patients were treated with PegIntron (0.5, 1.0 or 1.5 micrograms/kg/week) for one year with a follow-up period of six months. In addition, 303 patients received interferon alfa-2b (3 million International Units [MIU] three times a week) as a comparator. This study showed that PegIntron was superior to interferon alfa-2b (**Table 8**).

In the PegIntron combination trial, 1,530 naïve patients were treated for one year with one of the following combination regimens:

- PegIntron (1.5 micrograms/kg/week) + ribavirin (800 mg/day), (n = 511).
- PegIntron (1.5 micrograms/kg/week for one month followed by 0.5 microgram/kg/week for 11 months) + ribavirin (1,000/1,200 mg/day), (n = 514).
- Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day) (n = 505).

In this trial, the combination of PegIntron (1.5 micrograms/kg/week) and ribavirin was significantly more effective than the combination of interferon alfa-2b and ribavirin (**Table 7**), particularly in patients infected with Genotype 1 (**Table 8**). Sustained response was assessed by the response rate six months after the cessation of treatment.

HCV genotype and baseline virus load are prognostic factors which are known to affect response rates. However, response rates in this trial were shown to be dependent also on the dose of ribavirin administered in combination with PegIntron or interferon alfa-2b. In those patients that received > 10.6 mg/kg ribavirin (800 mg dose in typical 75 kg patient), regardless of genotype or viral load, response rates were significantly higher than in those patients that received ≤ 10.6 mg/kg ribavirin (**Table 8**), while response rates in patients that received > 13.2 mg/kg ribavirin were even higher.

<b>Table 7</b> Sustained virological response (% patients HCV negative)							
	<b>PegIntron monotherapy</b>				<b>PegIntron + ribavirin</b>		
<b>Treatment regimen</b>	<b>P 1.5</b>	<b>P 1.0</b>	<b>P 0.5</b>	<b>I</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>Number of patients</b>	304	297	315	303	511	514	505
<b>Response at end of treatment</b>	49 %	41 %	33 %	24 %	65 %	56 %	54 %
<b>Sustained response</b>	<b>23 %*</b>	<b>25 %</b>	<b>18 %</b>	<b>12 %</b>	<b>54 %**</b>	<b>47 %</b>	<b>47 %</b>



P 1.5	PegIntron 1.5 micrograms/kg
P 1.0	PegIntron 1.0 microgram/kg
P 0.5	PegIntron 0.5 microgram/kg
I	Interferon alfa-2b 3 MIU
P 1.5/R	PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)
P 0.5/R	PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)
I/R	Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)
*	p < 0.001 P 1.5 vs. I
**	p = 0.0143 P 1.5/R vs. I/R

<b>Table 8</b> Sustained response rates with PegIntron + ribavirin (by ribavirin dose, genotype and viral load)				
HCV Genotype	Ribavirin dose (mg/kg)	P 1.5/R	P 0.5/R	I/R
<b>All Genotypes</b>	<b>All</b>	<b>54 %</b>	<b>47 %</b>	<b>47 %</b>
	≤ 10.6	50 %	41 %	27 %
	> 10.6	61 %	48 %	47 %
<b>Genotype 1</b>	<b>All</b>	<b>42 %</b>	<b>34 %</b>	<b>33 %</b>
	≤ 10.6	38 %	25 %	20 %
	> 10.6	48 %	34 %	34 %
Genotype 1 ≤ 600,000 IU/ml	All	73 %	51 %	45 %
	≤ 10.6	74 %	25 %	33 %
	> 10.6	71 %	52 %	45 %
Genotype 1 > 600,000 IU/ml	All	30 %	27 %	29 %
	≤ 10.6	27 %	25 %	17 %
	> 10.6	37 %	27 %	29 %
<b>Genotype 2/3</b>	<b>All</b>	<b>82 %</b>	<b>80 %</b>	<b>79 %</b>
	≤ 10.6	79 %	73 %	50 %
	> 10.6	88 %	80 %	80 %

P 1.5/R	PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)
P 0.5/R	PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)
I/R	Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)

In the PegIntron monotherapy study, the Quality of Life was generally less affected by 0.5 microgram/kg of PegIntron than by either 1.0 microgram/kg of PegIntron once weekly or 3 MIU of interferon alfa-2b three times a week.

In a separate trial, 224 patients with genotype 2 or 3 received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with ribavirin 800 mg –1,400 mg p.o. for 6 months (based on body weight, only three patients weighing > 105 kg, received the 1,400 mg dose, which has not yet been validated) (**Table 9**). Twenty-four % had bridging fibrosis or cirrhosis (Knodell 3/4).

**Table 9. Virologic Response at End of Treatment, Sustained Virologic Response and Relapse by HCV Genotype and Viral Load\***

	PegIntron 1.5 µg/kg once weekly plus Ribavirin 800-1,400 mg/day		
	End of Treatment Response	Sustained Virologic Response	Relapse
All Subjects	94 % (211/224)	81 % (182/224)	12 % (27/224)
HCV 2 ≤ 600,000 IU/ml > 600,000 IU/ml	100 % (42/42)	93 % (39/42)	7 % (3/42)
	100 % (20/20)	95 % (19/20)	5 % (1/20)
	100 % (22/22)	91 % (20/22)	9 % (2/22)
HCV 3	93 % (169/182)	79 % (143/182)	14 % (24/166)

≤ 600,000 IU/ml	93 % (92/99)	86 % (85/99)	8 % (7/91)
> 600,000 IU/ml	93 % (77/83)	70 % (58/83)	23 % (17/75)

\* Any subject with an undetectable HCV-RNA level at the Follow-Up Week 12 visit and missing data at the Follow-Up Week 24 visit was considered a sustained responder. Any subject with missing data in and after the Follow-Up Week 12 window was considered to be a non-responder at Week 24 of follow-up.

The 6 month treatment duration in this trial was better tolerated than one year of treatment in the pivotal combination trial; for discontinuation 5 % vs. 14 %, for dose modification 18 % vs. 49 %.

In a non-comparative trial, 235 patients with genotype 1 and low viral load (< 600,000 IU/ml) received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with weight adjusted ribavirin. The overall sustained response rate after a 24-week treatment duration was 50 %. Forty-one percent of subjects (97/235) had nondetectable plasma HCV-RNA levels at Week 4 and Week 24 of therapy. In this subgroup, there was a 92 % (89/97) sustained virological response rate. The high sustained response rate in this subgroup of patients was identified in an interim analysis (n=49) and prospectively confirmed (n=48).

Limited historical data indicate that treatment for 48 weeks might be associated with a higher sustained response rate (11/11) and with a lower risk of relapse (0/11 as compared to 7/96 following 24 weeks of treatment).

### Predictability of sustained virological response

Virological response by week 12, defined as a 2-log viral load decrease or undetectable levels of HCV-RNA has been shown to be predictive for sustained response (**Table 10**).

<b>Table 10</b> Predictability of sustained response by viral response at week 12 and genotype*				
Treatment	Genotype	Viral response at week 12	Sustained response	Negative predictive value
PegIntron 1.5 + ribavirin (> 10.6 mg/kg) 48-week treatment	1	Yes 75 % (82/110)	71 % (58/82)	----
		No 25 % (28/110)	0 % (0/28)	100 %
PegIntron 1.5 + ribavirin 800-1,400 mg 24-week treatment	2 and 3	Yes 99 % (213/215)	83 % (177/213)	----
		No 1 % (2/215)	50 % (1/2)	50 %

\* reflects patients with 12 week data available

The negative predictive value for sustained response in patients treated with PegIntron in monotherapy was 98 %.

### HCV/HIV Co-infected patients

Two trials have been conducted in patients co-infected with HIV and HCV. The response to treatment in both of these trials is presented in **Table 11**. Study 1 (RIBAVIC; P01017) was a randomized, multicentre study which enrolled 412 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (1.5 µg/kg/week) plus ribavirin (800 mg/day) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800 mg/day) for 48 weeks with a follow-up period of 6 months. Study 2 (P02080) was a randomized, single centre study that enrolled 95 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (100 or 150 µg /week based on weight) plus ribavirin (800-1,200 mg/day based on weight) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800-1,200 mg/day based on weight). The duration of therapy was 48 weeks with a follow-up

period of 6 months except for patients infected with genotypes 2 or 3 and viral load < 800,000 IU/ml (Amplicor) who were treated for 24 weeks with a 6-month follow-up period.

<b>Table 11</b> Sustained virological response based on genotype after PegIntron in combination with Ribavirin in HCV/HIV Co-infected patients						
	<b>Study 1<sup>1</sup></b>			<b>Study 2<sup>2</sup></b>		
	PegIntron (1.5 µg/kg/week) + ribavirin (800 mg)	Interferon alfa-2b (3 MIU TIW) + ribavirin (800 mg)	p value <sup>a</sup>	PegIntron (100 or 150 <sup>c</sup> µg/week) + ribavirin (800-1,200 mg) <sup>d</sup>	Interferon alfa-2b (3 MIU TIW) + ribavirin (800-1,200 mg) <sup>d</sup>	p value <sup>b</sup>
All	27 % (56/205)	20 % (41/205)	0.047	44 % (23/52)	21 % (9/43)	0.017
Genotype 1, 4	17 % (21/125)	6 % (8/129)	0.006	38 % (12/32)	7 % (2/27)	0.007
Genotype 2, 3	44 % (35/80)	43 % (33/76)	0.88	53 % (10/19)	47 % (7/15)	0.730

MIU = million international units; TIW = three times a week.

a: p value based on Cochran-Mantel Haenszel Chi square test.

b: p value based on chi-square test.

c: subjects < 75 kg received 100 µg/week PegIntron and subjects ≥ 75 kg received 150 µg/week PegIntron.

d: ribavirin dosing was 800 mg for patients < 60 kg, 1,000 mg for patients 60-75 kg, and 1,200 mg for patients > 75 kg.

<sup>1</sup>Carrato F, Bani-Sadir F, Pol S et al. JAMA 2004; 292(23): 2839-2848.

<sup>2</sup>Laguno M, Murillas J, Blanco J et al. AIDS 2004; 18(13): F27-F36.

### *Histological response*

Liver biopsies were obtained before and after treatment in Study 1 and were available for 210 of the 412 subjects (51 %). Both the Metavir score and Ishak grade decreased among subjects treated with PegIntron in combination with ribavirin. This decline was significant among responders (-0.3 for Metavir and -1.2 for Ishak) and stable (-0.1 for Metavir and -0.2 for Ishak) among non-responders. In terms of activity, about one-third of sustained responders showed improvement and none showed worsening. There was no improvement in terms of fibrosis observed in this study. Steatosis was significantly improved in patients infected with HCV Genotype 3.

## **5.2 Pharmacokinetic properties**

PegIntron is a well characterized polyethylene glycol-modified (“pegylated”) derivative of interferon alfa-2b and is predominantly composed of monopegylated species. The plasma half-life of PegIntron is prolonged compared with non-pegylated interferon alfa-2b. PegIntron has a potential to depegylate to free interferon alfa-2b. The biologic activity of the pegylated isomers is qualitatively similar, but weaker than free interferon alfa-2b.

Following subcutaneous administration, maximal serum concentrations occur between 15-44 hours post-dose, and are sustained for up to 48-72 hours post-dose.

PegIntron C<sub>max</sub> and AUC measurements increase in a dose-related manner. Mean apparent volume of distribution is 0.99 l/kg.

Upon multiple dosing, there is an accumulation of immunoreactive interferons. There is, however, only a modest increase in biologic activity as measured by a bioassay.

Mean (SD) PegIntron elimination half-life is approximately 40 hours (13.3 hours), with apparent clearance of 22.0 ml/hr·kg. The mechanisms involved in clearance of interferons in man have not yet been fully elucidated. However, renal elimination may account for a minority (approximately 30 %) of PegIntron apparent clearance.

**Renal function:** Renal clearance appears to account for 30 % of total clearance of PegIntron. In a single dose study (1.0 microgram/kg) in patients with impaired renal function,  $C_{max}$ , AUC, and half-life increased in relation to the degree of renal impairment.

Following multiple dosing of PegIntron (1.0 microgram/kg subcutaneously administered every week for four weeks) the clearance of PegIntron is reduced by a mean of 17 % in patients with moderate renal impairment (creatinine clearance 30-49 ml/minute) and by a mean of 44 % in patients with severe renal impairment (creatinine clearance 15-29 ml/minute) compared to subjects with normal renal function. Based on single dose data, clearance was similar in patients with severe renal impairment not on dialysis and in patients who were receiving hemodialysis. The dose of PegIntron for monotherapy should be reduced in patients with moderate or severe renal impairment (see sections 4.2 and 4.4). Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see section 4.3).

Because of marked inter-subject variability in interferon pharmacokinetics, it is recommended that patients with severe renal impairment be closely monitored during treatment with PegIntron (see section 4.2)

**Hepatic function:** The pharmacokinetics of PegIntron have not been evaluated in patients with severe hepatic dysfunction.

**Elderly patients  $\geq$  65 years of age:** The pharmacokinetics of PegIntron following a single subcutaneous dose of 1.0 microgram/kg were not affected by age. The data suggest that no alteration in PegIntron dosage is necessary based on advancing age.

**Patients under the age of 18 years:** Specific pharmacokinetic evaluations have not been performed on these patients. PegIntron is indicated for the treatment of chronic hepatitis C only in patients 18 years of age or older.

**Interferon neutralising factors:** Interferon neutralising factor assays were performed on serum samples of patients who received PegIntron in the clinical trial. Interferon neutralising factors are antibodies which neutralise the antiviral activity of interferon. The clinical incidence of neutralising factors in patients who received PegIntron 0.5 micrograms/kg is 1.1 %.

### 5.3 Preclinical safety data

**PegIntron:** Adverse events not observed in clinical trials were not seen in toxicity studies in monkeys. These studies were limited to four weeks due to the appearance of anti-interferon antibodies in most monkeys.

Reproduction studies of PegIntron have not been performed. Interferon alfa-2b has been shown to be an abortifacient in primates. PegIntron is likely to also cause this effect. Effects on fertility have not been determined. It is not known whether the components of this medicinal product are excreted into experimental animal or human milk (see section 4.6 for relevant human data on pregnancy and lactation). PegIntron showed no genotoxic potential.

The relative non-toxicity of monomethoxy-polyethylene glycol (mPEG), which is liberated from PegIntron by metabolism *in vivo* has been demonstrated in preclinical acute and subchronic toxicity studies in rodents and monkeys, standard embryo-foetal development studies and in *in vitro* mutagenicity assays.

**PegIntron plus ribavirin:** When used in combination with ribavirin, PegIntron did not cause any effects not previously seen with either active substance alone. The major treatment-related change was a reversible, mild to moderate anaemia, the severity of which was greater than that produced by either active substance alone.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Powder for solution for injection:

Disodium phosphate, anhydrous,  
Sodium dihydrogen phosphate dihydrate,  
Sucrose,  
Polysorbate 80.

Solvent for parenteral use:

Water for injections.

### **6.2 Incompatibilities**

This medicinal product should only be reconstituted with the solvent provided (see section 6.6). In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

3 years

After reconstitution:

- Chemical and physical in-use stability has been demonstrated for 24 hours at 2°C - 8°C.
- From a microbiological point of view, the product is to be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C - 8°C.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C - 8°C).

### **6.5 Nature and contents of container**

The powder is contained in a 2 ml vial, Type I flint glass, with a butyl rubber stopper in an aluminium flip-off seal with a polypropylene bonnet. The solvent is presented in a 2 ml ampoule, Type I flint glass. PegIntron 50 micrograms is supplied as:

- 1 vial of powder for solution for injection and 1 ampoule of solvent for parenteral use;
- 1 vial of powder for solution for injection, 1 ampoule of solvent for parenteral use, 1 injection syringe, 2 injection needles and 1 cleansing swab;
- 4 vials of powder for solution for injection and 4 ampoules of solvent for parenteral use;
- 4 vials of powder for solution for injection, 4 ampoules of solvent for parenteral use, 4 injection syringes, 8 injection needles and 4 cleansing swabs;
- 6 vials of powder for solution for injection and 6 ampoules of solvent for parenteral use.
- 12 vials of powder for solution for injection, 12 ampoules of solvent for parenteral use, 12 injection syringes, 24 injection needles and 12 cleansing swabs.

Not all pack sizes may be marketed

### **6.6 Special precautions for disposal**

PegIntron is supplied as a powder of peginterferon alfa-2b at a strength of 50 micrograms for single use. Each vial must be reconstituted with 0.7 ml of water for injections for administration of up to 0.5 ml of solution. A small volume is lost during preparation of PegIntron for injection when the dose is measured and injected. Therefore, each vial contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection. The reconstituted solution has a

concentration of 50 micrograms/0.5 ml.

Using a sterilised injection syringe and injection needle, inject 0.7 ml of water for injections into the vial of PegIntron. Agitate gently to complete dissolution of powder. The appropriate dose can then be withdrawn with a sterilised injection syringe and injected. A complete set of instructions is provided in the Annex to the Package Leaflet.

As for all parenteral medicinal products, inspect visually the reconstituted solution prior to administration. The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. Discard any unused material.

## **7. MARKETING AUTHORISATION HOLDER**

SP Europe  
73, rue de Stalle  
B-1180 Bruxelles  
Belgium

## **8. MARKETING AUTHORISATION NUMBERS**

EU/1/00/131/001  
EU/1/00/131/002  
EU/1/00/131/003  
EU/1/00/131/004  
EU/1/00/131/005  
EU/1/00/131/026

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 25 May 2000  
Date of last renewal: 25 May 2005

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.emea.europa.eu/>

## **1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 80 micrograms powder and solvent for solution for injection

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each vial of PegIntron, powder for solution for injection contains 80 micrograms of peginterferon alfa-2b as measured on a protein basis.

Each vial provides 80 micrograms/0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

The active substance is a covalent conjugate of recombinant interferon alfa-2b\* with monomethoxy polyethylene glycol. The potency of this product should not be compared to that of another pegylated or non-pegylated protein of the same therapeutic class. For more information, see section 5.1.

\*produced by rDNA technology in *E.coli* cells harbouring a genetically engineered plasmid hybrid encompassing an interferon alfa-2b gene from human leukocytes

For a full list of excipients, see section 6.1. PegIntron contains 40 mg of sucrose per 0.5 ml.

## **3. PHARMACEUTICAL FORM**

Powder and solvent for solution for injection

White powder.

Clear and colourless solvent.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

PegIntron is indicated for the treatment of adult patients with chronic hepatitis C who have elevated transaminases without liver decompensation and who are positive for serum HCV-RNA or anti-HCV, including naïve patients with clinically stable HIV co-infection (see section 4.4).

The best way to use PegIntron in this indication is in combination with ribavirin.

This combination is indicated in naïve patients including patients with clinically stable HIV co-infection. This combination of PegIntron with ribavirin is also indicated in patients who have previously responded (with normalisation of ALT at the end of treatment) to interferon alpha monotherapy but who have subsequently relapsed.

Interferon monotherapy, including PegIntron, is indicated mainly in case of intolerance or contraindication to ribavirin.

Please refer also to the ribavirin Summary of Product Characteristics (SPC) when PegIntron is to be used in combination with ribavirin.

### **4.2 Posology and method of administration**

Treatment should be initiated and monitored only by a physician experienced in the management of patients with hepatitis C.

#### **Dose to be administered**

PegIntron should be administered as a once weekly subcutaneous injection. The dose administered depends on whether it is used in combination with ribavirin or as monotherapy.

## Combination therapy

PegIntron 1.5 micrograms/kg/week in combination with ribavirin capsules.

The intended dose of 1.5 µg/kg of PegIntron to be used in combination with ribavirin may be delivered in weight categories with the pen/vial strengths according to **Table 1**. Ribavirin capsules are to be administered orally each day in two divided doses with food (morning and evening).

**Table 1 - Dosing for Combination Therapy**

Body Weight (kg)	PegIntron		Ribavirin Capsules	
	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Total Daily Dose (mg)	Number of Capsules (200 mg)
< 40	50	0.5	800	4 <sup>a</sup>
40-50	80	0.4	800	4 <sup>a</sup>
51-64	80	0.5	800	4 <sup>a</sup>
65-75	100	0.5	1,000	5 <sup>b</sup>
76-85	120	0.5	1,000	5 <sup>b</sup>
> 85	150	0.5	1,200	6 <sup>c</sup>

a: 2 morning, 2 evening

b: 2 morning, 3 evening

c: 3 morning, 3 evening

## Duration of treatment

**Predictability of sustained virological response:** Patients infected with virus genotype 1 who fail to achieve virological response at Week 12 are highly unlikely to become sustained virological responders (see also section 5.1).

- **Genotype 1:** For patients who exhibit virological response at week 12, treatment should be continued for another nine month period (i.e., a total of 48 weeks).  
In the subset of patients with genotype 1 infection and low viral load (< 600,000 IU/ml) who become HCV-RNA negative at treatment week 4 and remain HCV-RNA negative at week 24, the treatment could either be stopped after this 24 week treatment course or pursued for an additional 24 weeks (i.e. overall 48 weeks treatment duration). However, an overall 24 weeks treatment duration may be associated with a higher risk of relapse than a 48 weeks treatment duration (see section 5.1).
- **Genotypes 2 or 3:** It is recommended that all patients be treated for 24 weeks, except for HCV/HIV co-infected patients who should receive 48 weeks of treatment.
- **Genotype 4:** In general, patients infected with genotype 4 are considered harder to treat and limited study data (n=66) indicate they are compatible with a duration of treatment as for genotype 1.

## HCV/HIV Co-infection

The recommended duration of dosing for HCV/HIV co-infected patients is 48 weeks, regardless of genotype.

### *Predictability of response and non-response in HCV/HIV Co-infection*

Early virological response by Week 12, defined as a 2 log viral load decrease or undetectable levels of HCV-RNA, has been shown to be predictive for sustained response. The negative predictive value for sustained response in HCV/HIV co-infected patients treated with PegIntron in combination with ribavirin was 99 % (67/68; Study 1) (see section 5.1). A positive predictive value of 50 % (52/104; Study 1) was observed for HCV/HIV co-infected patients receiving combination therapy.



### PegIntron monotherapy

As monotherapy the PegIntron regimen is 0.5 or 1.0 microgram/kg/week. The lowest vial or pen strength available is 50 µg/0.5 ml; therefore for patients prescribed 0.5 µg/kg/week, doses must be adjusted by volume as shown in **Table 2**. For the 1.0 µg/kg dose, similar volume adjustments can be made or alternate vial strengths can be used as shown in **Table 2**. PegIntron monotherapy was not studied in HCV/HIV co-infected patients.

**Table 2- Monotherapy Dosing**

	0.5 µg/kg		1.0 µg/kg	
Body Weight (kg)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)
30-35	50*	0.15	50	0.3
36-45	50*	0.2	50	0.4
46-56	50*	0.25	50	0.5
57-72	50	0.3	80	0.4
73-88	50	0.4	80	0.5
89-106	50	0.5	100	0.5
> 106**	80	0.4	120	0.5
* Must use vial. Minimum delivery for pen is 0.3 ml.				
** For patients > 120 kg, use 80 µg/0.5 ml vial				

### Duration of treatment

For patients who exhibit virological response at Week 12, treatment should be continued for at least another three-month period (i.e., a total of six months). The decision to extend therapy to one year of treatment should be based on prognostic factors (e.g., genotype, age > 40 years, male gender, bridging fibrosis).

### Dose modification for all patients

If severe adverse reactions or laboratory abnormalities develop during treatment with PegIntron monotherapy or PegIntron in combination with ribavirin, modify the dosages of each product as appropriate, until the adverse reactions abate. As adherence might be of importance for outcome of therapy, the dose should be kept as close as possible to the recommended standard dose. Guidelines were developed in clinical trials for dose modification.

### Combination Therapy Dose Reduction Guidelines

<b>Table 2a</b> Dose modification guidelines for combination therapy (with ribavirin)			
Laboratory values	Reduce only ribavirin dose to <u>600 mg/day</u> * if:	Reduce only PegIntron dose to one-half dose if:	Discontinue combination therapy if:
Haemoglobin	< 10 g/dl	-	< 8.5 g/dl
Haemoglobin in: Patients with history of stable cardiac disease	≥ 2 g/dl decrease in haemoglobin during any four week period during treatment (permanent dose reduction)		< 12 g/dl after four weeks of dose reduction
White blood cells	-	< 1.5 x 10 <sup>9</sup> /l	< 1.0 x 10 <sup>9</sup> /l
Neutrophils	-	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	-	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l
Bilirubin – direct	-	-	2.5 x ULN**

Bilirubin - indirect	> 5 mg/dl	-	> 4 mg/dl (for > 4 weeks)
Creatinine	-	-	> 2.0 mg/dl
ALT/AST	-	-	2 x baseline and > 10 x ULN**

\* Patients whose dose of ribavirin is reduced to 600 mg daily receive one 200 mg capsule in the morning and two 200 mg capsules in the evening.

\*\* Upper limit of normal

Dose reduction of PegIntron may be accomplished by either reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 2b**.

<b>Table 2b – Reduced PegIntron Dosing for Combination Therapy</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5 ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
< 40	25	50*	0.25	25
40-50	32	50	0.3	30
51-64	40	50	0.4	40
65-75	50	50	0.5	50
76-85	60	80	0.4	64
> 85	75	100	0.4	80
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### **PegIntron Monotherapy Dose Reduction Guidelines**

Dose modification guidelines for patients who use PegIntron monotherapy are shown in **Table 3a**.

<b>Table 3a Dose modification guidelines for PegIntron monotherapy</b>		
<b>Laboratory values</b>	<b>Reduce PegIntron to one-half dose if:</b>	<b>Discontinue PegIntron if:</b>
Neutrophils	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l

Dose reduction for patients who use 0.5 µg/kg PegIntron monotherapy must be accomplished by reducing the prescribed volume by one-half. The 50 µg/0.5 ml vial must be used if necessary since the pen can only deliver a minimum volume of 0.3 ml.

For patients who use 1.0 µg/kg PegIntron monotherapy, dose reduction may be accomplished by reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 3b**.

<b>Table 3b – Reduced PegIntron Dose for the 1.0 µg/kg Monotherapy Regimen</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
30-35	15	50*	0.15	15
36-45	20	50*	0.20	20
46-56	25	50*	0.25	25
57-72	32	50	0.3	30
73-89	40	50	0.4	40
90-106	50	50	0.5	50
> 106	60	80	0.4	64
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### Special populations

**Use in renal impairment:** *Monotherapy:* PegIntron should be used with caution in patients with moderate to severe renal impairment. In patients with moderate renal dysfunction (creatinine clearance 30-50 ml/minute), the starting dose of PegIntron should be reduced by 25 %. Patients with severe renal dysfunction (creatinine clearance 15-29 ml/minute) should have the starting dose of PegIntron reduced by 50 %. Data are not available for the use of PegIntron in patients with creatinine clearance < 15 ml/minute (see section 5.2). Patients with severe renal impairment, including those on hemodialysis, should be closely monitored. If renal function decreases during treatment, PegIntron therapy should be discontinued.

*Combination therapy:* Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see ribavirin SPC). When administered in combination with ribavirin, subjects with impaired renal function should be more carefully monitored with respect to the development of anaemia.

**Use in hepatic impairment:** The safety and efficacy of PegIntron therapy has not been evaluated in patients with severe hepatic dysfunction, therefore PegIntron must not be used for these patients.

**Use in the elderly (≥ 65 years of age):** There are no apparent age-related effects on the pharmacokinetics of PegIntron. Data from elderly patients treated with a single dose of PegIntron suggest no alteration in PegIntron dose is necessary based on age (see section 5.2).

**Use in patients under the age of 18 years:** There is no experience in children (see section 5.2).

### 4.3 Contraindications

- Hypersensitivity to the active substance or to any interferon or to any of the excipients;
- A history of severe pre-existing cardiac disease, including unstable or uncontrolled cardiac disease in the previous six months (see section 4.4);
- Severe, debilitating medical conditions;
- Autoimmune hepatitis or a history of autoimmune disease;
- Severe hepatic dysfunction or decompensated cirrhosis of the liver;
- Pre-existing thyroid disease unless it can be controlled with conventional treatment;
- Epilepsy and/or compromised central nervous system (CNS) function;
- Initiation of PegIntron is contraindicated in HCV/HIV patients with cirrhosis and a Child-Pugh score ≥ 6.

*Combination therapy with ribavirin:* Also see ribavirin Summary of the Product Characteristics (SPC) if PegIntron is to be administered in combination with ribavirin in patients with chronic hepatitis C.

#### 4.4 Special warnings and precautions for use

**Psychiatric and Central Nervous System (CNS):** Severe CNS effects, particularly depression, suicidal ideation and attempted suicide have been observed in some patients during PegIntron therapy, and even after treatment discontinuation mainly during the 6-month follow-up period. Other CNS effects including aggressive behaviour (sometimes directed against others), confusion and alterations of mental status have been observed with alpha interferons. Patients should be closely monitored for any signs or symptoms of psychiatric disorders. If such symptoms appear, the potential seriousness of these undesirable effects must be borne in mind by the prescribing physician and the need for adequate therapeutic management should be considered. If psychiatric symptoms persist or worsen, or suicidal ideation is identified, it is recommended that treatment with PegIntron be discontinued, and the patient followed, with psychiatric intervention as appropriate.

*Patients with existence of, or history of severe psychiatric conditions:* If treatment with peginterferon alfa-2b is judged necessary in patients with existence or history of severe psychiatric conditions, this should only be initiated after having ensured appropriate individualised diagnostic and therapeutic management of the psychiatric condition.

There is no experience with PegIntron in combination with ribavirin in patients who have relapsed after interferon alpha + ribavirin therapy.

All patients in the chronic hepatitis C studies had a liver biopsy before inclusion, but in certain cases (i.e. patients with genotype 2 and 3), treatment may be possible without histological confirmation. Current treatment guidelines should be consulted as to whether a liver biopsy is needed prior to commencing treatment.

**Acute hypersensitivity:** Acute hypersensitivity reactions (e.g., urticaria, angioedema, bronchoconstriction, anaphylaxis) have been observed rarely during interferon alfa-2b therapy. If such a reaction develops during treatment with PegIntron, discontinue treatment and institute appropriate medical therapy immediately. Transient rashes do not necessitate interruption of treatment.

**Cardiovascular system:** As with interferon alfa-2b, patients with a history of congestive heart failure, myocardial infarction and/or previous or current arrhythmic disorders, receiving PegIntron therapy require close monitoring. It is recommended that patients who have pre-existing cardiac abnormalities have electrocardiograms taken prior to and during the course of treatment. Cardiac arrhythmias (primarily supraventricular) usually respond to conventional therapy but may require discontinuation of PegIntron therapy.

More significant obtundation and coma, including cases of encephalopathy, have been observed in some patients, usually elderly, treated at higher doses for oncology indications. While these effects are generally reversible, in a few patients full resolution took up to three weeks. Very rarely, seizures have occurred with high doses of interferon alpha.

**Liver function:** As with all interferons, discontinue treatment with PegIntron in patients who develop prolongation of coagulation markers which might indicate liver decompensation.

**Fever:** While fever may be associated with the flu-like syndrome reported commonly during interferon therapy, other causes of persistent fever must be ruled out.

**Hydration:** Adequate hydration must be maintained in patients undergoing PegIntron therapy since hypotension related to fluid depletion has been seen in some patients treated with alpha interferons. Fluid replacement may be necessary.

**Pulmonary changes:** Pulmonary infiltrates, pneumonitis, and pneumonia, occasionally resulting in fatality, have been observed rarely in interferon alpha treated patients. Any patient developing fever,

cough, dyspnea or other respiratory symptoms must have a chest X-ray taken. If the chest X-ray shows pulmonary infiltrates or there is evidence of pulmonary function impairment, the patient is to be monitored closely, and, if appropriate, discontinue interferon alpha. Prompt discontinuation of interferon alpha administration and treatment with corticosteroids appear to be associated with resolution of pulmonary adverse events.

**Autoimmune disease:** The development of auto-antibodies and autoimmune disorders has been reported during treatment with alpha interferons. Patients predisposed to the development of autoimmune disorders may be at increased risk. Patients with signs or symptoms compatible with autoimmune disorders should be evaluated carefully, and the benefit-risk of continued interferon therapy should be reassessed (see also section 4.4 Thyroid changes and 4.8).

**Ocular changes:** Ophthalmologic disorders, including retinal haemorrhages, cotton wool spots, and retinal artery or vein obstruction have been reported in rare instances after treatment with alpha interferons (see section 4.8). All patients should have a baseline eye examination. Any patient complaining of ocular symptoms, including loss of visual acuity or visual field must have a prompt and complete eye examination. Periodic visual examinations are recommended during PegIntron therapy, particularly in patients with disorders that may be associated with retinopathy, such as diabetes mellitus or hypertension. Discontinuation of PegIntron should be considered in patients who develop new or worsening ophthalmological disorders.

**Thyroid changes:** Infrequently, patients treated for chronic hepatitis C with interferon alpha have developed thyroid abnormalities, either hypothyroidism or hyperthyroidism. Determine thyroid stimulating hormone (TSH) levels if, during the course of therapy, a patient develops symptoms consistent with possible thyroid dysfunction. In the presence of thyroid dysfunction, PegIntron treatment may be continued if TSH levels can be maintained in the normal range by medication.

**Metabolic disturbances:** Hypertriglyceridemia and aggravation of hypertriglyceridemia, sometimes severe, have been observed. Monitoring of lipid levels is, therefore, recommended.

#### **HCV/HIV Co-infection**

Mitochondrial toxicity and lactic acidosis:

Patients co-infected with HIV and receiving Highly Active Anti-Retroviral Therapy (HAART) may be at increased risk of developing lactic acidosis. Caution should be used when adding PegIntron and ribavirin to HAART therapy (see ribavirin SPC).

Moreover, patients treated with PegIntron and ribavirin combination therapy and zidovudine are at increased risk of developing anaemia.

Hepatic decompensation in HCV/HIV co-infected patients with advanced cirrhosis:

Co-infected patients with advanced cirrhosis receiving HAART may be at increased risk of hepatic decompensation and death. Adding treatment with alfa interferons alone or in combination with ribavirin may increase the risk in this patient subset. Other baseline factors in co-infected patients that may be associated with a higher risk of hepatic decompensation include treatment with didanosine and elevated bilirubin serum concentration.

Co-infected patients receiving both antiretroviral (ARV) and anti-hepatitis treatment should be closely monitored, assessing their Child-Pugh score during treatment. Patients progressing to hepatic decompensation should have their anti-hepatitis treatment immediately discontinued and the ARV treatment reassessed.

Haematological abnormalities in HCV/HIV co-infected patients:

HCV/HIV co-infected patients receiving peginterferon alfa-2b/ribavirin treatment and HAART may be at increased risk to develop haematological abnormalities (as neutropenia, thrombocytopenia and anaemia) compared to HCV mono-infected patients. Although, the majority of them could be managed by dose reduction, close monitoring of haematological parameters should be undertaken in this population of patients (see section 4.2 and below “Laboratory tests” and section 4.8).

Patients with low CD4 counts:

In patients co-infected with HCV/HIV, limited efficacy and safety data (N = 25) are available in subjects with CD4 counts less than 200 cells/ $\mu$ l. Caution is therefore warranted in the treatment of patients with low CD4 counts.

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron and ribavirin.

**Dental and periodontal disorders:** Dental and periodontal disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and mucous membranes of the mouth during long-term treatment with the combination of PegIntron and ribavirin. Patients should brush their teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If this reaction occurs, they should be advised to rinse out their mouth thoroughly afterwards.

**Organ transplant recipients:** The safety and efficacy of PegIntron alone or in combination with ribavirin for the treatment of hepatitis C in liver or other organ transplant recipients have not been studied. Preliminary data indicates that interferon alpha therapy may be associated with an increased rate of kidney graft rejection. Liver graft rejection has also been reported.

**Other:** Due to reports of interferon alpha exacerbating pre-existing psoriatic disease and sarcoidosis, use of PegIntron in patients with psoriasis or sarcoidosis is recommended only if the potential benefit justifies the potential risk.

**Laboratory tests:** Standard haematologic tests, blood chemistry and a test of thyroid function must be conducted in all patients prior to initiating therapy. Acceptable baseline values that may be considered as a guideline prior to initiation of PegIntron therapy are:

- |                    |                              |
|--------------------|------------------------------|
| • Platelets        | $\geq 100,000/\text{mm}^3$   |
| • Neutrophil count | $\geq 1,500/\text{mm}^3$     |
| • TSH level        | must be within normal limits |

Laboratory evaluations are to be conducted at weeks 2 and 4 of therapy, and periodically thereafter as clinically appropriate.

**Important information about some of the ingredients of PegIntron:**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

Patients with rare hereditary problems of fructose intolerance, glucose galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Results from a multiple-dose probe study assessing P450 substrates in chronic hepatitis C patients receiving once weekly PegIntron (1.5  $\mu$ g/kg) for 4 weeks demonstrated an increase in activity of CYP2D6 and CYP2C8/9. No change in activity of CYP1A2, CYP3A4, or N-acetyltransferase was observed.

Caution should be used when administering peginterferon alfa-2b with medicines metabolised by CYP2D6 and CYP2C8/9, especially those with narrow therapeutic window, such as warfarin and phenytoin (CYP2C9) and flecainide (CYP2D6).

These findings may partly relate to improved metabolic capacity due to reduced hepatic inflammation in patients undergoing treatment with PegIntron. Caution is therefore advised when PegIntron

treatment is initiated for chronic hepatitis in patients treated with medication with a narrow therapeutic window and sensitive to mild metabolic impairment of the liver.

No pharmacokinetic interactions were noted between PegIntron and ribavirin in a multiple-dose pharmacokinetic study.

**Methadone:** In patients with chronic hepatitis C that were on stable methadone maintenance therapy and naïve to peginterferon alfa-2b, addition of 1.5 microgram/kg/week of PegIntron subcutaneously for 4 weeks increased R-methadone AUC by approximately 15 % (95 % CI for AUC ratio estimate 103 – 128 %). The clinical significance of this finding is unknown; however, patients should be monitored for signs and symptoms of increased sedative effect, as well as respiratory depression. Especially in patients on a high dose of methadone, the risk for QTc prolongation should be considered.

#### *HCV/HIV Co-infection*

Nucleoside analogs: ribavirin has been shown *in vitro* to inhibit phosphorylation of zidovudine and stavudine. The clinical significance of these findings is unknown. However, these *in vitro* findings raise the possibility that concurrent use of ribavirin with either zidovudine or stavudine might lead to increased HIV plasma viraemia. Therefore, it is recommended that plasma HIV RNA levels be closely monitored in patients treated with ribavirin concurrently with either of these two agents. If HIV RNA levels increase, the use of ribavirin concomitantly with reverse transcriptase inhibitors must be reviewed (see ribavirin SPC).

Use of nucleoside analogs, alone or in combination with other nucleosides, has resulted in lactic acidosis. Pharmacologically, ribavirin increases phosphorylated metabolites of purine nucleosides *in vitro*. This activity could potentiate the risk of lactic acidosis induced by purine nucleoside analogs (e.g. didanosine or abacavir). Co-administration of ribavirin and didanosine is not recommended. Reports of mitochondrial toxicity, in particular lactic acidosis and pancreatitis, of which some fatal, have been reported (see ribavirin SPC).

## **4.6 Pregnancy and lactation**

PegIntron is recommended for use in fertile women only when they are using effective contraception during the treatment.

There are no adequate data from the use of interferon alfa-2b in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3). Interferon alfa-2b has been shown to be abortifacient in primates. PegIntron is likely to also cause this effect.

The potential risk in humans is unknown. PegIntron is to be used during pregnancy only if the potential benefit justifies the potential risk to the foetus

**Lactation:** It is not known whether the components of this medicinal product are excreted in human milk. Because of the potential for adverse reactions in breast-fed infants, breast-feeding should be discontinued prior to initiation of treatment.

#### *Combination therapy with ribavirin:*

Ribavirin causes serious birth defects when administered during pregnancy. Ribavirin therapy is contraindicated in women who are pregnant. Extreme care must be taken to avoid pregnancy in female patients or in partners of male patients taking PegIntron in combination with ribavirin. Females of childbearing potential and their partners must each use an effective contraceptive during treatment and for 4 months after treatment has been concluded. Male patients and their female partners must each use an effective contraceptive during treatment and for 7 months after treatment has been concluded (see Ribavirin SPC).

## **4.7 Effects on ability to drive and use machines**

Patients who develop fatigue, somnolence or confusion during treatment with PegIntron are cautioned to

avoid driving or operating machinery.

#### 4.8 Undesirable effects

The safety of PegIntron is evaluated from data from two clinical trials: one with PegIntron monotherapy, one with PegIntron in combination with ribavirin. In both cases, patients were treated for one year.

**Table 4** describes the regimens and patient exposure for one year of treatment in patients with no previous exposure to interferon (interferon-naïve patients). Because of a significant overlap in the pattern of undesirable effects with PegIntron monotherapy, groups of patients have been brought together in **Table 5** to show the pattern of reported effects for all monotherapy groups.

<b>Table 4</b> Regimens and patient exposure		
<b>Treatment</b>	<b>Regimen</b>	<b>Number of patients treated for one year</b>
PegIntron + ribavirin	PegIntron (1.5 micrograms/kg/week) + ribavirin (> 10.6 mg/kg/day)	188
Interferon alfa-2b + ribavirin	Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day)	505
PegIntron monotherapy	PegIntron (0.5 microgram/kg/week)	315
	PegIntron (1.0 microgram/kg/week)	297
	PegIntron (1.5 micrograms/kg/week)	304



<b>Table 5</b> Undesirable effects very commonly reported in clinical trials > 10 % of patients in PegIntron + ribavirin group Very common (> 1/10)			
	<b>PegIntron + ribavirin</b>	<b>Interferon alfa- 2b + ribavirin</b>	<b>PegIntron monotherapy</b>
<b>Infections and infestations</b>			
Infection viral	10 %	5 %	4-5 %
<b>Metabolism and nutrition disorders</b>			
Weight decrease	30 %	19 %	8-18 %
<b>Psychiatric disorders</b>			
Depression	34 %	32 %	26 %
Irritability	32 %	34 %	19 %
Insomnia	37 %	41 %	16-19 %
Anxiety	14 %	14 %	8 %
Concentration impaired	18 %	21 %	9-10 %
Emotional lability	11 %	10 %	5 %
<b>Nervous system disorders</b>			
Headache	58 %	57 %	57-63 %
Mouth dry	10 %	8 %	4-8 %
<b>Respiratory, thoracic and mediastinal disorders</b>			
Pharyngitis	10 %	7 %	3 %
Coughing	14 %	11 %	4 %
Dyspnea	26 %	22 %	5 %
<b>Gastrointestinal disorders</b>			
Nausea	43 %	31 %	20-23 %
Anorexia	35 %	26 %	10-25 %
Diarrhoea	20 %	13 %	14-17 %
Abdominal pain	12 %	9 %	11 %
Vomiting	16 %	10 %	4-7 %
<b>Skin and subcutaneous tissue disorders</b>			
Alopecia	45 %	32 %	20-34 %
Pruritus	27 %	27 %	7-9 %
Skin dry	23 %	21 %	6-9 %
Rash	21 %	21 %	5-7 %
<b>Musculoskeletal and connective tissue disorders</b>			
Myalgia	49 %	49 %	46-60 %
Arthralgia	31 %	26 %	23-28 %
Musculoskeletal pain	15 %	11 %	11-13 %
<b>General disorders and administration site conditions</b>			
Injection site inflammation	20 %	17 %	39-44 %
Injection site reaction	54 %	36 %	7-9 %
Fatigue	56 %	59 %	43 %
Rigors	42 %	40 %	33-43 %
Fever	39 %	32 %	29-43 %
Flu-like symptoms	21 %	23 %	18-25 %
Asthenia	28 %	17 %	12-14 %
Dizziness	17 %	16 %	7-12 %

<b>Table 6</b> Undesirable effects commonly reported in clinical trials in patients treated with PegIntron + ribavirin or PegIntron monotherapy Common (> 1/100, < 1/10)		
<b>Body system</b>	<b>5-10%</b>	<b>1-&lt;5%</b>
Infections and infestations		otitis media, fungal infection, bacterial infection, herpes simplex
Blood and lymphatic system disorders	Anaemia, leukopaenia	Thrombocytopenia, lymphadenopathy
Endocrine disorders	hypothyroidism	hyperthyroidism
Metabolism and nutrition disorders		hyperuricemia, hypocalcemia, thirst
Psychiatric disorders	agitation, nervousness	aggressive behaviour, somnolence, behavior disorder, apathy, appetite increased, sleep disorder, dreaming abnormal, decreased libido
Nervous system disorders	paresthesia, increased sweating	hypoesthesia, hyperaesthesia, hypertonia, confusion, tremor, vertigo, migraine, ataxia, neuralgia
Eye disorders		blurred vision, conjunctivitis, lacrimal gland disorder, eye pain
Cardiac disorders		tachycardia, palpitation
Vascular disorders		hypotension, hypertension, syncope, flushing
Ear and labyrinth disorders		tinnitus, hearing impairment/loss
Respiratory, thoracic and mediastinal disorders		nonproductive cough, rhinitis, sinusitis, bronchitis, respiratory disorder, nasal congestion, rhinorrhea, dysphonia, epistaxis
Gastrointestinal disorders	dyspepsia	constipation, taste perversion, loose stools, stomatitis, ulcerative stomatitis, gingival bleeding, glossitis, flatulence, hemorrhoids, gastroesophageal reflux, gingivitis, dehydration
Hepatobiliary disorders		hepatomegaly, hyperbilirubinemia
Skin and subcutaneous tissue disorders		erythematous rash, eczema, photosensitivity reaction, maculopapular rash, abnormal hair texture, acne, dermatitis, furunculosis, nail disorder, psoriasis, urticaria, erythema, face or peripheral oedema
Musculoskeletal and connective tissue disorders		arthritis
Renal and urinary disorders		micturition frequency, urine abnormal
Reproductive system and breast disorders	menstrual disorder, menorrhagia	ovarian disorder, vaginal disorder, sexual dysfunction (not specified), impotence, breast pain, amenorrhoea, prostatitis
General disorders and administration site conditions	RUQ pain, malaise, chest pain	injection site pain

Most cases of neutropaenia and thrombocytopaenia were mild (WHO grades 1 or 2). There were some cases of more severe neutropenia in patients treated with the recommended doses of PegIntron in combination with ribavirin (WHO grade 3: 39 of 186 [21 %]; and WHO grade 4: 13 of 186 [7 %]).

In a clinical trial, approximately 1.2 % of patients treated with PegIntron or interferon alfa-2b in combination with ribavirin reported life-threatening psychiatric events during treatment. These events included suicidal ideation and attempted suicide (see section 4.4). Following marketing, psychosis and hallucination have been reported rarely.

Rarely ( $> 1/10,000$ ,  $< 1/1,000$ ) or very rarely ( $< 1/10,000$ ) reported events with interferon alfa-2b, including PegIntron, include:

**Blood and lymphatic system disorders:**

Very rarely interferon alfa-2b or PegIntron used alone or in combination with ribavirin may be associated with aplastic anaemia. Pure red cell aplasia has been reported.

**Immune system disorders:**

very rarely: sarcoidosis or exacerbation of sarcoidosis

**Endocrine disorders:**

rarely: diabetes

**Nervous system disorders:**

rarely: seizure, peripheral neuropathy

very rarely: cerebrovascular ischaemia, cerebrovascular haemorrhage, encephalopathy

**Cardiac disorders:**

rarely: arrhythmia

very rarely: cardiac ischaemia, myocardial infarction

**Respiratory, thoracic and mediastinal disorders:**

very rarely: interstitial lung disease

**Gastrointestinal disorders:**

rarely: pancreatitis

very rarely: ulcerative and ischaemic colitis

**Skin and subcutaneous tissue disorders:**

very rarely: erythema multiforme, Stevens Johnson syndrome, toxic epidermal necrolysis, injection site necrosis

**Musculoskeletal and connective tissue disorders:**

rarely: rhabdomyolysis, myositis

**Renal and urinary disorders:**

rarely: renal insufficiency and renal failure

Cardiovascular (CVS) adverse events, particularly arrhythmia, appeared to be correlated mostly with pre-existing CVS disease and prior therapy with cardiotoxic agents (see section 4.4). Cardiomyopathy, that may be reversible upon discontinuation of interferon alpha, has been reported rarely in patients without prior evidence of cardiac disease.

Ophthalmological disorders that have been reported rarely with alpha interferons include retinopathies (including macular oedema), retinal haemorrhages, retinal artery or vein obstruction, cotton wool spots, loss of visual acuity or visual field, optic neuritis, and papilloedema (see section 4.4).

A wide variety of autoimmune and immune-mediated disorders have been reported with alpha interferons including thyroid disorders, systemic lupus erythematosus, rheumatoid arthritis (new or aggravated), idiopathic and thrombotic thrombocytopenic purpura, vasculitis, neuropathies including mononeuropathies (see also section 4.4, **Autoimmune disorders**).

**HCV/HIV co-infected patients**

For HCV/HIV co-infected patients receiving PegIntron in combination with ribavirin, other undesirable effects (that were not reported in mono-infected patients) which have been reported in the larger studies with a frequency  $> 5\%$  were: oral candidiasis (14 %), lipodystrophy acquired (13 %), CD4 lymphocytes decreased (8 %), appetite decreased (8 %), gamma-glutamyltransferase increased (9 %), back pain (5 %), blood amylase increased (6 %), blood lactic acid increased (5 %), cytolytic hepatitis (6 %), lipase increased (6 %) and pain in limb (6 %).

*Mitochondrial toxicity:*

Mitochondrial toxicity and lactic acidosis have been reported in HIV-positive patients receiving NRTI

regimen and associated ribavirin for co-HCV infection (see section 4.4).

*Laboratory values for HCV/HIV co-infected patients:*

Although haematological toxicities of neutropenia, thrombocytopenia and anaemia occurred more frequently in HCV/HIV co-infected patients, the majority could be managed by dose modification and rarely required premature discontinuation of treatment (see section 4.4). Haematological abnormalities were more frequently reported in patients receiving PegIntron in combination with ribavirin when compared to patients receiving interferon alfa-2b in combination with ribavirin. In Study 1 (see section 5.1), decrease in absolute neutrophil count levels below 500 cells/mm<sup>3</sup> was observed in 4 % (8/194) of patients and decrease in platelets below 50,000/mm<sup>3</sup> was observed in 4 % (8/194) of patients receiving PegIntron in combination with ribavirin. Anaemia (hemoglobin < 9.4g/dl) was reported in 12% (23/194) of patients treated with PegIntron in combination with ribavirin.

*CD4 lymphocytes decrease:*

Treatment with PegIntron in combination with ribavirin was associated with decreases in absolute CD4+ cell counts within the first 4 weeks without a reduction in CD4+ cell percentage. The decrease in CD4+ cell counts was reversible upon dose reduction or cessation of therapy. The use of PegIntron in combination with ribavirin had no observable negative impact on the control of HIV viraemia during therapy or follow-up. Limited safety data (N= 25) are available in co-infected patients with CD4+ cell counts < 200/μl (see section 4.4).

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron in combination with ribavirin.

## **4.9 Overdose**

In clinical trials, cases of accidental overdose, at never more than twice the prescribed dose, were reported. There were no serious reactions. Undesirable effects resolved during continued administration of PegIntron.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Immunostimulants, Cytokines and immunomodulators, Interferons, Peginterferon alfa-2b, ATC code: L03A B10.

Recombinant interferon alfa-2b is covalently conjugated with monomethoxy polyethylene glycol at an average degree of substitution of 1 mole of polymer/mole of protein. The average molecular mass is approximately 31,300 daltons of which the protein moiety constitutes approximately 19,300.

#### **Interferon alfa-2b**

*In vitro* and *in vivo* studies suggest that the biological activity of PegIntron is derived from its interferon alfa-2b moiety.

Interferons exert their cellular activities by binding to specific membrane receptors on the cell surface. Studies with other interferons have demonstrated species specificity. However, certain monkey species, e.g., Rhesus monkeys, are susceptible to pharmacodynamic stimulation upon exposure to human type 1 interferons.

Once bound to the cell membrane, interferon initiates a complex sequence of intracellular events that include the induction of certain enzymes. It is thought that this process, at least in part, is responsible for the various cellular responses to interferon, including inhibition of virus replication in virus-infected cells, suppression of cell proliferation and such immunomodulating activities as enhancement of the phagocytic activity of macrophages and augmentation of the specific cytotoxicity of

lymphocytes for target cells. Any or all of these activities may contribute to interferon's therapeutic effects.

Recombinant interferon alfa-2b also inhibits viral replication *in vitro* and *in vivo*. Although the exact antiviral mode of action of recombinant interferon alfa-2b is unknown, it appears to alter the host cell metabolism. This action inhibits viral replication or if replication occurs, the progeny virions are unable to leave the cell.

### **PegIntron**

PegIntron pharmacodynamics were assessed in a rising single-dose trial in healthy subjects by examining changes in oral temperature, concentrations of effector proteins such as serum neopterin and 2'5'-oligoadenylate synthetase (2'5'-OAS), as well as white cell and neutrophil counts. Subjects treated with PegIntron showed mild dose-related elevations in body temperature. Following single doses of PegIntron between 0.25 and 2.0 micrograms/kg/week, serum neopterin concentration was increased in a dose-related manner. Neutrophil and white cell count reductions at the end of week 4 correlated with the dose of PegIntron.

### **PegIntron clinical trials**

Two pivotal trials have been conducted, one (C/I97-010) with PegIntron monotherapy; the other (C/I98-580) with PegIntron in combination with ribavirin. Eligible patients for these trials had chronic hepatitis C confirmed by a positive HCV-RNA polymerase chain reaction (PCR) assay (> 30 IU/ml), a liver biopsy consistent with a histological diagnosis of chronic hepatitis with no other cause for the chronic hepatitis, and abnormal serum ALT.

In the PegIntron monotherapy trial, a total of 916 naïve chronic hepatitis C patients were treated with PegIntron (0.5, 1.0 or 1.5 micrograms/kg/week) for one year with a follow-up period of six months. In addition, 303 patients received interferon alfa-2b (3 million International Units [MIU] three times a week) as a comparator. This study showed that PegIntron was superior to interferon alfa-2b (**Table 8**).

In the PegIntron combination trial, 1,530 naïve patients were treated for one year with one of the following combination regimens:

- PegIntron (1.5 micrograms/kg/week) + ribavirin (800 mg/day), (n = 511).
- PegIntron (1.5 micrograms/kg/week for one month followed by 0.5 microgram/kg/week for 11 months) + ribavirin (1,000/1,200 mg/day), (n = 514).
- Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day) (n = 505).

In this trial, the combination of PegIntron (1.5 micrograms/kg/week) and ribavirin was significantly more effective than the combination of interferon alfa-2b and ribavirin (**Table 7**), particularly in patients infected with Genotype 1 (**Table 8**). Sustained response was assessed by the response rate six months after the cessation of treatment.

HCV genotype and baseline virus load are prognostic factors which are known to affect response rates. However, response rates in this trial were shown to be dependent also on the dose of ribavirin administered in combination with PegIntron or interferon alfa-2b. In those patients that received > 10.6 mg/kg ribavirin (800 mg dose in typical 75 kg patient), regardless of genotype or viral load, response rates were significantly higher than in those patients that received ≤ 10.6 mg/kg ribavirin (**Table 8**), while response rates in patients that received > 13.2 mg/kg ribavirin were even higher.

<b>Table 7</b> Sustained virological response (% patients HCV negative)							
	<b>PegIntron monotherapy</b>				<b>PegIntron + ribavirin</b>		
<b>Treatment regimen</b>	<b>P 1.5</b>	<b>P 1.0</b>	<b>P 0.5</b>	<b>I</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>Number of patients</b>	304	297	315	303	511	514	505
<b>Response at end of treatment</b>	49 %	41 %	33 %	24 %	65 %	56 %	54 %

<b>Sustained response</b>	<b>23 %*</b>	<b>25 %</b>	<b>18 %</b>	<b>12 %</b>	<b>54 %**</b>	<b>47 %</b>	<b>47 %</b>
---------------------------	--------------	-------------	-------------	-------------	---------------	-------------	-------------

P 1.5 PegIntron 1.5 micrograms/kg  
 P 1.0 PegIntron 1.0 microgram/kg  
 P 0.5 PegIntron 0.5 microgram/kg  
 I Interferon alfa-2b 3 MIU  
 P 1.5/R PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)  
 P 0.5/R PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)  
 I/R Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)  
 \* p < 0.001 P 1.5 vs. I  
 \*\* p = 0.0143 P 1.5/R vs. I/R

<b>Table 8</b> Sustained response rates with PegIntron + ribavirin (by ribavirin dose, genotype and viral load)				
<b>HCV Genotype</b>	<b>Ribavirin dose (mg/kg)</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>All Genotypes</b>	<b>All</b>	<b>54 %</b>	<b>47 %</b>	<b>47 %</b>
	≤ 10.6	50 %	41 %	27 %
	> 10.6	61 %	48 %	47 %
<b>Genotype 1</b>	<b>All</b>	<b>42 %</b>	<b>34 %</b>	<b>33 %</b>
	≤ 10.6	38 %	25 %	20 %
	> 10.6	48 %	34 %	34 %
Genotype 1 ≤ 600,000 IU/ml	All	73 %	51 %	45 %
	≤ 10.6	74 %	25 %	33 %
	> 10.6	71 %	52 %	45 %
Genotype 1 > 600,000 IU/ml	All	30 %	27 %	29 %
	≤ 10.6	27 %	25 %	17 %
	> 10.6	37 %	27 %	29 %
<b>Genotype 2/3</b>	<b>All</b>	<b>82 %</b>	<b>80 %</b>	<b>79 %</b>
	≤ 10.6	79 %	73 %	50 %
	> 10.6	88 %	80 %	80 %

P 1.5/R PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)  
 P 0.5/R PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)  
 I/R Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)

In the PegIntron monotherapy study, the Quality of Life was generally less affected by 0.5 microgram/kg of PegIntron than by either 1.0 microgram/kg of PegIntron once weekly or 3 MIU of interferon alfa-2b three times a week.

In a separate trial, 224 patients with genotype 2 or 3 received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with ribavirin 800 mg –1,400 mg p.o. for 6 months (based on body weight, only three patients weighing > 105 kg, received the 1,400 mg dose, which has not yet been validated) (**Table 9**). Twenty-four % had bridging fibrosis or cirrhosis (Knodell 3/4).

**Table 9. Virologic Response at End of Treatment, Sustained Virologic Response and Relapse by HCV Genotype and Viral Load\***

	PegIntron 1.5 µg/kg once weekly plus Ribavirin 800-1,400 mg/day		
	End of Treatment Response	Sustained Virologic Response	Relapse
All Subjects	94 % (211/224)	81 % (182/224)	12 % (27/224)
HCV 2 ≤ 600,000 IU/ml > 600,000	100 % (42/42)	93 % (39/42)	7 % (3/42)
	100 % (20/20)	95 % (19/20)	5 % (1/20)
	100 % (22/22)	91 % (20/22)	9 % (2/22)

IU/ml			
HCV 3	93 % (169/182)	79 % (143/182)	14 % (24/166)
≤ 600,000 IU/ml	93 % (92/99)	86 % (85/99)	8 % (7/91)
> 600,000 IU/ml	93 % (77/83)	70 % (58/83)	23 % (17/75)

\* Any subject with an undetectable HCV-RNA level at the Follow-Up Week 12 visit and missing data at the Follow-Up Week 24 visit was considered a sustained responder. Any subject with missing data in and after the Follow-Up Week 12 window was considered to be a non-responder at Week 24 of follow-up.

The 6 month treatment duration in this trial was better tolerated than one year of treatment in the pivotal combination trial; for discontinuation 5 % vs. 14 %, for dose modification 18 % vs. 49 %.

In a non-comparative trial, 235 patients with genotype 1 and low viral load (< 600,000 IU/ml) received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with weight adjusted ribavirin. The overall sustained response rate after a 24-week treatment duration was 50 %. Forty-one percent of subjects (97/235) had nondetectable plasma HCV-RNA levels at Week 4 and Week 24 of therapy. In this subgroup, there was a 92 % (89/97) sustained virological response rate. The high sustained response rate in this subgroup of patients was identified in an interim analysis (n=49) and prospectively confirmed (n=48).

Limited historical data indicate that treatment for 48 weeks might be associated with a higher sustained response rate (11/11) and with a lower risk of relapse (0/11 as compared to 7/96 following 24 weeks of treatment).

### Predictability of sustained virological response

Virological response by week 12, defined as a 2-log viral load decrease or undetectable levels of HCV-RNA has been shown to be predictive for sustained response (**Table 10**).

<b>Table 10</b> Predictability of sustained response by viral response at week 12 and genotype*				
Treatment	Genotype	Viral response at week 12	Sustained response	Negative predictive value
PegIntron 1.5 + ribavirin (> 10.6 mg/kg) 48-week treatment	1	Yes 75 % (82/110)	71 % (58/82)	----
		No 25 % (28/110)	0 % (0/28)	100 %
PegIntron 1.5 + ribavirin 800-1,400 mg 24-week treatment	2 and 3	Yes 99 % (213/215)	83 % (177/213)	----
		No 1 % (2/215)	50 % (1/2)	50 %

\* reflects patients with 12 week data available

The negative predictive value for sustained response in patients treated with PegIntron in monotherapy was 98 %.

### HCV/HIV Co-infected patients

Two trials have been conducted in patients co-infected with HIV and HCV. The response to treatment in both of these trials is presented in **Table 11**. Study 1 (RIBAVIC; P01017) was a randomized, multicentre study which enrolled 412 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (1.5 µg/kg/week) plus ribavirin (800 mg/day) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800 mg/day) for 48 weeks with a follow-up period of 6 months. Study 2 (P02080) was a randomized, single centre

study that enrolled 95 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (100 or 150 µg /week based on weight) plus ribavirin (800-1,200 mg/day based on weight) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800-1,200 mg/day based on weight). The duration of therapy was 48 weeks with a follow-up period of 6 months except for patients infected with genotypes 2 or 3 and viral load < 800,000 IU/ml (Amplicor) who were treated for 24 weeks with a 6month follow-up period.

<b>Table 11</b> Sustained virological response based on genotype after PegIntron in combination with Ribavirin in HCV/HIV Co-infected patients						
	<b>Study 1<sup>1</sup></b>			<b>Study 2<sup>2</sup></b>		
	PegIntron (1.5 µg/kg/week) + ribavirin (800 mg)	Interferon alfa-2b (3 MIU TIW) + ribavirin (800 mg)	p value <sup>a</sup>	PegIntron (100 or 150 <sup>c</sup> µg/week) + ribavirin (800-1,200 mg) <sup>d</sup>	Interferon alfa-2b (3 MIU TIW) + ribavirin (800-1,200 mg) <sup>d</sup>	p value <sup>b</sup>
All	27 % (56/205)	20 % (41/205)	0.047	44 % (23/52)	21 % (9/43)	0.017
Genotype 1, 4	17 % (21/125)	6 % (8/129)	0.006	38 % (12/32)	7 % (2/27)	0.007
Genotype 2, 3	44 % (35/80)	43 % (33/76)	0.88	53 % (10/19)	47 % (7/15)	0.730

MIU = million international units; TIW = three times a week.

a: p value based on Cochran-Mantel Haenszel Chi square test.

b: p value based on chi-square test.

c: subjects < 75 kg received 100 µg/week PegIntron and subjects ≥ 75 kg received 150 µg/week PegIntron.

d: ribavirin dosing was 800 mg for patients < 60 kg, 1,000 mg for patients 60-75 kg, and 1,200 mg for patients > 75 kg.

<sup>1</sup>Carrato F, Bani-Sadir F, Pol S et al. JAMA 2004; 292(23): 2839-2848.

<sup>2</sup>Laguno M, Murillas J, Blanco J et al. AIDS 2004; 18(13): F27-F36.

### *Histological response*

Liver biopsies were obtained before and after treatment in Study 1 and were available for 210 of the 412 subjects (51 %). Both the Metavir score and Ishak grade decreased among subjects treated with PegIntron in combination with ribavirin. This decline was significant among responders (-0.3 for Metavir and -1.2 for Ishak) and stable (-0.1 for Metavir and -0.2 for Ishak) among non-responders. In terms of activity, about one-third of sustained responders showed improvement and none showed worsening. There was no improvement in terms of fibrosis observed in this study. Steatosis was significantly improved in patients infected with HCV Genotype 3.

## **5.2 Pharmacokinetic properties**

PegIntron is a well characterized polyethylene glycol-modified (“pegylated”) derivative of interferon alfa-2b and is predominantly composed of monopegylated species. The plasma half-life of PegIntron is prolonged compared with non-pegylated interferon alfa-2b. PegIntron has a potential to depegylate to free interferon alfa-2b. The biologic activity of the pegylated isomers is qualitatively similar, but weaker than free interferon alfa-2b.

Following subcutaneous administration, maximal serum concentrations occur between 15-44 hours post-dose, and are sustained for up to 48-72 hours post-dose.

PegIntron C<sub>max</sub> and AUC measurements increase in a dose-related manner. Mean apparent volume of distribution is 0.99 l/kg.

Upon multiple dosing, there is an accumulation of immunoreactive interferons. There is, however, only a modest increase in biologic activity as measured by a bioassay.



Mean (SD) PegIntron elimination half-life is approximately 40 hours (13.3 hours), with apparent clearance of 22.0 ml/hr·kg. The mechanisms involved in clearance of interferons in man have not yet been fully elucidated. However, renal elimination may account for a minority (approximately 30 %) of PegIntron apparent clearance.

**Renal function:** Renal clearance appears to account for 30 % of total clearance of PegIntron. In a single dose study (1.0 microgram/kg) in patients with impaired renal function,  $C_{max}$ , AUC, and half-life increased in relation to the degree of renal impairment.

Following multiple dosing of PegIntron (1.0 microgram/kg subcutaneously administered every week for four weeks) the clearance of PegIntron is reduced by a mean of 17 % in patients with moderate renal impairment (creatinine clearance 30-49 ml/minute) and by a mean of 44 % in patients with severe renal impairment (creatinine clearance 15-29 ml/minute) compared to subjects with normal renal function. Based on single dose data, clearance was similar in patients with severe renal impairment not on dialysis and in patients who were receiving hemodialysis. The dose of PegIntron for monotherapy should be reduced in patients with moderate or severe renal impairment (see sections 4.2 and 4.4). Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see section 4.3).

Because of marked inter-subject variability in interferon pharmacokinetics, it is recommended that patients with severe renal impairment be closely monitored during treatment with PegIntron (see section 4.2)

**Hepatic function:** The pharmacokinetics of PegIntron have not been evaluated in patients with severe hepatic dysfunction.

**Elderly patients ≥ 65 years of age:** The pharmacokinetics of PegIntron following a single subcutaneous dose of 1.0 microgram/kg were not affected by age. The data suggest that no alteration in PegIntron dosage is necessary based on advancing age.

**Patients under the age of 18 years:** Specific pharmacokinetic evaluations have not been performed on these patients. PegIntron is indicated for the treatment of chronic hepatitis C only in patients 18 years of age or older.

**Interferon neutralising factors:** Interferon neutralising factor assays were performed on serum samples of patients who received PegIntron in the clinical trial. Interferon neutralising factors are antibodies which neutralise the antiviral activity of interferon. The clinical incidence of neutralising factors in patients who received PegIntron 0.5 micrograms/kg is 1.1 %.

### 5.3 Preclinical safety data

**PegIntron:** Adverse events not observed in clinical trials were not seen in toxicity studies in monkeys. These studies were limited to four weeks due to the appearance of anti-interferon antibodies in most monkeys.

Reproduction studies of PegIntron have not been performed. Interferon alfa-2b has been shown to be an abortifacient in primates. PegIntron is likely to also cause this effect. Effects on fertility have not been determined. It is not known whether the components of this medicinal product are excreted into experimental animal or human milk (see section 4.6 for relevant human data on pregnancy and lactation). PegIntron showed no genotoxic potential.

The relative non-toxicity of monomethoxy-polyethylene glycol (mPEG), which is liberated from PegIntron by metabolism *in vivo* has been demonstrated in preclinical acute and subchronic toxicity studies in rodents and monkeys, standard embryo-foetal development studies and in *in vitro* mutagenicity assays.

**PegIntron plus ribavirin:** When used in combination with ribavirin, PegIntron did not cause any effects not previously seen with either active substance alone. The major treatment-related change was a reversible, mild to moderate anaemia, the severity of which was greater than that produced by either active substance alone.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Powder for solution for injection:

Disodium phosphate, anhydrous,  
Sodium dihydrogen phosphate dihydrate,  
Sucrose,  
Polysorbate 80.

Solvent for parenteral use:

Water for injections.

### **6.2 Incompatibilities**

This medicinal product should only be reconstituted with the solvent provided (see section 6.6). In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

3 years

After reconstitution:

- Chemical and physical in-use stability has been demonstrated for 24 hours at 2°C - 8°C.
- From a microbiological point of view, the product is to be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C - 8°C.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C - 8°C).

### **6.5 Nature and contents of container**

The powder is contained in a 2 ml vial, Type I flint glass, with a butyl rubber stopper in an aluminium flip-off seal with a polypropylene bonnet. The solvent is presented in a 2 ml ampoule, Type I flint glass. PegIntron 80 micrograms is supplied as:

- 1 vial of powder for solution for injection and 1 ampoule of solvent for parenteral use;
- 1 vial of powder for solution for injection, 1 ampoule of solvent for parenteral use, 1 injection syringe, 2 injection needles and 1 cleansing swab;
- 4 vials of powder for solution for injection and 4 ampoules of solvent for parenteral use;
- 4 vials of powder for solution for injection, 4 ampoules of solvent for parenteral use, 4 injection syringes, 8 injection needles and 4 cleansing swabs;
- 6 vials of powder for solution for injection and 6 ampoules of solvent for parenteral use.
- 12 vials of powder for solution for injection, 12 ampoules of solvent for parenteral use, 12 injection syringes, 24 injection needles and 12 cleansing swabs.

Not all pack sizes may be marketed

### **6.6 Special precautions for disposal**

PegIntron is supplied as a powder of peginterferon alfa-2b at a strength of 80 micrograms for single use. Each vial must be reconstituted with 0.7 ml of water for injections for administration of up to 0.5 ml of solution. A small volume is lost during preparation of PegIntron for injection when the dose is measured and injected. Therefore, each vial contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection. The reconstituted solution has a concentration of 80 micrograms/0.5 ml.

Using a sterilised injection syringe and injection needle, inject 0.7 ml of water for injections into the vial of PegIntron. Agitate gently to complete dissolution of powder. The appropriate dose can then be withdrawn with a sterilised injection syringe and injected. A complete set of instructions is provided in the Annex to the Package Leaflet.

As for all parenteral medicinal products, inspect visually the reconstituted solution prior to administration. The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. Discard any unused material.

## **7. MARKETING AUTHORISATION HOLDER**

SP Europe  
73, rue de Stalle  
B-1180 Bruxelles  
Belgium

## **8. MARKETING AUTHORISATION NUMBERS**

EU/1/00/131/006  
EU/1/00/131/007  
EU/1/00/131/008  
EU/1/00/131/009  
EU/1/00/131/010  
EU/1/00/131/027

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 25 May 2000  
Date of last renewal: 25 May 2005

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>

## **1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 100 micrograms powder and solvent for solution for injection

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each vial of PegIntron, powder for solution for injection contains 100 micrograms of peginterferon alfa-2b as measured on a protein basis.

Each vial provides 100 micrograms/0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

The active substance is a covalent conjugate of recombinant interferon alfa-2b\* with monomethoxy polyethylene glycol. The potency of this product should not be compared to that of another pegylated or non-pegylated protein of the same therapeutic class. For more information, see section 5.1.

\*produced by rDNA technology in *E.coli* cells harbouring a genetically engineered plasmid hybrid encompassing an interferon alfa-2b gene from human leukocytes

For a full list of excipients, see section 6.1. PegIntron contains 40 mg of sucrose per 0.5 ml.

## **3. PHARMACEUTICAL FORM**

Powder and solvent for solution for injection

White powder.

Clear and colourless solvent.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

PegIntron is indicated for the treatment of adult patients with chronic hepatitis C who have elevated transaminases without liver decompensation and who are positive for serum HCV-RNA or anti-HCV, including naïve patients with clinically stable HIV co-infection (see section 4.4).

The best way to use PegIntron in this indication is in combination with ribavirin.

This combination is indicated in naïve patients including patients with clinically stable HIV co-infection. This combination of PegIntron with ribavirin is also indicated in patients who have previously responded (with normalisation of ALT at the end of treatment) to interferon alpha monotherapy but who have subsequently relapsed.

Interferon monotherapy, including PegIntron, is indicated mainly in case of intolerance or contraindication to ribavirin.

Please refer also to the ribavirin Summary of Product Characteristics (SPC) when PegIntron is to be used in combination with ribavirin.

### **4.2 Posology and method of administration**

Treatment should be initiated and monitored only by a physician experienced in the management of patients with hepatitis C.

#### **Dose to be administered**

PegIntron should be administered as a once weekly subcutaneous injection. The dose administered depends on whether it is used in combination with ribavirin or as monotherapy.

## Combination therapy

PegIntron 1.5 micrograms/kg/week in combination with ribavirin capsules.

The intended dose of 1.5 µg/kg of PegIntron to be used in combination with ribavirin may be delivered in weight categories with the pen/vial strengths according to **Table 1**. Ribavirin capsules are to be administered orally each day in two divided doses with food (morning and evening).

**Table 1 - Dosing for Combination Therapy**

Body Weight (kg)	PegIntron		Ribavirin Capsules	
	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Total Daily Dose (mg)	Number of Capsules (200 mg)
< 40	50	0.5	800	4 <sup>a</sup>
40-50	80	0.4	800	4 <sup>a</sup>
51-64	80	0.5	800	4 <sup>a</sup>
65-75	100	0.5	1,000	5 <sup>b</sup>
76-85	120	0.5	1,000	5 <sup>b</sup>
> 85	150	0.5	1,200	6 <sup>c</sup>

a: 2 morning, 2 evening

b: 2 morning, 3 evening

c: 3 morning, 3 evening

## Duration of treatment

**Predictability of sustained virological response:** Patients infected with virus genotype 1 who fail to achieve virological response at Week 12 are highly unlikely to become sustained virological responders (see also section 5.1).

- **Genotype 1:** For patients who exhibit virological response at week 12, treatment should be continued for another nine month period (i.e., a total of 48 weeks).  
In the subset of patients with genotype 1 infection and low viral load (< 600,000 IU/ml) who become HCV-RNA negative at treatment week 4 and remain HCV-RNA negative at week 24, the treatment could either be stopped after this 24 week treatment course or pursued for an additional 24 weeks (i.e. overall 48 weeks treatment duration). However, an overall 24 weeks treatment duration may be associated with a higher risk of relapse than a 48 weeks treatment duration (see section 5.1).
- **Genotypes 2 or 3:** It is recommended that all patients be treated for 24 weeks, except for HCV/HIV co-infected patients who should receive 48 weeks of treatment.
- **Genotype 4:** In general, patients infected with genotype 4 are considered harder to treat and limited study data (n=66) indicate they are compatible with a duration of treatment as for genotype 1.

## HCV/HIV Co-infection

The recommended duration of dosing for HCV/HIV co-infected patients is 48 weeks, regardless of genotype.

### *Predictability of response and non-response in HCV/HIV Co-infection*

Early virological response by Week 12, defined as a 2 log viral load decrease or undetectable levels of HCV-RNA, has been shown to be predictive for sustained response. The negative predictive value for sustained response in HCV/HIV co-infected patients treated with PegIntron in combination with ribavirin was 99 % (67/68; Study 1) (see section 5.1). A positive predictive value of 50 % (52/104; Study 1) was observed for HCV/HIV co-infected patients receiving combination therapy.

### PegIntron monotherapy

As monotherapy the PegIntron regimen is 0.5 or 1.0 microgram/kg/week. The lowest vial or pen strength available is 50 µg/0.5 ml; therefore for patients prescribed 0.5 µg/kg/week, doses must be adjusted by volume as shown in **Table 2**. For the 1.0 µg/kg dose, similar volume adjustments can be made or alternate vial strengths can be used as shown in **Table 2**. PegIntron monotherapy was not studied in HCV/HIV co-infected patients.

**Table 2- Monotherapy Dosing**

	0.5 µg/kg		1.0 µg/kg	
Body Weight (kg)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)
30-35	50*	0.15	50	0.3
36-45	50*	0.2	50	0.4
46-56	50*	0.25	50	0.5
57-72	50	0.3	80	0.4
73-88	50	0.4	80	0.5
89-106	50	0.5	100	0.5
> 106**	80	0.4	120	0.5
* Must use vial. Minimum delivery for pen is 0.3 ml.				
** For patients > 120 kg, use 80 µg/0.5 ml vial				

### Duration of treatment

For patients who exhibit virological response at Week 12, treatment should be continued for at least another three-month period (i.e., a total of six months). The decision to extend therapy to one year of treatment should be based on prognostic factors (e.g., genotype, age > 40 years, male gender, bridging fibrosis).

### Dose modification for all patients

If severe adverse reactions or laboratory abnormalities develop during treatment with PegIntron monotherapy or PegIntron in combination with ribavirin, modify the dosages of each product as appropriate, until the adverse reactions abate. As adherence might be of importance for outcome of therapy, the dose should be kept as close as possible to the recommended standard dose. Guidelines were developed in clinical trials for dose modification.

### Combination Therapy Dose Reduction Guidelines

<b>Table 2a</b> Dose modification guidelines for combination therapy (with ribavirin)			
Laboratory values	Reduce only ribavirin dose to <u>600 mg/day</u> * if:	Reduce only PegIntron dose to one-half dose if:	Discontinue combination therapy if:
Haemoglobin	< 10 g/dl	-	< 8.5 g/dl
Haemoglobin in: Patients with history of stable cardiac disease	≥ 2 g/dl decrease in haemoglobin during any four week period during treatment (permanent dose reduction)		< 12 g/dl after four weeks of dose reduction
White blood cells	-	< 1.5 x 10 <sup>9</sup> /l	< 1.0 x 10 <sup>9</sup> /l
Neutrophils	-	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	-	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l

Bilirubin – direct	-	-	2.5 x ULN**
Bilirubin - indirect	> 5 mg/dl	-	> 4 mg/dl (for > 4 weeks)
Creatinine	-	-	> 2.0 mg/dl
ALT/AST	-	-	2 x baseline and > 10 x ULN**

\* Patients whose dose of ribavirin is reduced to 600 mg daily receive one 200 mg capsule in the morning and two 200 mg capsules in the evening.

\*\* Upper limit of normal

Dose reduction of PegIntron may be accomplished by either reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 2b**.

<b>Table 2b – Reduced PegIntron Dosing for Combination Therapy</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5 ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
< 40	25	50*	0.25	25
40-50	32	50	0.3	30
51-64	40	50	0.4	40
65-75	50	50	0.5	50
76-85	60	80	0.4	64
> 85	75	100	0.4	80
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### **PegIntron Monotherapy Dose Reduction Guidelines**

Dose modification guidelines for patients who use PegIntron monotherapy are shown in **Table 3a**.

<b>Table 3a</b> Dose modification guidelines for PegIntron monotherapy		
<b>Laboratory values</b>	<b>Reduce PegIntron to one-half dose if:</b>	<b>Discontinue PegIntron if:</b>
Neutrophils	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l

Dose reduction for patients who use 0.5 µg/kg PegIntron monotherapy must be accomplished by reducing the prescribed volume by one-half. The 50 µg/0.5 ml vial must be used if necessary since the pen can only deliver a minimum volume of 0.3 ml.

For patients who use 1.0 µg/kg PegIntron monotherapy, dose reduction may be accomplished by reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 3b**.

<b>Table 3b – Reduced PegIntron Dose for the 1.0 µg/kg Monotherapy Regimen</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
30-35	15	50*	0.15	15
36-45	20	50*	0.20	20
46-56	25	50*	0.25	25
57-72	32	50	0.3	30
73-89	40	50	0.4	40
90-106	50	50	0.5	50
> 106	60	80	0.4	64
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### Special populations

**Use in renal impairment:** *Monotherapy:* PegIntron should be used with caution in patients with moderate to severe renal impairment. In patients with moderate renal dysfunction (creatinine clearance 30-50 ml/minute), the starting dose of PegIntron should be reduced by 25 %. Patients with severe renal dysfunction (creatinine clearance 15-29 ml/minute) should have the starting dose of PegIntron reduced by 50 %. Data are not available for the use of PegIntron in patients with creatinine clearance < 15 ml/minute (see section 5.2). Patients with severe renal impairment, including those on hemodialysis, should be closely monitored. If renal function decreases during treatment, PegIntron therapy should be discontinued.

*Combination therapy:* Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see ribavirin SPC). When administered in combination with ribavirin, subjects with impaired renal function should be more carefully monitored with respect to the development of anaemia.

**Use in hepatic impairment:** The safety and efficacy of PegIntron therapy has not been evaluated in patients with severe hepatic dysfunction, therefore PegIntron must not be used for these patients.

**Use in the elderly (≥ 65 years of age):** There are no apparent age-related effects on the pharmacokinetics of PegIntron. Data from elderly patients treated with a single dose of PegIntron suggest no alteration in PegIntron dose is necessary based on age (see section 5.2).

**Use in patients under the age of 18 years:** There is no experience in children (see section 5.2).

### 4.3 Contraindications

- Hypersensitivity to the active substance or to any interferon or to any of the excipients;
- A history of severe pre-existing cardiac disease, including unstable or uncontrolled cardiac disease in the previous six months (see section 4.4);
- Severe, debilitating medical conditions;
- Autoimmune hepatitis or a history of autoimmune disease;
- Severe hepatic dysfunction or decompensated cirrhosis of the liver;
- Pre-existing thyroid disease unless it can be controlled with conventional treatment;
- Epilepsy and/or compromised central nervous system (CNS) function;
- Initiation of PegIntron is contraindicated in HCV/HIV patients with cirrhosis and a Child-Pugh score ≥ 6.

*Combination therapy with ribavirin:* Also see ribavirin Summary of the Product Characteristics (SPC) if PegIntron is to be administered in combination with ribavirin in patients with chronic hepatitis C.



#### 4.4 Special warnings and precautions for use

**Psychiatric and Central Nervous System (CNS):** Severe CNS effects, particularly depression, suicidal ideation and attempted suicide have been observed in some patients during PegIntron therapy, and even after treatment discontinuation mainly during the 6-month follow-up period. Other CNS effects including aggressive behaviour (sometimes directed against others), confusion and alterations of mental status have been observed with alpha interferons. Patients should be closely monitored for any signs or symptoms of psychiatric disorders. If such symptoms appear, the potential seriousness of these undesirable effects must be borne in mind by the prescribing physician and the need for adequate therapeutic management should be considered. If psychiatric symptoms persist or worsen, or suicidal ideation is identified, it is recommended that treatment with PegIntron be discontinued, and the patient followed, with psychiatric intervention as appropriate.

*Patients with existence of, or history of severe psychiatric conditions:* If treatment with peginterferon alfa-2b is judged necessary in patients with existence or history of severe psychiatric conditions, this should only be initiated after having ensured appropriate individualised diagnostic and therapeutic management of the psychiatric condition.

There is no experience with PegIntron in combination with ribavirin in patients who have relapsed after interferon alpha + ribavirin therapy.

All patients in the chronic hepatitis C studies had a liver biopsy before inclusion, but in certain cases (i.e. patients with genotype 2 and 3), treatment may be possible without histological confirmation. Current treatment guidelines should be consulted as to whether a liver biopsy is needed prior to commencing treatment.

**Acute hypersensitivity:** Acute hypersensitivity reactions (e.g., urticaria, angioedema, bronchoconstriction, anaphylaxis) have been observed rarely during interferon alfa-2b therapy. If such a reaction develops during treatment with PegIntron, discontinue treatment and institute appropriate medical therapy immediately. Transient rashes do not necessitate interruption of treatment.

**Cardiovascular system:** As with interferon alfa-2b, patients with a history of congestive heart failure, myocardial infarction and/or previous or current arrhythmic disorders, receiving PegIntron therapy require close monitoring. It is recommended that patients who have pre-existing cardiac abnormalities have electrocardiograms taken prior to and during the course of treatment. Cardiac arrhythmias (primarily supraventricular) usually respond to conventional therapy but may require discontinuation of PegIntron therapy.

More significant obtundation and coma, including cases of encephalopathy, have been observed in some patients, usually elderly, treated at higher doses for oncology indications. While these effects are generally reversible, in a few patients full resolution took up to three weeks. Very rarely, seizures have occurred with high doses of interferon alpha.

**Liver function:** As with all interferons, discontinue treatment with PegIntron in patients who develop prolongation of coagulation markers which might indicate liver decompensation.

**Fever:** While fever may be associated with the flu-like syndrome reported commonly during interferon therapy, other causes of persistent fever must be ruled out.

**Hydration:** Adequate hydration must be maintained in patients undergoing PegIntron therapy since hypotension related to fluid depletion has been seen in some patients treated with alpha interferons. Fluid replacement may be necessary.

**Pulmonary changes:** Pulmonary infiltrates, pneumonitis, and pneumonia, occasionally resulting in fatality, have been observed rarely in interferon alpha treated patients. Any patient developing fever, cough, dyspnea or other respiratory symptoms must have a chest X-ray taken. If the chest X-ray shows pulmonary infiltrates or there is evidence of pulmonary function impairment, the patient is to be

monitored closely, and, if appropriate, discontinue interferon alpha. Prompt discontinuation of interferon alpha administration and treatment with corticosteroids appear to be associated with resolution of pulmonary adverse events.

**Autoimmune disease:** The development of auto-antibodies and autoimmune disorders has been reported during treatment with alpha interferons. Patients predisposed to the development of autoimmune disorders may be at increased risk. Patients with signs or symptoms compatible with autoimmune disorders should be evaluated carefully, and the benefit-risk of continued interferon therapy should be reassessed (see also section 4.4 Thyroid changes and 4.8).

**Ocular changes:** Ophthalmologic disorders, including retinal haemorrhages, cotton wool spots, and retinal artery or vein obstruction have been reported in rare instances after treatment with alpha interferons (see section 4.8). All patients should have a baseline eye examination. Any patient complaining of ocular symptoms, including loss of visual acuity or visual field must have a prompt and complete eye examination. Periodic visual examinations are recommended during PegIntron therapy, particularly in patients with disorders that may be associated with retinopathy, such as diabetes mellitus or hypertension. Discontinuation of PegIntron should be considered in patients who develop new or worsening ophthalmological disorders.

**Thyroid changes:** Infrequently, patients treated for chronic hepatitis C with interferon alpha have developed thyroid abnormalities, either hypothyroidism or hyperthyroidism. Determine thyroid stimulating hormone (TSH) levels if, during the course of therapy, a patient develops symptoms consistent with possible thyroid dysfunction. In the presence of thyroid dysfunction, PegIntron treatment may be continued if TSH levels can be maintained in the normal range by medication.

**Metabolic disturbances:** Hypertriglyceridemia and aggravation of hypertriglyceridemia, sometimes severe, have been observed. Monitoring of lipid levels is, therefore, recommended.

### **HCV/HIV Co-infection**

Mitochondrial toxicity and lactic acidosis:

Patients co-infected with HIV and receiving Highly Active Anti-Retroviral Therapy (HAART) may be at increased risk of developing lactic acidosis. Caution should be used when adding PegIntron and ribavirin to HAART therapy (see ribavirin SPC).

Moreover, patients treated with PegIntron and ribavirin combination therapy and zidovudine are at increased risk of developing anaemia.

Hepatic decompensation in HCV/HIV co-infected patients with advanced cirrhosis:

Co-infected patients with advanced cirrhosis receiving HAART may be at increased risk of hepatic decompensation and death. Adding treatment with alfa interferons alone or in combination with ribavirin may increase the risk in this patient subset. Other baseline factors in co-infected patients that may be associated with a higher risk of hepatic decompensation include treatment with didanosine and elevated bilirubin serum concentration.

Co-infected patients receiving both antiretroviral (ARV) and anti-hepatitis treatment should be closely monitored, assessing their Child-Pugh score during treatment. Patients progressing to hepatic decompensation should have their anti-hepatitis treatment immediately discontinued and the ARV treatment reassessed.

Haematological abnormalities in HCV/HIV co-infected patients:

HCV/HIV co-infected patients receiving peginterferon alfa-2b/ribavirin treatment and HAART may be at increased risk to develop haematological abnormalities (as neutropenia, thrombocytopenia and anaemia) compared to HCV mono-infected patients. Although, the majority of them could be managed by dose reduction, close monitoring of haematological parameters should be undertaken in this population of patients (see section 4.2 and below “Laboratory tests” and section 4.8).

Patients with low CD4 counts:

In patients co-infected with HCV/HIV, limited efficacy and safety data (N = 25) are available in subjects with CD4 counts less than 200 cells/μl. Caution is therefore warranted in the treatment of

patients with low CD4 counts.

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron and ribavirin.

**Dental and periodontal disorders:** Dental and periodontal disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and mucous membranes of the mouth during long-term treatment with the combination of PegIntron and ribavirin. Patients should brush their teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If this reaction occurs, they should be advised to rinse out their mouth thoroughly afterwards.

**Organ transplant recipients:** The safety and efficacy of PegIntron alone or in combination with ribavirin for the treatment of hepatitis C in liver or other organ transplant recipients have not been studied. Preliminary data indicates that interferon alpha therapy may be associated with an increased rate of kidney graft rejection. Liver graft rejection has also been reported.

**Other:** Due to reports of interferon alpha exacerbating pre-existing psoriatic disease and sarcoidosis, use of PegIntron in patients with psoriasis or sarcoidosis is recommended only if the potential benefit justifies the potential risk.

**Laboratory tests:** Standard haematologic tests, blood chemistry and a test of thyroid function must be conducted in all patients prior to initiating therapy. Acceptable baseline values that may be considered as a guideline prior to initiation of PegIntron therapy are:

- |                    |                              |
|--------------------|------------------------------|
| • Platelets        | $\geq 100,000/\text{mm}^3$   |
| • Neutrophil count | $\geq 1,500/\text{mm}^3$     |
| • TSH level        | must be within normal limits |

Laboratory evaluations are to be conducted at weeks 2 and 4 of therapy, and periodically thereafter as clinically appropriate.

**Important information about some of the ingredients of PegIntron:**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

Patients with rare hereditary problems of fructose intolerance, glucose galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Results from a multiple-dose probe study assessing P450 substrates in chronic hepatitis C patients receiving once weekly PegIntron (1.5 µg/kg) for 4 weeks demonstrated an increase in activity of CYP2D6 and CYP2C8/9. No change in activity of CYP1A2, CYP3A4, or N-acetyltransferase was observed.

Caution should be used when administering peginterferon alfa-2b with medicines metabolised by CYP2D6 and CYP2C8/9, especially those with narrow therapeutic window, such as warfarin and phenytoin (CYP2C9) and flecainide (CYP2D6).

These findings may partly relate to improved metabolic capacity due to reduced hepatic inflammation in patients undergoing treatment with PegIntron. Caution is therefore advised when PegIntron treatment is initiated for chronic hepatitis in patients treated with medication with a narrow therapeutic window and sensitive to mild metabolic impairment of the liver.

No pharmacokinetic interactions were noted between PegIntron and ribavirin in a multiple-dose pharmacokinetic study.

**Methadone:** In patients with chronic hepatitis C that were on stable methadone maintenance therapy and naïve to peginterferon alfa-2b, addition of 1.5 microgram/kg/week of PegIntron subcutaneously for 4 weeks increased R-methadone AUC by approximately 15 % (95 % CI for AUC ratio estimate 103 – 128 %). The clinical significance of this finding is unknown; however, patients should be monitored for signs and symptoms of increased sedative effect, as well as respiratory depression. Especially in patients on a high dose of methadone, the risk for QTc prolongation should be considered.

#### *HCV/HIV Co-infection*

Nucleoside analogs: ribavirin has been shown *in vitro* to inhibit phosphorylation of zidovudine and stavudine. The clinical significance of these findings is unknown. However, these *in vitro* findings raise the possibility that concurrent use of ribavirin with either zidovudine or stavudine might lead to increased HIV plasma viraemia. Therefore, it is recommended that plasma HIV RNA levels be closely monitored in patients treated with ribavirin concurrently with either of these two agents. If HIV RNA levels increase, the use of ribavirin concomitantly with reverse transcriptase inhibitors must be reviewed (see ribavirin SPC).

Use of nucleoside analogs, alone or in combination with other nucleosides, has resulted in lactic acidosis. Pharmacologically, ribavirin increases phosphorylated metabolites of purine nucleosides *in vitro*. This activity could potentiate the risk of lactic acidosis induced by purine nucleoside analogs (e.g. didanosine or abacavir). Co-administration of ribavirin and didanosine is not recommended. Reports of mitochondrial toxicity, in particular lactic acidosis and pancreatitis, of which some fatal, have been reported (see ribavirin SPC).

#### **4.6 Pregnancy and lactation**

PegIntron is recommended for use in fertile women only when they are using effective contraception during the treatment.

There are no adequate data from the use of interferon alfa-2b in pregnant women.. Studies in animals have shown reproductive toxicity (see section 5.3). Interferon alfa-2b has been shown to be abortifacient in primates. PegIntron is likely to also cause this effect.

The potential risk in humans is unknown. PegIntron is to be used during pregnancy only if the potential benefit justifies the potential risk to the foetus

**Lactation:** It is not known whether the components of this medicinal product are excreted in human milk. Because of the potential for adverse reactions in breast-fed infants, breast-feeding should be discontinued prior to initiation of treatment.

#### *Combination therapy with ribavirin:*

Ribavirin causes serious birth defects when administered during pregnancy. Ribavirin therapy is contraindicated in women who are pregnant. Extreme care must be taken to avoid pregnancy in female patients or in partners of male patients taking PegIntron in combination with ribavirin. Females of childbearing potential and their partners must each use an effective contraceptive during treatment and for 4 months after treatment has been concluded. Male patients and their female partners must each use an effective contraceptive during treatment and for 7 months after treatment has been concluded (see Ribavirin SPC).

#### **4.7 Effects on ability to drive and use machines**

Patients who develop fatigue, somnolence or confusion during treatment with PegIntron are cautioned to avoid driving or operating machinery.

#### **4.8 Undesirable effects**

The safety of PegIntron is evaluated from data from two clinical trials: one with PegIntron monotherapy, one with PegIntron in combination with ribavirin. In both cases, patients were treated for one year.

**Table 4** describes the regimens and patient exposure for one year of treatment in patients with no previous exposure to interferon (interferon-naïve patients). Because of a significant overlap in the pattern of undesirable effects with PegIntron monotherapy, groups of patients have been brought together in **Table 5** to show the pattern of reported effects for all monotherapy groups.

<b>Table 4</b> Regimens and patient exposure		
<b>Treatment</b>	<b>Regimen</b>	<b>Number of patients treated for one year</b>
PegIntron + ribavirin	PegIntron (1.5 micrograms/kg/week) + ribavirin (> 10.6 mg/kg/day)	188
Interferon alfa-2b + ribavirin	Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day)	505
PegIntron monotherapy	PegIntron (0.5 microgram/kg/week)	315
	PegIntron (1.0 microgram/kg/week)	297
	PegIntron (1.5 micrograms/kg/week)	304

<b>Table 5</b> Undesirable effects very commonly reported in clinical trials > 10 % of patients in PegIntron + ribavirin group Very common (> 1/10)			
	<b>PegIntron + ribavirin</b>	<b>Interferon alfa- 2b + ribavirin</b>	<b>PegIntron monotherapy</b>
<b>Infections and infestations</b>			
Infection viral	10 %	5 %	4-5 %
<b>Metabolism and nutrition disorders</b>			
Weight decrease	30 %	19 %	8-18 %
<b>Psychiatric disorders</b>			
Depression	34 %	32 %	26 %
Irritability	32 %	34 %	19 %
Insomnia	37 %	41 %	16-19 %
Anxiety	14 %	14 %	8 %
Concentration impaired	18 %	21 %	9-10 %
Emotional lability	11 %	10 %	5 %
<b>Nervous system disorders</b>			
Headache	58 %	57 %	57-63 %
Mouth dry	10 %	8 %	4-8 %
<b>Respiratory, thoracic and mediastinal disorders</b>			
Pharyngitis	10 %	7 %	3 %
Coughing	14 %	11 %	4 %
Dyspnea	26 %	22 %	5 %
<b>Gastrointestinal disorders</b>			
Nausea	43 %	31 %	20-23 %
Anorexia	35 %	26 %	10-25 %
Diarrhoea	20 %	13 %	14-17 %
Abdominal pain	12 %	9 %	11 %
Vomiting	16 %	10 %	4-7 %
<b>Skin and subcutaneous tissue disorders</b>			
Alopecia	45 %	32 %	20-34 %
Pruritus	27 %	27 %	7-9 %
Skin dry	23 %	21 %	6-9 %
Rash	21 %	21 %	5-7 %
<b>Musculoskeletal and connective tissue disorders</b>			
Myalgia	49 %	49 %	46-60 %
Arthralgia	31 %	26 %	23-28 %
Musculoskeletal pain	15 %	11 %	11-13 %
<b>General disorders and administration site conditions</b>			
Injection site inflammation	20 %	17 %	39-44 %
Injection site reaction	54 %	36 %	7-9 %
Fatigue	56 %	59 %	43 %
Rigors	42 %	40 %	33-43 %
Fever	39 %	32 %	29-43 %
Flu-like symptoms	21 %	23 %	18-25 %
Asthenia	28 %	17 %	12-14 %
Dizziness	17 %	16 %	7-12 %

<b>Table 6</b> Undesirable effects commonly reported in clinical trials in patients treated with PegIntron + ribavirin or PegIntron monotherapy Common (> 1/100, < 1/10)		
<b>Body system</b>	<b>5-10%</b>	<b>1-&lt;5%</b>
Infections and infestations		otitis media, fungal infection, bacterial infection, herpes simplex
Blood and lymphatic system disorders	Anaemia, leukopaenia	Thrombocytopenia, lymphadenopathy
Endocrine disorders	hypothyroidism	hyperthyroidism
Metabolism and nutrition disorders		hyperuricemia, hypocalcemia, thirst
Psychiatric disorders	agitation, nervousness	aggressive behaviour, somnolence, behavior disorder, apathy, appetite increased, sleep disorder, dreaming abnormal, decreased libido
Nervous system disorders	paresthesia, increased sweating	hypoesthesia, hyperaesthesia, hypertonia, confusion, tremor, vertigo, migraine, ataxia, neuralgia
Eye disorders		blurred vision, conjunctivitis, lacrimal gland disorder, eye pain
Cardiac disorders		tachycardia, palpitation
Vascular disorders		hypotension, hypertension, syncope, flushing
Ear and labyrinth disorders		tinnitus, hearing impairment/loss
Respiratory, thoracic and mediastinal disorders		nonproductive cough, rhinitis, sinusitis, bronchitis, respiratory disorder, nasal congestion, rhinorrhea, dysphonia, epistaxis
Gastrointestinal disorders	dyspepsia	constipation, taste perversion, loose stools, stomatitis, ulcerative stomatitis, gingival bleeding, glossitis, flatulence, hemorrhoids, gastroesophageal reflux, gingivitis, dehydration
Hepatobiliary disorders		hepatomegaly, hyperbilirubinemia
Skin and subcutaneous tissue disorders		erythematous rash, eczema, photosensitivity reaction, maculopapular rash, abnormal hair texture, acne, dermatitis, furunculosis, nail disorder, psoriasis, urticaria, erythema, face or peripheral oedema
Musculoskeletal and connective tissue disorders		arthritis
Renal and urinary disorders		micturition frequency, urine abnormal
Reproductive system and breast disorders	menstrual disorder, menorrhagia	ovarian disorder, vaginal disorder, sexual dysfunction (not specified), impotence, breast pain, amenorrhoea, prostatitis
General disorders and administration site conditions	RUQ pain, malaise, chest pain	injection site pain

Most cases of neutropaenia and thrombocytopaenia were mild (WHO grades 1 or 2). There were some cases of more severe neutropenia in patients treated with the recommended doses of PegIntron in combination with ribavirin (WHO grade 3: 39 of 186 [21 %]; and WHO grade 4: 13 of 186 [7 %]).

In a clinical trial, approximately 1.2 % of patients treated with PegIntron or interferon alfa-2b in combination with ribavirin reported life-threatening psychiatric events during treatment. These events included suicidal ideation and attempted suicide (see section 4.4). Following marketing, psychosis and hallucination have been reported rarely.

Rarely ( $> 1/10,000$ ,  $< 1/1,000$ ) or very rarely ( $< 1/10,000$ ) reported events with interferon alfa-2b, including PegIntron, include:

**Blood and lymphatic system disorders:**

Very rarely interferon alfa-2b or PegIntron used alone or in combination with ribavirin may be associated with aplastic anaemia. Pure red cell aplasia has been reported.

**Immune system disorders:**

very rarely: sarcoidosis or exacerbation of sarcoidosis

**Endocrine disorders:**

rarely: diabetes

**Nervous system disorders:**

rarely: seizure, peripheral neuropathy

very rarely: cerebrovascular ischaemia, cerebrovascular haemorrhage, encephalopathy

**Cardiac disorders:**

rarely: arrhythmia

very rarely: cardiac ischaemia, myocardial infarction

**Respiratory, thoracic and mediastinal disorders:**

very rarely: interstitial lung disease

**Gastrointestinal disorders:**

rarely: pancreatitis

very rarely: ulcerative and ischaemic colitis

**Skin and subcutaneous tissue disorders:**

very rarely: erythema multiforme, Stevens Johnson syndrome, toxic epidermal necrolysis, injection site necrosis

**Musculoskeletal and connective tissue disorders:**

rarely: rhabdomyolysis, myositis

**Renal and urinary disorders:**

rarely: renal insufficiency and renal failure

Cardiovascular (CVS) adverse events, particularly arrhythmia, appeared to be correlated mostly with pre-existing CVS disease and prior therapy with cardiotoxic agents (see section 4.4). Cardiomyopathy, that may be reversible upon discontinuation of interferon alpha, has been reported rarely in patients without prior evidence of cardiac disease.

Ophthalmological disorders that have been reported rarely with alpha interferons include retinopathies (including macular oedema), retinal haemorrhages, retinal artery or vein obstruction, cotton wool spots, loss of visual acuity or visual field, optic neuritis, and papilloedema (see section 4.4).

A wide variety of autoimmune and immune-mediated disorders have been reported with alpha interferons including thyroid disorders, systemic lupus erythematosus, rheumatoid arthritis (new or aggravated), idiopathic and thrombotic thrombocytopenic purpura, vasculitis, neuropathies including mononeuropathies (see also section 4.4, **Autoimmune disorders**).

**HCV/HIV co-infected patients**

For HCV/HIV co-infected patients receiving PegIntron in combination with ribavirin, other undesirable effects (that were not reported in mono-infected patients) which have been reported in the larger studies with a frequency  $> 5\%$  were: oral candidiasis (14 %), lipodystrophy acquired (13 %), CD4 lymphocytes decreased (8 %), appetite decreased (8 %), gamma-glutamyltransferase increased (9 %), back pain (5 %), blood amylase increased (6 %), blood lactic acid increased (5 %), cytolytic hepatitis (6 %), lipase increased (6 %) and pain in limb (6 %).

*Mitochondrial toxicity:*

Mitochondrial toxicity and lactic acidosis have been reported in HIV-positive patients receiving NRTI



regimen and associated ribavirin for co-HCV infection (see section 4.4).

*Laboratory values for HCV/HIV co-infected patients:*

Although haematological toxicities of neutropenia, thrombocytopenia and anaemia occurred more frequently in HCV/HIV co-infected patients, the majority could be managed by dose modification and rarely required premature discontinuation of treatment (see section 4.4). Haematological abnormalities were more frequently reported in patients receiving PegIntron in combination with ribavirin when compared to patients receiving interferon alfa-2b in combination with ribavirin. In Study 1 (see section 5.1), decrease in absolute neutrophil count levels below 500 cells/mm<sup>3</sup> was observed in 4 % (8/194) of patients and decrease in platelets below 50,000/mm<sup>3</sup> was observed in 4 % (8/194) of patients receiving PegIntron in combination with ribavirin. Anaemia (hemoglobin < 9.4g/dl) was reported in 12% (23/194) of patients treated with PegIntron in combination with ribavirin.

*CD4 lymphocytes decrease:*

Treatment with PegIntron in combination with ribavirin was associated with decreases in absolute CD4+ cell counts within the first 4 weeks without a reduction in CD4+ cell percentage. The decrease in CD4+ cell counts was reversible upon dose reduction or cessation of therapy. The use of PegIntron in combination with ribavirin had no observable negative impact on the control of HIV viraemia during therapy or follow-up. Limited safety data (N= 25) are available in co-infected patients with CD4+ cell counts < 200/μl (see section 4.4).

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron in combination with ribavirin.

## **4.9 Overdose**

In clinical trials, cases of accidental overdose, at never more than twice the prescribed dose, were reported. There were no serious reactions. Undesirable effects resolved during continued administration of PegIntron.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Immunostimulants, Cytokines and immunomodulators, Interferons, Peginterferon alfa-2b, ATC code: L03A B10.

Recombinant interferon alfa-2b is covalently conjugated with monomethoxy polyethylene glycol at an average degree of substitution of 1 mole of polymer/mole of protein. The average molecular mass is approximately 31,300 daltons of which the protein moiety constitutes approximately 19,300.

#### **Interferon alfa-2b**

*In vitro* and *in vivo* studies suggest that the biological activity of PegIntron is derived from its interferon alfa-2b moiety.

Interferons exert their cellular activities by binding to specific membrane receptors on the cell surface. Studies with other interferons have demonstrated species specificity. However, certain monkey species, e.g., Rhesus monkeys, are susceptible to pharmacodynamic stimulation upon exposure to human type 1 interferons.

Once bound to the cell membrane, interferon initiates a complex sequence of intracellular events that include the induction of certain enzymes. It is thought that this process, at least in part, is responsible for the various cellular responses to interferon, including inhibition of virus replication in virus-infected cells, suppression of cell proliferation and such immunomodulating activities as enhancement of the phagocytic activity of macrophages and augmentation of the specific cytotoxicity of

lymphocytes for target cells. Any or all of these activities may contribute to interferon's therapeutic effects.

Recombinant interferon alfa-2b also inhibits viral replication *in vitro* and *in vivo*. Although the exact antiviral mode of action of recombinant interferon alfa-2b is unknown, it appears to alter the host cell metabolism. This action inhibits viral replication or if replication occurs, the progeny virions are unable to leave the cell.

### **PegIntron**

PegIntron pharmacodynamics were assessed in a rising single-dose trial in healthy subjects by examining changes in oral temperature, concentrations of effector proteins such as serum neopterin and 2'5'-oligoadenylate synthetase (2'5'-OAS), as well as white cell and neutrophil counts. Subjects treated with PegIntron showed mild dose-related elevations in body temperature. Following single doses of PegIntron between 0.25 and 2.0 micrograms/kg/week, serum neopterin concentration was increased in a dose-related manner. Neutrophil and white cell count reductions at the end of week 4 correlated with the dose of PegIntron.

### **PegIntron clinical trials**

Two pivotal trials have been conducted, one (C/I97-010) with PegIntron monotherapy; the other (C/I98-580) with PegIntron in combination with ribavirin. Eligible patients for these trials had chronic hepatitis C confirmed by a positive HCV-RNA polymerase chain reaction (PCR) assay (> 30 IU/ml), a liver biopsy consistent with a histological diagnosis of chronic hepatitis with no other cause for the chronic hepatitis, and abnormal serum ALT.

In the PegIntron monotherapy trial, a total of 916 naïve chronic hepatitis C patients were treated with PegIntron (0.5, 1.0 or 1.5 micrograms/kg/week) for one year with a follow-up period of six months. In addition, 303 patients received interferon alfa-2b (3 million International Units [MIU] three times a week) as a comparator. This study showed that PegIntron was superior to interferon alfa-2b (**Table 8**).

In the PegIntron combination trial, 1,530 naïve patients were treated for one year with one of the following combination regimens:

- PegIntron (1.5 micrograms/kg/week) + ribavirin (800 mg/day), (n = 511).
- PegIntron (1.5 micrograms/kg/week for one month followed by 0.5 microgram/kg/week for 11 months) + ribavirin (1,000/1,200 mg/day), (n = 514).
- Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day) (n = 505).

In this trial, the combination of PegIntron (1.5 micrograms/kg/week) and ribavirin was significantly more effective than the combination of interferon alfa-2b and ribavirin (**Table 7**), particularly in patients infected with Genotype 1 (**Table 8**). Sustained response was assessed by the response rate six months after the cessation of treatment.

HCV genotype and baseline virus load are prognostic factors which are known to affect response rates. However, response rates in this trial were shown to be dependent also on the dose of ribavirin administered in combination with PegIntron or interferon alfa-2b. In those patients that received > 10.6 mg/kg ribavirin (800 mg dose in typical 75 kg patient), regardless of genotype or viral load, response rates were significantly higher than in those patients that received ≤ 10.6 mg/kg ribavirin (**Table 8**), while response rates in patients that received > 13.2 mg/kg ribavirin were even higher.

<b>Table 7</b> Sustained virological response (% patients HCV negative)							
	<b>PegIntron monotherapy</b>				<b>PegIntron + ribavirin</b>		
<b>Treatment regimen</b>	<b>P 1.5</b>	<b>P 1.0</b>	<b>P 0.5</b>	<b>I</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>Number of patients</b>	304	297	315	303	511	514	505
<b>Response at end of treatment</b>	49 %	41 %	33 %	24 %	65 %	56 %	54 %

<b>Sustained response</b>	<b>23 %*</b>	<b>25 %</b>	<b>18 %</b>	<b>12 %</b>	<b>54 %**</b>	<b>47 %</b>	<b>47 %</b>
---------------------------	--------------	-------------	-------------	-------------	---------------	-------------	-------------

P 1.5 PegIntron 1.5 micrograms/kg  
 P 1.0 PegIntron 1.0 microgram/kg  
 P 0.5 PegIntron 0.5 microgram/kg  
 I Interferon alfa-2b 3 MIU  
 P 1.5/R PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)  
 P 0.5/R PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)  
 I/R Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)  
 \* p < 0.001 P 1.5 vs. I  
 \*\* p = 0.0143 P 1.5/R vs. I/R

<b>Table 8</b> Sustained response rates with PegIntron + ribavirin (by ribavirin dose, genotype and viral load)				
<b>HCV Genotype</b>	<b>Ribavirin dose (mg/kg)</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>All Genotypes</b>	<b>All</b>	<b>54 %</b>	<b>47 %</b>	<b>47 %</b>
	≤ 10.6	50 %	41 %	27 %
	> 10.6	61 %	48 %	47 %
<b>Genotype 1</b>	<b>All</b>	<b>42 %</b>	<b>34 %</b>	<b>33 %</b>
	≤ 10.6	38 %	25 %	20 %
	> 10.6	48 %	34 %	34 %
Genotype 1 ≤ 600,000 IU/ml	All	73 %	51 %	45 %
	≤ 10.6	74 %	25 %	33 %
	> 10.6	71 %	52 %	45 %
Genotype 1 > 600,000 IU/ml	All	30 %	27 %	29 %
	≤ 10.6	27 %	25 %	17 %
	> 10.6	37 %	27 %	29 %
<b>Genotype 2/3</b>	<b>All</b>	<b>82 %</b>	<b>80 %</b>	<b>79 %</b>
	≤ 10.6	79 %	73 %	50 %
	> 10.6	88 %	80 %	80 %

P 1.5/R PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)  
 P 0.5/R PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)  
 I/R Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)

In the PegIntron monotherapy study, the Quality of Life was generally less affected by 0.5 microgram/kg of PegIntron than by either 1.0 microgram/kg of PegIntron once weekly or 3 MIU of interferon alfa-2b three times a week.

In a separate trial, 224 patients with genotype 2 or 3 received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with ribavirin 800 mg –1,400 mg p.o. for 6 months (based on body weight, only three patients weighing > 105 kg, received the 1,400 mg dose, which has not yet been validated) (**Table 9**). Twenty-four % had bridging fibrosis or cirrhosis (Knodell 3/4).

**Table 9. Virologic Response at End of Treatment, Sustained Virologic Response and Relapse by HCV Genotype and Viral Load\***

	PegIntron 1.5 µg/kg once weekly plus Ribavirin 800-1,400 mg/day		
	End of Treatment Response	Sustained Virologic Response	Relapse
All Subjects	94 % (211/224)	81 % (182/224)	12 % (27/224)
HCV 2 ≤ 600,000 IU/ml > 600,000	100 % (42/42)	93 % (39/42)	7 % (3/42)
	100 % (20/20)	95 % (19/20)	5 % (1/20)
	100 % (22/22)	91 % (20/22)	9 % (2/22)

IU/ml			
HCV 3	93 % (169/182)	79 % (143/182)	14 % (24/166)
≤ 600,000 IU/ml	93 % (92/99)	86 % (85/99)	8 % (7/91)
> 600,000 IU/ml	93 % (77/83)	70 % (58/83)	23 % (17/75)

\* Any subject with an undetectable HCV-RNA level at the Follow-Up Week 12 visit and missing data at the Follow-Up Week 24 visit was considered a sustained responder. Any subject with missing data in and after the Follow-Up Week 12 window was considered to be a non-responder at Week 24 of follow-up.

The 6 month treatment duration in this trial was better tolerated than one year of treatment in the pivotal combination trial; for discontinuation 5 % vs. 14 %, for dose modification 18 % vs. 49 %.

In a non-comparative trial, 235 patients with genotype 1 and low viral load (< 600,000 IU/ml) received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with weight adjusted ribavirin. The overall sustained response rate after a 24-week treatment duration was 50 %. Forty-one percent of subjects (97/235) had nondetectable plasma HCV-RNA levels at Week 4 and Week 24 of therapy. In this subgroup, there was a 92 % (89/97) sustained virological response rate. The high sustained response rate in this subgroup of patients was identified in an interim analysis (n=49) and prospectively confirmed (n=48).

Limited historical data indicate that treatment for 48 weeks might be associated with a higher sustained response rate (11/11) and with a lower risk of relapse (0/11 as compared to 7/96 following 24 weeks of treatment).

### Predictability of sustained virological response

Virological response by week 12, defined as a 2-log viral load decrease or undetectable levels of HCV-RNA has been shown to be predictive for sustained response (**Table 10**).

<b>Table 10</b> Predictability of sustained response by viral response at week 12 and genotype*				
Treatment	Genotype	Viral response at week 12	Sustained response	Negative predictive value
PegIntron 1.5 + ribavirin (> 10.6 mg/kg) 48-week treatment	1	Yes 75 % (82/110)	71 % (58/82)	----
		No 25 % (28/110)	0 % (0/28)	100 %
PegIntron 1.5 + ribavirin 800-1,400 mg 24-week treatment	2 and 3	Yes 99 % (213/215)	83 % (177/213)	----
		No 1 % (2/215)	50 % (1/2)	50 %

\* reflects patients with 12 week data available

The negative predictive value for sustained response in patients treated with PegIntron in monotherapy was 98 %.

### HCV/HIV Co-infected patients

Two trials have been conducted in patients co-infected with HIV and HCV. The response to treatment in both of these trials is presented in **Table 11**. Study 1 (RIBAVIC; P01017) was a randomized, multicentre study which enrolled 412 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (1.5 µg/kg/week) plus ribavirin (800 mg/day) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800 mg/day) for 48 weeks with a follow-up period of 6 months. Study 2 (P02080) was a randomized, single centre

study that enrolled 95 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (100 or 150 µg /week based on weight) plus ribavirin (800-1,200 mg/day based on weight) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800-1,200 mg/day based on weight). The duration of therapy was 48 weeks with a follow-up period of 6 months except for patients infected with genotypes 2 or 3 and viral load < 800,000 IU/ml (Amplicor) who were treated for 24 weeks with a 6-month follow-up period.

<b>Table 11</b> Sustained virological response based on genotype after PegIntron in combination with Ribavirin in HCV/HIV Co-infected patients						
	<b>Study 1<sup>1</sup></b>			<b>Study 2<sup>2</sup></b>		
	PegIntron (1.5 µg/kg/week) + ribavirin (800 mg)	Interferon alfa-2b (3 MIU TIW) + ribavirin (800 mg)	p value <sup>a</sup>	PegIntron (100 or 150 <sup>c</sup> µg/week) + ribavirin (800-1,200 mg) <sup>d</sup>	Interferon alfa-2b (3 MIU TIW) + ribavirin (800-1,200 mg) <sup>d</sup>	p value <sup>b</sup>
All	27 % (56/205)	20 % (41/205)	0.047	44 % (23/52)	21 % (9/43)	0.017
Genotype 1, 4	17 % (21/125)	6 % (8/129)	0.006	38 % (12/32)	7 % (2/27)	0.007
Genotype 2, 3	44 % (35/80)	43 % (33/76)	0.88	53 % (10/19)	47 % (7/15)	0.730

MIU = million international units; TIW = three times a week.

a: p value based on Cochran-Mantel Haenszel Chi square test.

b: p value based on chi-square test.

c: subjects < 75 kg received 100 µg/week PegIntron and subjects ≥ 75 kg received 150 µg/week PegIntron.

d: ribavirin dosing was 800 mg for patients < 60 kg, 1,000 mg for patients 60-75 kg, and 1,200 mg for patients > 75 kg.

<sup>1</sup>Carrato F, Bani-Sadir F, Pol S et al. JAMA 2004; 292(23): 2839-2848.

<sup>2</sup>Laguno M, Murillas J, Blanco J et al. AIDS 2004; 18(13): F27-F36.

### *Histological response*

Liver biopsies were obtained before and after treatment in Study 1 and were available for 210 of the 412 subjects (51 %). Both the Metavir score and Ishak grade decreased among subjects treated with PegIntron in combination with ribavirin. This decline was significant among responders (-0.3 for Metavir and -1.2 for Ishak) and stable (-0.1 for Metavir and -0.2 for Ishak) among non-responders. In terms of activity, about one-third of sustained responders showed improvement and none showed worsening. There was no improvement in terms of fibrosis observed in this study. Steatosis was significantly improved in patients infected with HCV Genotype 3.

## **5.2 Pharmacokinetic properties**

PegIntron is a well characterized polyethylene glycol-modified (“pegylated”) derivative of interferon alfa-2b and is predominantly composed of monopegylated species. The plasma half-life of PegIntron is prolonged compared with non-pegylated interferon alfa-2b. PegIntron has a potential to depegylate to free interferon alfa-2b. The biologic activity of the pegylated isomers is qualitatively similar, but weaker than free interferon alfa-2b.

Following subcutaneous administration, maximal serum concentrations occur between 15-44 hours post-dose, and are sustained for up to 48-72 hours post-dose.

PegIntron C<sub>max</sub> and AUC measurements increase in a dose-related manner. Mean apparent volume of distribution is 0.99 l/kg.

Upon multiple dosing, there is an accumulation of immunoreactive interferons. There is, however, only a modest increase in biologic activity as measured by a bioassay.

Mean (SD) PegIntron elimination half-life is approximately 40 hours (13.3 hours), with apparent clearance of 22.0 ml/hr·kg. The mechanisms involved in clearance of interferons in man have not yet been fully elucidated. However, renal elimination may account for a minority (approximately 30 %) of PegIntron apparent clearance.

**Renal function:** Renal clearance appears to account for 30 % of total clearance of PegIntron. In a single dose study (1.0 microgram/kg) in patients with impaired renal function,  $C_{max}$ , AUC, and half-life increased in relation to the degree of renal impairment.

Following multiple dosing of PegIntron (1.0 microgram/kg subcutaneously administered every week for four weeks) the clearance of PegIntron is reduced by a mean of 17 % in patients with moderate renal impairment (creatinine clearance 30-49 ml/minute) and by a mean of 44 % in patients with severe renal impairment (creatinine clearance 15-29 ml/minute) compared to subjects with normal renal function. Based on single dose data, clearance was similar in patients with severe renal impairment not on dialysis and in patients who were receiving hemodialysis. The dose of PegIntron for monotherapy should be reduced in patients with moderate or severe renal impairment (see sections 4.2 and 4.4). Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see section 4.3).

Because of marked inter-subject variability in interferon pharmacokinetics, it is recommended that patients with severe renal impairment be closely monitored during treatment with PegIntron (see section 4.2)

**Hepatic function:** The pharmacokinetics of PegIntron have not been evaluated in patients with severe hepatic dysfunction.

**Elderly patients ≥ 65 years of age:** The pharmacokinetics of PegIntron following a single subcutaneous dose of 1.0 microgram/kg were not affected by age. The data suggest that no alteration in PegIntron dosage is necessary based on advancing age.

**Patients under the age of 18 years:** Specific pharmacokinetic evaluations have not been performed on these patients. PegIntron is indicated for the treatment of chronic hepatitis C only in patients 18 years of age or older.

**Interferon neutralising factors:** Interferon neutralising factor assays were performed on serum samples of patients who received PegIntron in the clinical trial. Interferon neutralising factors are antibodies which neutralise the antiviral activity of interferon. The clinical incidence of neutralising factors in patients who received PegIntron 0.5 micrograms/kg is 1.1 %.

### 5.3 Preclinical safety data

**PegIntron:** Adverse events not observed in clinical trials were not seen in toxicity studies in monkeys. These studies were limited to four weeks due to the appearance of anti-interferon antibodies in most monkeys.

Reproduction studies of PegIntron have not been performed. Interferon alfa-2b has been shown to be an abortifacient in primates. PegIntron is likely to also cause this effect. Effects on fertility have not been determined. It is not known whether the components of this medicinal product are excreted into experimental animal or human milk (see section 4.6 for relevant human data on pregnancy and lactation). PegIntron showed no genotoxic potential.

The relative non-toxicity of monomethoxy-polyethylene glycol (mPEG), which is liberated from PegIntron by metabolism *in vivo* has been demonstrated in preclinical acute and subchronic toxicity studies in rodents and monkeys, standard embryo-foetal development studies and in *in vitro* mutagenicity assays.

**PegIntron plus ribavirin:** When used in combination with ribavirin, PegIntron did not cause any effects not previously seen with either active substance alone. The major treatment-related change was a reversible, mild to moderate anaemia, the severity of which was greater than that produced by either active substance alone.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Powder for solution for injection:

Disodium phosphate, anhydrous,  
Sodium dihydrogen phosphate dihydrate,  
Sucrose,  
Polysorbate 80.

Solvent for parenteral use:

Water for injections.

### **6.2 Incompatibilities**

This medicinal product should only be reconstituted with the solvent provided (see section 6.6). In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

3 years

After reconstitution:

- Chemical and physical in-use stability has been demonstrated for 24 hours at 2°C - 8°C.
- From a microbiological point of view, the product is to be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C - 8°C.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C - 8°C).

### **6.5 Nature and contents of container**

The powder is contained in a 2 ml vial, Type I flint glass, with a butyl rubber stopper in an aluminium flip-off seal with a polypropylene bonnet. The solvent is presented in a 2 ml ampoule, Type I flint glass. PegIntron 100 micrograms is supplied as:

- 1 vial of powder for solution for injection and 1 ampoule of solvent for parenteral use;
- 1 vial of powder for solution for injection, 1 ampoule of solvent for parenteral use, 1 injection syringe, 2 injection needles and 1 cleansing swab;
- 4 vials of powder for solution for injection and 4 ampoules of solvent for parenteral use;
- 4 vials of powder for solution for injection, 4 ampoules of solvent for parenteral use, 4 injection syringes, 8 injection needles and 4 cleansing swabs;
- 6 vials of powder for solution for injection and 6 ampoules of solvent for parenteral use.
- 12 vials of powder for solution for injection, 12 ampoules of solvent for parenteral use, 12 injection syringes, 24 injection needles and 12 cleansing swabs.

Not all pack sizes may be marketed

### **6.6 Special precautions for disposal**

PegIntron is supplied as a powder of peginterferon alfa-2b at a strength of 100 micrograms for single use. Each vial must be reconstituted with 0.7 ml of water for injections for administration of up to 0.5 ml of solution. A small volume is lost during preparation of PegIntron for injection when the dose is measured and injected. Therefore, each vial contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection. The reconstituted solution has a concentration of 100 micrograms/0.5 ml.

Using a sterilised injection syringe and injection needle, inject 0.7 ml of water for injections into the vial of PegIntron. Agitate gently to complete dissolution of powder. The appropriate dose can then be withdrawn with a sterilised injection syringe and injected. A complete set of instructions is provided in the Annex to the Package Leaflet.

As for all parenteral medicinal products, inspect visually the reconstituted solution prior to administration. The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. Discard any unused material.

## **7. MARKETING AUTHORISATION HOLDER**

SP Europe  
73, rue de Stalle  
B-1180 Bruxelles  
Belgium

## **8. MARKETING AUTHORISATION NUMBERS**

EU/1/00/131/011  
EU/1/00/131/012  
EU/1/00/131/013  
EU/1/00/131/014  
EU/1/00/131/015  
EU/1/00/131/028

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 25 May 2000  
Date of last renewal: 25 May 2005

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>



## **1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 120 micrograms powder and solvent for solution for injection

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each vial of PegIntron, powder for solution for injection contains 120 micrograms of peginterferon alfa-2b as measured on a protein basis.

Each vial provides 120 micrograms/0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

The active substance is a covalent conjugate of recombinant interferon alfa-2b\* with monomethoxy polyethylene glycol. The potency of this product should not be compared to that of another pegylated or non-pegylated protein of the same therapeutic class. For more information, see section 5.1.

\*produced by rDNA technology in *E.coli* cells harbouring a genetically engineered plasmid hybrid encompassing an interferon alfa-2b gene from human leukocytes

For a full list of excipients, see section 6.1. PegIntron contains 40 mg of sucrose per 0.5 ml.

## **3. PHARMACEUTICAL FORM**

Powder and solvent for solution for injection

White powder.

Clear and colourless solvent.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

PegIntron is indicated for the treatment of adult patients with chronic hepatitis C who have elevated transaminases without liver decompensation and who are positive for serum HCV-RNA or anti-HCV, including naïve patients with clinically stable HIV co-infection (see section 4.4).

The best way to use PegIntron in this indication is in combination with ribavirin.

This combination is indicated in naïve patients including patients with clinically stable HIV co-infection. This combination of PegIntron with ribavirin is also indicated in patients who have previously responded (with normalisation of ALT at the end of treatment) to interferon alpha monotherapy but who have subsequently relapsed.

Interferon monotherapy, including PegIntron, is indicated mainly in case of intolerance or contraindication to ribavirin.

Please refer also to the ribavirin Summary of Product Characteristics (SPC) when PegIntron is to be used in combination with ribavirin.

### **4.2 Posology and method of administration**

Treatment should be initiated and monitored only by a physician experienced in the management of patients with hepatitis C.

#### **Dose to be administered**

PegIntron should be administered as a once weekly subcutaneous injection. The dose administered depends on whether it is used in combination with ribavirin or as monotherapy.

## Combination therapy

PegIntron 1.5 micrograms/kg/week in combination with ribavirin capsules.

The intended dose of 1.5 µg/kg of PegIntron to be used in combination with ribavirin may be delivered in weight categories with the pen/vial strengths according to **Table 1**. Ribavirin capsules are to be administered orally each day in two divided doses with food (morning and evening).

**Table 1 - Dosing for Combination Therapy**

Body Weight (kg)	PegIntron		Ribavirin Capsules	
	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Total Daily Dose (mg)	Number of Capsules (200 mg)
< 40	50	0.5	800	4 <sup>a</sup>
40-50	80	0.4	800	4 <sup>a</sup>
51-64	80	0.5	800	4 <sup>a</sup>
65-75	100	0.5	1,000	5 <sup>b</sup>
76-85	120	0.5	1,000	5 <sup>b</sup>
> 85	150	0.5	1,200	6 <sup>c</sup>

a: 2 morning, 2 evening

b: 2 morning, 3 evening

c: 3 morning, 3 evening

## Duration of treatment

**Predictability of sustained virological response:** Patients infected with virus genotype 1 who fail to achieve virological response at Week 12 are highly unlikely to become sustained virological responders (see also section 5.1).

- **Genotype 1:** For patients who exhibit virological response at week 12, treatment should be continued for another nine month period (i.e., a total of 48 weeks).  
In the subset of patients with genotype 1 infection and low viral load (< 600,000 IU/ml) who become HCV-RNA negative at treatment week 4 and remain HCV-RNA negative at week 24, the treatment could either be stopped after this 24 week treatment course or pursued for an additional 24 weeks (i.e. overall 48 weeks treatment duration). However, an overall 24 weeks treatment duration may be associated with a higher risk of relapse than a 48 weeks treatment duration (see section 5.1).
- **Genotypes 2 or 3:** It is recommended that all patients be treated for 24 weeks, except for HCV/HIV co-infected patients who should receive 48 weeks of treatment.
- **Genotype 4:** In general, patients infected with genotype 4 are considered harder to treat and limited study data (n=66) indicate they are compatible with a duration of treatment as for genotype 1.

## HCV/HIV Co-infection

The recommended duration of dosing for HCV/HIV co-infected patients is 48 weeks, regardless of genotype.

### *Predictability of response and non-response in HCV/HIV Co-infection*

Early virological response by Week 12, defined as a 2 log viral load decrease or undetectable levels of HCV-RNA, has been shown to be predictive for sustained response. The negative predictive value for sustained response in HCV/HIV co-infected patients treated with PegIntron in combination with ribavirin was 99 % (67/68; Study 1) (see section 5.1). A positive predictive value of 50 % (52/104; Study 1) was observed for HCV/HIV co-infected patients receiving combination therapy.

### PegIntron monotherapy

As monotherapy the PegIntron regimen is 0.5 or 1.0 microgram/kg/week. The lowest vial or pen strength available is 50 µg/0.5 ml; therefore for patients prescribed 0.5 µg/kg/week, doses must be adjusted by volume as shown in **Table 2**. For the 1.0 µg/kg dose, similar volume adjustments can be made or alternate vial strengths can be used as shown in **Table 2**. PegIntron monotherapy was not studied in HCV/HIV co-infected patients.

**Table 2- Monotherapy Dosing**

	0.5 µg/kg		1.0 µg/kg	
Body Weight (kg)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)
30-35	50*	0.15	50	0.3
36-45	50*	0.2	50	0.4
46-56	50*	0.25	50	0.5
57-72	50	0.3	80	0.4
73-88	50	0.4	80	0.5
89-106	50	0.5	100	0.5
> 106**	80	0.4	120	0.5
* Must use vial. Minimum delivery for pen is 0.3 ml.				
** For patients > 120 kg, use 80 µg/0.5 ml vial				

### Duration of treatment

For patients who exhibit virological response at Week 12, treatment should be continued for at least another three-month period (i.e., a total of six months). The decision to extend therapy to one year of treatment should be based on prognostic factors (e.g., genotype, age > 40 years, male gender, bridging fibrosis).

### Dose modification for all patients

If severe adverse reactions or laboratory abnormalities develop during treatment with PegIntron monotherapy or PegIntron in combination with ribavirin, modify the dosages of each product as appropriate, until the adverse reactions abate. As adherence might be of importance for outcome of therapy, the dose should be kept as close as possible to the recommended standard dose. Guidelines were developed in clinical trials for dose modification.

### Combination Therapy Dose Reduction Guidelines

<b>Table 2a</b> Dose modification guidelines for combination therapy (with ribavirin)			
Laboratory values	Reduce only ribavirin dose to <u>600 mg/day</u> * if:	Reduce only PegIntron dose to one-half dose if:	Discontinue combination therapy if:
Haemoglobin	< 10 g/dl	-	< 8.5 g/dl
Haemoglobin in: Patients with history of stable cardiac disease	≥ 2 g/dl decrease in haemoglobin during any four week period during treatment (permanent dose reduction)		< 12 g/dl after four weeks of dose reduction
White blood cells	-	< 1.5 x 10 <sup>9</sup> /l	< 1.0 x 10 <sup>9</sup> /l
Neutrophils	-	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	-	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l

Bilirubin – direct	-	-	2.5 x ULN**
Bilirubin - indirect	> 5 mg/dl	-	> 4 mg/dl (for > 4 weeks)
Creatinine	-	-	> 2.0 mg/dl
ALT/AST	-	-	2 x baseline and > 10 x ULN**

\* Patients whose dose of ribavirin is reduced to 600 mg daily receive one 200 mg capsule in the morning and two 200 mg capsules in the evening.

\*\* Upper limit of normal

Dose reduction of PegIntron may be accomplished by either reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 2b**.

<b>Table 2b – Reduced PegIntron Dosing for Combination Therapy</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5 ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
< 40	25	50*	0.25	25
40-50	32	50	0.3	30
51-64	40	50	0.4	40
65-75	50	50	0.5	50
76-85	60	80	0.4	64
> 85	75	100	0.4	80
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### **PegIntron Monotherapy Dose Reduction Guidelines**

Dose modification guidelines for patients who use PegIntron monotherapy are shown in **Table 3a**.

<b>Table 3a</b> Dose modification guidelines for PegIntron monotherapy		
<b>Laboratory values</b>	<b>Reduce PegIntron to one-half dose if:</b>	<b>Discontinue PegIntron if:</b>
Neutrophils	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l

Dose reduction for patients who use 0.5 µg/kg PegIntron monotherapy must be accomplished by reducing the prescribed volume by one-half. The 50 µg/0.5 ml vial must be used if necessary since the pen can only deliver a minimum volume of 0.3 ml.

For patients who use 1.0 µg/kg PegIntron monotherapy, dose reduction may be accomplished by reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 3b**.

<b>Table 3b – Reduced PegIntron Dose for the 1.0 µg/kg Monotherapy Regimen</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
30-35	15	50*	0.15	15
36-45	20	50*	0.20	20
46-56	25	50*	0.25	25
57-72	32	50	0.3	30
73-89	40	50	0.4	40
90-106	50	50	0.5	50
> 106	60	80	0.4	64
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### Special populations

**Use in renal impairment:** *Monotherapy:* PegIntron should be used with caution in patients with moderate to severe renal impairment. In patients with moderate renal dysfunction (creatinine clearance 30-50 ml/minute), the starting dose of PegIntron should be reduced by 25 %. Patients with severe renal dysfunction (creatinine clearance 15-29 ml/minute) should have the starting dose of PegIntron reduced by 50 %. Data are not available for the use of PegIntron in patients with creatinine clearance < 15 ml/minute (see section 5.2). Patients with severe renal impairment, including those on hemodialysis, should be closely monitored. If renal function decreases during treatment, PegIntron therapy should be discontinued.

*Combination therapy:* Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see ribavirin SPC). When administered in combination with ribavirin, subjects with impaired renal function should be more carefully monitored with respect to the development of anaemia.

**Use in hepatic impairment:** The safety and efficacy of PegIntron therapy has not been evaluated in patients with severe hepatic dysfunction, therefore PegIntron must not be used for these patients.

**Use in the elderly (≥ 65 years of age):** There are no apparent age-related effects on the pharmacokinetics of PegIntron. Data from elderly patients treated with a single dose of PegIntron suggest no alteration in PegIntron dose is necessary based on age (see section 5.2).

**Use in patients under the age of 18 years:** There is no experience in children (see section 5.2).

### 4.3 Contraindications

- Hypersensitivity to the active substance or to any interferon or to any of the excipients;
- A history of severe pre-existing cardiac disease, including unstable or uncontrolled cardiac disease in the previous six months (see section 4.4);
- Severe, debilitating medical conditions;
- Autoimmune hepatitis or a history of autoimmune disease;
- Severe hepatic dysfunction or decompensated cirrhosis of the liver;
- Pre-existing thyroid disease unless it can be controlled with conventional treatment;
- Epilepsy and/or compromised central nervous system (CNS) function;
- Initiation of PegIntron is contraindicated in HCV/HIV patients with cirrhosis and a Child-Pugh score ≥ 6.

*Combination therapy with ribavirin:* Also see ribavirin Summary of the Product Characteristics (SPC) if PegIntron is to be administered in combination with ribavirin in patients with chronic hepatitis C.

#### 4.4 Special warnings and precautions for use

**Psychiatric and Central Nervous System (CNS):** Severe CNS effects, particularly depression, suicidal ideation and attempted suicide have been observed in some patients during PegIntron therapy, and even after treatment discontinuation mainly during the 6-month follow-up period. Other CNS effects including aggressive behaviour (sometimes directed against others), confusion and alterations of mental status have been observed with alpha interferons. Patients should be closely monitored for any signs or symptoms of psychiatric disorders. If such symptoms appear, the potential seriousness of these undesirable effects must be borne in mind by the prescribing physician and the need for adequate therapeutic management should be considered. If psychiatric symptoms persist or worsen, or suicidal ideation is identified, it is recommended that treatment with PegIntron be discontinued, and the patient followed, with psychiatric intervention as appropriate.

*Patients with existence of, or history of severe psychiatric conditions:* If treatment with peginterferon alfa-2b is judged necessary in patients with existence or history of severe psychiatric conditions, this should only be initiated after having ensured appropriate individualised diagnostic and therapeutic management of the psychiatric condition.

There is no experience with PegIntron in combination with ribavirin in patients who have relapsed after interferon alpha + ribavirin therapy.

All patients in the chronic hepatitis C studies had a liver biopsy before inclusion, but in certain cases (i.e. patients with genotype 2 and 3), treatment may be possible without histological confirmation. Current treatment guidelines should be consulted as to whether a liver biopsy is needed prior to commencing treatment.

**Acute hypersensitivity:** Acute hypersensitivity reactions (e.g., urticaria, angioedema, bronchoconstriction, anaphylaxis) have been observed rarely during interferon alfa-2b therapy. If such a reaction develops during treatment with PegIntron, discontinue treatment and institute appropriate medical therapy immediately. Transient rashes do not necessitate interruption of treatment.

**Cardiovascular system:** As with interferon alfa-2b, patients with a history of congestive heart failure, myocardial infarction and/or previous or current arrhythmic disorders, receiving PegIntron therapy require close monitoring. It is recommended that patients who have pre-existing cardiac abnormalities have electrocardiograms taken prior to and during the course of treatment. Cardiac arrhythmias (primarily supraventricular) usually respond to conventional therapy but may require discontinuation of PegIntron therapy.

More significant obtundation and coma, including cases of encephalopathy, have been observed in some patients, usually elderly, treated at higher doses for oncology indications. While these effects are generally reversible, in a few patients full resolution took up to three weeks. Very rarely, seizures have occurred with high doses of interferon alpha.

**Liver function:** As with all interferons, discontinue treatment with PegIntron in patients who develop prolongation of coagulation markers which might indicate liver decompensation.

**Fever:** While fever may be associated with the flu-like syndrome reported commonly during interferon therapy, other causes of persistent fever must be ruled out.

**Hydration:** Adequate hydration must be maintained in patients undergoing PegIntron therapy since hypotension related to fluid depletion has been seen in some patients treated with alpha interferons. Fluid replacement may be necessary.

**Pulmonary changes:** Pulmonary infiltrates, pneumonitis, and pneumonia, occasionally resulting in fatality, have been observed rarely in interferon alpha treated patients. Any patient developing fever, cough, dyspnea or other respiratory symptoms must have a chest X-ray taken. If the chest X-ray shows pulmonary infiltrates or there is evidence of pulmonary function impairment, the patient is to be

monitored closely, and, if appropriate, discontinue interferon alpha. Prompt discontinuation of interferon alpha administration and treatment with corticosteroids appear to be associated with resolution of pulmonary adverse events.

**Autoimmune disease:** The development of auto-antibodies and autoimmune disorders has been reported during treatment with alpha interferons. Patients predisposed to the development of autoimmune disorders may be at increased risk. Patients with signs or symptoms compatible with autoimmune disorders should be evaluated carefully, and the benefit-risk of continued interferon therapy should be reassessed (see also section 4.4 Thyroid changes and 4.8).

**Ocular changes:** Ophthalmologic disorders, including retinal haemorrhages, cotton wool spots, and retinal artery or vein obstruction have been reported in rare instances after treatment with alpha interferons (see section 4.8). All patients should have a baseline eye examination. Any patient complaining of ocular symptoms, including loss of visual acuity or visual field must have a prompt and complete eye examination. Periodic visual examinations are recommended during PegIntron therapy, particularly in patients with disorders that may be associated with retinopathy, such as diabetes mellitus or hypertension. Discontinuation of PegIntron should be considered in patients who develop new or worsening ophthalmological disorders.

**Thyroid changes:** Infrequently, patients treated for chronic hepatitis C with interferon alpha have developed thyroid abnormalities, either hypothyroidism or hyperthyroidism. Determine thyroid stimulating hormone (TSH) levels if, during the course of therapy, a patient develops symptoms consistent with possible thyroid dysfunction. In the presence of thyroid dysfunction, PegIntron treatment may be continued if TSH levels can be maintained in the normal range by medication.

**Metabolic disturbances:** Hypertriglyceridemia and aggravation of hypertriglyceridemia, sometimes severe, have been observed. Monitoring of lipid levels is, therefore, recommended.

#### **HCV/HIV Co-infection**

Mitochondrial toxicity and lactic acidosis:

Patients co-infected with HIV and receiving Highly Active Anti-Retroviral Therapy (HAART) may be at increased risk of developing lactic acidosis. Caution should be used when adding PegIntron and ribavirin to HAART therapy (see ribavirin SPC).

Moreover, patients treated with PegIntron and ribavirin combination therapy and zidovudine are at increased risk of developing anaemia.

Hepatic decompensation in HCV/HIV co-infected patients with advanced cirrhosis:

Co-infected patients with advanced cirrhosis receiving HAART may be at increased risk of hepatic decompensation and death. Adding treatment with alfa interferons alone or in combination with ribavirin may increase the risk in this patient subset. Other baseline factors in co-infected patients that may be associated with a higher risk of hepatic decompensation include treatment with didanosine and elevated bilirubin serum concentration.

Co-infected patients receiving both antiretroviral (ARV) and anti-hepatitis treatment should be closely monitored, assessing their Child-Pugh score during treatment. Patients progressing to hepatic decompensation should have their anti-hepatitis treatment immediately discontinued and the ARV treatment reassessed.

Haematological abnormalities in HCV/HIV co-infected patients:

HCV/HIV co-infected patients receiving peginterferon alfa-2b/ribavirin treatment and HAART may be at increased risk to develop haematological abnormalities (as neutropenia, thrombocytopenia and anaemia) compared to HCV mono-infected patients. Although, the majority of them could be managed by dose reduction, close monitoring of haematological parameters should be undertaken in this population of patients (see section 4.2 and below “Laboratory tests” and section 4.8).

Patients with low CD4 counts:

In patients co-infected with HCV/HIV, limited efficacy and safety data (N = 25) are available in subjects with CD4 counts less than 200 cells/μl. Caution is therefore warranted in the treatment of

patients with low CD4 counts.

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron and ribavirin.

**Dental and periodontal disorders:** Dental and periodontal disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and mucous membranes of the mouth during long-term treatment with the combination of PegIntron and ribavirin. Patients should brush their teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If this reaction occurs, they should be advised to rinse out their mouth thoroughly afterwards.

**Organ transplant recipients:** The safety and efficacy of PegIntron alone or in combination with ribavirin for the treatment of hepatitis C in liver or other organ transplant recipients have not been studied. Preliminary data indicates that interferon alpha therapy may be associated with an increased rate of kidney graft rejection. Liver graft rejection has also been reported.

**Other:** Due to reports of interferon alpha exacerbating pre-existing psoriatic disease and sarcoidosis, use of PegIntron in patients with psoriasis or sarcoidosis is recommended only if the potential benefit justifies the potential risk.

**Laboratory tests:** Standard haematologic tests, blood chemistry and a test of thyroid function must be conducted in all patients prior to initiating therapy. Acceptable baseline values that may be considered as a guideline prior to initiation of PegIntron therapy are:

- |                    |                              |
|--------------------|------------------------------|
| • Platelets        | $\geq 100,000/\text{mm}^3$   |
| • Neutrophil count | $\geq 1,500/\text{mm}^3$     |
| • TSH level        | must be within normal limits |

Laboratory evaluations are to be conducted at weeks 2 and 4 of therapy, and periodically thereafter as clinically appropriate.

**Important information about some of the ingredients of PegIntron:**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

Patients with rare hereditary problems of fructose intolerance, glucose galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Results from a multiple-dose probe study assessing P450 substrates in chronic hepatitis C patients receiving once weekly PegIntron (1.5 µg/kg) for 4 weeks demonstrated an increase in activity of CYP2D6 and CYP2C8/9. No change in activity of CYP1A2, CYP3A4, or N-acetyltransferase was observed.

Caution should be used when administering peginterferon alfa-2b with medicines metabolised by CYP2D6 and CYP2C8/9, especially those with narrow therapeutic window, such as warfarin and phenytoin (CYP2C9) and flecainide (CYP2D6).

These findings may partly relate to improved metabolic capacity due to reduced hepatic inflammation in patients undergoing treatment with PegIntron. Caution is therefore advised when PegIntron treatment is initiated for chronic hepatitis in patients treated with medication with a narrow therapeutic window and sensitive to mild metabolic impairment of the liver.



No pharmacokinetic interactions were noted between PegIntron and ribavirin in a multiple-dose pharmacokinetic study.

**Methadone:** In patients with chronic hepatitis C that were on stable methadone maintenance therapy and naïve to peginterferon alfa-2b, addition of 1.5 microgram/kg/week of PegIntron subcutaneously for 4 weeks increased R-methadone AUC by approximately 15 % (95 % CI for AUC ratio estimate 103 – 128 %). The clinical significance of this finding is unknown; however, patients should be monitored for signs and symptoms of increased sedative effect, as well as respiratory depression. Especially in patients on a high dose of methadone, the risk for QTc prolongation should be considered.

#### *HCV/HIV Co-infection*

Nucleoside analogs: ribavirin has been shown *in vitro* to inhibit phosphorylation of zidovudine and stavudine. The clinical significance of these findings is unknown. However, these *in vitro* findings raise the possibility that concurrent use of ribavirin with either zidovudine or stavudine might lead to increased HIV plasma viraemia. Therefore, it is recommended that plasma HIV RNA levels be closely monitored in patients treated with ribavirin concurrently with either of these two agents. If HIV RNA levels increase, the use of ribavirin concomitantly with reverse transcriptase inhibitors must be reviewed (see ribavirin SPC).

Use of nucleoside analogs, alone or in combination with other nucleosides, has resulted in lactic acidosis. Pharmacologically, ribavirin increases phosphorylated metabolites of purine nucleosides *in vitro*. This activity could potentiate the risk of lactic acidosis induced by purine nucleoside analogs (e.g. didanosine or abacavir). Co-administration of ribavirin and didanosine is not recommended. Reports of mitochondrial toxicity, in particular lactic acidosis and pancreatitis, of which some fatal, have been reported (see ribavirin SPC).

## **4.6 Pregnancy and lactation**

PegIntron is recommended for use in fertile women only when they are using effective contraception during the treatment.

There are no adequate data from the use of interferon alfa-2b in pregnant women.. Studies in animals have shown reproductive toxicity (see section 5.3). Interferon alfa-2b has been shown to be abortifacient in primates. PegIntron is likely to also cause this effect.

The potential risk in humans is unknown. PegIntron is to be used during pregnancy only if the potential benefit justifies the potential risk to the foetus

**Lactation:** It is not known whether the components of this medicinal product are excreted in human milk. Because of the potential for adverse reactions in breast-fed infants, breast-feeding should be discontinued prior to initiation of treatment.

#### *Combination therapy with ribavirin:*

Ribavirin causes serious birth defects when administered during pregnancy. Ribavirin therapy is contraindicated in women who are pregnant. Extreme care must be taken to avoid pregnancy in female patients or in partners of male patients taking PegIntron in combination with ribavirin. Females of childbearing potential and their partners must each use an effective contraceptive during treatment and for 4 months after treatment has been concluded. Male patients and their female partners must each use an effective contraceptive during treatment and for 7 months after treatment has been concluded (see Ribavirin SPC).

## **4.7 Effects on ability to drive and use machines**

Patients who develop fatigue, somnolence or confusion during treatment with PegIntron are cautioned to avoid driving or operating machinery.

## 4.8 Undesirable effects

The safety of PegIntron is evaluated from data from two clinical trials: one with PegIntron monotherapy, one with PegIntron in combination with ribavirin. In both cases, patients were treated for one year.

**Table 4** describes the regimens and patient exposure for one year of treatment in patients with no previous exposure to interferon (interferon-naïve patients). Because of a significant overlap in the pattern of undesirable effects with PegIntron monotherapy, groups of patients have been brought together in **Table 5** to show the pattern of reported effects for all monotherapy groups.

<b>Table 4</b> Regimens and patient exposure		
<b>Treatment</b>	<b>Regimen</b>	<b>Number of patients treated for one year</b>
PegIntron + ribavirin	PegIntron (1.5 micrograms/kg/week) + ribavirin (> 10.6 mg/kg/day)	188
Interferon alfa-2b + ribavirin	Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day)	505
PegIntron monotherapy	PegIntron (0.5 microgram/kg/week)	315
	PegIntron (1.0 microgram/kg/week)	297
	PegIntron (1.5 micrograms/kg/week)	304

<b>Table 5</b> Undesirable effects very commonly reported in clinical trials > 10 % of patients in PegIntron + ribavirin group Very common (> 1/10)			
	<b>PegIntron + ribavirin</b>	<b>Interferon alfa- 2b + ribavirin</b>	<b>PegIntron monotherapy</b>
<b>Infections and infestations</b>			
Infection viral	10 %	5 %	4-5 %
<b>Metabolism and nutrition disorders</b>			
Weight decrease	30 %	19 %	8-18 %
<b>Psychiatric disorders</b>			
Depression	34 %	32 %	26 %
Irritability	32 %	34 %	19 %
Insomnia	37 %	41 %	16-19 %
Anxiety	14 %	14 %	8 %
Concentration impaired	18 %	21 %	9-10 %
Emotional lability	11 %	10 %	5 %
<b>Nervous system disorders</b>			
Headache	58 %	57 %	57-63 %
Mouth dry	10 %	8 %	4-8 %
<b>Respiratory, thoracic and mediastinal disorders</b>			
Pharyngitis	10 %	7 %	3 %
Coughing	14 %	11 %	4 %
Dyspnea	26 %	22 %	5 %
<b>Gastrointestinal disorders</b>			
Nausea	43 %	31 %	20-23 %
Anorexia	35 %	26 %	10-25 %
Diarrhoea	20 %	13 %	14-17 %
Abdominal pain	12 %	9 %	11 %
Vomiting	16 %	10 %	4-7 %
<b>Skin and subcutaneous tissue disorders</b>			
Alopecia	45 %	32 %	20-34 %
Pruritus	27 %	27 %	7-9 %
Skin dry	23 %	21 %	6-9 %
Rash	21 %	21 %	5-7 %
<b>Musculoskeletal and connective tissue disorders</b>			
Myalgia	49 %	49 %	46-60 %
Arthralgia	31 %	26 %	23-28 %
Musculoskeletal pain	15 %	11 %	11-13 %
<b>General disorders and administration site conditions</b>			
Injection site inflammation	20 %	17 %	39-44 %
Injection site reaction	54 %	36 %	7-9 %
Fatigue	56 %	59 %	43 %
Rigors	42 %	40 %	33-43 %
Fever	39 %	32 %	29-43 %
Flu-like symptoms	21 %	23 %	18-25 %
Asthenia	28 %	17 %	12-14 %
Dizziness	17 %	16 %	7-12 %

<b>Table 6</b> Undesirable effects commonly reported in clinical trials in patients treated with PegIntron + ribavirin or PegIntron monotherapy Common (> 1/100, < 1/10)		
<b>Body system</b>	<b>5-10%</b>	<b>1-&lt;5%</b>
Infections and infestations		otitis media, fungal infection, bacterial infection, herpes simplex
Blood and lymphatic system disorders	Anaemia, leukopaenia	Thrombocytopenia, lymphadenopathy
Endocrine disorders	hypothyroidism	hyperthyroidism
Metabolism and nutrition disorders		hyperuricemia, hypocalcemia, thirst
Psychiatric disorders	agitation, nervousness	aggressive behaviour, somnolence, behavior disorder, apathy, appetite increased, sleep disorder, dreaming abnormal, decreased libido
Nervous system disorders	paresthesia, increased sweating	hypoesthesia, hyperaesthesia, hypertonia, confusion, tremor, vertigo, migraine, ataxia, neuralgia
Eye disorders		blurred vision, conjunctivitis, lacrimal gland disorder, eye pain
Cardiac disorders		tachycardia, palpitation
Vascular disorders		hypotension, hypertension, syncope, flushing
Ear and labyrinth disorders		tinnitus, hearing impairment/loss
Respiratory, thoracic and mediastinal disorders		nonproductive cough, rhinitis, sinusitis, bronchitis, respiratory disorder, nasal congestion, rhinorrhea, dysphonia, epistaxis
Gastrointestinal disorders	dyspepsia	constipation, taste perversion, loose stools, stomatitis, ulcerative stomatitis, gingival bleeding, glossitis, flatulence, hemorrhoids, gastroesophageal reflux, gingivitis, dehydration
Hepatobiliary disorders		hepatomegaly, hyperbilirubinemia
Skin and subcutaneous tissue disorders		erythematous rash, eczema, photosensitivity reaction, maculopapular rash, abnormal hair texture, acne, dermatitis, furunculosis, nail disorder, psoriasis, urticaria, erythema, face or peripheral oedema
Musculoskeletal and connective tissue disorders		arthritis
Renal and urinary disorders		micturition frequency, urine abnormal
Reproductive system and breast disorders	menstrual disorder, menorrhagia	ovarian disorder, vaginal disorder, sexual dysfunction (not specified), impotence, breast pain, amenorrhoea, prostatitis
General disorders and administration site conditions	RUQ pain, malaise, chest pain	injection site pain

Most cases of neutropaenia and thrombocytopaenia were mild (WHO grades 1 or 2). There were some cases of more severe neutropenia in patients treated with the recommended doses of PegIntron in combination with ribavirin (WHO grade 3: 39 of 186 [21 %]; and WHO grade 4: 13 of 186 [7 %]).

In a clinical trial, approximately 1.2 % of patients treated with PegIntron or interferon alfa-2b in combination with ribavirin reported life-threatening psychiatric events during treatment. These events included suicidal ideation and attempted suicide (see section 4.4). Following marketing, psychosis and hallucination have been reported rarely.

Rarely ( $> 1/10,000$ ,  $< 1/1,000$ ) or very rarely ( $< 1/10,000$ ) reported events with interferon alfa-2b, including PegIntron, include:

**Blood and lymphatic system disorders:**

Very rarely interferon alfa-2b or PegIntron used alone or in combination with ribavirin may be associated with aplastic anaemia. Pure red cell aplasia has been reported.

**Immune system disorders:**

very rarely: sarcoidosis or exacerbation of sarcoidosis

**Endocrine disorders:**

rarely: diabetes

**Nervous system disorders:**

rarely: seizure, peripheral neuropathy

very rarely: cerebrovascular ischaemia, cerebrovascular haemorrhage, encephalopathy

**Cardiac disorders:**

rarely: arrhythmia

very rarely: cardiac ischaemia, myocardial infarction

**Respiratory, thoracic and mediastinal disorders:**

very rarely: interstitial lung disease

**Gastrointestinal disorders:**

rarely: pancreatitis

very rarely: ulcerative and ischaemic colitis

**Skin and subcutaneous tissue disorders:**

very rarely: erythema multiforme, Stevens Johnson syndrome, toxic epidermal necrolysis, injection site necrosis

**Musculoskeletal and connective tissue disorders:**

rarely: rhabdomyolysis, myositis

**Renal and urinary disorders:**

rarely: renal insufficiency and renal failure

Cardiovascular (CVS) adverse events, particularly arrhythmia, appeared to be correlated mostly with pre-existing CVS disease and prior therapy with cardiotoxic agents (see section 4.4). Cardiomyopathy, that may be reversible upon discontinuation of interferon alpha, has been reported rarely in patients without prior evidence of cardiac disease.

Ophthalmological disorders that have been reported rarely with alpha interferons include retinopathies (including macular oedema), retinal haemorrhages, retinal artery or vein obstruction, cotton wool spots, loss of visual acuity or visual field, optic neuritis, and papilloedema (see section 4.4).

A wide variety of autoimmune and immune-mediated disorders have been reported with alpha interferons including thyroid disorders, systemic lupus erythematosus, rheumatoid arthritis (new or aggravated), idiopathic and thrombotic thrombocytopenic purpura, vasculitis, neuropathies including mononeuropathies (see also section 4.4, **Autoimmune disorders**).

**HCV/HIV co-infected patients**

For HCV/HIV co-infected patients receiving PegIntron in combination with ribavirin, other undesirable effects (that were not reported in mono-infected patients) which have been reported in the larger studies with a frequency  $>5\%$  were: oral candidiasis (14%), lipodystrophy acquired (13%), CD4 lymphocytes decreased (8%), appetite decreased (8%), gamma-glutamyltransferase increased (9%), back pain (5%), blood amylase increased (6%), blood lactic acid increased (5%), cytolytic hepatitis (6%), lipase increased (6%) and pain in limb (6%).

*Mitochondrial toxicity:*

Mitochondrial toxicity and lactic acidosis have been reported in HIV-positive patients receiving NRTI regimen and associated ribavirin for co-HCV infection (see section 4.4).

*Laboratory values for HCV/HIV co-infected patients:*

Although haematological toxicities of neutropenia, thrombocytopenia and anaemia occurred more frequently in HCV/HIV co-infected patients, the majority could be managed by dose modification and rarely required premature discontinuation of treatment (see section 4.4). Haematological abnormalities were more frequently reported in patients receiving PegIntron in combination with ribavirin when compared to patients receiving interferon alfa-2b in combination with ribavirin. In Study 1 (see section 5.1), decrease in absolute neutrophil count levels below 500 cells/mm<sup>3</sup> was observed in 4 % (8/194) of patients and decrease in platelets below 50,000/mm<sup>3</sup> was observed in 4 % (8/194) of patients receiving PegIntron in combination with ribavirin. Anaemia (hemoglobin < 9.4g/dl) was reported in 12% (23/194) of patients treated with PegIntron in combination with ribavirin.

*CD4 lymphocytes decrease:*

Treatment with PegIntron in combination with ribavirin was associated with decreases in absolute CD4+ cell counts within the first 4 weeks without a reduction in CD4+ cell percentage. The decrease in CD4+ cell counts was reversible upon dose reduction or cessation of therapy. The use of PegIntron in combination with ribavirin had no observable negative impact on the control of HIV viraemia during therapy or follow-up. Limited safety data (N= 25) are available in co-infected patients with CD4+ cell counts < 200/μl (see section 4.4).

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron in combination with ribavirin.

## **4.9 Overdose**

In clinical trials, cases of accidental overdose, at never more than twice the prescribed dose, were reported. There were no serious reactions. Undesirable effects resolved during continued administration of PegIntron.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Immunostimulants, Cytokines and immunomodulators, Interferons, Peginterferon alfa-2b, ATC code: L03A B10.

Recombinant interferon alfa-2b is covalently conjugated with monomethoxy polyethylene glycol at an average degree of substitution of 1 mole of polymer/mole of protein. The average molecular mass is approximately 31,300 daltons of which the protein moiety constitutes approximately 19,300.

#### **Interferon alfa-2b**

*In vitro* and *in vivo* studies suggest that the biological activity of PegIntron is derived from its interferon alfa-2b moiety.

Interferons exert their cellular activities by binding to specific membrane receptors on the cell surface. Studies with other interferons have demonstrated species specificity. However, certain monkey species, e.g., Rhesus monkeys, are susceptible to pharmacodynamic stimulation upon exposure to human type 1 interferons.

Once bound to the cell membrane, interferon initiates a complex sequence of intracellular events that include the induction of certain enzymes. It is thought that this process, at least in part, is responsible for the various cellular responses to interferon, including inhibition of virus replication in virus-infected cells, suppression of cell proliferation and such immunomodulating activities as enhancement

of the phagocytic activity of macrophages and augmentation of the specific cytotoxicity of lymphocytes for target cells. Any or all of these activities may contribute to interferon's therapeutic effects.

Recombinant interferon alfa-2b also inhibits viral replication *in vitro* and *in vivo*. Although the exact antiviral mode of action of recombinant interferon alfa-2b is unknown, it appears to alter the host cell metabolism. This action inhibits viral replication or if replication occurs, the progeny virions are unable to leave the cell.

### **PegIntron**

PegIntron pharmacodynamics were assessed in a rising single-dose trial in healthy subjects by examining changes in oral temperature, concentrations of effector proteins such as serum neopterin and 2'5'-oligoadenylate synthetase (2'5'-OAS), as well as white cell and neutrophil counts. Subjects treated with PegIntron showed mild dose-related elevations in body temperature. Following single doses of PegIntron between 0.25 and 2.0 micrograms/kg/week, serum neopterin concentration was increased in a dose-related manner. Neutrophil and white cell count reductions at the end of week 4 correlated with the dose of PegIntron.

### **PegIntron clinical trials**

Two pivotal trials have been conducted, one (C/I97-010) with PegIntron monotherapy; the other (C/I98-580) with PegIntron in combination with ribavirin. Eligible patients for these trials had chronic hepatitis C confirmed by a positive HCV-RNA polymerase chain reaction (PCR) assay (> 30 IU/ml), a liver biopsy consistent with a histological diagnosis of chronic hepatitis with no other cause for the chronic hepatitis, and abnormal serum ALT.

In the PegIntron monotherapy trial, a total of 916 naïve chronic hepatitis C patients were treated with PegIntron (0.5, 1.0 or 1.5 micrograms/kg/week) for one year with a follow-up period of six months. In addition, 303 patients received interferon alfa-2b (3 million International Units [MIU] three times a week) as a comparator. This study showed that PegIntron was superior to interferon alfa-2b (**Table 8**).

In the PegIntron combination trial, 1,530 naïve patients were treated for one year with one of the following combination regimens:

- PegIntron (1.5 micrograms/kg/week) + ribavirin (800 mg/day), (n = 511).
- PegIntron (1.5 micrograms/kg/week for one month followed by 0.5 microgram/kg/week for 11 months) + ribavirin (1,000/1,200 mg/day), (n = 514).
- Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day) (n = 505).

In this trial, the combination of PegIntron (1.5 micrograms/kg/week) and ribavirin was significantly more effective than the combination of interferon alfa-2b and ribavirin (**Table 7**), particularly in patients infected with Genotype 1 (**Table 8**). Sustained response was assessed by the response rate six months after the cessation of treatment.

HCV genotype and baseline virus load are prognostic factors which are known to affect response rates. However, response rates in this trial were shown to be dependent also on the dose of ribavirin administered in combination with PegIntron or interferon alfa-2b. In those patients that received > 10.6 mg/kg ribavirin (800 mg dose in typical 75 kg patient), regardless of genotype or viral load, response rates were significantly higher than in those patients that received ≤ 10.6 mg/kg ribavirin (**Table 8**), while response rates in patients that received > 13.2 mg/kg ribavirin were even higher.

<b>Table 7</b> Sustained virological response (% patients HCV negative)							
	<b>PegIntron monotherapy</b>				<b>PegIntron + ribavirin</b>		
<b>Treatment regimen</b>	<b>P 1.5</b>	<b>P 1.0</b>	<b>P 0.5</b>	<b>I</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>Number of patients</b>	304	297	315	303	511	514	505
<b>Response at end of treatment</b>	49 %	41 %	33 %	24 %	65 %	56 %	54 %

<b>Sustained response</b>	<b>23 %*</b>	<b>25 %</b>	<b>18 %</b>	<b>12 %</b>	<b>54 %**</b>	<b>47 %</b>	<b>47 %</b>
---------------------------	--------------	-------------	-------------	-------------	---------------	-------------	-------------

P 1.5 PegIntron 1.5 micrograms/kg  
 P 1.0 PegIntron 1.0 microgram/kg  
 P 0.5 PegIntron 0.5 microgram/kg  
 I Interferon alfa-2b 3 MIU  
 P 1.5/R PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)  
 P 0.5/R PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)  
 I/R Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)  
 \* p < 0.001 P 1.5 vs. I  
 \*\* p = 0.0143 P 1.5/R vs. I/R

<b>Table 8</b> Sustained response rates with PegIntron + ribavirin (by ribavirin dose, genotype and viral load)				
<b>HCV Genotype</b>	<b>Ribavirin dose (mg/kg)</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>All Genotypes</b>	<b>All</b>	<b>54 %</b>	<b>47 %</b>	<b>47 %</b>
	≤ 10.6	50 %	41 %	27 %
	> 10.6	61 %	48 %	47 %
<b>Genotype 1</b>	<b>All</b>	<b>42 %</b>	<b>34 %</b>	<b>33 %</b>
	≤ 10.6	38 %	25 %	20 %
	> 10.6	48 %	34 %	34 %
Genotype 1 ≤ 600,000 IU/ml	All	73 %	51 %	45 %
	≤ 10.6	74 %	25 %	33 %
	> 10.6	71 %	52 %	45 %
Genotype 1 > 600,000 IU/ml	All	30 %	27 %	29 %
	≤ 10.6	27 %	25 %	17 %
	> 10.6	37 %	27 %	29 %
<b>Genotype 2/3</b>	<b>All</b>	<b>82 %</b>	<b>80 %</b>	<b>79 %</b>
	≤ 10.6	79 %	73 %	50 %
	> 10.6	88 %	80 %	80 %

P 1.5/R PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)  
 P 0.5/R PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)  
 I/R Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)

In the PegIntron monotherapy study, the Quality of Life was generally less affected by 0.5 microgram/kg of PegIntron than by either 1.0 microgram/kg of PegIntron once weekly or 3 MIU of interferon alfa-2b three times a week.

In a separate trial, 224 patients with genotype 2 or 3 received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with ribavirin 800 mg –1,400 mg p.o. for 6 months (based on body weight, only three patients weighing > 105 kg, received the 1,400 mg dose, which has not yet been validated) (**Table 9**). Twenty-four % had bridging fibrosis or cirrhosis (Knodell 3/4).

**Table 9. Virologic Response at End of Treatment, Sustained Virologic Response and Relapse by HCV Genotype and Viral Load\***

	PegIntron 1.5 µg/kg once weekly plus Ribavirin 800-1,400 mg/day		
	End of Treatment Response	Sustained Virologic Response	Relapse
All Subjects	94 % (211/224)	81 % (182/224)	12 % (27/224)
HCV 2 ≤ 600,000 IU/ml > 600,000	100 % (42/42)	93 % (39/42)	7 % (3/42)
	100 % (20/20)	95 % (19/20)	5 % (1/20)
	100 % (22/22)	91 % (20/22)	9 % (2/22)



IU/ml			
HCV 3	93 % (169/182)	79 % (143/182)	14 % (24/166)
≤ 600,000 IU/ml	93 % (92/99)	86 % (85/99)	8 % (7/91)
> 600,000 IU/ml	93 % (77/83)	70 % (58/83)	23 % (17/75)

\* Any subject with an undetectable HCV-RNA level at the Follow-Up Week 12 visit and missing data at the Follow-Up Week 24 visit was considered a sustained responder. Any subject with missing data in and after the Follow-Up Week 12 window was considered to be a non-responder at Week 24 of follow-up.

The 6 month treatment duration in this trial was better tolerated than one year of treatment in the pivotal combination trial; for discontinuation 5 % vs. 14 %, for dose modification 18 % vs. 49 %.

In a non-comparative trial, 235 patients with genotype 1 and low viral load (< 600,000 IU/ml) received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with weight adjusted ribavirin. The overall sustained response rate after a 24-week treatment duration was 50 %. Forty-one percent of subjects (97/235) had nondetectable plasma HCV-RNA levels at Week 4 and Week 24 of therapy. In this subgroup, there was a 92 % (89/97) sustained virological response rate. The high sustained response rate in this subgroup of patients was identified in an interim analysis (n=49) and prospectively confirmed (n=48).

Limited historical data indicate that treatment for 48 weeks might be associated with a higher sustained response rate (11/11) and with a lower risk of relapse (0/11 as compared to 7/96 following 24 weeks of treatment).

### Predictability of sustained virological response

Virological response by week 12, defined as a 2-log viral load decrease or undetectable levels of HCV-RNA has been shown to be predictive for sustained response (**Table 10**).

<b>Table 10</b> Predictability of sustained response by viral response at week 12 and genotype*				
Treatment	Genotype	Viral response at week 12	Sustained response	Negative predictive value
PegIntron 1.5 + ribavirin (> 10.6 mg/kg) 48-week treatment	1	Yes 75 % (82/110)	71 % (58/82)	----
		No 25 % (28/110)	0 % (0/28)	100 %
PegIntron 1.5 + ribavirin 800-1,400 mg 24-week treatment	2 and 3	Yes 99 % (213/215)	83 % (177/213)	----
		No 1 % (2/215)	50 % (1/2)	50 %

\* reflects patients with 12 week data available

The negative predictive value for sustained response in patients treated with PegIntron in monotherapy was 98 %.

### HCV/HIV Co-infected patients

Two trials have been conducted in patients co-infected with HIV and HCV. The response to treatment in both of these trials is presented in **Table 11**. Study 1 (RIBAVIC; P01017) was a randomized, multicentre study which enrolled 412 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (1.5 µg/kg/week) plus ribavirin (800 mg/day) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800 mg/day) for 48 weeks with a follow-up period of 6 months. Study 2 (P02080) was a randomized, single centre

study that enrolled 95 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (100 or 150 µg /week based on weight) plus ribavirin (800-1,200 mg/day based on weight) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800-1,200 mg/day based on weight). The duration of therapy was 48 weeks with a follow-up period of 6 months except for patients infected with genotypes 2 or 3 and viral load < 800,000 IU/ml (Amplicor) who were treated for 24 weeks with a 6-month follow-up period.

<b>Table 11</b> Sustained virological response based on genotype after PegIntron in combination with Ribavirin in HCV/HIV Co-infected patients						
	<b>Study 1<sup>1</sup></b>			<b>Study 2<sup>2</sup></b>		
	PegIntron (1.5 µg/kg/week) + ribavirin (800 mg)	Interferon alfa-2b (3 MIU TIW) + ribavirin (800 mg)	p value <sup>a</sup>	PegIntron (100 or 150 <sup>c</sup> µg/week) + ribavirin (800-1,200 mg) <sup>d</sup>	Interferon alfa-2b (3 MIU TIW) + ribavirin (800-1,200 mg) <sup>d</sup>	p value <sup>b</sup>
All	27 % (56/205)	20 % (41/205)	0.047	44 % (23/52)	21 % (9/43)	0.017
Genotype 1, 4	17 % (21/125)	6 % (8/129)	0.006	38 % (12/32)	7 % (2/27)	0.007
Genotype 2, 3	44 % (35/80)	43 % (33/76)	0.88	53 % (10/19)	47 % (7/15)	0.730

MIU = million international units; TIW = three times a week.

a: p value based on Cochran-Mantel Haenszel Chi square test.

b: p value based on chi-square test.

c: subjects < 75 kg received 100 µg/week PegIntron and subjects ≥ 75 kg received 150 µg/week PegIntron.

d: ribavirin dosing was 800 mg for patients < 60 kg, 1,000 mg for patients 60-75 kg, and 1,200 mg for patients > 75 kg.

<sup>1</sup>Carrato F, Bani-Sadir F, Pol S et al. JAMA 2004; 292(23): 2839-2848.

<sup>2</sup>Laguno M, Murillas J, Blanco J et al. AIDS 2004; 18(13): F27-F36.

### *Histological response*

Liver biopsies were obtained before and after treatment in Study 1 and were available for 210 of the 412 subjects (51 %). Both the Metavir score and Ishak grade decreased among subjects treated with PegIntron in combination with ribavirin. This decline was significant among responders (-0.3 for Metavir and -1.2 for Ishak) and stable (-0.1 for Metavir and -0.2 for Ishak) among non-responders. In terms of activity, about one-third of sustained responders showed improvement and none showed worsening. There was no improvement in terms of fibrosis observed in this study. Steatosis was significantly improved in patients infected with HCV Genotype 3.

## **5.2 Pharmacokinetic properties**

PegIntron is a well characterized polyethylene glycol-modified (“pegylated”) derivative of interferon alfa-2b and is predominantly composed of monopegylated species. The plasma half-life of PegIntron is prolonged compared with non-pegylated interferon alfa-2b. PegIntron has a potential to depegylate to free interferon alfa-2b. The biologic activity of the pegylated isomers is qualitatively similar, but weaker than free interferon alfa-2b.

Following subcutaneous administration, maximal serum concentrations occur between 15-44 hours post-dose, and are sustained for up to 48-72 hours post-dose.

PegIntron C<sub>max</sub> and AUC measurements increase in a dose-related manner. Mean apparent volume of distribution is 0.99 l/kg.

Upon multiple dosing, there is an accumulation of immunoreactive interferons. There is, however, only a modest increase in biologic activity as measured by a bioassay.

Mean (SD) PegIntron elimination half-life is approximately 40 hours (13.3 hours), with apparent clearance of 22.0 ml/hr·kg. The mechanisms involved in clearance of interferons in man have not yet been fully elucidated. However, renal elimination may account for a minority (approximately 30 %) of PegIntron apparent clearance.

**Renal function:** Renal clearance appears to account for 30 % of total clearance of PegIntron. In a single dose study (1.0 microgram/kg) in patients with impaired renal function,  $C_{max}$ , AUC, and half-life increased in relation to the degree of renal impairment.

Following multiple dosing of PegIntron (1.0 microgram/kg subcutaneously administered every week for four weeks) the clearance of PegIntron is reduced by a mean of 17 % in patients with moderate renal impairment (creatinine clearance 30-49 ml/minute) and by a mean of 44 % in patients with severe renal impairment (creatinine clearance 15-29 ml/minute) compared to subjects with normal renal function. Based on single dose data, clearance was similar in patients with severe renal impairment not on dialysis and in patients who were receiving hemodialysis. The dose of PegIntron for monotherapy should be reduced in patients with moderate or severe renal impairment (see sections 4.2 and 4.4). Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see section 4.3).

Because of marked inter-subject variability in interferon pharmacokinetics, it is recommended that patients with severe renal impairment be closely monitored during treatment with PegIntron (see section 4.2)

**Hepatic function:** The pharmacokinetics of PegIntron have not been evaluated in patients with severe hepatic dysfunction.

**Elderly patients ≥ 65 years of age:** The pharmacokinetics of PegIntron following a single subcutaneous dose of 1.0 microgram/kg were not affected by age. The data suggest that no alteration in PegIntron dosage is necessary based on advancing age.

**Patients under the age of 18 years:** Specific pharmacokinetic evaluations have not been performed on these patients. PegIntron is indicated for the treatment of chronic hepatitis C only in patients 18 years of age or older.

**Interferon neutralising factors:** Interferon neutralising factor assays were performed on serum samples of patients who received PegIntron in the clinical trial. Interferon neutralising factors are antibodies which neutralise the antiviral activity of interferon. The clinical incidence of neutralising factors in patients who received PegIntron 0.5 micrograms/kg is 1.1 %.

### 5.3 Preclinical safety data

**PegIntron:** Adverse events not observed in clinical trials were not seen in toxicity studies in monkeys. These studies were limited to four weeks due to the appearance of anti-interferon antibodies in most monkeys.

Reproduction studies of PegIntron have not been performed. Interferon alfa-2b has been shown to be an abortifacient in primates. PegIntron is likely to also cause this effect. Effects on fertility have not been determined. It is not known whether the components of this medicinal product are excreted into experimental animal or human milk (see section 4.6 for relevant human data on pregnancy and lactation). PegIntron showed no genotoxic potential.

The relative non-toxicity of monomethoxy-polyethylene glycol (mPEG), which is liberated from PegIntron by metabolism *in vivo* has been demonstrated in preclinical acute and subchronic toxicity studies in rodents and monkeys, standard embryo-foetal development studies and in *in vitro* mutagenicity assays.

**PegIntron plus ribavirin:** When used in combination with ribavirin, PegIntron did not cause any effects not previously seen with either active substance alone. The major treatment-related change was a reversible, mild to moderate anaemia, the severity of which was greater than that produced by either active substance alone.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Powder for solution for injection:

Disodium phosphate, anhydrous,  
Sodium dihydrogen phosphate dihydrate,  
Sucrose,  
Polysorbate 80.

Solvent for parenteral use:

Water for injections.

### **6.2 Incompatibilities**

This medicinal product should only be reconstituted with the solvent provided (see section 6.6). In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

3 years

After reconstitution:

- Chemical and physical in-use stability has been demonstrated for 24 hours at 2°C - 8°C.
- From a microbiological point of view, the product is to be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C - 8°C.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C - 8°C).

### **6.5 Nature and contents of container**

The powder is contained in a 2 ml vial, Type I flint glass, with a butyl rubber stopper in an aluminium flip-off seal with a polypropylene bonnet. The solvent is presented in a 2 ml ampoule, Type I flint glass. PegIntron 120 micrograms is supplied as:

- 1 vial of powder for solution for injection and 1 ampoule of solvent for parenteral use;
- 1 vial of powder for solution for injection, 1 ampoule of solvent for parenteral use, 1 injection syringe, 2 injection needles and 1 cleansing swab;
- 4 vials of powder for solution for injection and 4 ampoules of solvent for parenteral use;
- 4 vials of powder for solution for injection, 4 ampoules of solvent for parenteral use, 4 injection syringes, 8 injection needles and 4 cleansing swabs;
- 6 vials of powder for solution for injection and 6 ampoules of solvent for parenteral use.
- 12 vials of powder for solution for injection, 12 ampoules of solvent for parenteral use, 12 injection syringes, 24 injection needles and 12 cleansing swabs.

Not all pack sizes may be marketed

### **6.6 Special precautions for disposal**

PegIntron is supplied as a powder of peginterferon alfa-2b at a strength of 120 micrograms for single use. Each vial must be reconstituted with 0.7 ml of water for injections for administration of up to 0.5 ml of solution. A small volume is lost during preparation of PegIntron for injection when the dose is measured and injected. Therefore, each vial contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection. The reconstituted solution has a concentration of 120 micrograms/0.5 ml.

Using a sterilised injection syringe and injection needle, inject 0.7 ml of water for injections into the vial of PegIntron. Agitate gently to complete dissolution of powder. The appropriate dose can then be withdrawn with a sterilised injection syringe and injected. A complete set of instructions is provided in the Annex to the Package Leaflet.

As for all parenteral medicinal products, inspect visually the reconstituted solution prior to administration. The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. Discard any unused material.

## **7. MARKETING AUTHORISATION HOLDER**

SP Europe  
73, rue de Stalle  
B-1180 Bruxelles  
Belgium

## **8. MARKETING AUTHORISATION NUMBERS**

EU/1/00/131/016  
EU/1/00/131/017  
EU/1/00/131/018  
EU/1/00/131/019  
EU/1/00/131/020  
EU/1/00/131/029

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 25 May 2000  
Date of last renewal: 25 May 2005

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.emea.europa.eu/>

## **1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 150 micrograms powder and solvent for solution for injection

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each vial of PegIntron, powder for solution for injection contains 150 micrograms of peginterferon alfa-2b as measured on a protein basis.

Each vial provides 150 micrograms/0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

The active substance is a covalent conjugate of recombinant interferon alfa-2b\* with monomethoxy polyethylene glycol. The potency of this product should not be compared to that of another pegylated or non-pegylated protein of the same therapeutic class. For more information, see section 5.1.

\*produced by rDNA technology in *E.coli* cells harbouring a genetically engineered plasmid hybrid encompassing an interferon alfa-2b gene from human leukocytes

For a full list of excipients, see section 6.1. PegIntron contains 40 mg of sucrose per 0.5 ml.

## **3. PHARMACEUTICAL FORM**

Powder and solvent for solution for injection

White powder.

Clear and colourless solvent.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

PegIntron is indicated for the treatment of adult patients with chronic hepatitis C who have elevated transaminases without liver decompensation and who are positive for serum HCV-RNA or anti-HCV, including naïve patients with clinically stable HIV co-infection (see section 4.4).

The best way to use PegIntron in this indication is in combination with ribavirin.

This combination is indicated in naïve patients including patients with clinically stable HIV co-infection. This combination of PegIntron with ribavirin is also indicated in patients who have previously responded (with normalisation of ALT at the end of treatment) to interferon alpha monotherapy but who have subsequently relapsed.

Interferon monotherapy, including PegIntron, is indicated mainly in case of intolerance or contraindication to ribavirin.

Please refer also to the ribavirin Summary of Product Characteristics (SPC) when PegIntron is to be used in combination with ribavirin.

### **4.2 Posology and method of administration**

Treatment should be initiated and monitored only by a physician experienced in the management of patients with hepatitis C.

#### **Dose to be administered**

PegIntron should be administered as a once weekly subcutaneous injection. The dose administered depends on whether it is used in combination with ribavirin or as monotherapy.

## Combination therapy

PegIntron 1.5 micrograms/kg/week in combination with ribavirin capsules.

The intended dose of 1.5 µg/kg of PegIntron to be used in combination with ribavirin may be delivered in weight categories with the pen/vial strengths according to **Table 1**. Ribavirin capsules are to be administered orally each day in two divided doses with food (morning and evening).

**Table 1 - Dosing for Combination Therapy**

Body Weight (kg)	PegIntron		Ribavirin Capsules	
	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Total Daily Dose (mg)	Number of Capsules (200 mg)
< 40	50	0.5	800	4 <sup>a</sup>
40-50	80	0.4	800	4 <sup>a</sup>
51-64	80	0.5	800	4 <sup>a</sup>
65-75	100	0.5	1,000	5 <sup>b</sup>
76-85	120	0.5	1,000	5 <sup>b</sup>
> 85	150	0.5	1,200	6 <sup>c</sup>

a: 2 morning, 2 evening

b: 2 morning, 3 evening

c: 3 morning, 3 evening

## Duration of treatment

**Predictability of sustained virological response:** Patients infected with virus genotype 1 who fail to achieve virological response at Week 12 are highly unlikely to become sustained virological responders (see also section 5.1).

- **Genotype 1:** For patients who exhibit virological response at week 12, treatment should be continued for another nine month period (i.e., a total of 48 weeks).  
In the subset of patients with genotype 1 infection and low viral load (< 600,000 IU/ml) who become HCV-RNA negative at treatment week 4 and remain HCV-RNA negative at week 24, the treatment could either be stopped after this 24 week treatment course or pursued for an additional 24 weeks (i.e. overall 48 weeks treatment duration). However, an overall 24 weeks treatment duration may be associated with a higher risk of relapse than a 48 weeks treatment duration (see section 5.1).
- **Genotypes 2 or 3:** It is recommended that all patients be treated for 24 weeks, except for HCV/HIV co-infected patients who should receive 48 weeks of treatment.
- **Genotype 4:** In general, patients infected with genotype 4 are considered harder to treat and limited study data (n=66) indicate they are compatible with a duration of treatment as for genotype 1.

## HCV/HIV Co-infection

The recommended duration of dosing for HCV/HIV co-infected patients is 48 weeks, regardless of genotype.

### *Predictability of response and non-response in HCV/HIV Co-infection*

Early virological response by Week 12, defined as a 2 log viral load decrease or undetectable levels of HCV-RNA, has been shown to be predictive for sustained response. The negative predictive value for sustained response in HCV/HIV co-infected patients treated with PegIntron in combination with ribavirin was 99 % (67/68; Study 1) (see section 5.1). A positive predictive value of 50 % (52/104; Study 1) was observed for HCV/HIV co-infected patients receiving combination therapy.

### PegIntron monotherapy

As monotherapy the PegIntron regimen is 0.5 or 1.0 microgram/kg/week. The lowest vial or pen strength available is 50 µg/0.5 ml; therefore for patients prescribed 0.5 µg/kg/week, doses must be adjusted by volume as shown in **Table 2**. For the 1.0 µg/kg dose, similar volume adjustments can be made or alternate vial strengths can be used as shown in **Table 2**. PegIntron monotherapy was not studied in HCV/HIV co-infected patients.

**Table 2- Monotherapy Dosing**

	0.5 µg/kg		1.0 µg/kg	
Body Weight (kg)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)
30-35	50*	0.15	50	0.3
36-45	50*	0.2	50	0.4
46-56	50*	0.25	50	0.5
57-72	50	0.3	80	0.4
73-88	50	0.4	80	0.5
89-106	50	0.5	100	0.5
> 106**	80	0.4	120	0.5
* Must use vial. Minimum delivery for pen is 0.3 ml.				
** For patients > 120 kg, use 80 µg/0.5 ml vial				

### Duration of treatment

For patients who exhibit virological response at Week 12, treatment should be continued for at least another three-month period (i.e., a total of six months). The decision to extend therapy to one year of treatment should be based on prognostic factors (e.g., genotype, age > 40 years, male gender, bridging fibrosis).

### Dose modification for all patients

If severe adverse reactions or laboratory abnormalities develop during treatment with PegIntron monotherapy or PegIntron in combination with ribavirin, modify the dosages of each product as appropriate, until the adverse reactions abate. As adherence might be of importance for outcome of therapy, the dose should be kept as close as possible to the recommended standard dose. Guidelines were developed in clinical trials for dose modification.

### Combination Therapy Dose Reduction Guidelines

<b>Table 2a</b> Dose modification guidelines for combination therapy (with ribavirin)			
Laboratory values	Reduce only ribavirin dose to <u>600 mg/day</u> * if:	Reduce only PegIntron dose to one-half dose if:	Discontinue combination therapy if:
Haemoglobin	< 10 g/dl	-	< 8.5 g/dl
Haemoglobin in: Patients with history of stable cardiac disease	≥ 2 g/dl decrease in haemoglobin during any four week period during treatment (permanent dose reduction)		< 12 g/dl after four weeks of dose reduction
White blood cells	-	< 1.5 x 10 <sup>9</sup> /l	< 1.0 x 10 <sup>9</sup> /l
Neutrophils	-	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	-	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l
Bilirubin – direct	-	-	2.5 x ULN**



Bilirubin - indirect	> 5 mg/dl	-	> 4 mg/dl (for > 4 weeks)
Creatinine	-	-	> 2.0 mg/dl
ALT/AST	-	-	2 x baseline and > 10 x ULN**

\* Patients whose dose of ribavirin is reduced to 600 mg daily receive one 200 mg capsule in the morning and two 200 mg capsules in the evening.

\*\* Upper limit of normal

Dose reduction of PegIntron may be accomplished by either reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 2b**.

<b>Table 2b – Reduced PegIntron Dosing for Combination Therapy</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5 ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
< 40	25	50*	0.25	25
40-50	32	50	0.3	30
51-64	40	50	0.4	40
65-75	50	50	0.5	50
76-85	60	80	0.4	64
> 85	75	100	0.4	80
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### **PegIntron Monotherapy Dose Reduction Guidelines**

Dose modification guidelines for patients who use PegIntron monotherapy are shown in **Table 3a**.

<b>Table 3a Dose modification guidelines for PegIntron monotherapy</b>		
<b>Laboratory values</b>	<b>Reduce PegIntron to one-half dose if:</b>	<b>Discontinue PegIntron if:</b>
Neutrophils	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l

Dose reduction for patients who use 0.5 µg/kg PegIntron monotherapy must be accomplished by reducing the prescribed volume by one-half. The 50 µg/0.5 ml vial must be used if necessary since the pen can only deliver a minimum volume of 0.3 ml.

For patients who use 1.0 µg/kg PegIntron monotherapy, dose reduction may be accomplished by reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 3b**.

<b>Table 3b – Reduced PegIntron Dose for the 1.0 µg/kg Monotherapy Regimen</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>

30-35	15	50*	0.15	15
36-45	20	50*	0.20	20
46-56	25	50*	0.25	25
57-72	32	50	0.3	30
73-89	40	50	0.4	40
90-106	50	50	0.5	50
> 106	60	80	0.4	64
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### Special populations

**Use in renal impairment:** *Monotherapy:* PegIntron should be used with caution in patients with moderate to severe renal impairment. In patients with moderate renal dysfunction (creatinine clearance 30-50 ml/minute), the starting dose of PegIntron should be reduced by 25 %. Patients with severe renal dysfunction (creatinine clearance 15-29 ml/minute) should have the starting dose of PegIntron reduced by 50 %. Data are not available for the use of PegIntron in patients with creatinine clearance < 15 ml/minute (see section 5.2). Patients with severe renal impairment, including those on hemodialysis, should be closely monitored. If renal function decreases during treatment, PegIntron therapy should be discontinued.

*Combination therapy:* Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see ribavirin SPC). When administered in combination with ribavirin, subjects with impaired renal function should be more carefully monitored with respect to the development of anaemia.

**Use in hepatic impairment:** The safety and efficacy of PegIntron therapy has not been evaluated in patients with severe hepatic dysfunction, therefore PegIntron must not be used for these patients.

**Use in the elderly (≥ 65 years of age):** There are no apparent age-related effects on the pharmacokinetics of PegIntron. Data from elderly patients treated with a single dose of PegIntron suggest no alteration in PegIntron dose is necessary based on age (see section 5.2).

**Use in patients under the age of 18 years:** There is no experience in children (see section 5.2).

### 4.3 Contraindications

- Hypersensitivity to the active substance or to any interferon or to any of the excipients;
- A history of severe pre-existing cardiac disease, including unstable or uncontrolled cardiac disease in the previous six months (see section 4.4);
- Severe, debilitating medical conditions;
- Autoimmune hepatitis or a history of autoimmune disease;
- Severe hepatic dysfunction or decompensated cirrhosis of the liver;
- Pre-existing thyroid disease unless it can be controlled with conventional treatment;
- Epilepsy and/or compromised central nervous system (CNS) function;
- Initiation of PegIntron is contraindicated in HCV/HIV patients with cirrhosis and a Child-Pugh score ≥ 6.

*Combination therapy with ribavirin:* Also see ribavirin Summary of the Product Characteristics (SPC) if PegIntron is to be administered in combination with ribavirin in patients with chronic hepatitis C.

### 4.4 Special warnings and precautions for use

**Psychiatric and Central Nervous System (CNS):** Severe CNS effects, particularly depression, suicidal ideation and attempted suicide have been observed in some patients during PegIntron therapy, and even after treatment discontinuation mainly during the 6-month follow-up period. Other CNS effects including aggressive behaviour (sometimes directed against others), confusion and alterations of mental status have been observed with alpha interferons. Patients should be closely monitored for any signs or symptoms of psychiatric disorders. If such symptoms appear, the potential seriousness of these undesirable effects must be borne in mind by the prescribing physician and the need for adequate therapeutic management should be considered. If psychiatric symptoms persist or worsen, or suicidal ideation is identified, it is recommended that treatment with PegIntron be discontinued, and the patient followed, with psychiatric intervention as appropriate.

*Patients with existence of, or history of severe psychiatric conditions:* If treatment with peginterferon alfa-2b is judged necessary in patients with existence or history of severe psychiatric conditions, this should only be initiated after having ensured appropriate individualised diagnostic and therapeutic management of the psychiatric condition.

There is no experience with PegIntron in combination with ribavirin in patients who have relapsed after interferon alpha + ribavirin therapy.

All patients in the chronic hepatitis C studies had a liver biopsy before inclusion, but in certain cases (i.e. patients with genotype 2 and 3), treatment may be possible without histological confirmation. Current treatment guidelines should be consulted as to whether a liver biopsy is needed prior to commencing treatment.

**Acute hypersensitivity:** Acute hypersensitivity reactions (e.g., urticaria, angioedema, bronchoconstriction, anaphylaxis) have been observed rarely during interferon alfa-2b therapy. If such a reaction develops during treatment with PegIntron, discontinue treatment and institute appropriate medical therapy immediately. Transient rashes do not necessitate interruption of treatment.

**Cardiovascular system:** As with interferon alfa-2b, patients with a history of congestive heart failure, myocardial infarction and/or previous or current arrhythmic disorders, receiving PegIntron therapy require close monitoring. It is recommended that patients who have pre-existing cardiac abnormalities have electrocardiograms taken prior to and during the course of treatment. Cardiac arrhythmias (primarily supraventricular) usually respond to conventional therapy but may require discontinuation of PegIntron therapy.

More significant obtundation and coma, including cases of encephalopathy, have been observed in some patients, usually elderly, treated at higher doses for oncology indications. While these effects are generally reversible, in a few patients full resolution took up to three weeks. Very rarely, seizures have occurred with high doses of interferon alpha.

**Liver function:** As with all interferons, discontinue treatment with PegIntron in patients who develop prolongation of coagulation markers which might indicate liver decompensation.

**Fever:** While fever may be associated with the flu-like syndrome reported commonly during interferon therapy, other causes of persistent fever must be ruled out.

**Hydration:** Adequate hydration must be maintained in patients undergoing PegIntron therapy since hypotension related to fluid depletion has been seen in some patients treated with alpha interferons. Fluid replacement may be necessary.

**Pulmonary changes:** Pulmonary infiltrates, pneumonitis, and pneumonia, occasionally resulting in fatality, have been observed rarely in interferon alpha treated patients. Any patient developing fever, cough, dyspnea or other respiratory symptoms must have a chest X-ray taken. If the chest X-ray shows pulmonary infiltrates or there is evidence of pulmonary function impairment, the patient is to be monitored closely, and, if appropriate, discontinue interferon alpha. Prompt discontinuation of interferon alpha administration and treatment with corticosteroids appear to be associated with resolution of pulmonary adverse events.

**Autoimmune disease:** The development of auto-antibodies and autoimmune disorders has been reported during treatment with alpha interferons. Patients predisposed to the development of autoimmune disorders may be at increased risk. Patients with signs or symptoms compatible with autoimmune disorders should be evaluated carefully, and the benefit-risk of continued interferon therapy should be reassessed (see also section 4.4 Thyroid changes and 4.8).

**Ocular changes:** Ophthalmologic disorders, including retinal haemorrhages, cotton wool spots, and retinal artery or vein obstruction have been reported in rare instances after treatment with alpha interferons (see section 4.8). All patients should have a baseline eye examination. Any patient complaining of ocular symptoms, including loss of visual acuity or visual field must have a prompt and complete eye examination. Periodic visual examinations are recommended during PegIntron therapy, particularly in patients with disorders that may be associated with retinopathy, such as diabetes mellitus or hypertension. Discontinuation of PegIntron should be considered in patients who develop new or worsening ophthalmological disorders.

**Thyroid changes:** Infrequently, patients treated for chronic hepatitis C with interferon alpha have developed thyroid abnormalities, either hypothyroidism or hyperthyroidism. Determine thyroid stimulating hormone (TSH) levels if, during the course of therapy, a patient develops symptoms consistent with possible thyroid dysfunction. In the presence of thyroid dysfunction, PegIntron treatment may be continued if TSH levels can be maintained in the normal range by medication.

**Metabolic disturbances:** Hypertriglyceridemia and aggravation of hypertriglyceridemia, sometimes severe, have been observed. Monitoring of lipid levels is, therefore, recommended.

### **HCV/HIV Co-infection**

Mitochondrial toxicity and lactic acidosis:

Patients co-infected with HIV and receiving Highly Active Anti-Retroviral Therapy (HAART) may be at increased risk of developing lactic acidosis. Caution should be used when adding PegIntron and ribavirin to HAART therapy (see ribavirin SPC).

Moreover, patients treated with PegIntron and ribavirin combination therapy and zidovudine are at increased risk of developing anaemia.

Hepatic decompensation in HCV/HIV co-infected patients with advanced cirrhosis:

Co-infected patients with advanced cirrhosis receiving HAART may be at increased risk of hepatic decompensation and death. Adding treatment with alfa interferons alone or in combination with ribavirin may increase the risk in this patient subset. Other baseline factors in co-infected patients that may be associated with a higher risk of hepatic decompensation include treatment with didanosine and elevated bilirubin serum concentration.

Co-infected patients receiving both antiretroviral (ARV) and anti-hepatitis treatment should be closely monitored, assessing their Child-Pugh score during treatment. Patients progressing to hepatic decompensation should have their anti-hepatitis treatment immediately discontinued and the ARV treatment reassessed.

Haematological abnormalities in HCV/HIV co-infected patients:

HCV/HIV co-infected patients receiving peginterferon alfa-2b/ribavirin treatment and HAART may be at increased risk to develop haematological abnormalities (as neutropenia, thrombocytopenia and anaemia) compared to HCV mono-infected patients. Although, the majority of them could be managed by dose reduction, close monitoring of haematological parameters should be undertaken in this population of patients (see section 4.2 and below “Laboratory tests” and section 4.8).

Patients with low CD4 counts:

In patients co-infected with HCV/HIV, limited efficacy and safety data (N = 25) are available in subjects with CD4 counts less than 200 cells/ $\mu$ l. Caution is therefore warranted in the treatment of patients with low CD4 counts.

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal

products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron and ribavirin.

**Dental and periodontal disorders:** Dental and periodontal disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and mucous membranes of the mouth during long-term treatment with the combination of PegIntron and ribavirin. Patients should brush their teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If this reaction occurs, they should be advised to rinse out their mouth thoroughly afterwards.

**Organ transplant recipients:** The safety and efficacy of PegIntron alone or in combination with ribavirin for the treatment of hepatitis C in liver or other organ transplant recipients have not been studied. Preliminary data indicates that interferon alpha therapy may be associated with an increased rate of kidney graft rejection. Liver graft rejection has also been reported.

**Other:** Due to reports of interferon alpha exacerbating pre-existing psoriatic disease and sarcoidosis, use of PegIntron in patients with psoriasis or sarcoidosis is recommended only if the potential benefit justifies the potential risk.

**Laboratory tests:** Standard haematologic tests, blood chemistry and a test of thyroid function must be conducted in all patients prior to initiating therapy. Acceptable baseline values that may be considered as a guideline prior to initiation of PegIntron therapy are:

- |                    |                              |
|--------------------|------------------------------|
| • Platelets        | $\geq 100,000/\text{mm}^3$   |
| • Neutrophil count | $\geq 1,500/\text{mm}^3$     |
| • TSH level        | must be within normal limits |

Laboratory evaluations are to be conducted at weeks 2 and 4 of therapy, and periodically thereafter as clinically appropriate.

**Important information about some of the ingredients of PegIntron:**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

Patients with rare hereditary problems of fructose intolerance, glucose galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Results from a multiple-dose probe study assessing P450 substrates in chronic hepatitis C patients receiving once weekly PegIntron (1.5 µg/kg) for 4 weeks demonstrated an increase in activity of CYP2D6 and CYP2C8/9. No change in activity of CYP1A2, CYP3A4, or N-acetyltransferase was observed.

Caution should be used when administering peginterferon alfa-2b with medicines metabolised by CYP2D6 and CYP2C8/9, especially those with narrow therapeutic window, such as warfarin and phenytoin (CYP2C9) and flecainide (CYP2D6).

These findings may partly relate to improved metabolic capacity due to reduced hepatic inflammation in patients undergoing treatment with PegIntron. Caution is therefore advised when PegIntron treatment is initiated for chronic hepatitis in patients treated with medication with a narrow therapeutic window and sensitive to mild metabolic impairment of the liver.

No pharmacokinetic interactions were noted between PegIntron and ribavirin in a multiple-dose pharmacokinetic study.

**Methadone:** In patients with chronic hepatitis C that were on stable methadone maintenance therapy and naïve to peginterferon alfa-2b, addition of 1.5 microgram/kg/week of PegIntron subcutaneously for 4 weeks increased R-methadone AUC by approximately 15 % (95 % CI for AUC ratio estimate 103 – 128 %). The clinical significance of this finding is unknown; however, patients should be monitored for signs and symptoms of increased sedative effect, as well as respiratory depression. Especially in patients on a high dose of methadone, the risk for QTc prolongation should be considered.

#### *HCV/HIV Co-infection*

Nucleoside analogs: ribavirin has been shown *in vitro* to inhibit phosphorylation of zidovudine and stavudine. The clinical significance of these finding is unknown. However, these *in vitro* findings raise the possibility that concurrent use of ribavirin with either zidovudine or stavudine might lead to increased HIV plasma viraemia. Therefore, it is recommended that plasma HIV RNA levels be closely monitored in patients treated with ribavirin concurrently with either of these two agents. If HIV RNA levels increase, the use of ribavirin concomitantly with reverse transcriptase inhibitors must be reviewed (see ribavirin SPC).

Use of nucleoside analogs, alone or in combination with other nucleosides, has resulted in lactic acidosis. Pharmacologically, ribavirin increases phosphorylated metabolites of purine nucleosides *in vitro*. This activity could potentiate the risk of lactic acidosis induced by purine nucleoside analogs (e.g. didanosine or abacavir). Co-administration of ribavirin and didanosine is not recommended. Reports of mitochondrial toxicity, in particular lactic acidosis and pancreatitis, of which some fatal, have been reported (see ribavirin SPC).

## **4.6 Pregnancy and lactation**

PegIntron is recommended for use in fertile women only when they are using effective contraception during the treatment.

There are no adequate data from the use of interferon alfa-2b in pregnant women.. Studies in animals have shown reproductive toxicity (see section 5.3). Interferon alfa-2b has been shown to be abortifacient in primates. PegIntron is likely to also cause this effect.

The potential risk in humans is unknown. PegIntron is to be used during pregnancy only if the potential benefit justifies the potential risk to the foetus.

**Lactation:** It is not known whether the components of this medicinal product are excreted in human milk. Because of the potential for adverse reactions in breast-fed infants, breast-feeding should be discontinued prior to initiation of treatment.

#### *Combination therapy with ribavirin:*

Ribavirin causes serious birth defects when administered during pregnancy. Ribavirin therapy is contraindicated in women who are pregnant. Extreme care must be taken to avoid pregnancy in female patients or in partners of male patients taking PegIntron in combination with ribavirin. Females of childbearing potential and their partners must each use an effective contraceptive during treatment and for 4 months after treatment has been concluded. Male patients and their female partners must each use an effective contraceptive during treatment and for 7 months after treatment has been concluded (see Ribavirin SPC).

## **4.7 Effects on ability to drive and use machines**

Patients who develop fatigue, somnolence or confusion during treatment with PegIntron are cautioned to avoid driving or operating machinery.

## **4.8 Undesirable effects**

The safety of PegIntron is evaluated from data from two clinical trials: one with PegIntron monotherapy, one with PegIntron in combination with ribavirin. In both cases, patients were treated for one year.

**Table 4** describes the regimens and patient exposure for one year of treatment in patients with no previous exposure to interferon (interferon-naïve patients). Because of a significant overlap in the pattern of undesirable effects with PegIntron monotherapy, groups of patients have been brought together in **Table 5** to show the pattern of reported effects for all monotherapy groups.

<b>Table 4</b> Regimens and patient exposure		
<b>Treatment</b>	<b>Regimen</b>	<b>Number of patients treated for one year</b>
PegIntron + ribavirin	PegIntron (1.5 micrograms/kg/week) + ribavirin (> 10.6 mg/kg/day)	188
Interferon alfa-2b + ribavirin	Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day)	505
PegIntron monotherapy	PegIntron (0.5 microgram/kg/week)	315
	PegIntron (1.0 microgram/kg/week)	297
	PegIntron (1.5 micrograms/kg/week)	304

<b>Table 5</b> Undesirable effects very commonly reported in clinical trials > 10 % of patients in PegIntron + ribavirin group Very common (> 1/10)			
	<b>PegIntron + ribavirin</b>	<b>Interferon alfa- 2b + ribavirin</b>	<b>PegIntron monotherapy</b>
<b>Infections and infestations</b>			
Infection viral	10 %	5 %	4-5 %
<b>Metabolism and nutrition disorders</b>			
Weight decrease	30 %	19 %	8-18 %
<b>Psychiatric disorders</b>			
Depression	34 %	32 %	26 %
Irritability	32 %	34 %	19 %
Insomnia	37 %	41 %	16-19 %
Anxiety	14 %	14 %	8 %
Concentration impaired	18 %	21 %	9-10 %
Emotional lability	11 %	10 %	5 %
<b>Nervous system disorders</b>			
Headache	58 %	57 %	57-63 %
Mouth dry	10 %	8 %	4-8 %
<b>Respiratory, thoracic and mediastinal disorders</b>			
Pharyngitis	10 %	7 %	3 %
Coughing	14 %	11 %	4 %
Dyspnea	26 %	22 %	5 %
<b>Gastrointestinal disorders</b>			
Nausea	43 %	31 %	20-23 %
Anorexia	35 %	26 %	10-25 %
Diarrhoea	20 %	13 %	14-17 %
Abdominal pain	12 %	9 %	11 %
Vomiting	16 %	10 %	4-7 %
<b>Skin and subcutaneous tissue disorders</b>			
Alopecia	45 %	32 %	20-34 %
Pruritus	27 %	27 %	7-9 %
Skin dry	23 %	21 %	6-9 %
Rash	21 %	21 %	5-7 %
<b>Musculoskeletal and connective tissue disorders</b>			
Myalgia	49 %	49 %	46-60 %
Arthralgia	31 %	26 %	23-28 %
Musculoskeletal pain	15 %	11 %	11-13 %
<b>General disorders and administration site conditions</b>			
Injection site inflammation	20 %	17 %	39-44 %
Injection site reaction	54 %	36 %	7-9 %
Fatigue	56 %	59 %	43 %
Rigors	42 %	40 %	33-43 %
Fever	39 %	32 %	29-43 %
Flu-like symptoms	21 %	23 %	18-25 %
Asthenia	28 %	17 %	12-14 %
Dizziness	17 %	16 %	7-12 %



<b>Table 6</b> Undesirable effects commonly reported in clinical trials in patients treated with PegIntron + ribavirin or PegIntron monotherapy Common (> 1/100, < 1/10)		
<b>Body system</b>	<b>5-10%</b>	<b>1-&lt;5%</b>
Infections and infestations		otitis media, fungal infection, bacterial infection, herpes simplex
Blood and lymphatic system disorders	Anaemia, leukopaenia	Thrombocytopenia, lymphadenopathy
Endocrine disorders	hypothyroidism	hyperthyroidism
Metabolism and nutrition disorders		hyperuricemia, hypocalcemia, thirst
Psychiatric disorders	agitation, nervousness	aggressive behaviour, somnolence, behavior disorder, apathy, appetite increased, sleep disorder, dreaming abnormal, decreased libido
Nervous system disorders	paresthesia, increased sweating	hypoesthesia, hyperaesthesia, hypertonia, confusion, tremor, vertigo, migraine, ataxia, neuralgia
Eye disorders		blurred vision, conjunctivitis, lacrimal gland disorder, eye pain
Cardiac disorders		tachycardia, palpitation
Vascular disorders		hypotension, hypertension, syncope, flushing
Ear and labyrinth disorders		tinnitus, hearing impairment/loss
Respiratory, thoracic and mediastinal disorders		nonproductive cough, rhinitis, sinusitis, bronchitis, respiratory disorder, nasal congestion, rhinorrhea, dysphonia, epistaxis
Gastrointestinal disorders	dyspepsia	constipation, taste perversion, loose stools, stomatitis, ulcerative stomatitis, gingival bleeding, glossitis, flatulence, hemorrhoids, gastroesophageal reflux, gingivitis, dehydration
Hepatobiliary disorders		hepatomegaly, hyperbilirubinemia
Skin and subcutaneous tissue disorders		erythematous rash, eczema, photosensitivity reaction, maculopapular rash, abnormal hair texture, acne, dermatitis, furunculosis, nail disorder, psoriasis, urticaria, erythema, face or peripheral oedema
Musculoskeletal and connective tissue disorders		arthritis
Renal and urinary disorders		micturition frequency, urine abnormal
Reproductive system and breast disorders	menstrual disorder, menorrhagia	ovarian disorder, vaginal disorder, sexual dysfunction (not specified), impotence, breast pain, amenorrhoea, prostatitis
General disorders and administration site conditions	RUQ pain, malaise, chest pain	injection site pain

Most cases of neutropaenia and thrombocytopaenia were mild (WHO grades 1 or 2). There were some cases of more severe neutropenia in patients treated with the recommended doses of PegIntron in combination with ribavirin (WHO grade 3: 39 of 186 [21 %]; and WHO grade 4: 13 of 186 [7 %]).

In a clinical trial, approximately 1.2 % of patients treated with PegIntron or interferon alfa-2b in combination with ribavirin reported life-threatening psychiatric events during treatment. These events included suicidal ideation and attempted suicide (see section 4.4). Following marketing, psychosis and hallucination have been reported rarely.

Rarely ( $> 1/10,000$ ,  $< 1/1,000$ ) or very rarely ( $< 1/10,000$ ) reported events with interferon alfa-2b, including PegIntron, include:

**Blood and lymphatic system disorders:**

Very rarely interferon alfa-2b or PegIntron used alone or in combination with ribavirin may be associated with aplastic anaemia. Pure red cell aplasia has been reported.

**Immune system disorders:**

very rarely: sarcoidosis or exacerbation of sarcoidosis

**Endocrine disorders:**

rarely: diabetes

**Nervous system disorders:**

rarely: seizure, peripheral neuropathy

very rarely: cerebrovascular ischaemia, cerebrovascular haemorrhage, encephalopathy

**Cardiac disorders:**

rarely: arrhythmia

very rarely: cardiac ischaemia, myocardial infarction

**Respiratory, thoracic and mediastinal disorders:**

very rarely: interstitial lung disease

**Gastrointestinal disorders:**

rarely: pancreatitis

very rarely: ulcerative and ischaemic colitis

**Skin and subcutaneous tissue disorders:**

very rarely: erythema multiforme, Stevens Johnson syndrome, toxic epidermal necrolysis, injection site necrosis

**Musculoskeletal and connective tissue disorders:**

rarely: rhabdomyolysis, myositis

**Renal and urinary disorders:**

rarely: renal insufficiency and renal failure

Cardiovascular (CVS) adverse events, particularly arrhythmia, appeared to be correlated mostly with pre-existing CVS disease and prior therapy with cardiotoxic agents (see section 4.4). Cardiomyopathy, that may be reversible upon discontinuation of interferon alpha, has been reported rarely in patients without prior evidence of cardiac disease.

Ophthalmological disorders that have been reported rarely with alpha interferons include retinopathies (including macular oedema), retinal haemorrhages, retinal artery or vein obstruction, cotton wool spots, loss of visual acuity or visual field, optic neuritis, and papilloedema (see section 4.4).

A wide variety of autoimmune and immune-mediated disorders have been reported with alpha interferons including thyroid disorders, systemic lupus erythematosus, rheumatoid arthritis (new or aggravated), idiopathic and thrombotic thrombocytopenic purpura, vasculitis, neuropathies including mononeuropathies (see also section 4.4, **Autoimmune disorders**).

**HCV/HIV co-infected patients**

For HCV/HIV co-infected patients receiving PegIntron in combination with ribavirin, other undesirable effects (that were not reported in mono-infected patients) which have been reported in the larger studies with a frequency  $> 5\%$  were: oral candidiasis (14 %), lipodystrophy acquired (13 %), CD4 lymphocytes decreased (8 %), appetite decreased (8 %), gamma-glutamyltransferase increased (9 %), back pain (5 %), blood amylase increased (6 %), blood lactic acid increased (5 %), cytolytic hepatitis (6 %), lipase increased (6 %) and pain in limb (6 %).

*Mitochondrial toxicity:*

Mitochondrial toxicity and lactic acidosis have been reported in HIV-positive patients receiving NRTI

regimen and associated ribavirin for co-HCV infection (see section 4.4).

*Laboratory values for HCV/HIV co-infected patients:*

Although haematological toxicities of neutropenia, thrombocytopenia and anaemia occurred more frequently in HCV/HIV co-infected patients, the majority could be managed by dose modification and rarely required premature discontinuation of treatment (see section 4.4). Haematological abnormalities were more frequently reported in patients receiving PegIntron in combination with ribavirin when compared to patients receiving interferon alfa-2b in combination with ribavirin. In Study 1 (see section 5.1), decrease in absolute neutrophil count levels below 500 cells/mm<sup>3</sup> was observed in 4 % (8/194) of patients and decrease in platelets below 50,000/mm<sup>3</sup> was observed in 4 % (8/194) of patients receiving PegIntron in combination with ribavirin. Anaemia (hemoglobin < 9.4g/dl) was reported in 12% (23/194) of patients treated with PegIntron in combination with ribavirin.

*CD4 lymphocytes decrease:*

Treatment with PegIntron in combination with ribavirin was associated with decreases in absolute CD4+ cell counts within the first 4 weeks without a reduction in CD4+ cell percentage. The decrease in CD4+ cell counts was reversible upon dose reduction or cessation of therapy. The use of PegIntron in combination with ribavirin had no observable negative impact on the control of HIV viraemia during therapy or follow-up. Limited safety data (N= 25) are available in co-infected patients with CD4+ cell counts < 200/μl (see section 4.4).

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron in combination with ribavirin.

## **4.9 Overdose**

In clinical trials, cases of accidental overdose, at never more than twice the prescribed dose, were reported. There were no serious reactions. Undesirable effects resolved during continued administration of PegIntron.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Immunostimulants, Cytokines and immunomodulators, Interferons, Peginterferon alfa-2b, ATC code: L03A B10.

Recombinant interferon alfa-2b is covalently conjugated with monomethoxy polyethylene glycol at an average degree of substitution of 1 mole of polymer/mole of protein. The average molecular mass is approximately 31,300 daltons of which the protein moiety constitutes approximately 19,300.

#### **Interferon alfa-2b**

*In vitro* and *in vivo* studies suggest that the biological activity of PegIntron is derived from its interferon alfa-2b moiety.

Interferons exert their cellular activities by binding to specific membrane receptors on the cell surface. Studies with other interferons have demonstrated species specificity. However, certain monkey species, e.g., Rhesus monkeys, are susceptible to pharmacodynamic stimulation upon exposure to human type 1 interferons.

Once bound to the cell membrane, interferon initiates a complex sequence of intracellular events that include the induction of certain enzymes. It is thought that this process, at least in part, is responsible for the various cellular responses to interferon, including inhibition of virus replication in virus-infected cells, suppression of cell proliferation and such immunomodulating activities as enhancement of the phagocytic activity of macrophages and augmentation of the specific cytotoxicity of

lymphocytes for target cells. Any or all of these activities may contribute to interferon's therapeutic effects.

Recombinant interferon alfa-2b also inhibits viral replication *in vitro* and *in vivo*. Although the exact antiviral mode of action of recombinant interferon alfa-2b is unknown, it appears to alter the host cell metabolism. This action inhibits viral replication or if replication occurs, the progeny virions are unable to leave the cell.

### **PegIntron**

PegIntron pharmacodynamics were assessed in a rising single-dose trial in healthy subjects by examining changes in oral temperature, concentrations of effector proteins such as serum neopterin and 2'5'-oligoadenylate synthetase (2'5'-OAS), as well as white cell and neutrophil counts. Subjects treated with PegIntron showed mild dose-related elevations in body temperature. Following single doses of PegIntron between 0.25 and 2.0 micrograms/kg/week, serum neopterin concentration was increased in a dose-related manner. Neutrophil and white cell count reductions at the end of week 4 correlated with the dose of PegIntron.

### **PegIntron clinical trials**

Two pivotal trials have been conducted, one (C/I97-010) with PegIntron monotherapy; the other (C/I98-580) with PegIntron in combination with ribavirin. Eligible patients for these trials had chronic hepatitis C confirmed by a positive HCV-RNA polymerase chain reaction (PCR) assay (> 30 IU/ml), a liver biopsy consistent with a histological diagnosis of chronic hepatitis with no other cause for the chronic hepatitis, and abnormal serum ALT.

In the PegIntron monotherapy trial, a total of 916 naïve chronic hepatitis C patients were treated with PegIntron (0.5, 1.0 or 1.5 micrograms/kg/week) for one year with a follow-up period of six months. In addition, 303 patients received interferon alfa-2b (3 million International Units [MIU] three times a week) as a comparator. This study showed that PegIntron was superior to interferon alfa-2b (**Table 8**).

In the PegIntron combination trial, 1,530 naïve patients were treated for one year with one of the following combination regimens:

- PegIntron (1.5 micrograms/kg/week) + ribavirin (800 mg/day), (n = 511).
- PegIntron (1.5 micrograms/kg/week for one month followed by 0.5 microgram/kg/week for 11 months) + ribavirin (1,000/1,200 mg/day), (n = 514).
- Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day) (n = 505).

In this trial, the combination of PegIntron (1.5 micrograms/kg/week) and ribavirin was significantly more effective than the combination of interferon alfa-2b and ribavirin (**Table 7**), particularly in patients infected with Genotype 1 (**Table 8**). Sustained response was assessed by the response rate six months after the cessation of treatment.

HCV genotype and baseline virus load are prognostic factors which are known to affect response rates. However, response rates in this trial were shown to be dependent also on the dose of ribavirin administered in combination with PegIntron or interferon alfa-2b. In those patients that received > 10.6 mg/kg ribavirin (800 mg dose in typical 75 kg patient), regardless of genotype or viral load, response rates were significantly higher than in those patients that received ≤ 10.6 mg/kg ribavirin (**Table 8**), while response rates in patients that received > 13.2 mg/kg ribavirin were even higher.

<b>Table 7</b> Sustained virological response (% patients HCV negative)							
	<b>PegIntron monotherapy</b>				<b>PegIntron + ribavirin</b>		
<b>Treatment regimen</b>	<b>P 1.5</b>	<b>P 1.0</b>	<b>P 0.5</b>	<b>I</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>Number of patients</b>	304	297	315	303	511	514	505
<b>Response at end of treatment</b>	49 %	41 %	33 %	24 %	65 %	56 %	54 %

<b>Sustained response</b>	<b>23 %*</b>	<b>25 %</b>	<b>18 %</b>	<b>12 %</b>	<b>54 %**</b>	<b>47 %</b>	<b>47 %</b>
---------------------------	--------------	-------------	-------------	-------------	---------------	-------------	-------------

P 1.5 PegIntron 1.5 micrograms/kg  
 P 1.0 PegIntron 1.0 microgram/kg  
 P 0.5 PegIntron 0.5 microgram/kg  
 I Interferon alfa-2b 3 MIU  
 P 1.5/R PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)  
 P 0.5/R PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)  
 I/R Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)  
 \* p < 0.001 P 1.5 vs. I  
 \*\* p = 0.0143 P 1.5/R vs. I/R

<b>Table 8</b> Sustained response rates with PegIntron + ribavirin (by ribavirin dose, genotype and viral load)				
<b>HCV Genotype</b>	<b>Ribavirin dose (mg/kg)</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>All Genotypes</b>	<b>All</b>	<b>54 %</b>	<b>47 %</b>	<b>47 %</b>
	≤ 10.6	50 %	41 %	27 %
	> 10.6	61 %	48 %	47 %
<b>Genotype 1</b>	<b>All</b>	<b>42 %</b>	<b>34 %</b>	<b>33 %</b>
	≤ 10.6	38 %	25 %	20 %
	> 10.6	48 %	34 %	34 %
Genotype 1 ≤ 600,000 IU/ml	All	73 %	51 %	45 %
	≤ 10.6	74 %	25 %	33 %
	> 10.6	71 %	52 %	45 %
Genotype 1 > 600,000 IU/ml	All	30 %	27 %	29 %
	≤ 10.6	27 %	25 %	17 %
	> 10.6	37 %	27 %	29 %
<b>Genotype 2/3</b>	<b>All</b>	<b>82 %</b>	<b>80 %</b>	<b>79 %</b>
	≤ 10.6	79 %	73 %	50 %
	> 10.6	88 %	80 %	80 %

P 1.5/R PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)  
 P 0.5/R PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)  
 I/R Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)

In the PegIntron monotherapy study, the Quality of Life was generally less affected by 0.5 microgram/kg of PegIntron than by either 1.0 microgram/kg of PegIntron once weekly or 3 MIU of interferon alfa-2b three times a week.

In a separate trial, 224 patients with genotype 2 or 3 received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with ribavirin 800 mg –1,400 mg p.o. for 6 months (based on body weight, only three patients weighing > 105 kg, received the 1,400 mg dose, which has not yet been validated) (**Table 9**). Twenty-four % had bridging fibrosis or cirrhosis (Knodell 3/4).

**Table 9. Virologic Response at End of Treatment, Sustained Virologic Response and Relapse by HCV Genotype and Viral Load\***

	PegIntron 1.5 µg/kg once weekly plus Ribavirin 800-1,400 mg/day		
	End of Treatment Response	Sustained Virologic Response	Relapse
All Subjects	94 % (211/224)	81 % (182/224)	12 % (27/224)
HCV 2 ≤ 600,000 IU/ml > 600,000	100 % (42/42)	93 % (39/42)	7 % (3/42)
	100 % (20/20)	95 % (19/20)	5 % (1/20)
	100 % (22/22)	91 % (20/22)	9 % (2/22)

IU/ml			
HCV 3	93 % (169/182)	79 % (143/182)	14 % (24/166)
≤ 600,000 IU/ml	93 % (92/99)	86 % (85/99)	8 % (7/91)
> 600,000 IU/ml	93 % (77/83)	70 % (58/83)	23 % (17/75)

\* Any subject with an undetectable HCV-RNA level at the Follow-Up Week 12 visit and missing data at the Follow-Up Week 24 visit was considered a sustained responder. Any subject with missing data in and after the Follow-Up Week 12 window was considered to be a non-responder at Week 24 of follow-up.

The 6 month treatment duration in this trial was better tolerated than one year of treatment in the pivotal combination trial; for discontinuation 5 % vs. 14 %, for dose modification 18 % vs. 49 %.

In a non-comparative trial, 235 patients with genotype 1 and low viral load (< 600,000 IU/ml) received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with weight adjusted ribavirin. The overall sustained response rate after a 24-week treatment duration was 50 %. Forty-one percent of subjects (97/235) had nondetectable plasma HCV-RNA levels at Week 4 and Week 24 of therapy. In this subgroup, there was a 92 % (89/97) sustained virological response rate. The high sustained response rate in this subgroup of patients was identified in an interim analysis (n=49) and prospectively confirmed (n=48).

Limited historical data indicate that treatment for 48 weeks might be associated with a higher sustained response rate (11/11) and with a lower risk of relapse (0/11 as compared to 7/96 following 24 weeks of treatment).

### Predictability of sustained virological response

Virological response by week 12, defined as a 2-log viral load decrease or undetectable levels of HCV-RNA has been shown to be predictive for sustained response (**Table 10**).

<b>Table 10</b> Predictability of sustained response by viral response at week 12 and genotype*				
Treatment	Genotype	Viral response at week 12	Sustained response	Negative predictive value
PegIntron 1.5 + ribavirin (> 10.6 mg/kg) 48-week treatment	1	Yes 75 % (82/110)	71 % (58/82)	----
		No 25 % (28/110)	0 % (0/28)	100 %
PegIntron 1.5 + ribavirin 800-1,400 mg 24-week treatment	2 and 3	Yes 99 % (213/215)	83 % (177/213)	----
		No 1 % (2/215)	50 % (1/2)	50 %

\* reflects patients with 12 week data available

The negative predictive value for sustained response in patients treated with PegIntron in monotherapy was 98 %.

### HCV/HIV Co-infected patients

Two trials have been conducted in patients co-infected with HIV and HCV. The response to treatment in both of these trials is presented in **Table 11**. Study 1 (RIBAVIC; P01017) was a randomized, multicentre study which enrolled 412 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (1.5 µg/kg/week) plus ribavirin (800 mg/day) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800 mg/day) for 48 weeks with a follow-up period of 6 months. Study 2 (P02080) was a randomized, single centre

study that enrolled 95 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (100 or 150 µg /week based on weight) plus ribavirin (800-1,200 mg/day based on weight) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800-1,200 mg/day based on weight). The duration of therapy was 48 weeks with a follow-up period of 6 months except for patients infected with genotypes 2 or 3 and viral load < 800,000 IU/ml (Amplicor) who were treated for 24 weeks with a 6-month follow-up period.

<b>Table 11</b> Sustained virological response based on genotype after PegIntron in combination with Ribavirin in HCV/HIV Co-infected patients						
	<b>Study 1<sup>1</sup></b>			<b>Study 2<sup>2</sup></b>		
	PegIntron (1.5 µg/kg/week) + ribavirin (800 mg)	Interferon alfa-2b (3 MIU TIW) + ribavirin (800 mg)	p value <sup>a</sup>	PegIntron (100 or 150 <sup>c</sup> µg/week) + ribavirin (800-1,200 mg) <sup>d</sup>	Interferon alfa-2b (3 MIU TIW) + ribavirin (800-1,200 mg) <sup>d</sup>	p value <sup>b</sup>
All	27 % (56/205)	20 % (41/205)	0.047	44 % (23/52)	21 % (9/43)	0.017
Genotype 1, 4	17 % (21/125)	6 % (8/129)	0.006	38 % (12/32)	7 % (2/27)	0.007
Genotype 2, 3	44 % (35/80)	43 % (33/76)	0.88	53 % (10/19)	47 % (7/15)	0.730

MIU = million international units; TIW = three times a week.

a: p value based on Cochran-Mantel Haenszel Chi square test.

b: p value based on chi-square test.

c: subjects < 75 kg received 100 µg/week PegIntron and subjects ≥ 75 kg received 150 µg/week PegIntron.

d: ribavirin dosing was 800 mg for patients < 60 kg, 1,000 mg for patients 60-75 kg, and 1,200 mg for patients > 75 kg.

<sup>1</sup>Carrato F, Bani-Sadir F, Pol S et al. JAMA 2004; 292(23): 2839-2848.

<sup>2</sup>Laguno M, Murillas J, Blanco J et al. AIDS 2004; 18(13): F27-F36.

### *Histological response*

Liver biopsies were obtained before and after treatment in Study 1 and were available for 210 of the 412 subjects (51 %). Both the Metavir score and Ishak grade decreased among subjects treated with PegIntron in combination with ribavirin. This decline was significant among responders (-0.3 for Metavir and -1.2 for Ishak) and stable (-0.1 for Metavir and -0.2 for Ishak) among non-responders. In terms of activity, about one-third of sustained responders showed improvement and none showed worsening. There was no improvement in terms of fibrosis observed in this study. Steatosis was significantly improved in patients infected with HCV Genotype 3.

## **5.2 Pharmacokinetic properties**

PegIntron is a well characterized polyethylene glycol-modified (“pegylated”) derivative of interferon alfa-2b and is predominantly composed of monopegylated species. The plasma half-life of PegIntron is prolonged compared with non-pegylated interferon alfa-2b. PegIntron has a potential to depegylate to free interferon alfa-2b. The biologic activity of the pegylated isomers is qualitatively similar, but weaker than free interferon alfa-2b.

Following subcutaneous administration, maximal serum concentrations occur between 15-44 hours post-dose, and are sustained for up to 48-72 hours post-dose.

PegIntron C<sub>max</sub> and AUC measurements increase in a dose-related manner. Mean apparent volume of distribution is 0.99 l/kg.

Upon multiple dosing, there is an accumulation of immunoreactive interferons. There is, however, only a modest increase in biologic activity as measured by a bioassay.

Mean (SD) PegIntron elimination half-life is approximately 40 hours (13.3 hours), with apparent clearance of 22.0 ml/hr·kg. The mechanisms involved in clearance of interferons in man have not yet been fully elucidated. However, renal elimination may account for a minority (approximately 30 %) of PegIntron apparent clearance.

**Renal function:** Renal clearance appears to account for 30 % of total clearance of PegIntron. In a single dose study (1.0 microgram/kg) in patients with impaired renal function,  $C_{max}$ , AUC, and half-life increased in relation to the degree of renal impairment.

Following multiple dosing of PegIntron (1.0 microgram/kg subcutaneously administered every week for four weeks) the clearance of PegIntron is reduced by a mean of 17 % in patients with moderate renal impairment (creatinine clearance 30-49 ml/minute) and by a mean of 44 % in patients with severe renal impairment (creatinine clearance 15-29 ml/minute) compared to subjects with normal renal function. Based on single dose data, clearance was similar in patients with severe renal impairment not on dialysis and in patients who were receiving hemodialysis. The dose of PegIntron for monotherapy should be reduced in patients with moderate or severe renal impairment (see sections 4.2 and 4.4). Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see section 4.3).

Because of marked inter-subject variability in interferon pharmacokinetics, it is recommended that patients with severe renal impairment be closely monitored during treatment with PegIntron (see section 4.2)

**Hepatic function:** The pharmacokinetics of PegIntron have not been evaluated in patients with severe hepatic dysfunction.

**Elderly patients ≥ 65 years of age:** The pharmacokinetics of PegIntron following a single subcutaneous dose of 1.0 microgram/kg were not affected by age. The data suggest that no alteration in PegIntron dosage is necessary based on advancing age.

**Patients under the age of 18 years:** Specific pharmacokinetic evaluations have not been performed on these patients. PegIntron is indicated for the treatment of chronic hepatitis C only in patients 18 years of age or older.

**Interferon neutralising factors:** Interferon neutralising factor assays were performed on serum samples of patients who received PegIntron in the clinical trial. Interferon neutralising factors are antibodies which neutralise the antiviral activity of interferon. The clinical incidence of neutralising factors in patients who received PegIntron 0.5 micrograms/kg is 1.1 %.

### 5.3 Preclinical safety data

**PegIntron:** Adverse events not observed in clinical trials were not seen in toxicity studies in monkeys. These studies were limited to four weeks due to the appearance of anti-interferon antibodies in most monkeys.

Reproduction studies of PegIntron have not been performed. Interferon alfa-2b has been shown to be an abortifacient in primates. PegIntron is likely to also cause this effect. Effects on fertility have not been determined. It is not known whether the components of this medicinal product are excreted into experimental animal or human milk (see section 4.6 for relevant human data on pregnancy and lactation). PegIntron showed no genotoxic potential.

The relative non-toxicity of monomethoxy-polyethylene glycol (mPEG), which is liberated from PegIntron by metabolism *in vivo* has been demonstrated in preclinical acute and subchronic toxicity studies in rodents and monkeys, standard embryo-foetal development studies and in *in vitro* mutagenicity assays.



**PegIntron plus ribavirin:** When used in combination with ribavirin, PegIntron did not cause any effects not previously seen with either active substance alone. The major treatment-related change was a reversible, mild to moderate anaemia, the severity of which was greater than that produced by either active substance alone.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Powder for solution for injection:

Disodium phosphate, anhydrous,  
Sodium dihydrogen phosphate dihydrate,  
Sucrose,  
Polysorbate 80.

Solvent for parenteral use:

Water for injections.

### **6.2 Incompatibilities**

This medicinal product should only be reconstituted with the solvent provided (see section 6.6). In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

3 years

After reconstitution:

- Chemical and physical in-use stability has been demonstrated for 24 hours at 2°C - 8°C.
- From a microbiological point of view, the product is to be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C - 8°C.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C - 8°C).

### **6.5 Nature and contents of container**

The powder is contained in a 2 ml vial, Type I flint glass, with a butyl rubber stopper in an aluminium flip-off seal with a polypropylene bonnet. The solvent is presented in a 2 ml ampoule, Type I flint glass. PegIntron 150 micrograms is supplied as:

- 1 vial of powder for solution for injection and 1 ampoule of solvent for parenteral use;
- 1 vial of powder for solution for injection, 1 ampoule of solvent for parenteral use, 1 injection syringe, 2 injection needles and 1 cleansing swab;
- 4 vials of powder for solution for injection and 4 ampoules of solvent for parenteral use;
- 4 vials of powder for solution for injection, 4 ampoules of solvent for parenteral use, 4 injection syringes, 8 injection needles and 4 cleansing swabs;
- 6 vials of powder for solution for injection and 6 ampoules of solvent for parenteral use.
- 12 vials of powder for solution for injection, 12 ampoules of solvent for parenteral use, 12 injection syringes, 24 injection needles and 12 cleansing swabs.

Not all pack sizes may be marketed

### **6.6 Special precautions for disposal**

PegIntron is supplied as a powder of peginterferon alfa-2b at a strength of 150 micrograms for single use. Each vial must be reconstituted with 0.7 ml of water for injections for administration of up to 0.5 ml of solution. A small volume is lost during preparation of PegIntron for injection when the dose is measured and injected. Therefore, each vial contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection. The reconstituted solution has a concentration of 150 micrograms/0.5 ml.

Using a sterilised injection syringe and injection needle, inject 0.7 ml of water for injections into the vial of PegIntron. Agitate gently to complete dissolution of powder. The appropriate dose can then be withdrawn with a sterilised injection syringe and injected. A complete set of instructions is provided in the Annex to the Package Leaflet.

As for all parenteral medicinal products, inspect visually the reconstituted solution prior to administration. The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. Discard any unused material.

## **7. MARKETING AUTHORISATION HOLDER**

SP Europe  
73, rue de Stalle  
B-1180 Bruxelles  
Belgium

## **8. MARKETING AUTHORISATION NUMBERS**

EU/1/00/131/021  
EU/1/00/131/022  
EU/1/00/131/023  
EU/1/00/131/024  
EU/1/00/131/025  
EU/1/00/131/030

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 25 May 2000  
Date of last renewal: 25 May 2005

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>

## **1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 50 micrograms, powder and solvent for solution for injection in pre-filled pen

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each pre-filled pen of PegIntron 50 micrograms contains a sufficient amount of peginterferon alfa-2b as measured on a protein basis in a powder for solution for injection, and the corresponding amount of solvent, to provide 50 micrograms in 0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

The active substance is a covalent conjugate of recombinant interferon alfa-2b\* with monomethoxy polyethylene glycol. The potency of this product should not be compared to that of another pegylated or non-pegylated protein of the same therapeutic class. For more information, see section 5.1.

\*produced by rDNA technology in *E.coli* cells harbouring a genetically engineered plasmid hybrid encompassing an interferon alfa-2b gene from human leukocytes

For a full list of excipients, see section 6.1. PegIntron contains 40 mg of sucrose per 0.5 ml.

## **3. PHARMACEUTICAL FORM**

Powder and solvent for solution for injection in pre-filled pen

White powder.

Clear and colourless solvent.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

PegIntron is indicated for the treatment of adult patients with chronic hepatitis C who have elevated transaminases without liver decompensation and who are positive for serum HCV-RNA or anti-HCV, including naïve patients with clinically stable HIV co-infection (see section 4.4).

The best way to use PegIntron in this indication is in combination with ribavirin.

This combination is indicated in naïve patients including patients with clinically stable HIV co-infection. This combination of PegIntron with ribavirin is also indicated in patients who have previously responded (with normalisation of ALT at the end of treatment) to interferon alpha monotherapy but who have subsequently relapsed.

Interferon monotherapy, including PegIntron, is indicated mainly in case of intolerance or contraindication to ribavirin.

Please refer also to the ribavirin Summary of Product Characteristics (SPC) when PegIntron is to be used in combination with ribavirin.

### **4.2 Posology and method of administration**

Treatment should be initiated and monitored only by a physician experienced in the management of patients with hepatitis C.

#### **Dose to be administered**

PegIntron should be administered as a once weekly subcutaneous injection. The dose administered depends on whether it is used in combination with ribavirin or as monotherapy.

## Combination therapy

PegIntron 1.5 micrograms/kg/week in combination with ribavirin capsules.

The intended dose of 1.5 µg/kg of PegIntron to be used in combination with ribavirin may be delivered in weight categories with the pen/vial strengths according to **Table 1**. Ribavirin capsules are to be administered orally each day in two divided doses with food (morning and evening).

**Table 1 - Dosing for Combination Therapy**

Body Weight (kg)	PegIntron		Ribavirin Capsules	
	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Total Daily Dose (mg)	Number of Capsules (200 mg)
< 40	50	0.5	800	4 <sup>a</sup>
40-50	80	0.4	800	4 <sup>a</sup>
51-64	80	0.5	800	4 <sup>a</sup>
65-75	100	0.5	1,000	5 <sup>b</sup>
76-85	120	0.5	1,000	5 <sup>b</sup>
> 85	150	0.5	1,200	6 <sup>c</sup>

a: 2 morning, 2 evening

b: 2 morning, 3 evening

c: 3 morning, 3 evening

## Duration of treatment

**Predictability of sustained virological response:** Patients infected with virus genotype 1 who fail to achieve virological response at Week 12 are highly unlikely to become sustained virological responders (see also section 5.1).

- **Genotype 1:** For patients who exhibit virological response at week 12, treatment should be continued for another nine month period (i.e., a total of 48 weeks).  
In the subset of patients with genotype 1 infection and low viral load (< 600,000 IU/ml) who become HCV-RNA negative at treatment week 4 and remain HCV-RNA negative at week 24, the treatment could either be stopped after this 24 week treatment course or pursued for an additional 24 weeks (i.e. overall 48 weeks treatment duration). However, an overall 24 weeks treatment duration may be associated with a higher risk of relapse than a 48 weeks treatment duration (see section 5.1).
- **Genotypes 2 or 3:** It is recommended that all patients be treated for 24 weeks, except for HCV/HIV co-infected patients who should receive 48 weeks of treatment.
- **Genotype 4:** In general, patients infected with genotype 4 are considered harder to treat and limited study data (n=66) indicate they are compatible with a duration of treatment as for genotype 1.

## HCV/HIV Co-infection

The recommended duration of dosing for HCV/HIV co-infected patients is 48 weeks, regardless of genotype.

### *Predictability of response and non-response in HCV/HIV Co-infection*

Early virological response by Week 12, defined as a 2 log viral load decrease or undetectable levels of HCV-RNA, has been shown to be predictive for sustained response. The negative predictive value for sustained response in HCV/HIV co-infected patients treated with PegIntron in combination with ribavirin was 99 % (67/68; Study 1) (see section 5.1). A positive predictive value of 50 % (52/104; Study 1) was observed for HCV/HIV co-infected patients receiving combination therapy.

### PegIntron monotherapy

As monotherapy the PegIntron regimen is 0.5 or 1.0 microgram/kg/week. The lowest vial or pen strength available is 50 µg/0.5 ml; therefore for patients prescribed 0.5 µg/kg/week, doses must be adjusted by volume as shown in **Table 2**. For the 1.0 µg/kg dose, similar volume adjustments can be made or alternate vial strengths can be used as shown in **Table 2**. PegIntron monotherapy was not studied in HCV/HIV co-infected patients.

**Table 2- Monotherapy Dosing**

	0.5 µg/kg		1.0 µg/kg	
Body Weight (kg)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)
30-35	50*	0.15	50	0.3
36-45	50*	0.2	50	0.4
46-56	50*	0.25	50	0.5
57-72	50	0.3	80	0.4
73-88	50	0.4	80	0.5
89-106	50	0.5	100	0.5
> 106**	80	0.4	120	0.5
* Must use vial. Minimum delivery for pen is 0.3 ml.				
** For patients > 120 kg, use 80 µg/0.5 ml vial				

### Duration of treatment

For patients who exhibit virological response at Week 12, treatment should be continued for at least another three-month period (i.e., a total of six months). The decision to extend therapy to one year of treatment should be based on prognostic factors (e.g., genotype, age > 40 years, male gender, bridging fibrosis).

### Dose modification for all patients

If severe adverse reactions or laboratory abnormalities develop during treatment with PegIntron monotherapy or PegIntron in combination with ribavirin, modify the dosages of each product as appropriate, until the adverse reactions abate. As adherence might be of importance for outcome of therapy, the dose should be kept as close as possible to the recommended standard dose. Guidelines were developed in clinical trials for dose modification.

### Combination Therapy Dose Reduction Guidelines

<b>Table 2a</b> Dose modification guidelines for combination therapy (with ribavirin)			
Laboratory values	Reduce only ribavirin dose to 600 mg/day* if:	Reduce only PegIntron dose to one-half dose if:	Discontinue combination therapy if:
Haemoglobin	< 10 g/dl	-	< 8.5 g/dl
Haemoglobin in: Patients with history of stable cardiac disease	≥ 2 g/dl decrease in haemoglobin during any four week period during treatment (permanent dose reduction)		< 12 g/dl after four weeks of dose reduction
White blood cells	-	< 1.5 x 10 <sup>9</sup> /l	< 1.0 x 10 <sup>9</sup> /l
Neutrophils	-	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	-	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l
Bilirubin – direct	-	-	2.5 x ULN**

Bilirubin – indirect	> 5 mg/dl	-	> 4 mg/dl (for > 4 weeks)
Creatinine	-	-	> 2.0 mg/dl
ALT/AST	-	-	2 x baseline and > 10 x ULN**

\* Patients whose dose of ribavirin is reduced to 600 mg daily receive one 200 mg capsule in the morning and two 200 mg capsules in the evening.

\*\* Upper limit of normal

Dose reduction of PegIntron may be accomplished by either reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 2b**.

<b>Table 2b – Reduced PegIntron Dosing for Combination Therapy</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5 ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
< 40	25	50*	0.25	25
40-50	32	50	0.3	30
51-64	40	50	0.4	40
65-75	50	50	0.5	50
76-85	60	80	0.4	64
> 85	75	100	0.4	80
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### **PegIntron Monotherapy Dose Reduction Guidelines**

Dose modification guidelines for patients who use PegIntron monotherapy are shown in **Table 3a**.

<b>Table 3a Dose modification guidelines for PegIntron monotherapy</b>		
<b>Laboratory values</b>	<b>Reduce PegIntron to one-half dose if:</b>	<b>Discontinue PegIntron if:</b>
Neutrophils	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l

Dose reduction for patients who use 0.5 µg/kg PegIntron monotherapy must be accomplished by reducing the prescribed volume by one-half. The 50 µg/0.5 ml vial must be used if necessary since the pen can only deliver a minimum volume of 0.3 ml.

For patients who use 1.0 µg/kg PegIntron monotherapy, dose reduction may be accomplished by reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 3b**.

<b>Table 3b – Reduced PegIntron Dose for the 1.0 µg/kg Monotherapy Regimen</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
30-35	15	50*	0.15	15

36-45	20	50*	0.20	20
46-56	25	50*	0.25	25
57-72	32	50	0.3	30
73-89	40	50	0.4	40
90-106	50	50	0.5	50
> 106	60	80	0.4	64
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### Special populations

**Use in renal impairment:** *Monotherapy:* PegIntron should be used with caution in patients with moderate to severe renal impairment. In patients with moderate renal dysfunction (creatinine clearance 30-50 ml/minute), the starting dose of PegIntron should be reduced by 25 %. Patients with severe renal dysfunction (creatinine clearance 15-29 ml/minute) should have the starting dose of PegIntron reduced by 50 %. Data are not available for the use of PegIntron in patients with creatinine clearance < 15 ml/minute (see section 5.2). Patients with severe renal impairment, including those on hemodialysis, should be closely monitored. If renal function decreases during treatment, PegIntron therapy should be discontinued.

*Combination therapy:* Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see ribavirin SPC). When administered in combination with ribavirin, subjects with impaired renal function should be more carefully monitored with respect to the development of anaemia.

**Use in hepatic impairment:** The safety and efficacy of PegIntron therapy has not been evaluated in patients with severe hepatic dysfunction, therefore PegIntron must not be used for these patients.

**Use in the elderly (≥ 65 years of age):** There are no apparent age-related effects on the pharmacokinetics of PegIntron. Data from elderly patients treated with a single dose of PegIntron suggest no alteration in PegIntron dose is necessary based on age (see section 5.2).

**Use in patients under the age of 18 years:** There is no experience in children (see section 5.2).

### 4.3 Contraindications

- Hypersensitivity to the active substance or to any interferon or to any of the excipients;
- A history of severe pre-existing cardiac disease, including unstable or uncontrolled cardiac disease in the previous six months (see section 4.4);
- Severe, debilitating medical conditions;
- Autoimmune hepatitis or a history of autoimmune disease;
- Severe hepatic dysfunction or decompensated cirrhosis of the liver;
- Pre-existing thyroid disease unless it can be controlled with conventional treatment;
- Epilepsy and/or compromised central nervous system (CNS) function;
- Initiation of PegIntron is contraindicated in HCV/HIV patients with cirrhosis and a Child-Pugh score ≥ 6.

*Combination therapy with ribavirin:* Also see ribavirin Summary of the Product Characteristics (SPC) if PegIntron is to be administered in combination with ribavirin in patients with chronic hepatitis C.

### 4.4 Special warnings and precautions for use

**Psychiatric and Central Nervous System (CNS):** Severe CNS effects, particularly depression, suicidal ideation and attempted suicide have been observed in some patients during PegIntron therapy, and even after treatment discontinuation mainly during the 6-month follow-up period. Other CNS effects including

aggressive behaviour (sometimes directed against others), confusion and alterations of mental status have been observed with alpha interferons. Patients should be closely monitored for any signs or symptoms of psychiatric disorders. If such symptoms appear, the potential seriousness of these undesirable effects must be borne in mind by the prescribing physician and the need for adequate therapeutic management should be considered. If psychiatric symptoms persist or worsen, or suicidal ideation is identified, it is recommended that treatment with PegIntron be discontinued, and the patient followed, with psychiatric intervention as appropriate.

*Patients with existence of, or history of severe psychiatric conditions:* If treatment with peginterferon alfa-2b is judged necessary in patients with existence or history of severe psychiatric conditions, this should only be initiated after having ensured appropriate individualised diagnostic and therapeutic management of the psychiatric condition.

There is no experience with PegIntron in combination with ribavirin in patients who have relapsed after interferon alpha + ribavirin therapy.

All patients in the chronic hepatitis C studies had a liver biopsy before inclusion, but in certain cases (i.e. patients with genotype 2 and 3), treatment may be possible without histological confirmation. Current treatment guidelines should be consulted as to whether a liver biopsy is needed prior to commencing treatment.

**Acute hypersensitivity:** Acute hypersensitivity reactions (e.g., urticaria, angioedema, bronchoconstriction, anaphylaxis) have been observed rarely during interferon alfa-2b therapy. If such a reaction develops during treatment with PegIntron, discontinue treatment and institute appropriate medical therapy immediately. Transient rashes do not necessitate interruption of treatment.

**Cardiovascular system:** As with interferon alfa-2b, patients with a history of congestive heart failure, myocardial infarction and/or previous or current arrhythmic disorders, receiving PegIntron therapy require close monitoring. It is recommended that patients who have pre-existing cardiac abnormalities have electrocardiograms taken prior to and during the course of treatment. Cardiac arrhythmias (primarily supraventricular) usually respond to conventional therapy but may require discontinuation of PegIntron therapy.

More significant obtundation and coma, including cases of encephalopathy, have been observed in some patients, usually elderly, treated at higher doses for oncology indications. While these effects are generally reversible, in a few patients full resolution took up to three weeks. Very rarely, seizures have occurred with high doses of interferon alpha.

**Liver function:** As with all interferons, discontinue treatment with PegIntron in patients who develop prolongation of coagulation markers which might indicate liver decompensation.

**Fever:** While fever may be associated with the flu-like syndrome reported commonly during interferon therapy, other causes of persistent fever must be ruled out.

**Hydration:** Adequate hydration must be maintained in patients undergoing PegIntron therapy since hypotension related to fluid depletion has been seen in some patients treated with alpha interferons. Fluid replacement may be necessary.

**Pulmonary changes:** Pulmonary infiltrates, pneumonitis, and pneumonia, occasionally resulting in fatality, have been observed rarely in interferon alpha treated patients. Any patient developing fever, cough, dyspnea or other respiratory symptoms must have a chest X-ray taken. If the chest X-ray shows pulmonary infiltrates or there is evidence of pulmonary function impairment, the patient is to be monitored closely, and, if appropriate, discontinue interferon alpha. Prompt discontinuation of interferon alpha administration and treatment with corticosteroids appear to be associated with resolution of pulmonary adverse events.

**Autoimmune disease:** The development of auto-antibodies and autoimmune disorders has been reported during treatment with alpha interferons. Patients predisposed to the development of



autoimmune disorders may be at increased risk. Patients with signs or symptoms compatible with autoimmune disorders should be evaluated carefully, and the benefit-risk of continued interferon therapy should be reassessed (see also section 4.4 Thyroid changes and 4.8).

**Ocular changes:** Ophthalmologic disorders, including retinal haemorrhages, cotton wool spots, and retinal artery or vein obstruction have been reported in rare instances after treatment with alpha interferons (see section 4.8). All patients should have a baseline eye examination. Any patient complaining of ocular symptoms, including loss of visual acuity or visual field must have a prompt and complete eye examination. Periodic visual examinations are recommended during PegIntron therapy, particularly in patients with disorders that may be associated with retinopathy, such as diabetes mellitus or hypertension. Discontinuation of PegIntron should be considered in patients who develop new or worsening ophthalmological disorders.

**Thyroid changes:** Infrequently, patients treated for chronic hepatitis C with interferon alpha have developed thyroid abnormalities, either hypothyroidism or hyperthyroidism. Determine thyroid stimulating hormone (TSH) levels if, during the course of therapy, a patient develops symptoms consistent with possible thyroid dysfunction. In the presence of thyroid dysfunction, PegIntron treatment may be continued if TSH levels can be maintained in the normal range by medication.

**Metabolic disturbances:** Hypertriglyceridemia and aggravation of hypertriglyceridemia, sometimes severe, have been observed. Monitoring of lipid levels is, therefore, recommended.

### **HCV/HIV Co-infection**

Mitochondrial toxicity and lactic acidosis:

Patients co-infected with HIV and receiving Highly Active Anti-Retroviral Therapy (HAART) may be at increased risk of developing lactic acidosis. Caution should be used when adding PegIntron and ribavirin to HAART therapy (see ribavirin SPC).

Moreover, patients treated with PegIntron and ribavirin combination therapy and zidovudine are at increased risk of developing anaemia.

Hepatic decompensation in HCV/HIV co-infected patients with advanced cirrhosis:

Co-infected patients with advanced cirrhosis receiving HAART may be at increased risk of hepatic decompensation and death. Adding treatment with alfa interferons alone or in combination with ribavirin may increase the risk in this patient subset. Other baseline factors in co-infected patients that may be associated with a higher risk of hepatic decompensation include treatment with didanosine and elevated bilirubin serum concentration.

Co-infected patients receiving both antiretroviral (ARV) and anti-hepatitis treatment should be closely monitored, assessing their Child-Pugh score during treatment. Patients progressing to hepatic decompensation should have their anti-hepatitis treatment immediately discontinued and the ARV treatment reassessed.

Haematological abnormalities in HCV/HIV co-infected patients:

HCV/HIV co-infected patients receiving peginterferon alfa-2b/ribavirin treatment and HAART may be at increased risk to develop haematological abnormalities (as neutropenia, thrombocytopenia and anaemia) compared to HCV mono-infected patients. Although, the majority of them could be managed by dose reduction, close monitoring of haematological parameters should be undertaken in this population of patients (see section 4.2 and below “Laboratory tests” and section 4.8).

Patients with low CD4 counts:

In patients co-infected with HCV/HIV, limited efficacy and safety data (N = 25) are available in subjects with CD4 counts less than 200 cells/ $\mu$ l. Caution is therefore warranted in the treatment of patients with low CD4 counts.

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron and ribavirin.

**Dental and periodontal disorders:** Dental and periodontal disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and mucous membranes of the mouth during long-term treatment with the combination of PegIntron and ribavirin. Patients should brush their teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If this reaction occurs, they should be advised to rinse out their mouth thoroughly afterwards.

**Organ transplant recipients:** The safety and efficacy of PegIntron alone or in combination with ribavirin for the treatment of hepatitis C in liver or other organ transplant recipients have not been studied. Preliminary data indicates that interferon alpha therapy may be associated with an increased rate of kidney graft rejection. Liver graft rejection has also been reported.

**Other:** Due to reports of interferon alpha exacerbating pre-existing psoriatic disease and sarcoidosis, use of PegIntron in patients with psoriasis or sarcoidosis is recommended only if the potential benefit justifies the potential risk.

**Laboratory tests:** Standard haematologic tests, blood chemistry and a test of thyroid function must be conducted in all patients prior to initiating therapy. Acceptable baseline values that may be considered as a guideline prior to initiation of PegIntron therapy are:

- |                    |                              |
|--------------------|------------------------------|
| • Platelets        | $\geq 100,000/\text{mm}^3$   |
| • Neutrophil count | $\geq 1,500/\text{mm}^3$     |
| • TSH level        | must be within normal limits |

Laboratory evaluations are to be conducted at weeks 2 and 4 of therapy, and periodically thereafter as clinically appropriate.

**Important information about some of the ingredients of PegIntron:**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

Patients with rare hereditary problems of fructose intolerance, glucose galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Results from a multiple-dose probe study assessing P450 substrates in chronic hepatitis C patients receiving once weekly PegIntron (1.5  $\mu\text{g}/\text{kg}$ ) for 4 weeks demonstrated an increase in activity of CYP2D6 and CYP2C8/9. No change in activity of CYP1A2, CYP3A4, or N-acetyltransferase was observed.

Caution should be used when administering peginterferon alfa-2b with medicines metabolised by CYP2D6 and CYP2C8/9, especially those with narrow therapeutic window, such as warfarin and phenytoin (CYP2C9) and flecainide (CYP2D6).

These findings may partly relate to improved metabolic capacity due to reduced hepatic inflammation in patients undergoing treatment with PegIntron. Caution is therefore advised when PegIntron treatment is initiated for chronic hepatitis in patients treated with medication with a narrow therapeutic window and sensitive to mild metabolic impairment of the liver.

No pharmacokinetic interactions were noted between PegIntron and ribavirin in a multiple-dose pharmacokinetic study.

**Methadone:** In patients with chronic hepatitis C that were on stable methadone maintenance therapy and naïve to peginterferon alfa-2b, addition of 1.5 microgram/kg/week of PegIntron subcutaneously for 4 weeks increased R-methadone AUC by approximately 15 % (95 % CI for AUC ratio estimate 103 – 128 %). The clinical significance of this finding is unknown; however, patients should be

monitored for signs and symptoms of increased sedative effect, as well as respiratory depression. Especially in patients on a high dose of methadone, the risk for QTc prolongation should be considered.

#### *HCV/HIV Co-infection*

Nucleoside analogs: ribavirin has been shown *in vitro* to inhibit phosphorylation of zidovudine and stavudine. The clinical significance of these findings is unknown. However, these *in vitro* findings raise the possibility that concurrent use of ribavirin with either zidovudine or stavudine might lead to increased HIV plasma viraemia. Therefore, it is recommended that plasma HIV RNA levels be closely monitored in patients treated with ribavirin concurrently with either of these two agents. If HIV RNA levels increase, the use of ribavirin concomitantly with reverse transcriptase inhibitors must be reviewed (see ribavirin SPC).

Use of nucleoside analogs, alone or in combination with other nucleosides, has resulted in lactic acidosis. Pharmacologically, ribavirin increases phosphorylated metabolites of purine nucleosides *in vitro*. This activity could potentiate the risk of lactic acidosis induced by purine nucleoside analogs (e.g. didanosine or abacavir). Co-administration of ribavirin and didanosine is not recommended. Reports of mitochondrial toxicity, in particular lactic acidosis and pancreatitis, of which some fatal, have been reported (see ribavirin SPC).

## **4.6 Pregnancy and lactation**

PegIntron is recommended for use in fertile women only when they are using effective contraception during the treatment.

There are no adequate data from the use of interferon alfa-2b in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3). Interferon alfa-2b has been shown to be abortifacient in primates. PegIntron is likely to also cause this effect.

The potential risk in humans is unknown. PegIntron is to be used during pregnancy only if the potential benefit justifies the potential risk to the foetus

**Lactation:** It is not known whether the components of this medicinal product are excreted in human milk. Because of the potential for adverse reactions in breast-fed infants, breast-feeding should be discontinued prior to initiation of treatment.

#### *Combination therapy with ribavirin:*

Ribavirin causes serious birth defects when administered during pregnancy. Ribavirin therapy is contraindicated in women who are pregnant. Extreme care must be taken to avoid pregnancy in female patients or in partners of male patients taking PegIntron in combination with ribavirin. Females of childbearing potential and their partners must each use an effective contraceptive during treatment and for 4 months after treatment has been concluded. Male patients and their female partners must each use an effective contraceptive during treatment and for 7 months after treatment has been concluded (see Ribavirin SPC).

## **4.7 Effects on ability to drive and use machines**

Patients who develop fatigue, somnolence or confusion during treatment with PegIntron are cautioned to avoid driving or operating machinery.

## **4.8 Undesirable effects**

The safety of PegIntron is evaluated from data from two clinical trials: one with PegIntron monotherapy, one with PegIntron in combination with ribavirin. In both cases, patients were treated for one year.

**Table 4** describes the regimens and patient exposure for one year of treatment in patients with no previous exposure to interferon (interferon-naïve patients). Because of a significant overlap in the

pattern of undesirable effects with PegIntron monotherapy, groups of patients have been brought together in **Table 5** to show the pattern of reported effects for all monotherapy groups.

<b>Table 4</b> Regimens and patient exposure		
<b>Treatment</b>	<b>Regimen</b>	<b>Number of patients treated for one year</b>
PegIntron + ribavirin	PegIntron (1.5 micrograms/kg/week) + ribavirin (> 10.6 mg/kg/day)	188
Interferon alfa-2b + ribavirin	Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day)	505
PegIntron monotherapy	PegIntron (0.5 microgram/kg/week)	315
	PegIntron (1.0 microgram/kg/week)	297
	PegIntron (1.5 micrograms/kg/week)	304

<b>Table 5</b> Undesirable effects very commonly reported in clinical trials > 10 % of patients in PegIntron + ribavirin group Very common (> 1/10)			
	<b>PegIntron + ribavirin</b>	<b>Interferon alfa- 2b + ribavirin</b>	<b>PegIntron monotherapy</b>
<b>Infections and infestations</b>			
Infection viral	10 %	5 %	4-5 %
<b>Metabolism and nutrition disorders</b>			
Weight decrease	30 %	19 %	8-18 %
<b>Psychiatric disorders</b>			
Depression	34 %	32 %	26 %
Irritability	32 %	34 %	19 %
Insomnia	37 %	41 %	16-19 %
Anxiety	14 %	14 %	8 %
Concentration impaired	18 %	21 %	9-10 %
Emotional lability	11 %	10 %	5 %
<b>Nervous system disorders</b>			
Headache	58 %	57 %	57-63 %
Mouth dry	10 %	8 %	4-8 %
<b>Respiratory, thoracic and mediastinal disorders</b>			
Pharyngitis	10 %	7 %	3 %
Coughing	14 %	11 %	4 %
Dyspnea	26 %	22 %	5 %
<b>Gastrointestinal disorders</b>			
Nausea	43 %	31 %	20-23 %
Anorexia	35 %	26 %	10-25 %
Diarrhoea	20 %	13 %	14-17 %
Abdominal pain	12 %	9 %	11 %
Vomiting	16 %	10 %	4-7 %
<b>Skin and subcutaneous tissue disorders</b>			
Alopecia	45 %	32 %	20-34 %
Pruritus	27 %	27 %	7-9 %
Skin dry	23 %	21 %	6-9 %
Rash	21 %	21 %	5-7 %
<b>Musculoskeletal and connective tissue disorders</b>			
Myalgia	49 %	49 %	46-60 %
Arthralgia	31 %	26 %	23-28 %
Musculoskeletal pain	15 %	11 %	11-13 %
<b>General disorders and administration site conditions</b>			
Injection site inflammation	20 %	17 %	39-44 %
Injection site reaction	54 %	36 %	7-9 %
Fatigue	56 %	59 %	43 %
Rigors	42 %	40 %	33-43 %
Fever	39 %	32 %	29-43 %
Flu-like symptoms	21 %	23 %	18-25 %
Asthenia	28 %	17 %	12-14 %
Dizziness	17 %	16 %	7-12 %

<b>Table 6</b> Undesirable effects commonly reported in clinical trials in patients treated with PegIntron + ribavirin or PegIntron monotherapy Common (> 1/100, < 1/10)		
<b>Body system</b>	<b>5-10%</b>	<b>1-&lt;5%</b>
Infections and infestations		otitis media, fungal infection, bacterial infection, herpes simplex
Blood and lymphatic system disorders	Anaemia, leukopaenia	Thrombocytopenia, lymphadenopathy
Endocrine disorders	hypothyroidism	hyperthyroidism
Metabolism and nutrition disorders		hyperuricemia, hypocalcemia, thirst
Psychiatric disorders	agitation, nervousness	aggressive behaviour, somnolence, behavior disorder, apathy, appetite increased, sleep disorder, dreaming abnormal, decreased libido
Nervous system disorders	paresthesia, increased sweating	hypoesthesia, hyperaesthesia, hypertonia, confusion, tremor, vertigo, migraine, ataxia, neuralgia
Eye disorders		blurred vision, conjunctivitis, lacrimal gland disorder, eye pain
Cardiac disorders		tachycardia, palpitation
Vascular disorders		hypotension, hypertension, syncope, flushing
Ear and labyrinth disorders		tinnitus, hearing impairment/loss
Respiratory, thoracic and mediastinal disorders		nonproductive cough, rhinitis, sinusitis, bronchitis, respiratory disorder, nasal congestion, rhinorrhea, dysphonia, epistaxis
Gastrointestinal disorders	dyspepsia	constipation, taste perversion, loose stools, stomatitis, ulcerative stomatitis, gingival bleeding, glossitis, flatulence, hemorrhoids, gastroesophageal reflux, gingivitis, dehydration
Hepatobiliary disorders		hepatomegaly, hyperbilirubinemia
Skin and subcutaneous tissue disorders		erythematous rash, eczema, photosensitivity reaction, maculopapular rash, abnormal hair texture, acne, dermatitis, furunculosis, nail disorder, psoriasis, urticaria, erythema, face or peripheral oedema
Musculoskeletal and connective tissue disorders		arthritis
Renal and urinary disorders		micturition frequency, urine abnormal
Reproductive system and breast disorders	menstrual disorder, menorrhagia	ovarian disorder, vaginal disorder, sexual dysfunction (not specified), impotence, breast pain, amenorrhoea, prostatitis
General disorders and administration site conditions	RUQ pain, malaise, chest pain	injection site pain

Most cases of neutropaenia and thrombocytopaenia were mild (WHO grades 1 or 2). There were some cases of more severe neutropenia in patients treated with the recommended doses of PegIntron in combination with ribavirin (WHO grade 3: 39 of 186 [21 %]; and WHO grade 4: 13 of 186 [7 %]).

In a clinical trial, approximately 1.2 % of patients treated with PegIntron or interferon alfa-2b in combination with ribavirin reported life-threatening psychiatric events during treatment. These events included suicidal ideation and attempted suicide (see section 4.4). Following marketing, psychosis and hallucination have been reported rarely.

Rarely ( $> 1/10,000$ ,  $< 1/1,000$ ) or very rarely ( $< 1/10,000$ ) reported events with interferon alfa-2b, including PegIntron, include:

**Blood and lymphatic system disorders:**

Very rarely interferon alfa-2b or PegIntron used alone or in combination with ribavirin may be associated with aplastic anaemia. Pure red cell aplasia has been reported.

**Immune system disorders:**

very rarely: sarcoidosis or exacerbation of sarcoidosis

**Endocrine disorders:**

rarely: diabetes

**Nervous system disorders:**

rarely: seizure, peripheral neuropathy

very rarely: cerebrovascular ischaemia, cerebrovascular haemorrhage, encephalopathy

**Cardiac disorders:**

rarely: arrhythmia

very rarely: cardiac ischaemia, myocardial infarction

**Respiratory, thoracic and mediastinal disorders:**

very rarely: interstitial lung disease

**Gastrointestinal disorders:**

rarely: pancreatitis

very rarely: ulcerative and ischaemic colitis

**Skin and subcutaneous tissue disorders:**

very rarely: erythema multiforme, Stevens Johnson syndrome, toxic epidermal necrolysis, injection site necrosis

**Musculoskeletal and connective tissue disorders:**

rarely: rhabdomyolysis, myositis

**Renal and urinary disorders:**

rarely: renal insufficiency and renal failure

Cardiovascular (CVS) adverse events, particularly arrhythmia, appeared to be correlated mostly with pre-existing CVS disease and prior therapy with cardiotoxic agents (see section 4.4). Cardiomyopathy, that may be reversible upon discontinuation of interferon alpha, has been reported rarely in patients without prior evidence of cardiac disease.

Ophthalmological disorders that have been reported rarely with alpha interferons include retinopathies (including macular oedema), retinal haemorrhages, retinal artery or vein obstruction, cotton wool spots, loss of visual acuity or visual field, optic neuritis, and papilloedema (see section 4.4).

A wide variety of autoimmune and immune-mediated disorders have been reported with alpha interferons including thyroid disorders, systemic lupus erythematosus, rheumatoid arthritis (new or aggravated), idiopathic and thrombotic thrombocytopenic purpura, vasculitis, neuropathies including mononeuropathies (see also section 4.4, **Autoimmune disorders**).

**HCV/HIV co-infected patients**

For HCV/HIV co-infected patients receiving PegIntron in combination with ribavirin, other undesirable effects (that were not reported in mono-infected patients) which have been reported in the larger studies with a frequency  $> 5\%$  were: oral candidiasis (14 %), lipodystrophy acquired (13 %), CD4 lymphocytes decreased (8 %), appetite decreased (8 %), gamma-glutamyltransferase increased (9 %), back pain (5 %), blood amylase increased (6 %), blood lactic acid increased (5 %), cytolytic hepatitis (6 %), lipase increased (6 %) and pain in limb (6 %).

*Mitochondrial toxicity:*

Mitochondrial toxicity and lactic acidosis have been reported in HIV-positive patients receiving NRTI

regimen and associated ribavirin for co-HCV infection (see section 4.4).

*Laboratory values for HCV/HIV co-infected patients:*

Although haematological toxicities of neutropenia, thrombocytopenia and anaemia occurred more frequently in HCV/HIV co-infected patients, the majority could be managed by dose modification and rarely required premature discontinuation of treatment (see section 4.4). Haematological abnormalities were more frequently reported in patients receiving PegIntron in combination with ribavirin when compared to patients receiving interferon alfa-2b in combination with ribavirin. In Study 1 (see section 5.1), decrease in absolute neutrophil count levels below 500 cells/mm<sup>3</sup> was observed in 4 % (8/194) of patients and decrease in platelets below 50,000/mm<sup>3</sup> was observed in 4 % (8/194) of patients receiving PegIntron in combination with ribavirin. Anaemia (hemoglobin < 9.4g/dl) was reported in 12% (23/194) of patients treated with PegIntron in combination with ribavirin.

*CD4 lymphocytes decrease:*

Treatment with PegIntron in combination with ribavirin was associated with decreases in absolute CD4+ cell counts within the first 4 weeks without a reduction in CD4+ cell percentage. The decrease in CD4+ cell counts was reversible upon dose reduction or cessation of therapy. The use of PegIntron in combination with ribavirin had no observable negative impact on the control of HIV viraemia during therapy or follow-up. Limited safety data (N= 25) are available in co-infected patients with CD4+ cell counts < 200/μl (see section 4.4).

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron in combination with ribavirin.

## **4.9 Overdose**

In clinical trials, cases of accidental overdose, at never more than twice the prescribed dose, were reported. There were no serious reactions. Undesirable effects resolved during continued administration of PegIntron.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Immunostimulants, Cytokines and immunomodulators, Interferons, Peginterferon alfa-2b, ATC code: L03A B10.

Recombinant interferon alfa-2b is covalently conjugated with monomethoxy polyethylene glycol at an average degree of substitution of 1 mole of polymer/mole of protein. The average molecular mass is approximately 31,300 daltons of which the protein moiety constitutes approximately 19,300.

#### **Interferon alfa-2b**

*In vitro* and *in vivo* studies suggest that the biological activity of PegIntron is derived from its interferon alfa-2b moiety.

Interferons exert their cellular activities by binding to specific membrane receptors on the cell surface. Studies with other interferons have demonstrated species specificity. However, certain monkey species, e.g., Rhesus monkeys, are susceptible to pharmacodynamic stimulation upon exposure to human type 1 interferons.

Once bound to the cell membrane, interferon initiates a complex sequence of intracellular events that include the induction of certain enzymes. It is thought that this process, at least in part, is responsible for the various cellular responses to interferon, including inhibition of virus replication in virus-infected cells, suppression of cell proliferation and such immunomodulating activities as enhancement of the phagocytic activity of macrophages and augmentation of the specific cytotoxicity of



lymphocytes for target cells. Any or all of these activities may contribute to interferon's therapeutic effects.

Recombinant interferon alfa-2b also inhibits viral replication *in vitro* and *in vivo*. Although the exact antiviral mode of action of recombinant interferon alfa-2b is unknown, it appears to alter the host cell metabolism. This action inhibits viral replication or if replication occurs, the progeny virions are unable to leave the cell.

### **PegIntron**

PegIntron pharmacodynamics were assessed in a rising single-dose trial in healthy subjects by examining changes in oral temperature, concentrations of effector proteins such as serum neopterin and 2'5'-oligoadenylate synthetase (2'5'-OAS), as well as white cell and neutrophil counts. Subjects treated with PegIntron showed mild dose-related elevations in body temperature. Following single doses of PegIntron between 0.25 and 2.0 micrograms/kg/week, serum neopterin concentration was increased in a dose-related manner. Neutrophil and white cell count reductions at the end of week 4 correlated with the dose of PegIntron.

### **PegIntron clinical trials**

Two pivotal trials have been conducted, one (C/I97-010) with PegIntron monotherapy; the other (C/I98-580) with PegIntron in combination with ribavirin. Eligible patients for these trials had chronic hepatitis C confirmed by a positive HCV-RNA polymerase chain reaction (PCR) assay (> 30 IU/ml), a liver biopsy consistent with a histological diagnosis of chronic hepatitis with no other cause for the chronic hepatitis, and abnormal serum ALT.

In the PegIntron monotherapy trial, a total of 916 naïve chronic hepatitis C patients were treated with PegIntron (0.5, 1.0 or 1.5 micrograms/kg/week) for one year with a follow-up period of six months. In addition, 303 patients received interferon alfa-2b (3 million International Units [MIU] three times a week) as a comparator. This study showed that PegIntron was superior to interferon alfa-2b (**Table 8**).

In the PegIntron combination trial, 1,530 naïve patients were treated for one year with one of the following combination regimens:

- PegIntron (1.5 micrograms/kg/week) + ribavirin (800 mg/day), (n = 511).
- PegIntron (1.5 micrograms/kg/week for one month followed by 0.5 microgram/kg/week for 11 months) + ribavirin (1,000/1,200 mg/day), (n = 514).
- Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day) (n = 505).

In this trial, the combination of PegIntron (1.5 micrograms/kg/week) and ribavirin was significantly more effective than the combination of interferon alfa-2b and ribavirin (**Table 7**), particularly in patients infected with Genotype 1 (**Table 8**). Sustained response was assessed by the response rate six months after the cessation of treatment.

HCV genotype and baseline virus load are prognostic factors which are known to affect response rates. However, response rates in this trial were shown to be dependent also on the dose of ribavirin administered in combination with PegIntron or interferon alfa-2b. In those patients that received > 10.6 mg/kg ribavirin (800 mg dose in typical 75 kg patient), regardless of genotype or viral load, response rates were significantly higher than in those patients that received ≤ 10.6 mg/kg ribavirin (**Table 8**), while response rates in patients that received > 13.2 mg/kg ribavirin were even higher.

<b>Table 7</b> Sustained virological response (% patients HCV negative)							
	<b>PegIntron monotherapy</b>				<b>PegIntron + ribavirin</b>		
<b>Treatment regimen</b>	<b>P 1.5</b>	<b>P 1.0</b>	<b>P 0.5</b>	<b>I</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>Number of patients</b>	304	297	315	303	511	514	505
<b>Response at end of treatment</b>	49 %	41 %	33 %	24 %	65 %	56 %	54 %

Sustained response	23 %*	25 %	18 %	12 %	54 %**	47 %	47 %
P 1.5	PegIntron 1.5 micrograms/kg						
P 1.0	PegIntron 1.0 microgram/kg						
P 0.5	PegIntron 0.5 microgram/kg						
I	Interferon alfa-2b 3 MIU						
P 1.5/R	PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)						
P 0.5/R	PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)						
I/R	Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)						
*	p < 0.001 P 1.5 vs. I						
**	p = 0.0143 P 1.5/R vs. I/R						

<b>Table 8</b> Sustained response rates with PegIntron + ribavirin (by ribavirin dose, genotype and viral load)				
HCV Genotype	Ribavirin dose (mg/kg)	P 1.5/R	P 0.5/R	I/R
<b>All Genotypes</b>	<b>All</b>	<b>54 %</b>	<b>47 %</b>	<b>47 %</b>
	≤ 10.6	50 %	41 %	27 %
	> 10.6	61 %	48 %	47 %
<b>Genotype 1</b>	<b>All</b>	<b>42 %</b>	<b>34 %</b>	<b>33 %</b>
	≤ 10.6	38 %	25 %	20 %
	> 10.6	48 %	34 %	34 %
Genotype 1 ≤ 600,000 IU/ml	All	73 %	51 %	45 %
	≤ 10.6	74 %	25 %	33 %
	> 10.6	71 %	52 %	45 %
Genotype 1 > 600,000 IU/ml	All	30 %	27 %	29 %
	≤ 10.6	27 %	25 %	17 %
	> 10.6	37 %	27 %	29 %
<b>Genotype 2/3</b>	<b>All</b>	<b>82 %</b>	<b>80 %</b>	<b>79 %</b>
	≤ 10.6	79 %	73 %	50 %
	> 10.6	88 %	80 %	80 %

P 1.5/R	PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)
P 0.5/R	PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)
I/R	Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)

In the PegIntron monotherapy study, the Quality of Life was generally less affected by 0.5 microgram/kg of PegIntron than by either 1.0 microgram/kg of PegIntron once weekly or 3 MIU of interferon alfa-2b three times a week.

In a separate trial, 224 patients with genotype 2 or 3 received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with ribavirin 800 mg –1,400 mg p.o. for 6 months (based on body weight, only three patients weighing > 105 kg, received the 1,400 mg dose, which has not yet been validated) (**Table 9**). Twenty-four % had bridging fibrosis or cirrhosis (Knodell 3/4).

**Table 9. Virologic Response at End of Treatment, Sustained Virologic Response and Relapse by HCV Genotype and Viral Load\***

	PegIntron 1.5 µg/kg once weekly plus Ribavirin 800-1,400 mg/day		
	End of Treatment Response	Sustained Virologic Response	Relapse
All Subjects	94 % (211/224)	81 % (182/224)	12 % (27/224)
HCV 2 ≤ 600,000 IU/ml > 600,000 IU/ml	100 % (42/42)	93 % (39/42)	7 % (3/42)
	100 % (20/20)	95 % (19/20)	5 % (1/20)
	100 % (22/22)	91 % (20/22)	9 % (2/22)
HCV 3	93 % (169/182)	79 % (143/182)	14 %

≤ 600,000 IU/ml	93 % (92/99)	86 % (85/99)	(24/166) 8 % (7/91)
> 600,000 IU/ml	93 % (77/83)	70 % (58/83)	23 % (17/75)

\* Any subject with an undetectable HCV-RNA level at the Follow-Up Week 12 visit and missing data at the Follow-Up Week 24 visit was considered a sustained responder. Any subject with missing data in and after the Follow-Up Week 12 window was considered to be a non-responder at Week 24 of follow-up.

The 6 month treatment duration in this trial was better tolerated than one year of treatment in the pivotal combination trial; for discontinuation 5 % vs. 14 %, for dose modification 18 % vs. 49 %.

In a non-comparative trial, 235 patients with genotype 1 and low viral load (< 600,000 IU/ml) received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with weight adjusted ribavirin. The overall sustained response rate after a 24-week treatment duration was 50 %. Forty-one percent of subjects (97/235) had nondetectable plasma HCV-RNA levels at Week 4 and Week 24 of therapy. In this subgroup, there was a 92 % (89/97) sustained virological response rate. The high sustained response rate in this subgroup of patients was identified in an interim analysis (n=49) and prospectively confirmed (n=48).

Limited historical data indicate that treatment for 48 weeks might be associated with a higher sustained response rate (11/11) and with a lower risk of relapse (0/11 as compared to 7/96 following 24 weeks of treatment).

#### Predictability of sustained virological response

Virological response by week 12, defined as a 2-log viral load decrease or undetectable levels of HCV-RNA has been shown to be predictive for sustained response (**Table 10**).

<b>Table 10</b> Predictability of sustained response by viral response at week 12 and genotype*				
Treatment	Genotype	Viral response at week 12	Sustained response	Negative predictive value
PegIntron 1.5 + ribavirin (> 10.6 mg/kg) 48-week treatment	1	Yes 75 % (82/110)	71 % (58/82)	----
		No 25 % (28/110)	0 % (0/28)	100 %
PegIntron 1.5 + ribavirin 800-1,400 mg 24-week treatment	2 and 3	Yes 99 % (213/215)	83 % (177/213)	----
		No 1 % (2/215)	50 % (1/2)	50 %

\* reflects patients with 12 week data available

The negative predictive value for sustained response in patients treated with PegIntron in monotherapy was 98 %.

#### HCV/HIV Co-infected patients

Two trials have been conducted in patients co-infected with HIV and HCV. The response to treatment in both of these trials is presented in **Table 11**. Study 1 (RIBAVIC; P01017) was a randomized, multicentre study which enrolled 412 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (1.5 µg/kg/week) plus ribavirin (800 mg/day) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800 mg/day) for 48 weeks with a follow-up period of 6 months. Study 2 (P02080) was a randomized, single centre study that enrolled 95 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (100 or 150 µg /week based on weight) plus ribavirin (800-1,200 mg/day based on weight) or interferon alfa-2b (3 MIU TIW) plus

ribavirin (800-1,200 mg/day based on weight). The duration of therapy was 48 weeks with a follow-up period of 6 months except for patients infected with genotypes 2 or 3 and viral load < 800,000 IU/ml (Amplicor) who were treated for 24 weeks with a 6-month follow-up period.

<b>Table 11</b> Sustained virological response based on genotype after PegIntron in combination with Ribavirin in HCV/HIV Co-infected patients						
	<b>Study 1<sup>1</sup></b>			<b>Study 2<sup>2</sup></b>		
	PegIntron (1.5 µg/kg/week) + ribavirin (800 mg)	Interferon alfa-2b (3 MIU TIW) + ribavirin (800 mg)	p value <sup>a</sup>	PegIntron (100 or 150 <sup>c</sup> µg/week) + ribavirin (800-1,200 mg) <sup>d</sup>	Interferon alfa-2b (3 MIU TIW) + ribavirin (800-1,200 mg) <sup>d</sup>	p value <sup>b</sup>
All	27 % (56/205)	20 % (41/205)	0.047	44 % (23/52)	21 % (9/43)	0.017
Genotype 1, 4	17 % (21/125)	6 % (8/129)	0.006	38 % (12/32)	7 % (2/27)	0.007
Genotype 2, 3	44 % (35/80)	43 % (33/76)	0.88	53 % (10/19)	47 % (7/15)	0.730

MIU = million international units; TIW = three times a week.

a: p value based on Cochran-Mantel Haenszel Chi square test.

b: p value based on chi-square test.

c: subjects < 75 kg received 100 µg/week PegIntron and subjects ≥ 75 kg received 150 µg/week PegIntron.

d: ribavirin dosing was 800 mg for patients < 60 kg, 1,000 mg for patients 60-75 kg, and 1,200 mg for patients > 75 kg.

<sup>1</sup>Carrato F, Bani-Sadir F, Pol S et al. JAMA 2004; 292(23): 2839-2848.

<sup>2</sup>Laguno M, Murillas J, Blanco J et al. AIDS 2004; 18(13): F27-F36.

### *Histological response*

Liver biopsies were obtained before and after treatment in Study 1 and were available for 210 of the 412 subjects (51 %). Both the Metavir score and Ishak grade decreased among subjects treated with PegIntron in combination with ribavirin. This decline was significant among responders (-0.3 for Metavir and -1.2 for Ishak) and stable (-0.1 for Metavir and -0.2 for Ishak) among non-responders. In terms of activity, about one-third of sustained responders showed improvement and none showed worsening. There was no improvement in terms of fibrosis observed in this study. Steatosis was significantly improved in patients infected with HCV Genotype 3.

## **5.2 Pharmacokinetic properties**

PegIntron is a well characterized polyethylene glycol-modified (“pegylated”) derivative of interferon alfa-2b and is predominantly composed of monopegylated species. The plasma half-life of PegIntron is prolonged compared with non-pegylated interferon alfa-2b. PegIntron has a potential to depegylate to free interferon alfa-2b. The biologic activity of the pegylated isomers is qualitatively similar, but weaker than free interferon alfa-2b.

Following subcutaneous administration, maximal serum concentrations occur between 15-44 hours post-dose, and are sustained for up to 48-72 hours post-dose.

PegIntron C<sub>max</sub> and AUC measurements increase in a dose-related manner. Mean apparent volume of distribution is 0.99 l/kg.

Upon multiple dosing, there is an accumulation of immunoreactive interferons. There is, however, only a modest increase in biologic activity as measured by a bioassay.

Mean (SD) PegIntron elimination half-life is approximately 40 hours (13.3 hours), with apparent clearance of 22.0 ml/hr·kg. The mechanisms involved in clearance of interferons in man have not yet been

fully elucidated. However, renal elimination may account for a minority (approximately 30 %) of PegIntron apparent clearance.

**Renal function:** Renal clearance appears to account for 30 % of total clearance of PegIntron. In a single dose study (1.0 microgram/kg) in patients with impaired renal function,  $C_{max}$ , AUC, and half-life increased in relation to the degree of renal impairment.

Following multiple dosing of PegIntron (1.0 microgram/kg subcutaneously administered every week for four weeks) the clearance of PegIntron is reduced by a mean of 17 % in patients with moderate renal impairment (creatinine clearance 30-49 ml/minute) and by a mean of 44 % in patients with severe renal impairment (creatinine clearance 15-29 ml/minute) compared to subjects with normal renal function. Based on single dose data, clearance was similar in patients with severe renal impairment not on dialysis and in patients who were receiving hemodialysis. The dose of PegIntron for monotherapy should be reduced in patients with moderate or severe renal impairment (see sections 4.2 and 4.4). Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see section 4.3).

Because of marked inter-subject variability in interferon pharmacokinetics, it is recommended that patients with severe renal impairment be closely monitored during treatment with PegIntron (see section 4.2)

**Hepatic function:** The pharmacokinetics of PegIntron have not been evaluated in patients with severe hepatic dysfunction.

**Elderly patients  $\geq 65$  years of age:** The pharmacokinetics of PegIntron following a single subcutaneous dose of 1.0 microgram/kg were not affected by age. The data suggest that no alteration in PegIntron dosage is necessary based on advancing age.

**Patients under the age of 18 years:** Specific pharmacokinetic evaluations have not been performed on these patients. PegIntron is indicated for the treatment of chronic hepatitis C only in patients 18 years of age or older.

**Interferon neutralising factors:** Interferon neutralising factor assays were performed on serum samples of patients who received PegIntron in the clinical trial. Interferon neutralising factors are antibodies which neutralise the antiviral activity of interferon. The clinical incidence of neutralising factors in patients who received PegIntron 0.5 micrograms/kg is 1.1 %.

### 5.3 Preclinical safety data

**PegIntron:** Adverse events not observed in clinical trials were not seen in toxicity studies in monkeys. These studies were limited to four weeks due to the appearance of anti-interferon antibodies in most monkeys.

Reproduction studies of PegIntron have not been performed. Interferon alfa-2b has been shown to be an abortifacient in primates. PegIntron is likely to also cause this effect. Effects on fertility have not been determined. It is not known whether the components of this medicinal product are excreted into experimental animal or human milk (see section 4.6 for relevant human data on pregnancy and lactation). PegIntron showed no genotoxic potential.

The relative non-toxicity of monomethoxy-polyethylene glycol (mPEG), which is liberated from PegIntron by metabolism *in vivo* has been demonstrated in preclinical acute and subchronic toxicity studies in rodents and monkeys, standard embryo-foetal development studies and in *in vitro* mutagenicity assays.

**PegIntron plus ribavirin:** When used in combination with ribavirin, PegIntron did not cause any effects not previously seen with either active substance alone. The major treatment-related change was

a reversible, mild to moderate anaemia, the severity of which was greater than that produced by either active substance alone.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Powder for solution for injection:

Disodium phosphate, anhydrous,  
Sodium dihydrogen phosphate dihydrate,  
Sucrose,  
Polysorbate 80.

Solvent for parenteral use:

Water for injections.

Deliverable volume from pen = 0.5 ml.

### **6.2 Incompatibilities**

This medicinal product should only be reconstituted with the solvent provided (see section 6.6). In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

3 years

After reconstitution:

- Chemical and physical in-use stability has been demonstrated for 24 hours at 2°C - 8°C.
- From a microbiological point of view, the product is to be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C - 8°C.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C - 8°C). Do not freeze.

### **6.5 Nature and contents of container**

The powder and solvent are both contained in a two-chamber cartridge, Type I flint glass, separated by a bromobutyl rubber plunger. The cartridge is sealed at one end with a polypropylene cap containing a bromobutyl rubber liner and at the other end by a bromobutyl rubber plunger.

PegIntron 50 micrograms is supplied as:

- 1 pen containing powder and solvent for solution for injection, 1 injection needle and 2 cleansing swabs;
- 4 pens containing powder and solvent for solution for injection, 4 injection needles and 8 cleansing swabs;
- 6 pens containing powder and solvent for solution for injection, 6 injection needles and 12 cleansing swabs;
- 12 pens containing powder and solvent for solution for injection, 12 injection needles and 24 cleansing swabs.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

PegIntron pre-filled pen contains a powder of peginterferon alfa-2b and a solvent for solution at a strength of 50 micrograms for single use. Each pen is reconstituted with the solvent provided in the two-chamber cartridge (water for injections) for administration of up to 0.5 ml of solution. A small volume is lost during preparation of PegIntron for injection when the dose is measured and injected. Therefore, each pen contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection. The reconstituted solution has a concentration of 50 micrograms in 0.5 ml.

PegIntron is injected subcutaneously after reconstituting the powder as instructed, attaching an injection needle and setting the prescribed dose. A complete and illustrated set of instructions is provided in the Annex to the Package Leaflet.

Remove PegIntron pre-filled pen from the refrigerator before administration to allow the solvent to reach room temperature (not more than 25°C).

As for all parenteral medicinal products, inspect visually the reconstituted solution prior to administration. The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. After administering the dose, discard the PegIntron pre-filled pen and any unused solution contained in it.

## **7. MARKETING AUTHORISATION HOLDER**

SP Europe  
73, rue de Stalle  
B-1180 Bruxelles  
Belgium

## **8. MARKETING AUTHORISATION NUMBERS**

EU/1/00/131/031  
EU/1/00/131/032  
EU/1/00/131/033  
EU/1/00/131/034

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 6 February 2002  
Date of last renewal: 25 May 2005

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.emea.europa.eu/>

## **1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 80 micrograms, powder and solvent for solution for injection in pre-filled pen

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each pre-filled pen of PegIntron 80 micrograms contains a sufficient amount of peginterferon alfa-2b as measured on a protein basis in a powder for solution for injection, and the corresponding amount of solvent, to provide 80 micrograms in 0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

The active substance is a covalent conjugate of recombinant interferon alfa-2b\* with monomethoxy polyethylene glycol. The potency of this product should not be compared to that of another pegylated or non-pegylated protein of the same therapeutic class. For more information, see section 5.1.

\*produced by rDNA technology in *E.coli* cells harbouring a genetically engineered plasmid hybrid encompassing an interferon alfa-2b gene from human leukocytes

For a full list of excipients, see section 6.1. PegIntron contains 40 mg of sucrose per 0.5 ml.

## **3. PHARMACEUTICAL FORM**

Powder and solvent for solution for injection in pre-filled pen

White powder.

Clear and colourless solvent.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

PegIntron is indicated for the treatment of adult patients with chronic hepatitis C who have elevated transaminases without liver decompensation and who are positive for serum HCV-RNA or anti-HCV, including naïve patients with clinically stable HIV co-infection (see section 4.4).

The best way to use PegIntron in this indication is in combination with ribavirin.

This combination is indicated in naïve patients including patients with clinically stable HIV co-infection. This combination of PegIntron with ribavirin is also indicated in patients who have previously responded (with normalisation of ALT at the end of treatment) to interferon alpha monotherapy but who have subsequently relapsed.

Interferon monotherapy, including PegIntron, is indicated mainly in case of intolerance or contraindication to ribavirin.

Please refer also to the ribavirin Summary of Product Characteristics (SPC) when PegIntron is to be used in combination with ribavirin.

### **4.2 Posology and method of administration**

Treatment should be initiated and monitored only by a physician experienced in the management of patients with hepatitis C.

#### **Dose to be administered**

PegIntron should be administered as a once weekly subcutaneous injection. The dose administered depends on whether it is used in combination with ribavirin or as monotherapy.



## Combination therapy

PegIntron 1.5 micrograms/kg/week in combination with ribavirin capsules.

The intended dose of 1.5 µg/kg of PegIntron to be used in combination with ribavirin may be delivered in weight categories with the pen/vial strengths according to **Table 1**. Ribavirin capsules are to be administered orally each day in two divided doses with food (morning and evening).

**Table 1 - Dosing for Combination Therapy**

Body Weight (kg)	PegIntron		Ribavirin Capsules	
	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Total Daily Dose (mg)	Number of Capsules (200 mg)
< 40	50	0.5	800	4 <sup>a</sup>
40-50	80	0.4	800	4 <sup>a</sup>
51-64	80	0.5	800	4 <sup>a</sup>
65-75	100	0.5	1,000	5 <sup>b</sup>
76-85	120	0.5	1,000	5 <sup>b</sup>
> 85	150	0.5	1,200	6 <sup>c</sup>

a: 2 morning, 2 evening

b: 2 morning, 3 evening

c: 3 morning, 3 evening

## Duration of treatment

**Predictability of sustained virological response:** Patients infected with virus genotype 1 who fail to achieve virological response at Week 12 are highly unlikely to become sustained virological responders (see also section 5.1).

- **Genotype 1:** For patients who exhibit virological response at week 12, treatment should be continued for another nine month period (i.e., a total of 48 weeks).  
In the subset of patients with genotype 1 infection and low viral load (< 600,000 IU/ml) who become HCV-RNA negative at treatment week 4 and remain HCV-RNA negative at week 24, the treatment could either be stopped after this 24 week treatment course or pursued for an additional 24 weeks (i.e. overall 48 weeks treatment duration). However, an overall 24 weeks treatment duration may be associated with a higher risk of relapse than a 48 weeks treatment duration (see section 5.1).
- **Genotypes 2 or 3:** It is recommended that all patients be treated for 24 weeks, except for HCV/HIV co-infected patients who should receive 48 weeks of treatment.
- **Genotype 4:** In general, patients infected with genotype 4 are considered harder to treat and limited study data (n=66) indicate they are compatible with a duration of treatment as for genotype 1.

## HCV/HIV Co-infection

The recommended duration of dosing for HCV/HIV co-infected patients is 48 weeks, regardless of genotype.

### *Predictability of response and non-response in HCV/HIV Co-infection*

Early virological response by Week 12, defined as a 2 log viral load decrease or undetectable levels of HCV-RNA, has been shown to be predictive for sustained response. The negative predictive value for sustained response in HCV/HIV co-infected patients treated with PegIntron in combination with ribavirin was 99 % (67/68; Study 1) (see section 5.1). A positive predictive value of 50 % (52/104; Study 1) was observed for HCV/HIV co-infected patients receiving combination therapy.

## PegIntron monotherapy

As monotherapy the PegIntron regimen is 0.5 or 1.0 microgram/kg/week. The lowest vial or pen strength available is 50 µg/0.5 ml; therefore for patients prescribed 0.5 µg/kg/week, doses must be adjusted by volume as shown in **Table 2**. For the 1.0 µg/kg dose, similar volume adjustments can be made or alternate vial strengths can be used as shown in **Table 2**. PegIntron monotherapy was not studied in HCV/HIV co-infected patients.

**Table 2- Monotherapy Dosing**

	0.5 µg/kg		1.0 µg/kg	
Body Weight (kg)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)
30-35	50*	0.15	50	0.3
36-45	50*	0.2	50	0.4
46-56	50*	0.25	50	0.5
57-72	50	0.3	80	0.4
73-88	50	0.4	80	0.5
89-106	50	0.5	100	0.5
> 106**	80	0.4	120	0.5
* Must use vial. Minimum delivery for pen is 0.3 ml.				
** For patients > 120 kg, use 80 µg/0.5 ml vial				

### Duration of treatment

For patients who exhibit virological response at Week 12, treatment should be continued for at least another three-month period (i.e., a total of six months). The decision to extend therapy to one year of treatment should be based on prognostic factors (e.g., genotype, age > 40 years, male gender, bridging fibrosis).

### Dose modification for all patients

If severe adverse reactions or laboratory abnormalities develop during treatment with PegIntron monotherapy or PegIntron in combination with ribavirin, modify the dosages of each product as appropriate, until the adverse reactions abate. As adherence might be of importance for outcome of therapy, the dose should be kept as close as possible to the recommended standard dose. Guidelines were developed in clinical trials for dose modification.

### Combination Therapy Dose Reduction Guidelines

<b>Table 2a</b> Dose modification guidelines for combination therapy (with ribavirin)			
Laboratory values	Reduce only ribavirin dose to 600 mg/day* if:	Reduce only PegIntron dose to one-half dose if:	Discontinue combination therapy if:
Haemoglobin	< 10 g/dl	-	< 8.5 g/dl
Haemoglobin in: Patients with history of stable cardiac disease	≥ 2 g/dl decrease in haemoglobin during any four week period during treatment (permanent dose reduction)		< 12 g/dl after four weeks of dose reduction
White blood cells	-	< 1.5 x 10 <sup>9</sup> /l	< 1.0 x 10 <sup>9</sup> /l
Neutrophils	-	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	-	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l
Bilirubin – direct	-	-	2.5 x ULN**
Bilirubin – indirect	> 5 mg/dl	-	> 4 mg/dl

			(for > 4 weeks)
Creatinine	-	-	> 2.0 mg/dl
ALT/AST	-	-	2 x baseline and > 10 x ULN**

\* Patients whose dose of ribavirin is reduced to 600 mg daily receive one 200 mg capsule in the morning and two 200 mg capsules in the evening.

\*\* Upper limit of normal

Dose reduction of PegIntron may be accomplished by either reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 2b**.

<b>Table 2b – Reduced PegIntron Dosing for Combination Therapy</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5 ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
< 40	25	50*	0.25	25
40-50	32	50	0.3	30
51-64	40	50	0.4	40
65-75	50	50	0.5	50
76-85	60	80	0.4	64
> 85	75	100	0.4	80
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### **PegIntron Monotherapy Dose Reduction Guidelines**

Dose modification guidelines for patients who use PegIntron monotherapy are shown in **Table 3a**.

<b>Table 3a Dose modification guidelines for PegIntron monotherapy</b>		
<b>Laboratory values</b>	<b>Reduce PegIntron to one-half dose if:</b>	<b>Discontinue PegIntron if:</b>
Neutrophils	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l

Dose reduction for patients who use 0.5 µg/kg PegIntron monotherapy must be accomplished by reducing the prescribed volume by one-half. The 50 µg/0.5 ml vial must be used if necessary since the pen can only deliver a minimum volume of 0.3 ml.

For patients who use 1.0 µg/kg PegIntron monotherapy, dose reduction may be accomplished by reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 3b**.

<b>Table 3b – Reduced PegIntron Dose for the 1.0 µg/kg Monotherapy Regimen</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
30-35	15	50*	0.15	15

36-45	20	50*	0.20	20
46-56	25	50*	0.25	25
57-72	32	50	0.3	30
73-89	40	50	0.4	40
90-106	50	50	0.5	50
> 106	60	80	0.4	64
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### Special populations

**Use in renal impairment:** *Monotherapy:* PegIntron should be used with caution in patients with moderate to severe renal impairment. In patients with moderate renal dysfunction (creatinine clearance 30-50 ml/minute), the starting dose of PegIntron should be reduced by 25 %. Patients with severe renal dysfunction (creatinine clearance 15-29 ml/minute) should have the starting dose of PegIntron reduced by 50 %. Data are not available for the use of PegIntron in patients with creatinine clearance < 15 ml/minute (see section 5.2). Patients with severe renal impairment, including those on hemodialysis, should be closely monitored. If renal function decreases during treatment, PegIntron therapy should be discontinued.

*Combination therapy:* Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see ribavirin SPC). When administered in combination with ribavirin, subjects with impaired renal function should be more carefully monitored with respect to the development of anaemia.

**Use in hepatic impairment:** The safety and efficacy of PegIntron therapy has not been evaluated in patients with severe hepatic dysfunction, therefore PegIntron must not be used for these patients.

**Use in the elderly (≥ 65 years of age):** There are no apparent age-related effects on the pharmacokinetics of PegIntron. Data from elderly patients treated with a single dose of PegIntron suggest no alteration in PegIntron dose is necessary based on age (see section 5.2).

**Use in patients under the age of 18 years:** There is no experience in children (see section 5.2).

### 4.3 Contraindications

- Hypersensitivity to the active substance or to any interferon or to any of the excipients;
- A history of severe pre-existing cardiac disease, including unstable or uncontrolled cardiac disease in the previous six months (see section 4.4);
- Severe, debilitating medical conditions;
- Autoimmune hepatitis or a history of autoimmune disease;
- Severe hepatic dysfunction or decompensated cirrhosis of the liver;
- Pre-existing thyroid disease unless it can be controlled with conventional treatment;
- Epilepsy and/or compromised central nervous system (CNS) function;
- Initiation of PegIntron is contraindicated in HCV/HIV patients with cirrhosis and a Child-Pugh score ≥ 6.

*Combination therapy with ribavirin:* Also see ribavirin Summary of the Product Characteristics (SPC) if PegIntron is to be administered in combination with ribavirin in patients with chronic hepatitis C.

### 4.4 Special warnings and precautions for use

**Psychiatric and Central Nervous System (CNS):** Severe CNS effects, particularly depression, suicidal ideation and attempted suicide have been observed in some patients during PegIntron therapy, and even after treatment discontinuation mainly during the 6-month follow-up period. Other CNS effects including

aggressive behaviour (sometimes directed against others), confusion and alterations of mental status have been observed with alpha interferons. Patients should be closely monitored for any signs or symptoms of psychiatric disorders. If such symptoms appear, the potential seriousness of these undesirable effects must be borne in mind by the prescribing physician and the need for adequate therapeutic management should be considered. If psychiatric symptoms persist or worsen, or suicidal ideation is identified, it is recommended that treatment with PegIntron be discontinued, and the patient followed, with psychiatric intervention as appropriate.

*Patients with existence of, or history of severe psychiatric conditions:* If treatment with peginterferon alfa-2b is judged necessary in patients with existence or history of severe psychiatric conditions, this should only be initiated after having ensured appropriate individualised diagnostic and therapeutic management of the psychiatric condition.

There is no experience with PegIntron in combination with ribavirin in patients who have relapsed after interferon alpha + ribavirin therapy.

All patients in the chronic hepatitis C studies had a liver biopsy before inclusion, but in certain cases (i.e. patients with genotype 2 and 3), treatment may be possible without histological confirmation. Current treatment guidelines should be consulted as to whether a liver biopsy is needed prior to commencing treatment.

**Acute hypersensitivity:** Acute hypersensitivity reactions (e.g., urticaria, angioedema, bronchoconstriction, anaphylaxis) have been observed rarely during interferon alfa-2b therapy. If such a reaction develops during treatment with PegIntron, discontinue treatment and institute appropriate medical therapy immediately. Transient rashes do not necessitate interruption of treatment.

**Cardiovascular system:** As with interferon alfa-2b, patients with a history of congestive heart failure, myocardial infarction and/or previous or current arrhythmic disorders, receiving PegIntron therapy require close monitoring. It is recommended that patients who have pre-existing cardiac abnormalities have electrocardiograms taken prior to and during the course of treatment. Cardiac arrhythmias (primarily supraventricular) usually respond to conventional therapy but may require discontinuation of PegIntron therapy.

More significant obtundation and coma, including cases of encephalopathy, have been observed in some patients, usually elderly, treated at higher doses for oncology indications. While these effects are generally reversible, in a few patients full resolution took up to three weeks. Very rarely, seizures have occurred with high doses of interferon alpha.

**Liver function:** As with all interferons, discontinue treatment with PegIntron in patients who develop prolongation of coagulation markers which might indicate liver decompensation.

**Fever:** While fever may be associated with the flu-like syndrome reported commonly during interferon therapy, other causes of persistent fever must be ruled out.

**Hydration:** Adequate hydration must be maintained in patients undergoing PegIntron therapy since hypotension related to fluid depletion has been seen in some patients treated with alpha interferons. Fluid replacement may be necessary.

**Pulmonary changes:** Pulmonary infiltrates, pneumonitis, and pneumonia, occasionally resulting in fatality, have been observed rarely in interferon alpha treated patients. Any patient developing fever, cough, dyspnea or other respiratory symptoms must have a chest X-ray taken. If the chest X-ray shows pulmonary infiltrates or there is evidence of pulmonary function impairment, the patient is to be monitored closely, and, if appropriate, discontinue interferon alpha. Prompt discontinuation of interferon alpha administration and treatment with corticosteroids appear to be associated with resolution of pulmonary adverse events.

**Autoimmune disease:** The development of auto-antibodies and autoimmune disorders has been reported during treatment with alpha interferons. Patients predisposed to the development of

autoimmune disorders may be at increased risk. Patients with signs or symptoms compatible with autoimmune disorders should be evaluated carefully, and the benefit-risk of continued interferon therapy should be reassessed (see also section 4.4 Thyroid changes and 4.8).

**Ocular changes:** Ophthalmologic disorders, including retinal haemorrhages, cotton wool spots, and retinal artery or vein obstruction have been reported in rare instances after treatment with alpha interferons (see section 4.8). All patients should have a baseline eye examination. Any patient complaining of ocular symptoms, including loss of visual acuity or visual field must have a prompt and complete eye examination. Periodic visual examinations are recommended during PegIntron therapy, particularly in patients with disorders that may be associated with retinopathy, such as diabetes mellitus or hypertension. Discontinuation of PegIntron should be considered in patients who develop new or worsening ophthalmological disorders.

**Thyroid changes:** Infrequently, patients treated for chronic hepatitis C with interferon alpha have developed thyroid abnormalities, either hypothyroidism or hyperthyroidism. Determine thyroid stimulating hormone (TSH) levels if, during the course of therapy, a patient develops symptoms consistent with possible thyroid dysfunction. In the presence of thyroid dysfunction, PegIntron treatment may be continued if TSH levels can be maintained in the normal range by medication.

**Metabolic disturbances:** Hypertriglyceridemia and aggravation of hypertriglyceridemia, sometimes severe, have been observed. Monitoring of lipid levels is, therefore, recommended.

### **HCV/HIV Co-infection**

Mitochondrial toxicity and lactic acidosis:

Patients co-infected with HIV and receiving Highly Active Anti-Retroviral Therapy (HAART) may be at increased risk of developing lactic acidosis. Caution should be used when adding PegIntron and ribavirin to HAART therapy (see ribavirin SPC).

Moreover, patients treated with PegIntron and ribavirin combination therapy and zidovudine are at increased risk of developing anaemia.

Hepatic decompensation in HCV/HIV co-infected patients with advanced cirrhosis:

Co-infected patients with advanced cirrhosis receiving HAART may be at increased risk of hepatic decompensation and death. Adding treatment with alfa interferons alone or in combination with ribavirin may increase the risk in this patient subset. Other baseline factors in co-infected patients that may be associated with a higher risk of hepatic decompensation include treatment with didanosine and elevated bilirubin serum concentration.

Co-infected patients receiving both antiretroviral (ARV) and anti-hepatitis treatment should be closely monitored, assessing their Child-Pugh score during treatment. Patients progressing to hepatic decompensation should have their anti-hepatitis treatment immediately discontinued and the ARV treatment reassessed.

Haematological abnormalities in HCV/HIV co-infected patients:

HCV/HIV co-infected patients receiving peginterferon alfa-2b/ribavirin treatment and HAART may be at increased risk to develop haematological abnormalities (as neutropenia, thrombocytopenia and anaemia) compared to HCV mono-infected patients. Although, the majority of them could be managed by dose reduction, close monitoring of haematological parameters should be undertaken in this population of patients (see section 4.2 and below “Laboratory tests” and section 4.8).

Patients with low CD4 counts:

In patients co-infected with HCV/HIV, limited efficacy and safety data (N = 25) are available in subjects with CD4 counts less than 200 cells/ $\mu$ l. Caution is therefore warranted in the treatment of patients with low CD4 counts.

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron and ribavirin.

**Dental and periodontal disorders:** Dental and periodontal disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and mucous membranes of the mouth during long-term treatment with the combination of PegIntron and ribavirin. Patients should brush their teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If this reaction occurs, they should be advised to rinse out their mouth thoroughly afterwards.

**Organ transplant recipients:** The safety and efficacy of PegIntron alone or in combination with ribavirin for the treatment of hepatitis C in liver or other organ transplant recipients have not been studied. Preliminary data indicates that interferon alpha therapy may be associated with an increased rate of kidney graft rejection. Liver graft rejection has also been reported.

**Other:** Due to reports of interferon alpha exacerbating pre-existing psoriatic disease and sarcoidosis, use of PegIntron in patients with psoriasis or sarcoidosis is recommended only if the potential benefit justifies the potential risk.

**Laboratory tests:** Standard haematologic tests, blood chemistry and a test of thyroid function must be conducted in all patients prior to initiating therapy. Acceptable baseline values that may be considered as a guideline prior to initiation of PegIntron therapy are:

- Platelets  $\geq 100,000/\text{mm}^3$
- Neutrophil count  $\geq 1,500/\text{mm}^3$
- TSH level must be within normal limits

Laboratory evaluations are to be conducted at weeks 2 and 4 of therapy, and periodically thereafter as clinically appropriate.

**Important information about some of the ingredients of PegIntron:**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

Patients with rare hereditary problems of fructose intolerance, glucose galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Results from a multiple-dose probe study assessing P450 substrates in chronic hepatitis C patients receiving once weekly PegIntron (1.5  $\mu\text{g/kg}$ ) for 4 weeks demonstrated an increase in activity of CYP2D6 and CYP2C8/9. No change in activity of CYP1A2, CYP3A4, or N-acetyltransferase was observed.

Caution should be used when administering peginterferon alfa-2b with medicines metabolised by CYP2D6 and CYP2C8/9, especially those with narrow therapeutic window, such as warfarin and phenytoin (CYP2C9) and flecainide (CYP2D6).

These findings may partly relate to improved metabolic capacity due to reduced hepatic inflammation in patients undergoing treatment with PegIntron. Caution is therefore advised when PegIntron treatment is initiated for chronic hepatitis in patients treated with medication with a narrow therapeutic window and sensitive to mild metabolic impairment of the liver.

No pharmacokinetic interactions were noted between PegIntron and ribavirin in a multiple-dose pharmacokinetic study.

**Methadone:** In patients with chronic hepatitis C that were on stable methadone maintenance therapy and naïve to peginterferon alfa-2b, addition of 1.5 microgram/kg/week of PegIntron subcutaneously for 4 weeks increased R-methadone AUC by approximately 15 % (95 % CI for AUC ratio estimate

103 – 128 %). The clinical significance of this finding is unknown; however, patients should be monitored for signs and symptoms of increased sedative effect, as well as respiratory depression. Especially in patients on a high dose of methadone, the risk for QTc prolongation should be considered.

#### *HCV/HIV Co-infection*

Nucleoside analogs: ribavirin has been shown *in vitro* to inhibit phosphorylation of zidovudine and stavudine. The clinical significance of these finding is unknown. However, these *in vitro* findings raise the possibility that concurrent use of ribavirin with either zidovudine or stavudine might lead to increased HIV plasma viraemia. Therefore, it is recommended that plasma HIV RNA levels be closely monitored in patients treated with ribavirin concurrently with either of these two agents. If HIV RNA levels increase, the use of ribavirin concomitantly with reverse transcriptase inhibitors must be reviewed (see ribavirin SPC).

Use of nucleoside analogs, alone or in combination with other nucleosides, has resulted in lactic acidosis. Pharmacologically, ribavirin increases phosphorylated metabolites of purine nucleosides *in vitro*. This activity could potentiate the risk of lactic acidosis induced by purine nucleoside analogs (e.g. didanosine or abacavir). Co-administration of ribavirin and didanosine is not recommended. Reports of mitochondrial toxicity, in particular lactic acidosis and pancreatitis, of which some fatal, have been reported (see ribavirin SPC).

## **4.6 Pregnancy and lactation**

PegIntron is recommended for use in fertile women only when they are using effective contraception during the treatment.

There are no adequate data from the use of interferon alfa-2b in pregnant women.. Studies in animals have shown reproductive toxicity (see section 5.3). Interferon alfa-2b has been shown to be abortifacient in primates. PegIntron is likely to also cause this effect.

The potential risk in humans is unknown. PegIntron is to be used during pregnancy only if the potential benefit justifies the potential risk to the foetus

**Lactation:** It is not known whether the components of this medicinal product are excreted in human milk. Because of the potential for adverse reactions in breast-fed infants, breast-feeding should be discontinued prior to initiation of treatment.

#### *Combination therapy with ribavirin:*

Ribavirin causes serious birth defects when administered during pregnancy. Ribavirin therapy is contraindicated in women who are pregnant. Extreme care must be taken to avoid pregnancy in female patients or in partners of male patients taking PegIntron in combination with ribavirin. Females of childbearing potential and their partners must each use an effective contraceptive during treatment and for 4 months after treatment has been concluded. Male patients and their female partners must each use an effective contraceptive during treatment and for 7 months after treatment has been concluded (see Ribavirin SPC).

## **4.7 Effects on ability to drive and use machines**

Patients who develop fatigue, somnolence or confusion during treatment with PegIntron are cautioned to avoid driving or operating machinery.

## **4.8 Undesirable effects**

The safety of PegIntron is evaluated from data from two clinical trials: one with PegIntron monotherapy, one with PegIntron in combination with ribavirin. In both cases, patients were treated for one year.



**Table 4** describes the regimens and patient exposure for one year of treatment in patients with no previous exposure to interferon (interferon-naïve patients). Because of a significant overlap in the pattern of undesirable effects with PegIntron monotherapy, groups of patients have been brought together in **Table 5** to show the pattern of reported effects for all monotherapy groups.

<b>Table 4</b> Regimens and patient exposure		
<b>Treatment</b>	<b>Regimen</b>	<b>Number of patients treated for one year</b>
PegIntron + ribavirin	PegIntron (1.5 micrograms/kg/week) + ribavirin (> 10.6 mg/kg/day)	188
Interferon alfa-2b + ribavirin	Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day)	505
PegIntron monotherapy	PegIntron (0.5 microgram/kg/week)	315
	PegIntron (1.0 microgram/kg/week)	297
	PegIntron (1.5 micrograms/kg/week)	304

<b>Table 5</b> Undesirable effects very commonly reported in clinical trials > 10 % of patients in PegIntron + ribavirin group Very common (> 1/10)			
	<b>PegIntron + ribavirin</b>	<b>Interferon alfa- 2b + ribavirin</b>	<b>PegIntron monotherapy</b>
<b>Infections and infestations</b>			
Infection viral	10 %	5 %	4-5 %
<b>Metabolism and nutrition disorders</b>			
Weight decrease	30 %	19 %	8-18 %
<b>Psychiatric disorders</b>			
Depression	34 %	32 %	26 %
Irritability	32 %	34 %	19 %
Insomnia	37 %	41 %	16-19 %
Anxiety	14 %	14 %	8 %
Concentration impaired	18 %	21 %	9-10 %
Emotional lability	11 %	10 %	5 %
<b>Nervous system disorders</b>			
Headache	58 %	57 %	57-63 %
Mouth dry	10 %	8 %	4-8 %
<b>Respiratory, thoracic and mediastinal disorders</b>			
Pharyngitis	10 %	7 %	3 %
Coughing	14 %	11 %	4 %
Dyspnea	26 %	22 %	5 %
<b>Gastrointestinal disorders</b>			
Nausea	43 %	31 %	20-23 %
Anorexia	35 %	26 %	10-25 %
Diarrhoea	20 %	13 %	14-17 %
Abdominal pain	12 %	9 %	11 %
Vomiting	16 %	10 %	4-7 %
<b>Skin and subcutaneous tissue disorders</b>			
Alopecia	45 %	32 %	20-34 %
Pruritus	27 %	27 %	7-9 %
Skin dry	23 %	21 %	6-9 %
Rash	21 %	21 %	5-7 %
<b>Musculoskeletal and connective tissue disorders</b>			
Myalgia	49 %	49 %	46-60 %
Arthralgia	31 %	26 %	23-28 %
Musculoskeletal pain	15 %	11 %	11-13 %
<b>General disorders and administration site conditions</b>			
Injection site inflammation	20 %	17 %	39-44 %
Injection site reaction	54 %	36 %	7-9 %
Fatigue	56 %	59 %	43 %
Rigors	42 %	40 %	33-43 %
Fever	39 %	32 %	29-43 %
Flu-like symptoms	21 %	23 %	18-25 %
Asthenia	28 %	17 %	12-14 %
Dizziness	17 %	16 %	7-12 %

<b>Table 6</b> Undesirable effects commonly reported in clinical trials in patients treated with PegIntron + ribavirin or PegIntron monotherapy Common (> 1/100, < 1/10)		
<b>Body system</b>	<b>5-10%</b>	<b>1-&lt;5%</b>
Infections and infestations		otitis media, fungal infection, bacterial infection, herpes simplex
Blood and lymphatic system disorders	Anaemia, leukopaenia	Thrombocytopenia, lymphadenopathy
Endocrine disorders	hypothyroidism	hyperthyroidism
Metabolism and nutrition disorders		hyperuricemia, hypocalcemia, thirst
Psychiatric disorders	agitation, nervousness	aggressive behaviour, somnolence, behavior disorder, apathy, appetite increased, sleep disorder, dreaming abnormal, decreased libido
Nervous system disorders	paresthesia, increased sweating	hypoesthesia, hyperaesthesia, hypertonia, confusion, tremor, vertigo, migraine, ataxia, neuralgia
Eye disorders		blurred vision, conjunctivitis, lacrimal gland disorder, eye pain
Cardiac disorders		tachycardia, palpitation
Vascular disorders		hypotension, hypertension, syncope, flushing
Ear and labyrinth disorders		tinnitus, hearing impairment/loss
Respiratory, thoracic and mediastinal disorders		nonproductive cough, rhinitis, sinusitis, bronchitis, respiratory disorder, nasal congestion, rhinorrhea, dysphonia, epistaxis
Gastrointestinal disorders	dyspepsia	constipation, taste perversion, loose stools, stomatitis, ulcerative stomatitis, gingival bleeding, glossitis, flatulence, hemorrhoids, gastroesophageal reflux, gingivitis, dehydration
Hepatobiliary disorders		hepatomegaly, hyperbilirubinemia
Skin and subcutaneous tissue disorders		erythematous rash, eczema, photosensitivity reaction, maculopapular rash, abnormal hair texture, acne, dermatitis, furunculosis, nail disorder, psoriasis, urticaria, erythema, face or peripheral oedema
Musculoskeletal and connective tissue disorders		arthritis
Renal and urinary disorders		micturition frequency, urine abnormal
Reproductive system and breast disorders	menstrual disorder, menorrhagia	ovarian disorder, vaginal disorder, sexual dysfunction (not specified), impotence, breast pain, amenorrhoea, prostatitis
General disorders and administration site conditions	RUQ pain, malaise, chest pain	injection site pain

Most cases of neutropaenia and thrombocytopaenia were mild (WHO grades 1 or 2). There were some cases of more severe neutropenia in patients treated with the recommended doses of PegIntron in combination with ribavirin (WHO grade 3: 39 of 186 [21 %]; and WHO grade 4: 13 of 186 [7 %]).

In a clinical trial, approximately 1.2 % of patients treated with PegIntron or interferon alfa-2b in combination with ribavirin reported life-threatening psychiatric events during treatment. These events included suicidal ideation and attempted suicide (see section 4.4). Following marketing, psychosis and hallucination have been reported rarely.

Rarely ( $> 1/10,000$ ,  $< 1/1,000$ ) or very rarely ( $< 1/10,000$ ) reported events with interferon alfa-2b, including PegIntron, include:

**Blood and lymphatic system disorders:**

Very rarely interferon alfa-2b or PegIntron used alone or in combination with ribavirin may be associated with aplastic anaemia. Pure red cell aplasia has been reported.

**Immune system disorders:**

very rarely: sarcoidosis or exacerbation of sarcoidosis

**Endocrine disorders:**

rarely: diabetes

**Nervous system disorders:**

rarely: seizure, peripheral neuropathy

very rarely: cerebrovascular ischaemia, cerebrovascular haemorrhage, encephalopathy

**Cardiac disorders:**

rarely: arrhythmia

very rarely: cardiac ischaemia, myocardial infarction

**Respiratory, thoracic and mediastinal disorders:**

very rarely: interstitial lung disease

**Gastrointestinal disorders:**

rarely: pancreatitis

very rarely: ulcerative and ischaemic colitis

**Skin and subcutaneous tissue disorders:**

very rarely: erythema multiforme, Stevens Johnson syndrome, toxic epidermal necrolysis, injection site necrosis

**Musculoskeletal and connective tissue disorders:**

rarely: rhabdomyolysis, myositis

**Renal and urinary disorders:**

rarely: renal insufficiency and renal failure

Cardiovascular (CVS) adverse events, particularly arrhythmia, appeared to be correlated mostly with pre-existing CVS disease and prior therapy with cardiotoxic agents (see section 4.4). Cardiomyopathy, that may be reversible upon discontinuation of interferon alpha, has been reported rarely in patients without prior evidence of cardiac disease.

Ophthalmological disorders that have been reported rarely with alpha interferons include retinopathies (including macular oedema), retinal haemorrhages, retinal artery or vein obstruction, cotton wool spots, loss of visual acuity or visual field, optic neuritis, and papilloedema (see section 4.4).

A wide variety of autoimmune and immune-mediated disorders have been reported with alpha interferons including thyroid disorders, systemic lupus erythematosus, rheumatoid arthritis (new or aggravated), idiopathic and thrombotic thrombocytopenic purpura, vasculitis, neuropathies including mononeuropathies (see also section 4.4, **Autoimmune disorders**).

**HCV/HIV co-infected patients**

For HCV/HIV co-infected patients receiving PegIntron in combination with ribavirin, other undesirable effects (that were not reported in mono-infected patients) which have been reported in the larger studies with a frequency  $> 5\%$  were: oral candidiasis (14 %), lipodystrophy acquired (13 %), CD4 lymphocytes decreased (8 %), appetite decreased (8 %), gamma-glutamyltransferase increased (9 %), back pain (5 %), blood amylase increased (6 %), blood lactic acid increased (5 %), cytolytic hepatitis (6 %), lipase increased (6 %) and pain in limb (6 %).

*Mitochondrial toxicity:*

Mitochondrial toxicity and lactic acidosis have been reported in HIV-positive patients receiving NRTI

regimen and associated ribavirin for co-HCV infection (see section 4.4).

*Laboratory values for HCV/HIV co-infected patients:*

Although haematological toxicities of neutropenia, thrombocytopenia and anaemia occurred more frequently in HCV/HIV co-infected patients, the majority could be managed by dose modification and rarely required premature discontinuation of treatment (see section 4.4). Haematological abnormalities were more frequently reported in patients receiving PegIntron in combination with ribavirin when compared to patients receiving interferon alfa-2b in combination with ribavirin. In Study 1 (see section 5.1), decrease in absolute neutrophil count levels below 500 cells/mm<sup>3</sup> was observed in 4 % (8/194) of patients and decrease in platelets below 50,000/mm<sup>3</sup> was observed in 4 % (8/194) of patients receiving PegIntron in combination with ribavirin. Anaemia (hemoglobin < 9.4g/dl) was reported in 12% (23/194) of patients treated with PegIntron in combination with ribavirin.

*CD4 lymphocytes decrease:*

Treatment with PegIntron in combination with ribavirin was associated with decreases in absolute CD4+ cell counts within the first 4 weeks without a reduction in CD4+ cell percentage. The decrease in CD4+ cell counts was reversible upon dose reduction or cessation of therapy. The use of PegIntron in combination with ribavirin had no observable negative impact on the control of HIV viraemia during therapy or follow-up. Limited safety data (N= 25) are available in co-infected patients with CD4+ cell counts < 200/μl (see section 4.4).

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron in combination with ribavirin.

## **4.9 Overdose**

In clinical trials, cases of accidental overdose, at never more than twice the prescribed dose, were reported. There were no serious reactions. Undesirable effects resolved during continued administration of PegIntron.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Immunostimulants, Cytokines and immunomodulators, Interferons, Peginterferon alfa-2b, ATC code: L03A B10.

Recombinant interferon alfa-2b is covalently conjugated with monomethoxy polyethylene glycol at an average degree of substitution of 1 mole of polymer/mole of protein. The average molecular mass is approximately 31,300 daltons of which the protein moiety constitutes approximately 19,300.

#### **Interferon alfa-2b**

*In vitro* and *in vivo* studies suggest that the biological activity of PegIntron is derived from its interferon alfa-2b moiety.

Interferons exert their cellular activities by binding to specific membrane receptors on the cell surface. Studies with other interferons have demonstrated species specificity. However, certain monkey species, e.g., Rhesus monkeys, are susceptible to pharmacodynamic stimulation upon exposure to human type 1 interferons.

Once bound to the cell membrane, interferon initiates a complex sequence of intracellular events that include the induction of certain enzymes. It is thought that this process, at least in part, is responsible for the various cellular responses to interferon, including inhibition of virus replication in virus-infected cells, suppression of cell proliferation and such immunomodulating activities as enhancement of the phagocytic activity of macrophages and augmentation of the specific cytotoxicity of

lymphocytes for target cells. Any or all of these activities may contribute to interferon's therapeutic effects.

Recombinant interferon alfa-2b also inhibits viral replication *in vitro* and *in vivo*. Although the exact antiviral mode of action of recombinant interferon alfa-2b is unknown, it appears to alter the host cell metabolism. This action inhibits viral replication or if replication occurs, the progeny virions are unable to leave the cell.

### **PegIntron**

PegIntron pharmacodynamics were assessed in a rising single-dose trial in healthy subjects by examining changes in oral temperature, concentrations of effector proteins such as serum neopterin and 2'5'-oligoadenylate synthetase (2'5'-OAS), as well as white cell and neutrophil counts. Subjects treated with PegIntron showed mild dose-related elevations in body temperature. Following single doses of PegIntron between 0.25 and 2.0 micrograms/kg/week, serum neopterin concentration was increased in a dose-related manner. Neutrophil and white cell count reductions at the end of week 4 correlated with the dose of PegIntron.

### **PegIntron clinical trials**

Two pivotal trials have been conducted, one (C/I97-010) with PegIntron monotherapy; the other (C/I98-580) with PegIntron in combination with ribavirin. Eligible patients for these trials had chronic hepatitis C confirmed by a positive HCV-RNA polymerase chain reaction (PCR) assay (> 30 IU/ml), a liver biopsy consistent with a histological diagnosis of chronic hepatitis with no other cause for the chronic hepatitis, and abnormal serum ALT.

In the PegIntron monotherapy trial, a total of 916 naïve chronic hepatitis C patients were treated with PegIntron (0.5, 1.0 or 1.5 micrograms/kg/week) for one year with a follow-up period of six months. In addition, 303 patients received interferon alfa-2b (3 million International Units [MIU] three times a week) as a comparator. This study showed that PegIntron was superior to interferon alfa-2b (**Table 8**).

In the PegIntron combination trial, 1,530 naïve patients were treated for one year with one of the following combination regimens:

- PegIntron (1.5 micrograms/kg/week) + ribavirin (800 mg/day), (n = 511).
- PegIntron (1.5 micrograms/kg/week for one month followed by 0.5 microgram/kg/week for 11 months) + ribavirin (1,000/1,200 mg/day), (n = 514).
- Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day) (n = 505).

In this trial, the combination of PegIntron (1.5 micrograms/kg/week) and ribavirin was significantly more effective than the combination of interferon alfa-2b and ribavirin (**Table 7**), particularly in patients infected with Genotype 1 (**Table 8**). Sustained response was assessed by the response rate six months after the cessation of treatment.

HCV genotype and baseline virus load are prognostic factors which are known to affect response rates. However, response rates in this trial were shown to be dependent also on the dose of ribavirin administered in combination with PegIntron or interferon alfa-2b. In those patients that received > 10.6 mg/kg ribavirin (800 mg dose in typical 75 kg patient), regardless of genotype or viral load, response rates were significantly higher than in those patients that received ≤ 10.6 mg/kg ribavirin (**Table 8**), while response rates in patients that received > 13.2 mg/kg ribavirin were even higher.

<b>Table 7</b> Sustained virological response (% patients HCV negative)							
	<b>PegIntron monotherapy</b>				<b>PegIntron + ribavirin</b>		
<b>Treatment regimen</b>	<b>P 1.5</b>	<b>P 1.0</b>	<b>P 0.5</b>	<b>I</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>Number of patients</b>	304	297	315	303	511	514	505
<b>Response at end of treatment</b>	49 %	41 %	33 %	24 %	65 %	56 %	54 %

Sustained response	23 %*	25 %	18 %	12 %	54 %**	47 %	47 %
P 1.5	PegIntron 1.5 micrograms/kg						
P 1.0	PegIntron 1.0 microgram/kg						
P 0.5	PegIntron 0.5 microgram/kg						
I	Interferon alfa-2b 3 MIU						
P 1.5/R	PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)						
P 0.5/R	PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)						
I/R	Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)						
*	p < 0.001 P 1.5 vs. I						
**	p = 0.0143 P 1.5/R vs. I/R						

<b>Table 8</b> Sustained response rates with PegIntron + ribavirin (by ribavirin dose, genotype and viral load)				
HCV Genotype	Ribavirin dose (mg/kg)	P 1.5/R	P 0.5/R	I/R
<b>All Genotypes</b>	<b>All</b>	<b>54 %</b>	<b>47 %</b>	<b>47 %</b>
	≤ 10.6	50 %	41 %	27 %
	> 10.6	61 %	48 %	47 %
<b>Genotype 1</b>	<b>All</b>	<b>42 %</b>	<b>34 %</b>	<b>33 %</b>
	≤ 10.6	38 %	25 %	20 %
	> 10.6	48 %	34 %	34 %
Genotype 1 ≤ 600,000 IU/ml	All	73 %	51 %	45 %
	≤ 10.6	74 %	25 %	33 %
	> 10.6	71 %	52 %	45 %
Genotype 1 > 600,000 IU/ml	All	30 %	27 %	29 %
	≤ 10.6	27 %	25 %	17 %
	> 10.6	37 %	27 %	29 %
<b>Genotype 2/3</b>	<b>All</b>	<b>82 %</b>	<b>80 %</b>	<b>79 %</b>
	≤ 10.6	79 %	73 %	50 %
	> 10.6	88 %	80 %	80 %

P 1.5/R PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)  
P 0.5/R PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)  
I/R Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)

In the PegIntron monotherapy study, the Quality of Life was generally less affected by 0.5 microgram/kg of PegIntron than by either 1.0 microgram/kg of PegIntron once weekly or 3 MIU of interferon alfa-2b three times a week.

In a separate trial, 224 patients with genotype 2 or 3 received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with ribavirin 800 mg –1,400 mg p.o. for 6 months (based on body weight, only three patients weighing > 105 kg, received the 1,400 mg dose, which has not yet been validated) (**Table 9**). Twenty-four % had bridging fibrosis or cirrhosis (Knodell 3/4).

**Table 9. Virologic Response at End of Treatment, Sustained Virologic Response and Relapse by HCV Genotype and Viral Load\***

	PegIntron 1.5 µg/kg once weekly plus Ribavirin 800-1,400 mg/day		
	End of Treatment Response	Sustained Virologic Response	Relapse
All Subjects	94 % (211/224)	81 % (182/224)	12 % (27/224)
HCV 2 ≤ 600,000 IU/ml > 600,000 IU/ml	100 % (42/42)	93 % (39/42)	7 % (3/42)
	100 % (20/20)	95 % (19/20)	5 % (1/20)
	100 % (22/22)	91 % (20/22)	9 % (2/22)
HCV 3	93 % (169/182)	79 % (143/182)	14 %

≤ 600,000 IU/ml	93 % (92/99)	86 % (85/99)	(24/166) 8 % (7/91)
> 600,000 IU/ml	93 % (77/83)	70 % (58/83)	23 % (17/75)

\* Any subject with an undetectable HCV-RNA level at the Follow-Up Week 12 visit and missing data at the Follow-Up Week 24 visit was considered a sustained responder. Any subject with missing data in and after the Follow-Up Week 12 window was considered to be a non-responder at Week 24 of follow-up.

The 6 month treatment duration in this trial was better tolerated than one year of treatment in the pivotal combination trial; for discontinuation 5 % vs. 14 %, for dose modification 18 % vs. 49 %.

In a non-comparative trial, 235 patients with genotype 1 and low viral load (< 600,000 IU/ml) received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with weight adjusted ribavirin. The overall sustained response rate after a 24-week treatment duration was 50 %. Forty-one percent of subjects (97/235) had nondetectable plasma HCV-RNA levels at Week 4 and Week 24 of therapy. In this subgroup, there was a 92 % (89/97) sustained virological response rate. The high sustained response rate in this subgroup of patients was identified in an interim analysis (n=49) and prospectively confirmed (n=48).

Limited historical data indicate that treatment for 48 weeks might be associated with a higher sustained response rate (11/11) and with a lower risk of relapse (0/11 as compared to 7/96 following 24 weeks of treatment).

#### Predictability of sustained virological response

Virological response by week 12, defined as a 2-log viral load decrease or undetectable levels of HCV-RNA has been shown to be predictive for sustained response (**Table 10**).

<b>Table 10</b> Predictability of sustained response by viral response at week 12 and genotype*				
Treatment	Genotype	Viral response at week 12	Sustained response	Negative predictive value
PegIntron 1.5 + ribavirin (> 10.6 mg/kg) 48-week treatment	1	Yes 75 % (82/110)	71 % (58/82)	----
		No 25 % (28/110)	0 % (0/28)	100 %
PegIntron 1.5 + ribavirin 800-1,400 mg 24-week treatment	2 and 3	Yes 99 % (213/215)	83 % (177/213)	----
		No 1 % (2/215)	50 % (1/2)	50 %

\* reflects patients with 12 week data available

The negative predictive value for sustained response in patients treated with PegIntron in monotherapy was 98 %.

#### HCV/HIV Co-infected patients

Two trials have been conducted in patients co-infected with HIV and HCV. The response to treatment in both of these trials is presented in **Table 11**. Study 1 (RIBAVIC; P01017) was a randomized, multicentre study which enrolled 412 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (1.5 µg/kg/week) plus ribavirin (800 mg/day) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800 mg/day) for 48 weeks with a follow-up period of 6 months. Study 2 (P02080) was a randomized, single centre study that enrolled 95 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (100 or 150 µg /week based



on weight) plus ribavirin (800-1,200 mg/day based on weight) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800-1,200 mg/day based on weight). The duration of therapy was 48 weeks with a follow-up period of 6 months except for patients infected with genotypes 2 or 3 and viral load < 800,000 IU/ml (Amplicor) who were treated for 24 weeks with a 6-month follow-up period.

<b>Table 11</b> Sustained virological response based on genotype after PegIntron in combination with Ribavirin in HCV/HIV Co-infected patients						
	<b>Study 1<sup>1</sup></b>			<b>Study 2<sup>2</sup></b>		
	PegIntron (1.5 µg/kg/week) + ribavirin (800 mg)	Interferon alfa-2b (3 MIU TIW) + ribavirin (800 mg)	p value <sup>a</sup>	PegIntron (100 or 150 <sup>c</sup> µg/week) + ribavirin (800-1,200 mg) <sup>d</sup>	Interferon alfa-2b (3 MIU TIW) + ribavirin (800-1,200 mg) <sup>d</sup>	p value <sup>b</sup>
All	27 % (56/205)	20 % (41/205)	0.047	44 % (23/52)	21 % (9/43)	0.017
Genotype 1, 4	17 % (21/125)	6 % (8/129)	0.006	38 % (12/32)	7 % (2/27)	0.007
Genotype 2, 3	44 % (35/80)	43 % (33/76)	0.88	53 % (10/19)	47 % (7/15)	0.730

MIU = million international units; TIW = three times a week.

a: p value based on Cochran-Mantel Haenszel Chi square test.

b: p value based on chi-square test.

c: subjects < 75 kg received 100 µg/week PegIntron and subjects ≥ 75 kg received 150 µg/week PegIntron.

d: ribavirin dosing was 800 mg for patients < 60 kg, 1,000 mg for patients 60-75 kg, and 1,200 mg for patients > 75 kg.

<sup>1</sup>Carrato F, Bani-Sadir F, Pol S et al. JAMA 2004; 292(23): 2839-2848.

<sup>2</sup>Laguno M, Murillas J, Blanco J et al. AIDS 2004; 18(13): F27-F36.

### *Histological response*

Liver biopsies were obtained before and after treatment in Study 1 and were available for 210 of the 412 subjects (51 %). Both the Metavir score and Ishak grade decreased among subjects treated with PegIntron in combination with ribavirin. This decline was significant among responders (-0.3 for Metavir and -1.2 for Ishak) and stable (-0.1 for Metavir and -0.2 for Ishak) among non-responders. In terms of activity, about one-third of sustained responders showed improvement and none showed worsening. There was no improvement in terms of fibrosis observed in this study. Steatosis was significantly improved in patients infected with HCV Genotype 3.

## **5.2 Pharmacokinetic properties**

PegIntron is a well characterized polyethylene glycol-modified (“pegylated”) derivative of interferon alfa-2b and is predominantly composed of monopegylated species. The plasma half-life of PegIntron is prolonged compared with non-pegylated interferon alfa-2b. PegIntron has a potential to depegylate to free interferon alfa-2b. The biologic activity of the pegylated isomers is qualitatively similar, but weaker than free interferon alfa-2b.

Following subcutaneous administration, maximal serum concentrations occur between 15-44 hours post-dose, and are sustained for up to 48-72 hours post-dose.

PegIntron C<sub>max</sub> and AUC measurements increase in a dose-related manner. Mean apparent volume of distribution is 0.99 l/kg.

Upon multiple dosing, there is an accumulation of immunoreactive interferons. There is, however, only a modest increase in biologic activity as measured by a bioassay.

Mean (SD) PegIntron elimination half-life is approximately 40 hours (13.3 hours), with apparent clearance of 22.0 ml/hr·kg. The mechanisms involved in clearance of interferons in man have not yet been

fully elucidated. However, renal elimination may account for a minority (approximately 30 %) of PegIntron apparent clearance.

**Renal function:** Renal clearance appears to account for 30 % of total clearance of PegIntron. In a single dose study (1.0 microgram/kg) in patients with impaired renal function,  $C_{max}$ , AUC, and half-life increased in relation to the degree of renal impairment.

Following multiple dosing of PegIntron (1.0 microgram/kg subcutaneously administered every week for four weeks) the clearance of PegIntron is reduced by a mean of 17 % in patients with moderate renal impairment (creatinine clearance 30-49 ml/minute) and by a mean of 44 % in patients with severe renal impairment (creatinine clearance 15-29 ml/minute) compared to subjects with normal renal function. Based on single dose data, clearance was similar in patients with severe renal impairment not on dialysis and in patients who were receiving hemodialysis. The dose of PegIntron for monotherapy should be reduced in patients with moderate or severe renal impairment (see sections 4.2 and 4.4). Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see section 4.3).

Because of marked inter-subject variability in interferon pharmacokinetics, it is recommended that patients with severe renal impairment be closely monitored during treatment with PegIntron (see section 4.2)

**Hepatic function:** The pharmacokinetics of PegIntron have not been evaluated in patients with severe hepatic dysfunction.

**Elderly patients  $\geq$  65 years of age:** The pharmacokinetics of PegIntron following a single subcutaneous dose of 1.0 microgram/kg were not affected by age. The data suggest that no alteration in PegIntron dosage is necessary based on advancing age.

**Patients under the age of 18 years:** Specific pharmacokinetic evaluations have not been performed on these patients. PegIntron is indicated for the treatment of chronic hepatitis C only in patients 18 years of age or older.

**Interferon neutralising factors:** Interferon neutralising factor assays were performed on serum samples of patients who received PegIntron in the clinical trial. Interferon neutralising factors are antibodies which neutralise the antiviral activity of interferon. The clinical incidence of neutralising factors in patients who received PegIntron 0.5 micrograms/kg is 1.1 %.

### 5.3 Preclinical safety data

**PegIntron:** Adverse events not observed in clinical trials were not seen in toxicity studies in monkeys. These studies were limited to four weeks due to the appearance of anti-interferon antibodies in most monkeys.

Reproduction studies of PegIntron have not been performed. Interferon alfa-2b has been shown to be an abortifacient in primates. PegIntron is likely to also cause this effect. Effects on fertility have not been determined. It is not known whether the components of this medicinal product are excreted into experimental animal or human milk (see section 4.6 for relevant human data on pregnancy and lactation). PegIntron showed no genotoxic potential.

The relative non-toxicity of monomethoxy-polyethylene glycol (mPEG), which is liberated from PegIntron by metabolism *in vivo* has been demonstrated in preclinical acute and subchronic toxicity studies in rodents and monkeys, standard embryo-foetal development studies and in *in vitro* mutagenicity assays.

**PegIntron plus ribavirin:** When used in combination with ribavirin, PegIntron did not cause any effects not previously seen with either active substance alone. The major treatment-related change was

a reversible, mild to moderate anaemia, the severity of which was greater than that produced by either active substance alone.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Powder for solution for injection:

Disodium phosphate, anhydrous,  
Sodium dihydrogen phosphate dihydrate,  
Sucrose,  
Polysorbate 80.

Solvent for parenteral use:

Water for injections.

Deliverable volume from pen = 0.5 ml.

### **6.2 Incompatibilities**

This medicinal product should only be reconstituted with the solvent provided (see section 6.6). In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

3 years

After reconstitution:

- Chemical and physical in-use stability has been demonstrated for 24 hours at 2°C - 8°C.
- From a microbiological point of view, the product is to be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C - 8°C.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C - 8°C). Do not freeze.

### **6.5 Nature and contents of container**

The powder and solvent are both contained in a two-chamber cartridge, Type I flint glass, separated by a bromobutyl rubber plunger. The cartridge is sealed at one end with a polypropylene cap containing a bromobutyl rubber liner and at the other end by a bromobutyl rubber plunger.

PegIntron 80 micrograms is supplied as:

- 1 pen containing powder and solvent for solution for injection, 1 injection needle and 2 cleansing swabs;
- 4 pens containing powder and solvent for solution for injection, 4 injection needles and 8 cleansing swabs;
- 6 pens containing powder and solvent for solution for injection, 6 injection needles and 12 cleansing swabs;
- 12 pens containing powder and solvent for solution for injection, 12 injection needles and 24 cleansing swabs.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

PegIntron pre-filled pen contains a powder of peginterferon alfa-2b and a solvent for solution at a strength of 80 micrograms for single use. Each pen is reconstituted with the solvent provided in the two-chamber cartridge (water for injections) for administration of up to 0.5 ml of solution. A small volume is lost during preparation of PegIntron for injection when the dose is measured and injected. Therefore, each pen contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection. The reconstituted solution has a concentration of 80 micrograms in 0.5 ml.

PegIntron is injected subcutaneously after reconstituting the powder as instructed, attaching an injection needle and setting the prescribed dose. A complete and illustrated set of instructions is provided in the Annex to the Package Leaflet.

Remove PegIntron pre-filled pen from the refrigerator before administration to allow the solvent to reach room temperature (not more than 25°C).

As for all parenteral medicinal products, inspect visually the reconstituted solution prior to administration. The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. After administering the dose, discard the PegIntron pre-filled pen and any unused solution contained in it.

## **7. MARKETING AUTHORISATION HOLDER**

SP Europe  
73, rue de Stalle  
B-1180 Bruxelles  
Belgium

## **8. MARKETING AUTHORISATION NUMBERS**

EU/1/00/131/035  
EU/1/00/131/036  
EU/1/00/131/037  
EU/1/00/131/038

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 6 February 2002  
Date of last renewal: 25 May 2005

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.emea.europa.eu/>

## **1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 100 micrograms, powder and solvent for solution for injection in pre-filled pen

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each pre-filled pen of PegIntron 100 micrograms contains a sufficient amount of peginterferon alfa-2b as measured on a protein basis in a powder for solution for injection, and the corresponding amount of solvent, to provide 100 micrograms in 0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

The active substance is a covalent conjugate of recombinant interferon alfa-2b\* with monomethoxy polyethylene glycol. The potency of this product should not be compared to that of another pegylated or non-pegylated protein of the same therapeutic class. For more information, see section 5.1.

\*produced by rDNA technology in *E.coli* cells harbouring a genetically engineered plasmid hybrid encompassing an interferon alfa-2b gene from human leukocytes

For a full list of excipients, see section 6.1. PegIntron contains 40 mg of sucrose per 0.5 ml.

## **3. PHARMACEUTICAL FORM**

Powder and solvent for solution for injection in pre-filled pen

White powder.

Clear and colourless solvent.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

PegIntron is indicated for the treatment of adult patients with chronic hepatitis C who have elevated transaminases without liver decompensation and who are positive for serum HCV-RNA or anti-HCV, including naïve patients with clinically stable HIV co-infection (see section 4.4).

The best way to use PegIntron in this indication is in combination with ribavirin.

This combination is indicated in naïve patients including patients with clinically stable HIV co-infection. This combination of PegIntron with ribavirin is also indicated in patients who have previously responded (with normalisation of ALT at the end of treatment) to interferon alpha monotherapy but who have subsequently relapsed.

Interferon monotherapy, including PegIntron, is indicated mainly in case of intolerance or contraindication to ribavirin.

Please refer also to the ribavirin Summary of Product Characteristics (SPC) when PegIntron is to be used in combination with ribavirin.

### **4.2 Posology and method of administration**

Treatment should be initiated and monitored only by a physician experienced in the management of patients with hepatitis C.

#### **Dose to be administered**

PegIntron should be administered as a once weekly subcutaneous injection. The dose administered depends on whether it is used in combination with ribavirin or as monotherapy.

## Combination therapy

PegIntron 1.5 micrograms/kg/week in combination with ribavirin capsules.

The intended dose of 1.5 µg/kg of PegIntron to be used in combination with ribavirin may be delivered in weight categories with the pen/vial strengths according to **Table 1**. Ribavirin capsules are to be administered orally each day in two divided doses with food (morning and evening).

**Table 1 - Dosing for Combination Therapy**

Body Weight (kg)	PegIntron		Ribavirin Capsules	
	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Total Daily Dose (mg)	Number of Capsules (200 mg)
< 40	50	0.5	800	4 <sup>a</sup>
40-50	80	0.4	800	4 <sup>a</sup>
51-64	80	0.5	800	4 <sup>a</sup>
65-75	100	0.5	1,000	5 <sup>b</sup>
76-85	120	0.5	1,000	5 <sup>b</sup>
> 85	150	0.5	1,200	6 <sup>c</sup>

a: 2 morning, 2 evening

b: 2 morning, 3 evening

c: 3 morning, 3 evening

## Duration of treatment

**Predictability of sustained virological response:** Patients infected with virus genotype 1 who fail to achieve virological response at Week 12 are highly unlikely to become sustained virological responders (see also section 5.1).

- **Genotype 1:** For patients who exhibit virological response at week 12, treatment should be continued for another nine month period (i.e., a total of 48 weeks).  
In the subset of patients with genotype 1 infection and low viral load (< 600,000 IU/ml) who become HCV-RNA negative at treatment week 4 and remain HCV-RNA negative at week 24, the treatment could either be stopped after this 24 week treatment course or pursued for an additional 24 weeks (i.e. overall 48 weeks treatment duration). However, an overall 24 weeks treatment duration may be associated with a higher risk of relapse than a 48 weeks treatment duration (see section 5.1).
- **Genotypes 2 or 3:** It is recommended that all patients be treated for 24 weeks, except for HCV/HIV co-infected patients who should receive 48 weeks of treatment.
- **Genotype 4:** In general, patients infected with genotype 4 are considered harder to treat and limited study data (n=66) indicate they are compatible with a duration of treatment as for genotype 1.

## HCV/HIV Co-infection

The recommended duration of dosing for HCV/HIV co-infected patients is 48 weeks, regardless of genotype.

### *Predictability of response and non-response in HCV/HIV Co-infection*

Early virological response by Week 12, defined as a 2 log viral load decrease or undetectable levels of HCV-RNA, has been shown to be predictive for sustained response. The negative predictive value for sustained response in HCV/HIV co-infected patients treated with PegIntron in combination with ribavirin was 99 % (67/68; Study 1) (see section 5.1). A positive predictive value of 50 % (52/104; Study 1) was observed for HCV/HIV co-infected patients receiving combination therapy.

### PegIntron monotherapy

As monotherapy the PegIntron regimen is 0.5 or 1.0 microgram/kg/week. The lowest vial or pen strength available is 50 µg/0.5 ml; therefore for patients prescribed 0.5 µg/kg/week, doses must be adjusted by volume as shown in **Table 2**. For the 1.0 µg/kg dose, similar volume adjustments can be made or alternate vial strengths can be used as shown in **Table 2**. PegIntron monotherapy was not studied in HCV/HIV co-infected patients.

**Table 2- Monotherapy Dosing**

	0.5 µg/kg		1.0 µg/kg	
Body Weight (kg)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)
30-35	50*	0.15	50	0.3
36-45	50*	0.2	50	0.4
46-56	50*	0.25	50	0.5
57-72	50	0.3	80	0.4
73-88	50	0.4	80	0.5
89-106	50	0.5	100	0.5
> 106**	80	0.4	120	0.5
* Must use vial. Minimum delivery for pen is 0.3 ml.				
** For patients > 120 kg, use 80 µg/0.5 ml vial				

### Duration of treatment

For patients who exhibit virological response at Week 12, treatment should be continued for at least another three-month period (i.e., a total of six months). The decision to extend therapy to one year of treatment should be based on prognostic factors (e.g., genotype, age > 40 years, male gender, bridging fibrosis).

### Dose modification for all patients

If severe adverse reactions or laboratory abnormalities develop during treatment with PegIntron monotherapy or PegIntron in combination with ribavirin, modify the dosages of each product as appropriate, until the adverse reactions abate. As adherence might be of importance for outcome of therapy, the dose should be kept as close as possible to the recommended standard dose. Guidelines were developed in clinical trials for dose modification.

### Combination Therapy Dose Reduction Guidelines

<b>Table 2a</b> Dose modification guidelines for combination therapy (with ribavirin)			
Laboratory values	Reduce only ribavirin dose to 600 mg/day* if:	Reduce only PegIntron dose to one-half dose if:	Discontinue combination therapy if:
Haemoglobin	< 10 g/dl	-	< 8.5 g/dl
Haemoglobin in: Patients with history of stable cardiac disease	≥ 2 g/dl decrease in haemoglobin during any four week period during treatment (permanent dose reduction)		< 12 g/dl after four weeks of dose reduction
White blood cells	-	< 1.5 x 10 <sup>9</sup> /l	< 1.0 x 10 <sup>9</sup> /l
Neutrophils	-	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	-	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l
Bilirubin – direct	-	-	2.5 x ULN**

Bilirubin – indirect	> 5 mg/dl	-	> 4 mg/dl (for > 4 weeks)
Creatinine	-	-	> 2.0 mg/dl
ALT/AST	-	-	2 x baseline and > 10 x ULN**

\* Patients whose dose of ribavirin is reduced to 600 mg daily receive one 200 mg capsule in the morning and two 200 mg capsules in the evening.

\*\* Upper limit of normal

Dose reduction of PegIntron may be accomplished by either reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 2b**.

<b>Table 2b – Reduced PegIntron Dosing for Combination Therapy</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5 ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
< 40	25	50*	0.25	25
40-50	32	50	0.3	30
51-64	40	50	0.4	40
65-75	50	50	0.5	50
76-85	60	80	0.4	64
> 85	75	100	0.4	80
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### **PegIntron Monotherapy Dose Reduction Guidelines**

Dose modification guidelines for patients who use PegIntron monotherapy are shown in **Table 3a**.

<b>Table 3a Dose modification guidelines for PegIntron monotherapy</b>		
<b>Laboratory values</b>	<b>Reduce PegIntron to one-half dose if:</b>	<b>Discontinue PegIntron if:</b>
Neutrophils	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l

Dose reduction for patients who use 0.5 µg/kg PegIntron monotherapy must be accomplished by reducing the prescribed volume by one-half. The 50 µg/0.5 ml vial must be used if necessary since the pen can only deliver a minimum volume of 0.3 ml.

For patients who use 1.0 µg/kg PegIntron monotherapy, dose reduction may be accomplished by reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 3b**.

<b>Table 3b – Reduced PegIntron Dose for the 1.0 µg/kg Monotherapy Regimen</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
30-35	15	50*	0.15	15



36-45	20	50*	0.20	20
46-56	25	50*	0.25	25
57-72	32	50	0.3	30
73-89	40	50	0.4	40
90-106	50	50	0.5	50
> 106	60	80	0.4	64
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### Special populations

**Use in renal impairment:** *Monotherapy:* PegIntron should be used with caution in patients with moderate to severe renal impairment. In patients with moderate renal dysfunction (creatinine clearance 30-50 ml/minute), the starting dose of PegIntron should be reduced by 25 %. Patients with severe renal dysfunction (creatinine clearance 15-29 ml/minute) should have the starting dose of PegIntron reduced by 50 %. Data are not available for the use of PegIntron in patients with creatinine clearance < 15 ml/minute (see section 5.2). Patients with severe renal impairment, including those on hemodialysis, should be closely monitored. If renal function decreases during treatment, PegIntron therapy should be discontinued.

*Combination therapy:* Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see ribavirin SPC). When administered in combination with ribavirin, subjects with impaired renal function should be more carefully monitored with respect to the development of anaemia.

**Use in hepatic impairment:** The safety and efficacy of PegIntron therapy has not been evaluated in patients with severe hepatic dysfunction, therefore PegIntron must not be used for these patients.

**Use in the elderly (≥ 65 years of age):** There are no apparent age-related effects on the pharmacokinetics of PegIntron. Data from elderly patients treated with a single dose of PegIntron suggest no alteration in PegIntron dose is necessary based on age (see section 5.2).

**Use in patients under the age of 18 years:** There is no experience in children (see section 5.2).

### 4.3 Contraindications

- Hypersensitivity to the active substance or to any interferon or to any of the excipients;
- A history of severe pre-existing cardiac disease, including unstable or uncontrolled cardiac disease in the previous six months (see section 4.4);
- Severe, debilitating medical conditions;
- Autoimmune hepatitis or a history of autoimmune disease;
- Severe hepatic dysfunction or decompensated cirrhosis of the liver;
- Pre-existing thyroid disease unless it can be controlled with conventional treatment;
- Epilepsy and/or compromised central nervous system (CNS) function;
- Initiation of PegIntron is contraindicated in HCV/HIV patients with cirrhosis and a Child-Pugh score ≥ 6.

*Combination therapy with ribavirin:* Also see ribavirin Summary of the Product Characteristics (SPC) if PegIntron is to be administered in combination with ribavirin in patients with chronic hepatitis C.

### 4.4 Special warnings and precautions for use

**Psychiatric and Central Nervous System (CNS):** Severe CNS effects, particularly depression, suicidal ideation and attempted suicide have been observed in some patients during PegIntron therapy, and even after treatment discontinuation mainly during the 6-month follow-up period. Other CNS effects including

aggressive behaviour (sometimes directed against others), confusion and alterations of mental status have been observed with alpha interferons. Patients should be closely monitored for any signs or symptoms of psychiatric disorders. If such symptoms appear, the potential seriousness of these undesirable effects must be borne in mind by the prescribing physician and the need for adequate therapeutic management should be considered. If psychiatric symptoms persist or worsen, or suicidal ideation is identified, it is recommended that treatment with PegIntron be discontinued, and the patient followed, with psychiatric intervention as appropriate.

*Patients with existence of, or history of severe psychiatric conditions:* If treatment with peginterferon alfa-2b is judged necessary in patients with existence or history of severe psychiatric conditions, this should only be initiated after having ensured appropriate individualised diagnostic and therapeutic management of the psychiatric condition.

There is no experience with PegIntron in combination with ribavirin in patients who have relapsed after interferon alpha + ribavirin therapy.

All patients in the chronic hepatitis C studies had a liver biopsy before inclusion, but in certain cases (i.e. patients with genotype 2 and 3), treatment may be possible without histological confirmation. Current treatment guidelines should be consulted as to whether a liver biopsy is needed prior to commencing treatment.

**Acute hypersensitivity:** Acute hypersensitivity reactions (e.g., urticaria, angioedema, bronchoconstriction, anaphylaxis) have been observed rarely during interferon alfa-2b therapy. If such a reaction develops during treatment with PegIntron, discontinue treatment and institute appropriate medical therapy immediately. Transient rashes do not necessitate interruption of treatment.

**Cardiovascular system:** As with interferon alfa-2b, patients with a history of congestive heart failure, myocardial infarction and/or previous or current arrhythmic disorders, receiving PegIntron therapy require close monitoring. It is recommended that patients who have pre-existing cardiac abnormalities have electrocardiograms taken prior to and during the course of treatment. Cardiac arrhythmias (primarily supraventricular) usually respond to conventional therapy but may require discontinuation of PegIntron therapy.

More significant obtundation and coma, including cases of encephalopathy, have been observed in some patients, usually elderly, treated at higher doses for oncology indications. While these effects are generally reversible, in a few patients full resolution took up to three weeks. Very rarely, seizures have occurred with high doses of interferon alpha.

**Liver function:** As with all interferons, discontinue treatment with PegIntron in patients who develop prolongation of coagulation markers which might indicate liver decompensation.

**Fever:** While fever may be associated with the flu-like syndrome reported commonly during interferon therapy, other causes of persistent fever must be ruled out.

**Hydration:** Adequate hydration must be maintained in patients undergoing PegIntron therapy since hypotension related to fluid depletion has been seen in some patients treated with alpha interferons. Fluid replacement may be necessary.

**Pulmonary changes:** Pulmonary infiltrates, pneumonitis, and pneumonia, occasionally resulting in fatality, have been observed rarely in interferon alpha treated patients. Any patient developing fever, cough, dyspnea or other respiratory symptoms must have a chest X-ray taken. If the chest X-ray shows pulmonary infiltrates or there is evidence of pulmonary function impairment, the patient is to be monitored closely, and, if appropriate, discontinue interferon alpha. Prompt discontinuation of interferon alpha administration and treatment with corticosteroids appear to be associated with resolution of pulmonary adverse events.

**Autoimmune disease:** The development of auto-antibodies and autoimmune disorders has been reported during treatment with alpha interferons. Patients predisposed to the development of

autoimmune disorders may be at increased risk. Patients with signs or symptoms compatible with autoimmune disorders should be evaluated carefully, and the benefit-risk of continued interferon therapy should be reassessed (see also section 4.4 Thyroid changes and 4.8).

**Ocular changes:** Ophthalmologic disorders, including retinal haemorrhages, cotton wool spots, and retinal artery or vein obstruction have been reported in rare instances after treatment with alpha interferons (see section 4.8). All patients should have a baseline eye examination. Any patient complaining of ocular symptoms, including loss of visual acuity or visual field must have a prompt and complete eye examination. Periodic visual examinations are recommended during PegIntron therapy, particularly in patients with disorders that may be associated with retinopathy, such as diabetes mellitus or hypertension. Discontinuation of PegIntron should be considered in patients who develop new or worsening ophthalmological disorders.

**Thyroid changes:** Infrequently, patients treated for chronic hepatitis C with interferon alpha have developed thyroid abnormalities, either hypothyroidism or hyperthyroidism. Determine thyroid stimulating hormone (TSH) levels if, during the course of therapy, a patient develops symptoms consistent with possible thyroid dysfunction. In the presence of thyroid dysfunction, PegIntron treatment may be continued if TSH levels can be maintained in the normal range by medication.

**Metabolic disturbances:** Hypertriglyceridemia and aggravation of hypertriglyceridemia, sometimes severe, have been observed. Monitoring of lipid levels is, therefore, recommended.

#### **HCV/HIV Co-infection**

Mitochondrial toxicity and lactic acidosis:

Patients co-infected with HIV and receiving Highly Active Anti-Retroviral Therapy (HAART) may be at increased risk of developing lactic acidosis. Caution should be used when adding PegIntron and ribavirin to HAART therapy (see ribavirin SPC).

Moreover, patients treated with PegIntron and ribavirin combination therapy and zidovudine are at increased risk of developing anaemia.

Hepatic decompensation in HCV/HIV co-infected patients with advanced cirrhosis:

Co-infected patients with advanced cirrhosis receiving HAART may be at increased risk of hepatic decompensation and death. Adding treatment with alfa interferons alone or in combination with ribavirin may increase the risk in this patient subset. Other baseline factors in co-infected patients that may be associated with a higher risk of hepatic decompensation include treatment with didanosine and elevated bilirubin serum concentration.

Co-infected patients receiving both antiretroviral (ARV) and anti-hepatitis treatment should be closely monitored, assessing their Child-Pugh score during treatment. Patients progressing to hepatic decompensation should have their anti-hepatitis treatment immediately discontinued and the ARV treatment reassessed.

Haematological abnormalities in HCV/HIV co-infected patients:

HCV/HIV co-infected patients receiving peginterferon alfa-2b/ribavirin treatment and HAART may be at increased risk to develop haematological abnormalities (as neutropenia, thrombocytopenia and anaemia) compared to HCV mono-infected patients. Although, the majority of them could be managed by dose reduction, close monitoring of haematological parameters should be undertaken in this population of patients (see section 4.2 and below “Laboratory tests” and section 4.8).

Patients with low CD4 counts:

In patients co-infected with HCV/HIV, limited efficacy and safety data (N = 25) are available in subjects with CD4 counts less than 200 cells/ $\mu$ l. Caution is therefore warranted in the treatment of patients with low CD4 counts.

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron and ribavirin.

**Dental and periodontal disorders:** Dental and periodontal disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and mucous membranes of the mouth during long-term treatment with the combination of PegIntron and ribavirin. Patients should brush their teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If this reaction occurs, they should be advised to rinse out their mouth thoroughly afterwards.

**Organ transplant recipients:** The safety and efficacy of PegIntron alone or in combination with ribavirin for the treatment of hepatitis C in liver or other organ transplant recipients have not been studied. Preliminary data indicates that interferon alpha therapy may be associated with an increased rate of kidney graft rejection. Liver graft rejection has also been reported.

**Other:** Due to reports of interferon alpha exacerbating pre-existing psoriatic disease and sarcoidosis, use of PegIntron in patients with psoriasis or sarcoidosis is recommended only if the potential benefit justifies the potential risk.

**Laboratory tests:** Standard haematologic tests, blood chemistry and a test of thyroid function must be conducted in all patients prior to initiating therapy. Acceptable baseline values that may be considered as a guideline prior to initiation of PegIntron therapy are:

- |                    |                              |
|--------------------|------------------------------|
| • Platelets        | $\geq 100,000/\text{mm}^3$   |
| • Neutrophil count | $\geq 1,500/\text{mm}^3$     |
| • TSH level        | must be within normal limits |

Laboratory evaluations are to be conducted at weeks 2 and 4 of therapy, and periodically thereafter as clinically appropriate.

**Important information about some of the ingredients of PegIntron:**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

Patients with rare hereditary problems of fructose intolerance, glucose galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Results from a multiple-dose probe study assessing P450 substrates in chronic hepatitis C patients receiving once weekly PegIntron (1.5  $\mu\text{g}/\text{kg}$ ) for 4 weeks demonstrated an increase in activity of CYP2D6 and CYP2C8/9. No change in activity of CYP1A2, CYP3A4, or N-acetyltransferase was observed.

Caution should be used when administering peginterferon alfa-2b with medicines metabolised by CYP2D6 and CYP2C8/9, especially those with narrow therapeutic window, such as warfarin and phenytoin (CYP2C9) and flecainide (CYP2D6).

These findings may partly relate to improved metabolic capacity due to reduced hepatic inflammation in patients undergoing treatment with PegIntron. Caution is therefore advised when PegIntron treatment is initiated for chronic hepatitis in patients treated with medication with a narrow therapeutic window and sensitive to mild metabolic impairment of the liver.

No pharmacokinetic interactions were noted between PegIntron and ribavirin in a multiple-dose pharmacokinetic study.

**Methadone:** In patients with chronic hepatitis C that were on stable methadone maintenance therapy and naïve to peginterferon alfa-2b, addition of 1.5 microgram/kg/week of PegIntron subcutaneously for 4 weeks increased R-methadone AUC by approximately 15 % (95 % CI for AUC ratio estimate 103 – 128 %). The clinical significance of this finding is unknown; however, patients should be

monitored for signs and symptoms of increased sedative effect, as well as respiratory depression. Especially in patients on a high dose of methadone, the risk for QTc prolongation should be considered.

#### *HCV/HIV Co-infection*

Nucleoside analogs: ribavirin has been shown *in vitro* to inhibit phosphorylation of zidovudine and stavudine. The clinical significance of these findings is unknown. However, these *in vitro* findings raise the possibility that concurrent use of ribavirin with either zidovudine or stavudine might lead to increased HIV plasma viraemia. Therefore, it is recommended that plasma HIV RNA levels be closely monitored in patients treated with ribavirin concurrently with either of these two agents. If HIV RNA levels increase, the use of ribavirin concomitantly with reverse transcriptase inhibitors must be reviewed (see ribavirin SPC).

Use of nucleoside analogs, alone or in combination with other nucleosides, has resulted in lactic acidosis. Pharmacologically, ribavirin increases phosphorylated metabolites of purine nucleosides *in vitro*. This activity could potentiate the risk of lactic acidosis induced by purine nucleoside analogs (e.g. didanosine or abacavir). Co-administration of ribavirin and didanosine is not recommended. Reports of mitochondrial toxicity, in particular lactic acidosis and pancreatitis, of which some fatal, have been reported (see ribavirin SPC).

### **4.6 Pregnancy and lactation**

PegIntron is recommended for use in fertile women only when they are using effective contraception during the treatment.

There are no adequate data from the use of interferon alfa-2b in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3). Interferon alfa-2b has been shown to be abortifacient in primates. PegIntron is likely to also cause this effect.

The potential risk in humans is unknown. PegIntron is to be used during pregnancy only if the potential benefit justifies the potential risk to the foetus

**Lactation:** It is not known whether the components of this medicinal product are excreted in human milk. Because of the potential for adverse reactions in breast-fed infants, breast-feeding should be discontinued prior to initiation of treatment.

#### *Combination therapy with ribavirin:*

Ribavirin causes serious birth defects when administered during pregnancy. Ribavirin therapy is contraindicated in women who are pregnant. Extreme care must be taken to avoid pregnancy in female patients or in partners of male patients taking PegIntron in combination with ribavirin. Females of childbearing potential and their partners must each use an effective contraceptive during treatment and for 4 months after treatment has been concluded. Male patients and their female partners must each use an effective contraceptive during treatment and for 7 months after treatment has been concluded (see Ribavirin SPC).

### **4.7 Effects on ability to drive and use machines**

Patients who develop fatigue, somnolence or confusion during treatment with PegIntron are cautioned to avoid driving or operating machinery.

### **4.8 Undesirable effects**

The safety of PegIntron is evaluated from data from two clinical trials: one with PegIntron monotherapy, one with PegIntron in combination with ribavirin. In both cases, patients were treated for one year.

**Table 4** describes the regimens and patient exposure for one year of treatment in patients with no previous exposure to interferon (interferon-naïve patients). Because of a significant overlap in the

pattern of undesirable effects with PegIntron monotherapy, groups of patients have been brought together in **Table 5** to show the pattern of reported effects for all monotherapy groups.

<b>Table 4</b> Regimens and patient exposure		
<b>Treatment</b>	<b>Regimen</b>	<b>Number of patients treated for one year</b>
PegIntron + ribavirin	PegIntron (1.5 micrograms/kg/week) + ribavirin (> 10.6 mg/kg/day)	188
Interferon alfa-2b + ribavirin	Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day)	505
PegIntron monotherapy	PegIntron (0.5 microgram/kg/week)	315
	PegIntron (1.0 microgram/kg/week)	297
	PegIntron (1.5 micrograms/kg/week)	304

<b>Table 5</b> Undesirable effects very commonly reported in clinical trials > 10 % of patients in PegIntron + ribavirin group Very common (> 1/10)			
	<b>PegIntron + ribavirin</b>	<b>Interferon alfa- 2b + ribavirin</b>	<b>PegIntron monotherapy</b>
<b>Infections and infestations</b>			
Infection viral	10 %	5 %	4-5 %
<b>Metabolism and nutrition disorders</b>			
Weight decrease	30 %	19 %	8-18 %
<b>Psychiatric disorders</b>			
Depression	34 %	32 %	26 %
Irritability	32 %	34 %	19 %
Insomnia	37 %	41 %	16-19 %
Anxiety	14 %	14 %	8 %
Concentration impaired	18 %	21 %	9-10 %
Emotional lability	11 %	10 %	5 %
<b>Nervous system disorders</b>			
Headache	58 %	57 %	57-63 %
Mouth dry	10 %	8 %	4-8 %
<b>Respiratory, thoracic and mediastinal disorders</b>			
Pharyngitis	10 %	7 %	3 %
Coughing	14 %	11 %	4 %
Dyspnea	26 %	22 %	5 %
<b>Gastrointestinal disorders</b>			
Nausea	43 %	31 %	20-23 %
Anorexia	35 %	26 %	10-25 %
Diarrhoea	20 %	13 %	14-17 %
Abdominal pain	12 %	9 %	11 %
Vomiting	16 %	10 %	4-7 %
<b>Skin and subcutaneous tissue disorders</b>			
Alopecia	45 %	32 %	20-34 %
Pruritus	27 %	27 %	7-9 %
Skin dry	23 %	21 %	6-9 %
Rash	21 %	21 %	5-7 %
<b>Musculoskeletal and connective tissue disorders</b>			
Myalgia	49 %	49 %	46-60 %
Arthralgia	31 %	26 %	23-28 %
Musculoskeletal pain	15 %	11 %	11-13 %
<b>General disorders and administration site conditions</b>			
Injection site inflammation	20 %	17 %	39-44 %
Injection site reaction	54 %	36 %	7-9 %
Fatigue	56 %	59 %	43 %
Rigors	42 %	40 %	33-43 %
Fever	39 %	32 %	29-43 %
Flu-like symptoms	21 %	23 %	18-25 %
Asthenia	28 %	17 %	12-14 %
Dizziness	17 %	16 %	7-12 %

<b>Table 6</b> Undesirable effects commonly reported in clinical trials in patients treated with PegIntron + ribavirin or PegIntron monotherapy Common (> 1/100, < 1/10)		
<b>Body system</b>	<b>5-10%</b>	<b>1-&lt;5%</b>
Infections and infestations		otitis media, fungal infection, bacterial infection, herpes simplex
Blood and lymphatic system disorders	Anaemia, leukopaenia	Thrombocytopenia, lymphadenopathy
Endocrine disorders	hypothyroidism	hyperthyroidism
Metabolism and nutrition disorders		hyperuricemia, hypocalcemia, thirst
Psychiatric disorders	agitation, nervousness	aggressive behaviour, somnolence, behavior disorder, apathy, appetite increased, sleep disorder, dreaming abnormal, decreased libido
Nervous system disorders	paresthesia, increased sweating	hypoesthesia, hyperaesthesia, hypertonia, confusion, tremor, vertigo, migraine, ataxia, neuralgia
Eye disorders		blurred vision, conjunctivitis, lacrimal gland disorder, eye pain
Cardiac disorders		tachycardia, palpitation
Vascular disorders		hypotension, hypertension, syncope, flushing
Ear and labyrinth disorders		tinnitus, hearing impairment/loss
Respiratory, thoracic and mediastinal disorders		nonproductive cough, rhinitis, sinusitis, bronchitis, respiratory disorder, nasal congestion, rhinorrhea, dysphonia, epistaxis
Gastrointestinal disorders	dyspepsia	constipation, taste perversion, loose stools, stomatitis, ulcerative stomatitis, gingival bleeding, glossitis, flatulence, hemorrhoids, gastroesophageal reflux, gingivitis, dehydration
Hepatobiliary disorders		hepatomegaly, hyperbilirubinemia
Skin and subcutaneous tissue disorders		erythematous rash, eczema, photosensitivity reaction, maculopapular rash, abnormal hair texture, acne, dermatitis, furunculosis, nail disorder, psoriasis, urticaria, erythema, face or peripheral oedema
Musculoskeletal and connective tissue disorders		arthritis
Renal and urinary disorders		micturition frequency, urine abnormal
Reproductive system and breast disorders	menstrual disorder, menorrhagia	ovarian disorder, vaginal disorder, sexual dysfunction (not specified), impotence, breast pain, amenorrhoea, prostatitis
General disorders and administration site conditions	RUQ pain, malaise, chest pain	injection site pain

Most cases of neutropaenia and thrombocytopaenia were mild (WHO grades 1 or 2). There were some cases of more severe neutropenia in patients treated with the recommended doses of PegIntron in combination with ribavirin (WHO grade 3: 39 of 186 [21 %]; and WHO grade 4: 13 of 186 [7 %]).



In a clinical trial, approximately 1.2 % of patients treated with PegIntron or interferon alfa-2b in combination with ribavirin reported life-threatening psychiatric events during treatment. These events included suicidal ideation and attempted suicide (see section 4.4). Following marketing, psychosis and hallucination have been reported rarely.

Rarely ( $> 1/10,000$ ,  $< 1/1,000$ ) or very rarely ( $< 1/10,000$ ) reported events with interferon alfa-2b, including PegIntron, include:

**Blood and lymphatic system disorders:**

Very rarely interferon alfa-2b or PegIntron used alone or in combination with ribavirin may be associated with aplastic anaemia. Pure red cell aplasia has been reported.

**Immune system disorders:**

very rarely: sarcoidosis or exacerbation of sarcoidosis

**Endocrine disorders:**

rarely: diabetes

**Nervous system disorders:**

rarely: seizure, peripheral neuropathy

very rarely: cerebrovascular ischaemia, cerebrovascular haemorrhage, encephalopathy

**Cardiac disorders:**

rarely: arrhythmia

very rarely: cardiac ischaemia, myocardial infarction

**Respiratory, thoracic and mediastinal disorders:**

very rarely: interstitial lung disease

**Gastrointestinal disorders:**

rarely: pancreatitis

very rarely: ulcerative and ischaemic colitis

**Skin and subcutaneous tissue disorders:**

very rarely: erythema multiforme, Stevens Johnson syndrome, toxic epidermal necrolysis, injection site necrosis

**Musculoskeletal and connective tissue disorders:**

rarely: rhabdomyolysis, myositis

**Renal and urinary disorders:**

rarely: renal insufficiency and renal failure

Cardiovascular (CVS) adverse events, particularly arrhythmia, appeared to be correlated mostly with pre-existing CVS disease and prior therapy with cardiotoxic agents (see section 4.4). Cardiomyopathy, that may be reversible upon discontinuation of interferon alpha, has been reported rarely in patients without prior evidence of cardiac disease.

Ophthalmological disorders that have been reported rarely with alpha interferons include retinopathies (including macular oedema), retinal haemorrhages, retinal artery or vein obstruction, cotton wool spots, loss of visual acuity or visual field, optic neuritis, and papilloedema (see section 4.4).

A wide variety of autoimmune and immune-mediated disorders have been reported with alpha interferons including thyroid disorders, systemic lupus erythematosus, rheumatoid arthritis (new or aggravated), idiopathic and thrombotic thrombocytopenic purpura, vasculitis, neuropathies including mononeuropathies (see also section 4.4, **Autoimmune disorders**).

**HCV/HIV co-infected patients**

For HCV/HIV co-infected patients receiving PegIntron in combination with ribavirin, other undesirable effects (that were not reported in mono-infected patients) which have been reported in the larger studies with a frequency  $> 5\%$  were: oral candidiasis (14 %), lipodystrophy acquired (13 %), CD4 lymphocytes decreased (8 %), appetite decreased (8 %), gamma-glutamyltransferase increased (9 %), back pain (5 %), blood amylase increased (6 %), blood lactic acid increased (5 %), cytolytic hepatitis (6 %), lipase increased (6 %) and pain in limb (6 %).

*Mitochondrial toxicity:*

Mitochondrial toxicity and lactic acidosis have been reported in HIV-positive patients receiving NRTI

regimen and associated ribavirin for co-HCV infection (see section 4.4).

*Laboratory values for HCV/HIV co-infected patients:*

Although haematological toxicities of neutropenia, thrombocytopenia and anaemia occurred more frequently in HCV/HIV co-infected patients, the majority could be managed by dose modification and rarely required premature discontinuation of treatment (see section 4.4). Haematological abnormalities were more frequently reported in patients receiving PegIntron in combination with ribavirin when compared to patients receiving interferon alfa-2b in combination with ribavirin. In Study 1 (see section 5.1), decrease in absolute neutrophil count levels below 500 cells/mm<sup>3</sup> was observed in 4 % (8/194) of patients and decrease in platelets below 50,000/mm<sup>3</sup> was observed in 4 % (8/194) of patients receiving PegIntron in combination with ribavirin. Anaemia (hemoglobin < 9.4g/dl) was reported in 12% (23/194) of patients treated with PegIntron in combination with ribavirin.

*CD4 lymphocytes decrease:*

Treatment with PegIntron in combination with ribavirin was associated with decreases in absolute CD4+ cell counts within the first 4 weeks without a reduction in CD4+ cell percentage. The decrease in CD4+ cell counts was reversible upon dose reduction or cessation of therapy. The use of PegIntron in combination with ribavirin had no observable negative impact on the control of HIV viraemia during therapy or follow-up. Limited safety data (N= 25) are available in co-infected patients with CD4+ cell counts < 200/μl (see section 4.4).

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron in combination with ribavirin.

## **4.9 Overdose**

In clinical trials, cases of accidental overdose, at never more than twice the prescribed dose, were reported. There were no serious reactions. Undesirable effects resolved during continued administration of PegIntron.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Immunostimulants, Cytokines and immunomodulators, Interferons, Peginterferon alfa-2b, ATC code: L03A B10.

Recombinant interferon alfa-2b is covalently conjugated with monomethoxy polyethylene glycol at an average degree of substitution of 1 mole of polymer/mole of protein. The average molecular mass is approximately 31,300 daltons of which the protein moiety constitutes approximately 19,300.

#### **Interferon alfa-2b**

*In vitro* and *in vivo* studies suggest that the biological activity of PegIntron is derived from its interferon alfa-2b moiety.

Interferons exert their cellular activities by binding to specific membrane receptors on the cell surface. Studies with other interferons have demonstrated species specificity. However, certain monkey species, e.g., Rhesus monkeys, are susceptible to pharmacodynamic stimulation upon exposure to human type 1 interferons.

Once bound to the cell membrane, interferon initiates a complex sequence of intracellular events that include the induction of certain enzymes. It is thought that this process, at least in part, is responsible for the various cellular responses to interferon, including inhibition of virus replication in virus-infected cells, suppression of cell proliferation and such immunomodulating activities as enhancement of the phagocytic activity of macrophages and augmentation of the specific cytotoxicity of

lymphocytes for target cells. Any or all of these activities may contribute to interferon's therapeutic effects.

Recombinant interferon alfa-2b also inhibits viral replication *in vitro* and *in vivo*. Although the exact antiviral mode of action of recombinant interferon alfa-2b is unknown, it appears to alter the host cell metabolism. This action inhibits viral replication or if replication occurs, the progeny virions are unable to leave the cell.

### **PegIntron**

PegIntron pharmacodynamics were assessed in a rising single-dose trial in healthy subjects by examining changes in oral temperature, concentrations of effector proteins such as serum neopterin and 2'5'-oligoadenylate synthetase (2'5'-OAS), as well as white cell and neutrophil counts. Subjects treated with PegIntron showed mild dose-related elevations in body temperature. Following single doses of PegIntron between 0.25 and 2.0 micrograms/kg/week, serum neopterin concentration was increased in a dose-related manner. Neutrophil and white cell count reductions at the end of week 4 correlated with the dose of PegIntron.

### **PegIntron clinical trials**

Two pivotal trials have been conducted, one (C/I97-010) with PegIntron monotherapy; the other (C/I98-580) with PegIntron in combination with ribavirin. Eligible patients for these trials had chronic hepatitis C confirmed by a positive HCV-RNA polymerase chain reaction (PCR) assay (> 30 IU/ml), a liver biopsy consistent with a histological diagnosis of chronic hepatitis with no other cause for the chronic hepatitis, and abnormal serum ALT.

In the PegIntron monotherapy trial, a total of 916 naïve chronic hepatitis C patients were treated with PegIntron (0.5, 1.0 or 1.5 micrograms/kg/week) for one year with a follow-up period of six months. In addition, 303 patients received interferon alfa-2b (3 million International Units [MIU] three times a week) as a comparator. This study showed that PegIntron was superior to interferon alfa-2b (**Table 8**).

In the PegIntron combination trial, 1,530 naïve patients were treated for one year with one of the following combination regimens:

- PegIntron (1.5 micrograms/kg/week) + ribavirin (800 mg/day), (n = 511).
- PegIntron (1.5 micrograms/kg/week for one month followed by 0.5 microgram/kg/week for 11 months) + ribavirin (1,000/1,200 mg/day), (n = 514).
- Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day) (n = 505).

In this trial, the combination of PegIntron (1.5 micrograms/kg/week) and ribavirin was significantly more effective than the combination of interferon alfa-2b and ribavirin (**Table 7**), particularly in patients infected with Genotype 1 (**Table 8**). Sustained response was assessed by the response rate six months after the cessation of treatment.

HCV genotype and baseline virus load are prognostic factors which are known to affect response rates. However, response rates in this trial were shown to be dependent also on the dose of ribavirin administered in combination with PegIntron or interferon alfa-2b. In those patients that received > 10.6 mg/kg ribavirin (800 mg dose in typical 75 kg patient), regardless of genotype or viral load, response rates were significantly higher than in those patients that received ≤ 10.6 mg/kg ribavirin (**Table 8**), while response rates in patients that received > 13.2 mg/kg ribavirin were even higher.

<b>Table 7</b> Sustained virological response (% patients HCV negative)							
	<b>PegIntron monotherapy</b>				<b>PegIntron + ribavirin</b>		
<b>Treatment regimen</b>	<b>P 1.5</b>	<b>P 1.0</b>	<b>P 0.5</b>	<b>I</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>Number of patients</b>	304	297	315	303	511	514	505
<b>Response at end of treatment</b>	49 %	41 %	33 %	24 %	65 %	56 %	54 %

Sustained response	23 %*	25 %	18 %	12 %	54 %**	47 %	47 %
P 1.5	PegIntron 1.5 micrograms/kg						
P 1.0	PegIntron 1.0 microgram/kg						
P 0.5	PegIntron 0.5 microgram/kg						
I	Interferon alfa-2b 3 MIU						
P 1.5/R	PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)						
P 0.5/R	PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)						
I/R	Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)						
*	p < 0.001 P 1.5 vs. I						
**	p = 0.0143 P 1.5/R vs. I/R						

<b>Table 8</b> Sustained response rates with PegIntron + ribavirin (by ribavirin dose, genotype and viral load)				
HCV Genotype	Ribavirin dose (mg/kg)	P 1.5/R	P 0.5/R	I/R
<b>All Genotypes</b>	<b>All</b>	<b>54 %</b>	<b>47 %</b>	<b>47 %</b>
	≤ 10.6	50 %	41 %	27 %
	> 10.6	61 %	48 %	47 %
<b>Genotype 1</b>	<b>All</b>	<b>42 %</b>	<b>34 %</b>	<b>33 %</b>
	≤ 10.6	38 %	25 %	20 %
	> 10.6	48 %	34 %	34 %
Genotype 1 ≤ 600,000 IU/ml	All	73 %	51 %	45 %
	≤ 10.6	74 %	25 %	33 %
	> 10.6	71 %	52 %	45 %
Genotype 1 > 600,000 IU/ml	All	30 %	27 %	29 %
	≤ 10.6	27 %	25 %	17 %
	> 10.6	37 %	27 %	29 %
<b>Genotype 2/3</b>	<b>All</b>	<b>82 %</b>	<b>80 %</b>	<b>79 %</b>
	≤ 10.6	79 %	73 %	50 %
	> 10.6	88 %	80 %	80 %

P 1.5/R	PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)
P 0.5/R	PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)
I/R	Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)

In the PegIntron monotherapy study, the Quality of Life was generally less affected by 0.5 microgram/kg of PegIntron than by either 1.0 microgram/kg of PegIntron once weekly or 3 MIU of interferon alfa-2b three times a week.

In a separate trial, 224 patients with genotype 2 or 3 received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with ribavirin 800 mg –1,400 mg p.o. for 6 months (based on body weight, only three patients weighing > 105 kg, received the 1,400 mg dose, which has not yet been validated) (**Table 9**). Twenty-four % had bridging fibrosis or cirrhosis (Knodell 3/4).

**Table 9. Virologic Response at End of Treatment, Sustained Virologic Response and Relapse by HCV Genotype and Viral Load\***

	PegIntron 1.5 µg/kg once weekly plus Ribavirin 800-1,400 mg/day		
	End of Treatment Response	Sustained Virologic Response	Relapse
All Subjects	94 % (211/224)	81 % (182/224)	12 % (27/224)
HCV 2 ≤ 600,000 IU/ml > 600,000 IU/ml	100 % (42/42)	93 % (39/42)	7 % (3/42)
	100 % (20/20)	95 % (19/20)	5 % (1/20)
	100 % (22/22)	91 % (20/22)	9 % (2/22)
HCV 3	93 % (169/182)	79 % (143/182)	14 %

≤ 600,000 IU/ml	93 % (92/99)	86 % (85/99)	(24/166) 8 % (7/91)
> 600,000 IU/ml	93 % (77/83)	70 % (58/83)	23 % (17/75)

\* Any subject with an undetectable HCV-RNA level at the Follow-Up Week 12 visit and missing data at the Follow-Up Week 24 visit was considered a sustained responder. Any subject with missing data in and after the Follow-Up Week 12 window was considered to be a non-responder at Week 24 of follow-up.

The 6 month treatment duration in this trial was better tolerated than one year of treatment in the pivotal combination trial; for discontinuation 5 % vs. 14 %, for dose modification 18 % vs. 49 %.

In a non-comparative trial, 235 patients with genotype 1 and low viral load (< 600,000 IU/ml) received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with weight adjusted ribavirin. The overall sustained response rate after a 24-week treatment duration was 50 %. Forty-one percent of subjects (97/235) had nondetectable plasma HCV-RNA levels at Week 4 and Week 24 of therapy. In this subgroup, there was a 92 % (89/97) sustained virological response rate. The high sustained response rate in this subgroup of patients was identified in an interim analysis (n=49) and prospectively confirmed (n=48).

Limited historical data indicate that treatment for 48 weeks might be associated with a higher sustained response rate (11/11) and with a lower risk of relapse (0/11 as compared to 7/96 following 24 weeks of treatment).

#### Predictability of sustained virological response

Virological response by week 12, defined as a 2-log viral load decrease or undetectable levels of HCV-RNA has been shown to be predictive for sustained response (**Table 10**).

<b>Table 10</b> Predictability of sustained response by viral response at week 12 and genotype*				
Treatment	Genotype	Viral response at week 12	Sustained response	Negative predictive value
PegIntron 1.5 + ribavirin (> 10.6 mg/kg) 48-week treatment	1	Yes 75 % (82/110)	71 % (58/82)	----
		No 25 % (28/110)	0 % (0/28)	100 %
PegIntron 1.5 + ribavirin 800-1,400 mg 24-week treatment	2 and 3	Yes 99 % (213/215)	83 % (177/213)	----
		No 1 % (2/215)	50 % (1/2)	50 %

\* reflects patients with 12 week data available

The negative predictive value for sustained response in patients treated with PegIntron in monotherapy was 98 %.

#### HCV/HIV Co-infected patients

Two trials have been conducted in patients co-infected with HIV and HCV. The response to treatment in both of these trials is presented in **Table 11**. Study 1 (RIBAVIC; P01017) was a randomized, multicentre study which enrolled 412 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (1.5 µg/kg/week) plus ribavirin (800 mg/day) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800 mg/day) for 48 weeks with a follow-up period of 6 months. Study 2 (P02080) was a randomized, single centre study that enrolled 95 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (100 or 150 µg /week based on weight) plus ribavirin (800-1,200 mg/day based on weight) or interferon alfa-2b (3 MIU TIW) plus

ribavirin (800-1,200 mg/day based on weight). The duration of therapy was 48 weeks with a follow-up period of 6 months except for patients infected with genotypes 2 or 3 and viral load < 800,000 IU/ml (Amplicor) who were treated for 24 weeks with a 6-month follow-up period.

<b>Table 11</b> Sustained virological response based on genotype after PegIntron in combination with Ribavirin in HCV/HIV Co-infected patients						
	<b>Study 1<sup>1</sup></b>			<b>Study 2<sup>2</sup></b>		
	PegIntron (1.5 µg/kg/week) + ribavirin (800 mg)	Interferon alfa-2b (3 MIU TIW) + ribavirin (800 mg)	p value <sup>a</sup>	PegIntron (100 or 150 <sup>c</sup> µg/week) + ribavirin (800-1,200 mg) <sup>d</sup>	Interferon alfa-2b (3 MIU TIW) + ribavirin (800-1,200 mg) <sup>d</sup>	p value <sup>b</sup>
All	27 % (56/205)	20 % (41/205)	0.047	44 % (23/52)	21 % (9/43)	0.017
Genotype 1, 4	17 % (21/125)	6 % (8/129)	0.006	38 % (12/32)	7 % (2/27)	0.007
Genotype 2, 3	44 % (35/80)	43 % (33/76)	0.88	53 % (10/19)	47 % (7/15)	0.730

MIU = million international units; TIW = three times a week.

a: p value based on Cochran-Mantel Haenszel Chi square test.

b: p value based on chi-square test.

c: subjects < 75 kg received 100 µg/week PegIntron and subjects ≥ 75 kg received 150 µg/week PegIntron.

d: ribavirin dosing was 800 mg for patients < 60 kg, 1,000 mg for patients 60-75 kg, and 1,200 mg for patients > 75 kg.

<sup>1</sup>Carrato F, Bani-Sadir F, Pol S et al. JAMA 2004; 292(23): 2839-2848.

<sup>2</sup>Laguno M, Murillas J, Blanco J et al. AIDS 2004; 18(13): F27-F36.

### *Histological response*

Liver biopsies were obtained before and after treatment in Study 1 and were available for 210 of the 412 subjects (51 %). Both the Metavir score and Ishak grade decreased among subjects treated with PegIntron in combination with ribavirin. This decline was significant among responders (-0.3 for Metavir and -1.2 for Ishak) and stable (-0.1 for Metavir and -0.2 for Ishak) among non-responders. In terms of activity, about one-third of sustained responders showed improvement and none showed worsening. There was no improvement in terms of fibrosis observed in this study. Steatosis was significantly improved in patients infected with HCV Genotype 3.

## **5.2 Pharmacokinetic properties**

PegIntron is a well characterized polyethylene glycol-modified (“pegylated”) derivative of interferon alfa-2b and is predominantly composed of monopegylated species. The plasma half-life of PegIntron is prolonged compared with non-pegylated interferon alfa-2b. PegIntron has a potential to depegylate to free interferon alfa-2b. The biologic activity of the pegylated isomers is qualitatively similar, but weaker than free interferon alfa-2b.

Following subcutaneous administration, maximal serum concentrations occur between 15-44 hours post-dose, and are sustained for up to 48-72 hours post-dose.

PegIntron C<sub>max</sub> and AUC measurements increase in a dose-related manner. Mean apparent volume of distribution is 0.99 l/kg.

Upon multiple dosing, there is an accumulation of immunoreactive interferons. There is, however, only a modest increase in biologic activity as measured by a bioassay.

Mean (SD) PegIntron elimination half-life is approximately 40 hours (13.3 hours), with apparent clearance of 22.0 ml/hr·kg. The mechanisms involved in clearance of interferons in man have not yet been

fully elucidated. However, renal elimination may account for a minority (approximately 30 %) of PegIntron apparent clearance.

**Renal function:** Renal clearance appears to account for 30 % of total clearance of PegIntron. In a single dose study (1.0 microgram/kg) in patients with impaired renal function,  $C_{max}$ , AUC, and half-life increased in relation to the degree of renal impairment.

Following multiple dosing of PegIntron (1.0 microgram/kg subcutaneously administered every week for four weeks) the clearance of PegIntron is reduced by a mean of 17 % in patients with moderate renal impairment (creatinine clearance 30-49 ml/minute) and by a mean of 44 % in patients with severe renal impairment (creatinine clearance 15-29 ml/minute) compared to subjects with normal renal function. Based on single dose data, clearance was similar in patients with severe renal impairment not on dialysis and in patients who were receiving hemodialysis. The dose of PegIntron for monotherapy should be reduced in patients with moderate or severe renal impairment (see sections 4.2 and 4.4). Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see section 4.3).

Because of marked inter-subject variability in interferon pharmacokinetics, it is recommended that patients with severe renal impairment be closely monitored during treatment with PegIntron (see section 4.2)

**Hepatic function:** The pharmacokinetics of PegIntron have not been evaluated in patients with severe hepatic dysfunction.

**Elderly patients  $\geq 65$  years of age:** The pharmacokinetics of PegIntron following a single subcutaneous dose of 1.0 microgram/kg were not affected by age. The data suggest that no alteration in PegIntron dosage is necessary based on advancing age.

**Patients under the age of 18 years:** Specific pharmacokinetic evaluations have not been performed on these patients. PegIntron is indicated for the treatment of chronic hepatitis C only in patients 18 years of age or older.

**Interferon neutralising factors:** Interferon neutralising factor assays were performed on serum samples of patients who received PegIntron in the clinical trial. Interferon neutralising factors are antibodies which neutralise the antiviral activity of interferon. The clinical incidence of neutralising factors in patients who received PegIntron 0.5 micrograms/kg is 1.1 %.

### 5.3 Preclinical safety data

**PegIntron:** Adverse events not observed in clinical trials were not seen in toxicity studies in monkeys. These studies were limited to four weeks due to the appearance of anti-interferon antibodies in most monkeys.

Reproduction studies of PegIntron have not been performed. Interferon alfa-2b has been shown to be an abortifacient in primates. PegIntron is likely to also cause this effect. Effects on fertility have not been determined. It is not known whether the components of this medicinal product are excreted into experimental animal or human milk (see section 4.6 for relevant human data on pregnancy and lactation). PegIntron showed no genotoxic potential.

The relative non-toxicity of monomethoxy-polyethylene glycol (mPEG), which is liberated from PegIntron by metabolism *in vivo* has been demonstrated in preclinical acute and subchronic toxicity studies in rodents and monkeys, standard embryo-foetal development studies and in *in vitro* mutagenicity assays.

**PegIntron plus ribavirin:** When used in combination with ribavirin, PegIntron did not cause any effects not previously seen with either active substance alone. The major treatment-related change was

a reversible, mild to moderate anaemia, the severity of which was greater than that produced by either active substance alone.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Powder for solution for injection:

Disodium phosphate, anhydrous,  
Sodium dihydrogen phosphate dihydrate,  
Sucrose,  
Polysorbate 80.

Solvent for parenteral use:

Water for injections.

Deliverable volume from pen = 0.5 ml.

### **6.2 Incompatibilities**

This medicinal product should only be reconstituted with the solvent provided (see section 6.6). In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

3 years

After reconstitution:

- Chemical and physical in-use stability has been demonstrated for 24 hours at 2°C - 8°C.
- From a microbiological point of view, the product is to be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C - 8°C.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C - 8°C). Do not freeze.

### **6.5 Nature and contents of container**

The powder and solvent are both contained in a two-chamber cartridge, Type I flint glass, separated by a bromobutyl rubber plunger. The cartridge is sealed at one end with a polypropylene cap containing a bromobutyl rubber liner and at the other end by a bromobutyl rubber plunger.

PegIntron 100 micrograms is supplied as:

- 1 pen containing powder and solvent for solution for injection, 1 injection needle and 2 cleansing swabs;
- 4 pens containing powder and solvent for solution for injection, 4 injection needles and 8 cleansing swabs;
- 6 pens containing powder and solvent for solution for injection, 6 injection needles and 12 cleansing swabs;
- 12 pens containing powder and solvent for solution for injection, 12 injection needles and 24 cleansing swabs.

Not all pack sizes may be marketed.



## **6.6 Special precautions for disposal**

PegIntron pre-filled pen contains a powder of peginterferon alfa-2b and a solvent for solution at a strength of 100 micrograms for single use. Each pen is reconstituted with the solvent provided in the two-chamber cartridge (water for injections) for administration of up to 0.5 ml of solution. A small volume is lost during preparation of PegIntron for injection when the dose is measured and injected. Therefore, each pen contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection. The reconstituted solution has a concentration of 100 micrograms in 0.5 ml.

PegIntron is injected subcutaneously after reconstituting the powder as instructed, attaching an injection needle and setting the prescribed dose. A complete and illustrated set of instructions is provided in the Annex to the Package Leaflet.

Remove PegIntron pre-filled pen from the refrigerator before administration to allow the solvent to reach room temperature (not more than 25°C).

As for all parenteral medicinal products, inspect visually the reconstituted solution prior to administration. The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. After administering the dose, discard the PegIntron pre-filled pen and any unused solution contained in it.

## **7. MARKETING AUTHORISATION HOLDER**

SP Europe  
73, rue de Stalle  
B-1180 Bruxelles  
Belgium

## **8. MARKETING AUTHORISATION NUMBERS**

EU/1/00/131/039  
EU/1/00/131/040  
EU/1/00/131/041  
EU/1/00/131/042

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 6 February 2002  
Date of last renewal: 25 May 2005

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.emea.europa.eu/>

## **1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 120 micrograms, powder and solvent for solution for injection in pre-filled pen

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each pre-filled pen of PegIntron 120 micrograms contains a sufficient amount of peginterferon alfa-2b as measured on a protein basis in a powder for solution for injection, and the corresponding amount of solvent, to provide 120 micrograms in 0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

The active substance is a covalent conjugate of recombinant interferon alfa-2b\* with monomethoxy polyethylene glycol. The potency of this product should not be compared to that of another pegylated or non-pegylated protein of the same therapeutic class. For more information, see section 5.1.

\*produced by rDNA technology in *E.coli* cells harbouring a genetically engineered plasmid hybrid encompassing an interferon alfa-2b gene from human leukocytes

For a full list of excipients, see section 6.1. PegIntron contains 40 mg of sucrose per 0.5 ml.

## **3. PHARMACEUTICAL FORM**

Powder and solvent for solution for injection in pre-filled pen

White powder.

Clear and colourless solvent.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

PegIntron is indicated for the treatment of adult patients with chronic hepatitis C who have elevated transaminases without liver decompensation and who are positive for serum HCV-RNA or anti-HCV, including naïve patients with clinically stable HIV co-infection (see section 4.4).

The best way to use PegIntron in this indication is in combination with ribavirin.

This combination is indicated in naïve patients including patients with clinically stable HIV co-infection. This combination of PegIntron with ribavirin is also indicated in patients who have previously responded (with normalisation of ALT at the end of treatment) to interferon alpha monotherapy but who have subsequently relapsed.

Interferon monotherapy, including PegIntron, is indicated mainly in case of intolerance or contraindication to ribavirin.

Please refer also to the ribavirin Summary of Product Characteristics (SPC) when PegIntron is to be used in combination with ribavirin.

### **4.2 Posology and method of administration**

Treatment should be initiated and monitored only by a physician experienced in the management of patients with hepatitis C.

#### **Dose to be administered**

PegIntron should be administered as a once weekly subcutaneous injection. The dose administered depends on whether it is used in combination with ribavirin or as monotherapy.

## Combination therapy

PegIntron 1.5 micrograms/kg/week in combination with ribavirin capsules.

The intended dose of 1.5 µg/kg of PegIntron to be used in combination with ribavirin may be delivered in weight categories with the pen/vial strengths according to **Table 1**. Ribavirin capsules are to be administered orally each day in two divided doses with food (morning and evening).

**Table 1 - Dosing for Combination Therapy**

Body Weight (kg)	PegIntron		Ribavirin Capsules	
	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Total Daily Dose (mg)	Number of Capsules (200 mg)
< 40	50	0.5	800	4 <sup>a</sup>
40-50	80	0.4	800	4 <sup>a</sup>
51-64	80	0.5	800	4 <sup>a</sup>
65-75	100	0.5	1,000	5 <sup>b</sup>
76-85	120	0.5	1,000	5 <sup>b</sup>
> 85	150	0.5	1,200	6 <sup>c</sup>

a: 2 morning, 2 evening

b: 2 morning, 3 evening

c: 3 morning, 3 evening

## Duration of treatment

**Predictability of sustained virological response:** Patients infected with virus genotype 1 who fail to achieve virological response at Week 12 are highly unlikely to become sustained virological responders (see also section 5.1).

- **Genotype 1:** For patients who exhibit virological response at week 12, treatment should be continued for another nine month period (i.e., a total of 48 weeks).  
In the subset of patients with genotype 1 infection and low viral load (< 600,000 IU/ml) who become HCV-RNA negative at treatment week 4 and remain HCV-RNA negative at week 24, the treatment could either be stopped after this 24 week treatment course or pursued for an additional 24 weeks (i.e. overall 48 weeks treatment duration). However, an overall 24 weeks treatment duration may be associated with a higher risk of relapse than a 48 weeks treatment duration (see section 5.1).
- **Genotypes 2 or 3:** It is recommended that all patients be treated for 24 weeks, except for HCV/HIV co-infected patients who should receive 48 weeks of treatment.
- **Genotype 4:** In general, patients infected with genotype 4 are considered harder to treat and limited study data (n=66) indicate they are compatible with a duration of treatment as for genotype 1.

## HCV/HIV Co-infection

The recommended duration of dosing for HCV/HIV co-infected patients is 48 weeks, regardless of genotype.

### *Predictability of response and non-response in HCV/HIV Co-infection*

Early virological response by Week 12, defined as a 2 log viral load decrease or undetectable levels of HCV-RNA, has been shown to be predictive for sustained response. The negative predictive value for sustained response in HCV/HIV co-infected patients treated with PegIntron in combination with ribavirin was 99 % (67/68; Study 1) (see section 5.1). A positive predictive value of 50 % (52/104; Study 1) was observed for HCV/HIV co-infected patients receiving combination therapy.

### PegIntron monotherapy

As monotherapy the PegIntron regimen is 0.5 or 1.0 microgram/kg/week. The lowest vial or pen strength available is 50 µg/0.5 ml; therefore for patients prescribed 0.5 µg/kg/week, doses must be adjusted by volume as shown in **Table 2**. For the 1.0 µg/kg dose, similar volume adjustments can be made or alternate vial strengths can be used as shown in **Table 2**. PegIntron monotherapy was not studied in HCV/HIV co-infected patients.

**Table 2- Monotherapy Dosing**

	0.5 µg/kg		1.0 µg/kg	
Body Weight (kg)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)
30-35	50*	0.15	50	0.3
36-45	50*	0.2	50	0.4
46-56	50*	0.25	50	0.5
57-72	50	0.3	80	0.4
73-88	50	0.4	80	0.5
89-106	50	0.5	100	0.5
> 106**	80	0.4	120	0.5
* Must use vial. Minimum delivery for pen is 0.3 ml.				
** For patients > 120 kg, use 80 µg/0.5 ml vial				

### Duration of treatment

For patients who exhibit virological response at Week 12, treatment should be continued for at least another three-month period (i.e., a total of six months). The decision to extend therapy to one year of treatment should be based on prognostic factors (e.g., genotype, age > 40 years, male gender, bridging fibrosis).

### Dose modification for all patients

If severe adverse reactions or laboratory abnormalities develop during treatment with PegIntron monotherapy or PegIntron in combination with ribavirin, modify the dosages of each product as appropriate, until the adverse reactions abate. As adherence might be of importance for outcome of therapy, the dose should be kept as close as possible to the recommended standard dose. Guidelines were developed in clinical trials for dose modification.

### Combination Therapy Dose Reduction Guidelines

<b>Table 2a</b> Dose modification guidelines for combination therapy (with ribavirin)			
Laboratory values	Reduce only ribavirin dose to 600 mg/day* if:	Reduce only PegIntron dose to one-half dose if:	Discontinue combination therapy if:
Haemoglobin	< 10 g/dl	-	< 8.5 g/dl
Haemoglobin in: Patients with history of stable cardiac disease	≥ 2 g/dl decrease in haemoglobin during any four week period during treatment (permanent dose reduction)		< 12 g/dl after four weeks of dose reduction
White blood cells	-	< 1.5 x 10 <sup>9</sup> /l	< 1.0 x 10 <sup>9</sup> /l
Neutrophils	-	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	-	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l
Bilirubin – direct	-	-	2.5 x ULN**

Bilirubin – indirect	> 5 mg/dl	-	> 4 mg/dl (for > 4 weeks)
Creatinine	-	-	> 2.0 mg/dl
ALT/AST	-	-	2 x baseline and > 10 x ULN**

\* Patients whose dose of ribavirin is reduced to 600 mg daily receive one 200 mg capsule in the morning and two 200 mg capsules in the evening.

\*\* Upper limit of normal

Dose reduction of PegIntron may be accomplished by either reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 2b**.

<b>Table 2b – Reduced PegIntron Dosing for Combination Therapy</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5 ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
< 40	25	50*	0.25	25
40-50	32	50	0.3	30
51-64	40	50	0.4	40
65-75	50	50	0.5	50
76-85	60	80	0.4	64
> 85	75	100	0.4	80
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### **PegIntron Monotherapy Dose Reduction Guidelines**

Dose modification guidelines for patients who use PegIntron monotherapy are shown in **Table 3a**.

<b>Table 3a Dose modification guidelines for PegIntron monotherapy</b>		
<b>Laboratory values</b>	<b>Reduce PegIntron to one-half dose if:</b>	<b>Discontinue PegIntron if:</b>
Neutrophils	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l

Dose reduction for patients who use 0.5 µg/kg PegIntron monotherapy must be accomplished by reducing the prescribed volume by one-half. The 50 µg/0.5 ml vial must be used if necessary since the pen can only deliver a minimum volume of 0.3 ml.

For patients who use 1.0 µg/kg PegIntron monotherapy, dose reduction may be accomplished by reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 3b**.

<b>Table 3b – Reduced PegIntron Dose for the 1.0 µg/kg Monotherapy Regimen</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
30-35	15	50*	0.15	15

36-45	20	50*	0.20	20
46-56	25	50*	0.25	25
57-72	32	50	0.3	30
73-89	40	50	0.4	40
90-106	50	50	0.5	50
> 106	60	80	0.4	64
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### Special populations

**Use in renal impairment:** *Monotherapy:* PegIntron should be used with caution in patients with moderate to severe renal impairment. In patients with moderate renal dysfunction (creatinine clearance 30-50 ml/minute), the starting dose of PegIntron should be reduced by 25 %. Patients with severe renal dysfunction (creatinine clearance 15-29 ml/minute) should have the starting dose of PegIntron reduced by 50 %. Data are not available for the use of PegIntron in patients with creatinine clearance < 15 ml/minute (see section 5.2). Patients with severe renal impairment, including those on hemodialysis, should be closely monitored. If renal function decreases during treatment, PegIntron therapy should be discontinued.

*Combination therapy:* Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see ribavirin SPC). When administered in combination with ribavirin, subjects with impaired renal function should be more carefully monitored with respect to the development of anaemia.

**Use in hepatic impairment:** The safety and efficacy of PegIntron therapy has not been evaluated in patients with severe hepatic dysfunction, therefore PegIntron must not be used for these patients.

**Use in the elderly (≥ 65 years of age):** There are no apparent age-related effects on the pharmacokinetics of PegIntron. Data from elderly patients treated with a single dose of PegIntron suggest no alteration in PegIntron dose is necessary based on age (see section 5.2).

**Use in patients under the age of 18 years:** There is no experience in children (see section 5.2).

### 4.3 Contraindications

- Hypersensitivity to the active substance or to any interferon or to any of the excipients;
- A history of severe pre-existing cardiac disease, including unstable or uncontrolled cardiac disease in the previous six months (see section 4.4);
- Severe, debilitating medical conditions;
- Autoimmune hepatitis or a history of autoimmune disease;
- Severe hepatic dysfunction or decompensated cirrhosis of the liver;
- Pre-existing thyroid disease unless it can be controlled with conventional treatment;
- Epilepsy and/or compromised central nervous system (CNS) function;
- Initiation of PegIntron is contraindicated in HCV/HIV patients with cirrhosis and a Child-Pugh score ≥ 6.

*Combination therapy with ribavirin:* Also see ribavirin Summary of the Product Characteristics (SPC) if PegIntron is to be administered in combination with ribavirin in patients with chronic hepatitis C.

### 4.4 Special warnings and precautions for use

**Psychiatric and Central Nervous System (CNS):** Severe CNS effects, particularly depression, suicidal ideation and attempted suicide have been observed in some patients during PegIntron therapy, and even after treatment discontinuation mainly during the 6-month follow-up period. Other CNS effects including

aggressive behaviour (sometimes directed against others), confusion and alterations of mental status have been observed with alpha interferons. Patients should be closely monitored for any signs or symptoms of psychiatric disorders. If such symptoms appear, the potential seriousness of these undesirable effects must be borne in mind by the prescribing physician and the need for adequate therapeutic management should be considered. If psychiatric symptoms persist or worsen, or suicidal ideation is identified, it is recommended that treatment with PegIntron be discontinued, and the patient followed, with psychiatric intervention as appropriate.

*Patients with existence of, or history of severe psychiatric conditions:* If treatment with peginterferon alfa-2b is judged necessary in patients with existence or history of severe psychiatric conditions, this should only be initiated after having ensured appropriate individualised diagnostic and therapeutic management of the psychiatric condition.

There is no experience with PegIntron in combination with ribavirin in patients who have relapsed after interferon alpha + ribavirin therapy.

All patients in the chronic hepatitis C studies had a liver biopsy before inclusion, but in certain cases (i.e. patients with genotype 2 and 3), treatment may be possible without histological confirmation. Current treatment guidelines should be consulted as to whether a liver biopsy is needed prior to commencing treatment.

**Acute hypersensitivity:** Acute hypersensitivity reactions (e.g., urticaria, angioedema, bronchoconstriction, anaphylaxis) have been observed rarely during interferon alfa-2b therapy. If such a reaction develops during treatment with PegIntron, discontinue treatment and institute appropriate medical therapy immediately. Transient rashes do not necessitate interruption of treatment.

**Cardiovascular system:** As with interferon alfa-2b, patients with a history of congestive heart failure, myocardial infarction and/or previous or current arrhythmic disorders, receiving PegIntron therapy require close monitoring. It is recommended that patients who have pre-existing cardiac abnormalities have electrocardiograms taken prior to and during the course of treatment. Cardiac arrhythmias (primarily supraventricular) usually respond to conventional therapy but may require discontinuation of PegIntron therapy.

More significant obtundation and coma, including cases of encephalopathy, have been observed in some patients, usually elderly, treated at higher doses for oncology indications. While these effects are generally reversible, in a few patients full resolution took up to three weeks. Very rarely, seizures have occurred with high doses of interferon alpha.

**Liver function:** As with all interferons, discontinue treatment with PegIntron in patients who develop prolongation of coagulation markers which might indicate liver decompensation.

**Fever:** While fever may be associated with the flu-like syndrome reported commonly during interferon therapy, other causes of persistent fever must be ruled out.

**Hydration:** Adequate hydration must be maintained in patients undergoing PegIntron therapy since hypotension related to fluid depletion has been seen in some patients treated with alpha interferons. Fluid replacement may be necessary.

**Pulmonary changes:** Pulmonary infiltrates, pneumonitis, and pneumonia, occasionally resulting in fatality, have been observed rarely in interferon alpha treated patients. Any patient developing fever, cough, dyspnea or other respiratory symptoms must have a chest X-ray taken. If the chest X-ray shows pulmonary infiltrates or there is evidence of pulmonary function impairment, the patient is to be monitored closely, and, if appropriate, discontinue interferon alpha. Prompt discontinuation of interferon alpha administration and treatment with corticosteroids appear to be associated with resolution of pulmonary adverse events.

**Autoimmune disease:** The development of auto-antibodies and autoimmune disorders has been reported during treatment with alpha interferons. Patients predisposed to the development of

autoimmune disorders may be at increased risk. Patients with signs or symptoms compatible with autoimmune disorders should be evaluated carefully, and the benefit-risk of continued interferon therapy should be reassessed (see also section 4.4 Thyroid changes and 4.8).

**Ocular changes:** Ophthalmologic disorders, including retinal haemorrhages, cotton wool spots, and retinal artery or vein obstruction have been reported in rare instances after treatment with alpha interferons (see section 4.8). All patients should have a baseline eye examination. Any patient complaining of ocular symptoms, including loss of visual acuity or visual field must have a prompt and complete eye examination. Periodic visual examinations are recommended during PegIntron therapy, particularly in patients with disorders that may be associated with retinopathy, such as diabetes mellitus or hypertension. Discontinuation of PegIntron should be considered in patients who develop new or worsening ophthalmological disorders.

**Thyroid changes:** Infrequently, patients treated for chronic hepatitis C with interferon alpha have developed thyroid abnormalities, either hypothyroidism or hyperthyroidism. Determine thyroid stimulating hormone (TSH) levels if, during the course of therapy, a patient develops symptoms consistent with possible thyroid dysfunction. In the presence of thyroid dysfunction, PegIntron treatment may be continued if TSH levels can be maintained in the normal range by medication.

**Metabolic disturbances:** Hypertriglyceridemia and aggravation of hypertriglyceridemia, sometimes severe, have been observed. Monitoring of lipid levels is, therefore, recommended.

#### **HCV/HIV Co-infection**

Mitochondrial toxicity and lactic acidosis:

Patients co-infected with HIV and receiving Highly Active Anti-Retroviral Therapy (HAART) may be at increased risk of developing lactic acidosis. Caution should be used when adding PegIntron and ribavirin to HAART therapy (see ribavirin SPC).

Moreover, patients treated with PegIntron and ribavirin combination therapy and zidovudine are at increased risk of developing anaemia.

Hepatic decompensation in HCV/HIV co-infected patients with advanced cirrhosis:

Co-infected patients with advanced cirrhosis receiving HAART may be at increased risk of hepatic decompensation and death. Adding treatment with alfa interferons alone or in combination with ribavirin may increase the risk in this patient subset. Other baseline factors in co-infected patients that may be associated with a higher risk of hepatic decompensation include treatment with didanosine and elevated bilirubin serum concentration.

Co-infected patients receiving both antiretroviral (ARV) and anti-hepatitis treatment should be closely monitored, assessing their Child-Pugh score during treatment. Patients progressing to hepatic decompensation should have their anti-hepatitis treatment immediately discontinued and the ARV treatment reassessed.

Haematological abnormalities in HCV/HIV co-infected patients:

HCV/HIV co-infected patients receiving peginterferon alfa-2b/ribavirin treatment and HAART may be at increased risk to develop haematological abnormalities (as neutropenia, thrombocytopenia and anaemia) compared to HCV mono-infected patients. Although, the majority of them could be managed by dose reduction, close monitoring of haematological parameters should be undertaken in this population of patients (see section 4.2 and below “Laboratory tests” and section 4.8).

Patients with low CD4 counts:

In patients co-infected with HCV/HIV, limited efficacy and safety data (N = 25) are available in subjects with CD4 counts less than 200 cells/ $\mu$ l. Caution is therefore warranted in the treatment of patients with low CD4 counts.

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron and ribavirin.



**Dental and periodontal disorders:** Dental and periodontal disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and mucous membranes of the mouth during long-term treatment with the combination of PegIntron and ribavirin. Patients should brush their teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If this reaction occurs, they should be advised to rinse out their mouth thoroughly afterwards.

**Organ transplant recipients:** The safety and efficacy of PegIntron alone or in combination with ribavirin for the treatment of hepatitis C in liver or other organ transplant recipients have not been studied. Preliminary data indicates that interferon alpha therapy may be associated with an increased rate of kidney graft rejection. Liver graft rejection has also been reported.

**Other:** Due to reports of interferon alpha exacerbating pre-existing psoriatic disease and sarcoidosis, use of PegIntron in patients with psoriasis or sarcoidosis is recommended only if the potential benefit justifies the potential risk.

**Laboratory tests:** Standard haematologic tests, blood chemistry and a test of thyroid function must be conducted in all patients prior to initiating therapy. Acceptable baseline values that may be considered as a guideline prior to initiation of PegIntron therapy are:

- |                    |                              |
|--------------------|------------------------------|
| • Platelets        | $\geq 100,000/\text{mm}^3$   |
| • Neutrophil count | $\geq 1,500/\text{mm}^3$     |
| • TSH level        | must be within normal limits |

Laboratory evaluations are to be conducted at weeks 2 and 4 of therapy, and periodically thereafter as clinically appropriate.

**Important information about some of the ingredients of PegIntron:**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

Patients with rare hereditary problems of fructose intolerance, glucose galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Results from a multiple-dose probe study assessing P450 substrates in chronic hepatitis C patients receiving once weekly PegIntron (1.5  $\mu\text{g}/\text{kg}$ ) for 4 weeks demonstrated an increase in activity of CYP2D6 and CYP2C8/9. No change in activity of CYP1A2, CYP3A4, or N-acetyltransferase was observed.

Caution should be used when administering peginterferon alfa-2b with medicines metabolised by CYP2D6 and CYP2C8/9, especially those with narrow therapeutic window, such as warfarin and phenytoin (CYP2C9) and flecainide (CYP2D6).

These findings may partly relate to improved metabolic capacity due to reduced hepatic inflammation in patients undergoing treatment with PegIntron. Caution is therefore advised when PegIntron treatment is initiated for chronic hepatitis in patients treated with medication with a narrow therapeutic window and sensitive to mild metabolic impairment of the liver.

No pharmacokinetic interactions were noted between PegIntron and ribavirin in a multiple-dose pharmacokinetic study.

**Methadone:** In patients with chronic hepatitis C that were on stable methadone maintenance therapy and naïve to peginterferon alfa-2b, addition of 1.5 microgram/kg/week of PegIntron subcutaneously for 4 weeks increased R-methadone AUC by approximately 15 % (95 % CI for AUC ratio estimate 103 – 128 %). The clinical significance of this finding is unknown; however, patients should be

monitored for signs and symptoms of increased sedative effect, as well as respiratory depression. Especially in patients on a high dose of methadone, the risk for QTc prolongation should be considered.

#### *HCV/HIV Co-infection*

Nucleoside analogs: ribavirin has been shown *in vitro* to inhibit phosphorylation of zidovudine and stavudine. The clinical significance of these finding is unknown. However, these *in vitro* findings raise the possibility that concurrent use of ribavirin with either zidovudine or stavudine might lead to increased HIV plasma viraemia. Therefore, it is recommended that plasma HIV RNA levels be closely monitored in patients treated with ribavirin concurrently with either of these two agents. If HIV RNA levels increase, the use of ribavirin concomitantly with reverse transcriptase inhibitors must be reviewed (see ribavirin SPC).

Use of nucleoside analogs, alone or in combination with other nucleosides, has resulted in lactic acidosis. Pharmacologically, ribavirin increases phosphorylated metabolites of purine nucleosides *in vitro*. This activity could potentiate the risk of lactic acidosis induced by purine nucleoside analogs (e.g. didanosine or abacavir). Co-administration of ribavirin and didanosine is not recommended. Reports of mitochondrial toxicity, in particular lactic acidosis and pancreatitis, of which some fatal, have been reported (see ribavirin SPC).

### **4.6 Pregnancy and lactation**

PegIntron is recommended for use in fertile women only when they are using effective contraception during the treatment.

There are no adequate data from the use of interferon alfa-2b in pregnant women.. Studies in animals have shown reproductive toxicity (see section 5.3). Interferon alfa-2b has been shown to be abortifacient in primates. PegIntron is likely to also cause this effect.

The potential risk in humans is unknown. PegIntron is to be used during pregnancy only if the potential benefit justifies the potential risk to the foetus

**Lactation:** It is not known whether the components of this medicinal product are excreted in human milk. Because of the potential for adverse reactions in breast-fed infants, breast-feeding should be discontinued prior to initiation of treatment.

#### *Combination therapy with ribavirin:*

Ribavirin causes serious birth defects when administered during pregnancy. Ribavirin therapy is contraindicated in women who are pregnant. Extreme care must be taken to avoid pregnancy in female patients or in partners of male patients taking PegIntron in combination with ribavirin. Females of childbearing potential and their partners must each use an effective contraceptive during treatment and for 4 months after treatment has been concluded. Male patients and their female partners must each use an effective contraceptive during treatment and for 7 months after treatment has been concluded (see Ribavirin SPC).

### **4.7 Effects on ability to drive and use machines**

Patients who develop fatigue, somnolence or confusion during treatment with PegIntron are cautioned to avoid driving or operating machinery.

### **4.8 Undesirable effects**

The safety of PegIntron is evaluated from data from two clinical trials: one with PegIntron monotherapy, one with PegIntron in combination with ribavirin. In both cases, patients were treated for one year.

**Table 4** describes the regimens and patient exposure for one year of treatment in patients with no previous exposure to interferon (interferon-naïve patients). Because of a significant overlap in the

pattern of undesirable effects with PegIntron monotherapy, groups of patients have been brought together in **Table 5** to show the pattern of reported effects for all monotherapy groups.

<b>Table 4</b> Regimens and patient exposure		
<b>Treatment</b>	<b>Regimen</b>	<b>Number of patients treated for one year</b>
PegIntron + ribavirin	PegIntron (1.5 micrograms/kg/week) + ribavirin (> 10.6 mg/kg/day)	188
Interferon alfa-2b + ribavirin	Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day)	505
PegIntron monotherapy	PegIntron (0.5 microgram/kg/week)	315
	PegIntron (1.0 microgram/kg/week)	297
	PegIntron (1.5 micrograms/kg/week)	304

<b>Table 5</b> Undesirable effects very commonly reported in clinical trials > 10 % of patients in PegIntron + ribavirin group Very common (> 1/10)			
	<b>PegIntron + ribavirin</b>	<b>Interferon alfa- 2b + ribavirin</b>	<b>PegIntron monotherapy</b>
<b>Infections and infestations</b>			
Infection viral	10 %	5 %	4-5 %
<b>Metabolism and nutrition disorders</b>			
Weight decrease	30 %	19 %	8-18 %
<b>Psychiatric disorders</b>			
Depression	34 %	32 %	26 %
Irritability	32 %	34 %	19 %
Insomnia	37 %	41 %	16-19 %
Anxiety	14 %	14 %	8 %
Concentration impaired	18 %	21 %	9-10 %
Emotional lability	11 %	10 %	5 %
<b>Nervous system disorders</b>			
Headache	58 %	57 %	57-63 %
Mouth dry	10 %	8 %	4-8 %
<b>Respiratory, thoracic and mediastinal disorders</b>			
Pharyngitis	10 %	7 %	3 %
Coughing	14 %	11 %	4 %
Dyspnea	26 %	22 %	5 %
<b>Gastrointestinal disorders</b>			
Nausea	43 %	31 %	20-23 %
Anorexia	35 %	26 %	10-25 %
Diarrhoea	20 %	13 %	14-17 %
Abdominal pain	12 %	9 %	11 %
Vomiting	16 %	10 %	4-7 %
<b>Skin and subcutaneous tissue disorders</b>			
Alopecia	45 %	32 %	20-34 %
Pruritus	27 %	27 %	7-9 %
Skin dry	23 %	21 %	6-9 %
Rash	21 %	21 %	5-7 %
<b>Musculoskeletal and connective tissue disorders</b>			
Myalgia	49 %	49 %	46-60 %
Arthralgia	31 %	26 %	23-28 %
Musculoskeletal pain	15 %	11 %	11-13 %
<b>General disorders and administration site conditions</b>			
Injection site inflammation	20 %	17 %	39-44 %
Injection site reaction	54 %	36 %	7-9 %
Fatigue	56 %	59 %	43 %
Rigors	42 %	40 %	33-43 %
Fever	39 %	32 %	29-43 %
Flu-like symptoms	21 %	23 %	18-25 %
Asthenia	28 %	17 %	12-14 %
Dizziness	17 %	16 %	7-12 %

<b>Table 6</b> Undesirable effects commonly reported in clinical trials in patients treated with PegIntron + ribavirin or PegIntron monotherapy Common (> 1/100, < 1/10)		
<b>Body system</b>	<b>5-10%</b>	<b>1-&lt;5%</b>
Infections and infestations		otitis media, fungal infection, bacterial infection, herpes simplex
Blood and lymphatic system disorders	Anaemia, leukopaenia	Thrombocytopenia, lymphadenopathy
Endocrine disorders	hypothyroidism	hyperthyroidism
Metabolism and nutrition disorders		hyperuricemia, hypocalcemia, thirst
Psychiatric disorders	agitation, nervousness	aggressive behaviour, somnolence, behavior disorder, apathy, appetite increased, sleep disorder, dreaming abnormal, decreased libido
Nervous system disorders	paresthesia, increased sweating	hypoesthesia, hyperaesthesia, hypertonia, confusion, tremor, vertigo, migraine, ataxia, neuralgia
Eye disorders		blurred vision, conjunctivitis, lacrimal gland disorder, eye pain
Cardiac disorders		tachycardia, palpitation
Vascular disorders		hypotension, hypertension, syncope, flushing
Ear and labyrinth disorders		tinnitus, hearing impairment/loss
Respiratory, thoracic and mediastinal disorders		nonproductive cough, rhinitis, sinusitis, bronchitis, respiratory disorder, nasal congestion, rhinorrhea, dysphonia, epistaxis
Gastrointestinal disorders	dyspepsia	constipation, taste perversion, loose stools, stomatitis, ulcerative stomatitis, gingival bleeding, glossitis, flatulence, hemorrhoids, gastroesophageal reflux, gingivitis, dehydration
Hepatobiliary disorders		hepatomegaly, hyperbilirubinemia
Skin and subcutaneous tissue disorders		erythematous rash, eczema, photosensitivity reaction, maculopapular rash, abnormal hair texture, acne, dermatitis, furunculosis, nail disorder, psoriasis, urticaria, erythema, face or peripheral oedema
Musculoskeletal and connective tissue disorders		arthritis
Renal and urinary disorders		micturition frequency, urine abnormal
Reproductive system and breast disorders	menstrual disorder, menorrhagia	ovarian disorder, vaginal disorder, sexual dysfunction (not specified), impotence, breast pain, amenorrhoea, prostatitis
General disorders and administration site conditions	RUQ pain, malaise, chest pain	injection site pain

Most cases of neutropaenia and thrombocytopaenia were mild (WHO grades 1 or 2). There were some cases of more severe neutropenia in patients treated with the recommended doses of PegIntron in combination with ribavirin (WHO grade 3: 39 of 186 [21 %]; and WHO grade 4: 13 of 186 [7 %]).

In a clinical trial, approximately 1.2 % of patients treated with PegIntron or interferon alfa-2b in combination with ribavirin reported life-threatening psychiatric events during treatment. These events included suicidal ideation and attempted suicide (see section 4.4). Following marketing, psychosis and hallucination have been reported rarely.

Rarely ( $> 1/10,000$ ,  $< 1/1,000$ ) or very rarely ( $< 1/10,000$ ) reported events with interferon alfa-2b, including PegIntron, include:

**Blood and lymphatic system disorders:**

Very rarely interferon alfa-2b or PegIntron used alone or in combination with ribavirin may be associated with aplastic anaemia. Pure red cell aplasia has been reported.

**Immune system disorders:**

very rarely: sarcoidosis or exacerbation of sarcoidosis

**Endocrine disorders:**

rarely: diabetes

**Nervous system disorders:**

rarely: seizure, peripheral neuropathy

very rarely: cerebrovascular ischaemia, cerebrovascular haemorrhage, encephalopathy

**Cardiac disorders:**

rarely: arrhythmia

very rarely: cardiac ischaemia, myocardial infarction

**Respiratory, thoracic and mediastinal disorders:**

very rarely: interstitial lung disease

**Gastrointestinal disorders:**

rarely: pancreatitis

very rarely: ulcerative and ischaemic colitis

**Skin and subcutaneous tissue disorders:**

very rarely: erythema multiforme, Stevens Johnson syndrome, toxic epidermal necrolysis, injection site necrosis

**Musculoskeletal and connective tissue disorders:**

rarely: rhabdomyolysis, myositis

**Renal and urinary disorders:**

rarely: renal insufficiency and renal failure

Cardiovascular (CVS) adverse events, particularly arrhythmia, appeared to be correlated mostly with pre-existing CVS disease and prior therapy with cardiotoxic agents (see section 4.4). Cardiomyopathy, that may be reversible upon discontinuation of interferon alpha, has been reported rarely in patients without prior evidence of cardiac disease.

Ophthalmological disorders that have been reported rarely with alpha interferons include retinopathies (including macular oedema), retinal haemorrhages, retinal artery or vein obstruction, cotton wool spots, loss of visual acuity or visual field, optic neuritis, and papilloedema (see section 4.4).

A wide variety of autoimmune and immune-mediated disorders have been reported with alpha interferons including thyroid disorders, systemic lupus erythematosus, rheumatoid arthritis (new or aggravated), idiopathic and thrombotic thrombocytopenic purpura, vasculitis, neuropathies including mononeuropathies (see also section 4.4, **Autoimmune disorders**).

**HCV/HIV co-infected patients**

For HCV/HIV co-infected patients receiving PegIntron in combination with ribavirin, other undesirable effects (that were not reported in mono-infected patients) which have been reported in the larger studies with a frequency  $> 5\%$  were: oral candidiasis (14 %), lipodystrophy acquired (13 %), CD4 lymphocytes decreased (8 %), appetite decreased (8 %), gamma-glutamyltransferase increased (9 %), back pain (5 %), blood amylase increased (6 %), blood lactic acid increased (5 %), cytolytic hepatitis (6 %), lipase increased (6 %) and pain in limb (6 %).

*Mitochondrial toxicity:*

Mitochondrial toxicity and lactic acidosis have been reported in HIV-positive patients receiving NRTI

regimen and associated ribavirin for co-HCV infection (see section 4.4).

*Laboratory values for HCV/HIV co-infected patients:*

Although haematological toxicities of neutropenia, thrombocytopenia and anaemia occurred more frequently in HCV/HIV co-infected patients, the majority could be managed by dose modification and rarely required premature discontinuation of treatment (see section 4.4). Haematological abnormalities were more frequently reported in patients receiving PegIntron in combination with ribavirin when compared to patients receiving interferon alfa-2b in combination with ribavirin. In Study 1 (see section 5.1), decrease in absolute neutrophil count levels below 500 cells/mm<sup>3</sup> was observed in 4 % (8/194) of patients and decrease in platelets below 50,000/mm<sup>3</sup> was observed in 4 % (8/194) of patients receiving PegIntron in combination with ribavirin. Anaemia (hemoglobin < 9.4g/dl) was reported in 12% (23/194) of patients treated with PegIntron in combination with ribavirin.

*CD4 lymphocytes decrease:*

Treatment with PegIntron in combination with ribavirin was associated with decreases in absolute CD4+ cell counts within the first 4 weeks without a reduction in CD4+ cell percentage. The decrease in CD4+ cell counts was reversible upon dose reduction or cessation of therapy. The use of PegIntron in combination with ribavirin had no observable negative impact on the control of HIV viraemia during therapy or follow-up. Limited safety data (N= 25) are available in co-infected patients with CD4+ cell counts < 200/μl (see section 4.4).

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron in combination with ribavirin.

## **4.9 Overdose**

In clinical trials, cases of accidental overdose, at never more than twice the prescribed dose, were reported. There were no serious reactions. Undesirable effects resolved during continued administration of PegIntron.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Immunostimulants, Cytokines and immunomodulators, Interferons, Peginterferon alfa-2b, ATC code: L03A B10.

Recombinant interferon alfa-2b is covalently conjugated with monomethoxy polyethylene glycol at an average degree of substitution of 1 mole of polymer/mole of protein. The average molecular mass is approximately 31,300 daltons of which the protein moiety constitutes approximately 19,300.

#### **Interferon alfa-2b**

*In vitro* and *in vivo* studies suggest that the biological activity of PegIntron is derived from its interferon alfa-2b moiety.

Interferons exert their cellular activities by binding to specific membrane receptors on the cell surface. Studies with other interferons have demonstrated species specificity. However, certain monkey species, e.g., Rhesus monkeys, are susceptible to pharmacodynamic stimulation upon exposure to human type 1 interferons.

Once bound to the cell membrane, interferon initiates a complex sequence of intracellular events that include the induction of certain enzymes. It is thought that this process, at least in part, is responsible for the various cellular responses to interferon, including inhibition of virus replication in virus-infected cells, suppression of cell proliferation and such immunomodulating activities as enhancement of the phagocytic activity of macrophages and augmentation of the specific cytotoxicity of

lymphocytes for target cells. Any or all of these activities may contribute to interferon's therapeutic effects.

Recombinant interferon alfa-2b also inhibits viral replication *in vitro* and *in vivo*. Although the exact antiviral mode of action of recombinant interferon alfa-2b is unknown, it appears to alter the host cell metabolism. This action inhibits viral replication or if replication occurs, the progeny virions are unable to leave the cell.

### **PegIntron**

PegIntron pharmacodynamics were assessed in a rising single-dose trial in healthy subjects by examining changes in oral temperature, concentrations of effector proteins such as serum neopterin and 2'5'-oligoadenylate synthetase (2'5'-OAS), as well as white cell and neutrophil counts. Subjects treated with PegIntron showed mild dose-related elevations in body temperature. Following single doses of PegIntron between 0.25 and 2.0 micrograms/kg/week, serum neopterin concentration was increased in a dose-related manner. Neutrophil and white cell count reductions at the end of week 4 correlated with the dose of PegIntron.

### **PegIntron clinical trials**

Two pivotal trials have been conducted, one (C/I97-010) with PegIntron monotherapy; the other (C/I98-580) with PegIntron in combination with ribavirin. Eligible patients for these trials had chronic hepatitis C confirmed by a positive HCV-RNA polymerase chain reaction (PCR) assay (> 30 IU/ml), a liver biopsy consistent with a histological diagnosis of chronic hepatitis with no other cause for the chronic hepatitis, and abnormal serum ALT.

In the PegIntron monotherapy trial, a total of 916 naïve chronic hepatitis C patients were treated with PegIntron (0.5, 1.0 or 1.5 micrograms/kg/week) for one year with a follow-up period of six months. In addition, 303 patients received interferon alfa-2b (3 million International Units [MIU] three times a week) as a comparator. This study showed that PegIntron was superior to interferon alfa-2b (**Table 8**).

In the PegIntron combination trial, 1,530 naïve patients were treated for one year with one of the following combination regimens:

- PegIntron (1.5 micrograms/kg/week) + ribavirin (800 mg/day), (n = 511).
- PegIntron (1.5 micrograms/kg/week for one month followed by 0.5 microgram/kg/week for 11 months) + ribavirin (1,000/1,200 mg/day), (n = 514).
- Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day) (n = 505).

In this trial, the combination of PegIntron (1.5 micrograms/kg/week) and ribavirin was significantly more effective than the combination of interferon alfa-2b and ribavirin (**Table 7**), particularly in patients infected with Genotype 1 (**Table 8**). Sustained response was assessed by the response rate six months after the cessation of treatment.

HCV genotype and baseline virus load are prognostic factors which are known to affect response rates. However, response rates in this trial were shown to be dependent also on the dose of ribavirin administered in combination with PegIntron or interferon alfa-2b. In those patients that received > 10.6 mg/kg ribavirin (800 mg dose in typical 75 kg patient), regardless of genotype or viral load, response rates were significantly higher than in those patients that received ≤ 10.6 mg/kg ribavirin (**Table 8**), while response rates in patients that received > 13.2 mg/kg ribavirin were even higher.

<b>Table 7</b> Sustained virological response (% patients HCV negative)							
	<b>PegIntron monotherapy</b>				<b>PegIntron + ribavirin</b>		
<b>Treatment regimen</b>	<b>P 1.5</b>	<b>P 1.0</b>	<b>P 0.5</b>	<b>I</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>Number of patients</b>	304	297	315	303	511	514	505
<b>Response at end of treatment</b>	49 %	41 %	33 %	24 %	65 %	56 %	54 %



Sustained response	23 %*	25 %	18 %	12 %	54 %**	47 %	47 %
P 1.5	PegIntron 1.5 micrograms/kg						
P 1.0	PegIntron 1.0 microgram/kg						
P 0.5	PegIntron 0.5 microgram/kg						
I	Interferon alfa-2b 3 MIU						
P 1.5/R	PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)						
P 0.5/R	PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)						
I/R	Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)						
*	p < 0.001 P 1.5 vs. I						
**	p = 0.0143 P 1.5/R vs. I/R						

<b>Table 8</b> Sustained response rates with PegIntron + ribavirin (by ribavirin dose, genotype and viral load)				
HCV Genotype	Ribavirin dose (mg/kg)	P 1.5/R	P 0.5/R	I/R
<b>All Genotypes</b>	<b>All</b>	<b>54 %</b>	<b>47 %</b>	<b>47 %</b>
	≤ 10.6	50 %	41 %	27 %
	> 10.6	61 %	48 %	47 %
<b>Genotype 1</b>	<b>All</b>	<b>42 %</b>	<b>34 %</b>	<b>33 %</b>
	≤ 10.6	38 %	25 %	20 %
	> 10.6	48 %	34 %	34 %
Genotype 1 ≤ 600,000 IU/ml	All	73 %	51 %	45 %
	≤ 10.6	74 %	25 %	33 %
	> 10.6	71 %	52 %	45 %
Genotype 1 > 600,000 IU/ml	All	30 %	27 %	29 %
	≤ 10.6	27 %	25 %	17 %
	> 10.6	37 %	27 %	29 %
<b>Genotype 2/3</b>	<b>All</b>	<b>82 %</b>	<b>80 %</b>	<b>79 %</b>
	≤ 10.6	79 %	73 %	50 %
	> 10.6	88 %	80 %	80 %

P 1.5/R	PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)
P 0.5/R	PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)
I/R	Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)

In the PegIntron monotherapy study, the Quality of Life was generally less affected by 0.5 microgram/kg of PegIntron than by either 1.0 microgram/kg of PegIntron once weekly or 3 MIU of interferon alfa-2b three times a week.

In a separate trial, 224 patients with genotype 2 or 3 received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with ribavirin 800 mg –1,400 mg p.o. for 6 months (based on body weight, only three patients weighing > 105 kg, received the 1,400 mg dose, which has not yet been validated) (**Table 9**). Twenty-four % had bridging fibrosis or cirrhosis (Knodell 3/4).

**Table 9. Virologic Response at End of Treatment, Sustained Virologic Response and Relapse by HCV Genotype and Viral Load\***

	PegIntron 1.5 µg/kg once weekly plus Ribavirin 800-1,400 mg/day		
	End of Treatment Response	Sustained Virologic Response	Relapse
All Subjects	94 % (211/224)	81 % (182/224)	12 % (27/224)
HCV 2 ≤ 600,000 IU/ml > 600,000 IU/ml	100 % (42/42)	93 % (39/42)	7 % (3/42)
	100 % (20/20)	95 % (19/20)	5 % (1/20)
	100 % (22/22)	91 % (20/22)	9 % (2/22)
HCV 3	93 % (169/182)	79 % (143/182)	14 %

≤ 600,000 IU/ml	93 % (92/99)	86 % (85/99)	(24/166) 8 % (7/91)
> 600,000 IU/ml	93 % (77/83)	70 % (58/83)	23 % (17/75)

\* Any subject with an undetectable HCV-RNA level at the Follow-Up Week 12 visit and missing data at the Follow-Up Week 24 visit was considered a sustained responder. Any subject with missing data in and after the Follow-Up Week 12 window was considered to be a non-responder at Week 24 of follow-up.

The 6 month treatment duration in this trial was better tolerated than one year of treatment in the pivotal combination trial; for discontinuation 5 % vs. 14 %, for dose modification 18 % vs. 49 %.

In a non-comparative trial, 235 patients with genotype 1 and low viral load (< 600,000 IU/ml) received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with weight adjusted ribavirin. The overall sustained response rate after a 24-week treatment duration was 50 %. Forty-one percent of subjects (97/235) had nondetectable plasma HCV-RNA levels at Week 4 and Week 24 of therapy. In this subgroup, there was a 92 % (89/97) sustained virological response rate. The high sustained response rate in this subgroup of patients was identified in an interim analysis (n=49) and prospectively confirmed (n=48).

Limited historical data indicate that treatment for 48 weeks might be associated with a higher sustained response rate (11/11) and with a lower risk of relapse (0/11 as compared to 7/96 following 24 weeks of treatment).

#### Predictability of sustained virological response

Virological response by week 12, defined as a 2-log viral load decrease or undetectable levels of HCV-RNA has been shown to be predictive for sustained response (**Table 10**).

<b>Table 10</b> Predictability of sustained response by viral response at week 12 and genotype*				
Treatment	Genotype	Viral response at week 12	Sustained response	Negative predictive value
PegIntron 1.5 + ribavirin (> 10.6 mg/kg) 48-week treatment	1	Yes 75 % (82/110)	71 % (58/82)	----
		No 25 % (28/110)	0 % (0/28)	100 %
PegIntron 1.5 + ribavirin 800-1,400 mg 24-week treatment	2 and 3	Yes 99 % (213/215)	83 % (177/213)	----
		No 1 % (2/215)	50 % (1/2)	50 %

\* reflects patients with 12 week data available

The negative predictive value for sustained response in patients treated with PegIntron in monotherapy was 98 %.

#### HCV/HIV Co-infected patients

Two trials have been conducted in patients co-infected with HIV and HCV. The response to treatment in both of these trials is presented in **Table 11**. Study 1 (RIBAVIC; P01017) was a randomized, multicentre study which enrolled 412 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (1.5 µg/kg/week) plus ribavirin (800 mg/day) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800 mg/day) for 48 weeks with a follow-up period of 6 months. Study 2 (P02080) was a randomized, single centre study that enrolled 95 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (100 or 150 µg /week based

on weight) plus ribavirin (800-1,200 mg/day based on weight) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800-1,200 mg/day based on weight). The duration of therapy was 48 weeks with a follow-up period of 6 months except for patients infected with genotypes 2 or 3 and viral load < 800,000 IU/ml (Amplicor) who were treated for 24 weeks with a 6-month follow-up period.

<b>Table 11</b> Sustained virological response based on genotype after PegIntron in combination with Ribavirin in HCV/HIV Co-infected patients						
	<b>Study 1<sup>1</sup></b>			<b>Study 2<sup>2</sup></b>		
	PegIntron (1.5 µg/kg/week) + ribavirin (800 mg)	Interferon alfa-2b (3 MIU TIW) + ribavirin (800 mg)	p value <sup>a</sup>	PegIntron (100 or 150 <sup>c</sup> µg/week) + ribavirin (800-1,200 mg) <sup>d</sup>	Interferon alfa-2b (3 MIU TIW) + ribavirin (800-1,200 mg) <sup>d</sup>	p value <sup>b</sup>
All	27 % (56/205)	20 % (41/205)	0.047	44 % (23/52)	21 % (9/43)	0.017
Genotype 1, 4	17 % (21/125)	6 % (8/129)	0.006	38 % (12/32)	7 % (2/27)	0.007
Genotype 2, 3	44 % (35/80)	43 % (33/76)	0.88	53 % (10/19)	47 % (7/15)	0.730

MIU = million international units; TIW = three times a week.

a: p value based on Cochran-Mantel Haenszel Chi square test.

b: p value based on chi-square test.

c: subjects < 75 kg received 100 µg/week PegIntron and subjects ≥ 75 kg received 150 µg/week PegIntron.

d: ribavirin dosing was 800 mg for patients < 60 kg, 1,000 mg for patients 60-75 kg, and 1,200 mg for patients > 75 kg.

<sup>1</sup>Carrato F, Bani-Sadir F, Pol S et al. JAMA 2004; 292(23): 2839-2848.

<sup>2</sup>Laguno M, Murillas J, Blanco J et al. AIDS 2004; 18(13): F27-F36.

### *Histological response*

Liver biopsies were obtained before and after treatment in Study 1 and were available for 210 of the 412 subjects (51 %). Both the Metavir score and Ishak grade decreased among subjects treated with PegIntron in combination with ribavirin. This decline was significant among responders (-0.3 for Metavir and -1.2 for Ishak) and stable (-0.1 for Metavir and -0.2 for Ishak) among non-responders. In terms of activity, about one-third of sustained responders showed improvement and none showed worsening. There was no improvement in terms of fibrosis observed in this study. Steatosis was significantly improved in patients infected with HCV Genotype 3.

## **5.2 Pharmacokinetic properties**

PegIntron is a well characterized polyethylene glycol-modified (“pegylated”) derivative of interferon alfa-2b and is predominantly composed of monopegylated species. The plasma half-life of PegIntron is prolonged compared with non-pegylated interferon alfa-2b. PegIntron has a potential to depegylate to free interferon alfa-2b. The biologic activity of the pegylated isomers is qualitatively similar, but weaker than free interferon alfa-2b.

Following subcutaneous administration, maximal serum concentrations occur between 15-44 hours post-dose, and are sustained for up to 48-72 hours post-dose.

PegIntron C<sub>max</sub> and AUC measurements increase in a dose-related manner. Mean apparent volume of distribution is 0.99 l/kg.

Upon multiple dosing, there is an accumulation of immunoreactive interferons. There is, however, only a modest increase in biologic activity as measured by a bioassay.

Mean (SD) PegIntron elimination half-life is approximately 40 hours (13.3 hours), with apparent clearance of 22.0 ml/hr·kg. The mechanisms involved in clearance of interferons in man have not yet been

fully elucidated. However, renal elimination may account for a minority (approximately 30 %) of PegIntron apparent clearance.

**Renal function:** Renal clearance appears to account for 30 % of total clearance of PegIntron. In a single dose study (1.0 microgram/kg) in patients with impaired renal function,  $C_{max}$ , AUC, and half-life increased in relation to the degree of renal impairment.

Following multiple dosing of PegIntron (1.0 microgram/kg subcutaneously administered every week for four weeks) the clearance of PegIntron is reduced by a mean of 17 % in patients with moderate renal impairment (creatinine clearance 30-49 ml/minute) and by a mean of 44 % in patients with severe renal impairment (creatinine clearance 15-29 ml/minute) compared to subjects with normal renal function. Based on single dose data, clearance was similar in patients with severe renal impairment not on dialysis and in patients who were receiving hemodialysis. The dose of PegIntron for monotherapy should be reduced in patients with moderate or severe renal impairment (see sections 4.2 and 4.4). Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see section 4.3).

Because of marked inter-subject variability in interferon pharmacokinetics, it is recommended that patients with severe renal impairment be closely monitored during treatment with PegIntron (see section 4.2)

**Hepatic function:** The pharmacokinetics of PegIntron have not been evaluated in patients with severe hepatic dysfunction.

**Elderly patients  $\geq 65$  years of age:** The pharmacokinetics of PegIntron following a single subcutaneous dose of 1.0 microgram/kg were not affected by age. The data suggest that no alteration in PegIntron dosage is necessary based on advancing age.

**Patients under the age of 18 years:** Specific pharmacokinetic evaluations have not been performed on these patients. PegIntron is indicated for the treatment of chronic hepatitis C only in patients 18 years of age or older.

**Interferon neutralising factors:** Interferon neutralising factor assays were performed on serum samples of patients who received PegIntron in the clinical trial. Interferon neutralising factors are antibodies which neutralise the antiviral activity of interferon. The clinical incidence of neutralising factors in patients who received PegIntron 0.5 micrograms/kg is 1.1 %.

### 5.3 Preclinical safety data

**PegIntron:** Adverse events not observed in clinical trials were not seen in toxicity studies in monkeys. These studies were limited to four weeks due to the appearance of anti-interferon antibodies in most monkeys.

Reproduction studies of PegIntron have not been performed. Interferon alfa-2b has been shown to be an abortifacient in primates. PegIntron is likely to also cause this effect. Effects on fertility have not been determined. It is not known whether the components of this medicinal product are excreted into experimental animal or human milk (see section 4.6 for relevant human data on pregnancy and lactation). PegIntron showed no genotoxic potential.

The relative non-toxicity of monomethoxy-polyethylene glycol (mPEG), which is liberated from PegIntron by metabolism *in vivo* has been demonstrated in preclinical acute and subchronic toxicity studies in rodents and monkeys, standard embryo-foetal development studies and in *in vitro* mutagenicity assays.

**PegIntron plus ribavirin:** When used in combination with ribavirin, PegIntron did not cause any effects not previously seen with either active substance alone. The major treatment-related change was

a reversible, mild to moderate anaemia, the severity of which was greater than that produced by either active substance alone.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Powder for solution for injection:

Disodium phosphate, anhydrous,  
Sodium dihydrogen phosphate dihydrate,  
Sucrose,  
Polysorbate 80.

Solvent for parenteral use:

Water for injections.

Deliverable volume from pen = 0.5 ml.

### **6.2 Incompatibilities**

This medicinal product should only be reconstituted with the solvent provided (see section 6.6). In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

3 years

After reconstitution:

- Chemical and physical in-use stability has been demonstrated for 24 hours at 2°C - 8°C.
- From a microbiological point of view, the product is to be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C - 8°C.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C - 8°C). Do not freeze.

### **6.5 Nature and contents of container**

The powder and solvent are both contained in a two-chamber cartridge, Type I flint glass, separated by a bromobutyl rubber plunger. The cartridge is sealed at one end with a polypropylene cap containing a bromobutyl rubber liner and at the other end by a bromobutyl rubber plunger.

PegIntron 120 micrograms is supplied as:

- 1 pen containing powder and solvent for solution for injection, 1 injection needle and 2 cleansing swabs;
- 4 pens containing powder and solvent for solution for injection, 4 injection needles and 8 cleansing swabs;
- 6 pens containing powder and solvent for solution for injection, 6 injection needles and 12 cleansing swabs;
- 12 pens containing powder and solvent for solution for injection, 12 injection needles and 24 cleansing swabs.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

PegIntron pre-filled pen contains a powder of peginterferon alfa-2b and a solvent for solution at a strength of 120 micrograms for single use. Each pen is reconstituted with the solvent provided in the two-chamber cartridge (water for injections) for administration of up to 0.5 ml of solution. A small volume is lost during preparation of PegIntron for injection when the dose is measured and injected. Therefore, each pen contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection. The reconstituted solution has a concentration of 120 micrograms in 0.5 ml.

PegIntron is injected subcutaneously after reconstituting the powder as instructed, attaching an injection needle and setting the prescribed dose. A complete and illustrated set of instructions is provided in the Annex to the Package Leaflet.

Remove PegIntron pre-filled pen from the refrigerator before administration to allow the solvent to reach room temperature (not more than 25°C).

As for all parenteral medicinal products, inspect visually the reconstituted solution prior to administration. The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. After administering the dose, discard the PegIntron pre-filled pen and any unused solution contained in it.

## **7. MARKETING AUTHORISATION HOLDER**

SP Europe  
73, rue de Stalle  
B-1180 Bruxelles  
Belgium

## **8. MARKETING AUTHORISATION NUMBERS**

EU/1/00/131/043  
EU/1/00/131/044  
EU/1/00/131/045  
EU/1/00/131/046

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 6 February 2002  
Date of last renewal: 25 May 2005

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.emea.europa.eu/>

## **1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 150 micrograms, powder and solvent for solution for injection in pre-filled pen

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each pre-filled pen of PegIntron 150 micrograms contains a sufficient amount of peginterferon alfa-2b as measured on a protein basis in a powder for solution for injection, and the corresponding amount of solvent, to provide 150 micrograms in 0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

The active substance is a covalent conjugate of recombinant interferon alfa-2b\* with monomethoxy polyethylene glycol. The potency of this product should not be compared to that of another pegylated or non-pegylated protein of the same therapeutic class. For more information, see section 5.1.

\*produced by rDNA technology in *E.coli* cells harbouring a genetically engineered plasmid hybrid encompassing an interferon alfa-2b gene from human leukocytes

For a full list of excipients, see section 6.1. PegIntron contains 40 mg of sucrose per 0.5 ml.

## **3. PHARMACEUTICAL FORM**

Powder and solvent for solution for injection in pre-filled pen

White powder.

Clear and colourless solvent.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

PegIntron is indicated for the treatment of adult patients with chronic hepatitis C who have elevated transaminases without liver decompensation and who are positive for serum HCV-RNA or anti-HCV, including naïve patients with clinically stable HIV co-infection (see section 4.4).

The best way to use PegIntron in this indication is in combination with ribavirin.

This combination is indicated in naïve patients including patients with clinically stable HIV co-infection. This combination of PegIntron with ribavirin is also indicated in patients who have previously responded (with normalisation of ALT at the end of treatment) to interferon alpha monotherapy but who have subsequently relapsed.

Interferon monotherapy, including PegIntron, is indicated mainly in case of intolerance or contraindication to ribavirin.

Please refer also to the ribavirin Summary of Product Characteristics (SPC) when PegIntron is to be used in combination with ribavirin.

### **4.2 Posology and method of administration**

Treatment should be initiated and monitored only by a physician experienced in the management of patients with hepatitis C.

#### **Dose to be administered**

PegIntron should be administered as a once weekly subcutaneous injection. The dose administered depends on whether it is used in combination with ribavirin or as monotherapy.

## Combination therapy

PegIntron 1.5 micrograms/kg/week in combination with ribavirin capsules.

The intended dose of 1.5 µg/kg of PegIntron to be used in combination with ribavirin may be delivered in weight categories with the pen/vial strengths according to **Table 1**. Ribavirin capsules are to be administered orally each day in two divided doses with food (morning and evening).

**Table 1 - Dosing for Combination Therapy**

Body Weight (kg)	PegIntron		Ribavirin Capsules	
	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Total Daily Dose (mg)	Number of Capsules (200 mg)
< 40	50	0.5	800	4 <sup>a</sup>
40-50	80	0.4	800	4 <sup>a</sup>
51-64	80	0.5	800	4 <sup>a</sup>
65-75	100	0.5	1,000	5 <sup>b</sup>
76-85	120	0.5	1,000	5 <sup>b</sup>
> 85	150	0.5	1,200	6 <sup>c</sup>

a: 2 morning, 2 evening

b: 2 morning, 3 evening

c: 3 morning, 3 evening

## Duration of treatment

**Predictability of sustained virological response:** Patients infected with virus genotype 1 who fail to achieve virological response at Week 12 are highly unlikely to become sustained virological responders (see also section 5.1).

- **Genotype 1:** For patients who exhibit virological response at week 12, treatment should be continued for another nine month period (i.e., a total of 48 weeks).  
In the subset of patients with genotype 1 infection and low viral load (< 600,000 IU/ml) who become HCV-RNA negative at treatment week 4 and remain HCV-RNA negative at week 24, the treatment could either be stopped after this 24 week treatment course or pursued for an additional 24 weeks (i.e. overall 48 weeks treatment duration). However, an overall 24 weeks treatment duration may be associated with a higher risk of relapse than a 48 weeks treatment duration (see section 5.1).
- **Genotypes 2 or 3:** It is recommended that all patients be treated for 24 weeks, except for HCV/HIV co-infected patients who should receive 48 weeks of treatment.
- **Genotype 4:** In general, patients infected with genotype 4 are considered harder to treat and limited study data (n=66) indicate they are compatible with a duration of treatment as for genotype 1.

## HCV/HIV Co-infection

The recommended duration of dosing for HCV/HIV co-infected patients is 48 weeks, regardless of genotype.

### *Predictability of response and non-response in HCV/HIV Co-infection*

Early virological response by Week 12, defined as a 2 log viral load decrease or undetectable levels of HCV-RNA, has been shown to be predictive for sustained response. The negative predictive value for sustained response in HCV/HIV co-infected patients treated with PegIntron in combination with ribavirin was 99 % (67/68; Study 1) (see section 5.1). A positive predictive value of 50 % (52/104; Study 1) was observed for HCV/HIV co-infected patients receiving combination therapy.



### PegIntron monotherapy

As monotherapy the PegIntron regimen is 0.5 or 1.0 microgram/kg/week. The lowest vial or pen strength available is 50 µg/0.5 ml; therefore for patients prescribed 0.5 µg/kg/week, doses must be adjusted by volume as shown in **Table 2**. For the 1.0 µg/kg dose, similar volume adjustments can be made or alternate vial strengths can be used as shown in **Table 2**. PegIntron monotherapy was not studied in HCV/HIV co-infected patients.

**Table 2- Monotherapy Dosing**

	0.5 µg/kg		1.0 µg/kg	
Body Weight (kg)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)	Vial/Pen Strength (µg/0.5ml)	Administer Once Weekly (ml)
30-35	50*	0.15	50	0.3
36-45	50*	0.2	50	0.4
46-56	50*	0.25	50	0.5
57-72	50	0.3	80	0.4
73-88	50	0.4	80	0.5
89-106	50	0.5	100	0.5
> 106**	80	0.4	120	0.5
* Must use vial. Minimum delivery for pen is 0.3 ml.				
** For patients > 120 kg, use 80 µg/0.5 ml vial				

### Duration of treatment

For patients who exhibit virological response at Week 12, treatment should be continued for at least another three-month period (i.e., a total of six months). The decision to extend therapy to one year of treatment should be based on prognostic factors (e.g., genotype, age > 40 years, male gender, bridging fibrosis).

### Dose modification for all patients

If severe adverse reactions or laboratory abnormalities develop during treatment with PegIntron monotherapy or PegIntron in combination with ribavirin, modify the dosages of each product as appropriate, until the adverse reactions abate. As adherence might be of importance for outcome of therapy, the dose should be kept as close as possible to the recommended standard dose. Guidelines were developed in clinical trials for dose modification.

### Combination Therapy Dose Reduction Guidelines

<b>Table 2a</b> Dose modification guidelines for combination therapy (with ribavirin)			
Laboratory values	Reduce only ribavirin dose to 600 mg/day* if:	Reduce only PegIntron dose to one-half dose if:	Discontinue combination therapy if:
Haemoglobin	< 10 g/dl	-	< 8.5 g/dl
Haemoglobin in: Patients with history of stable cardiac disease	≥ 2 g/dl decrease in haemoglobin during any four week period during treatment (permanent dose reduction)		< 12 g/dl after four weeks of dose reduction
White blood cells	-	< 1.5 x 10 <sup>9</sup> /l	< 1.0 x 10 <sup>9</sup> /l
Neutrophils	-	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	-	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l
Bilirubin – direct	-	-	2.5 x ULN**

Bilirubin – indirect	> 5 mg/dl	-	> 4 mg/dl (for > 4 weeks)
Creatinine	-	-	> 2.0 mg/dl
ALT/AST	-	-	2 x baseline and > 10 x ULN**

\* Patients whose dose of ribavirin is reduced to 600 mg daily receive one 200 mg capsule in the morning and two 200 mg capsules in the evening.

\*\* Upper limit of normal

Dose reduction of PegIntron may be accomplished by either reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 2b**.

<b>Table 2b – Reduced PegIntron Dosing for Combination Therapy</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5 ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
< 40	25	50*	0.25	25
40-50	32	50	0.3	30
51-64	40	50	0.4	40
65-75	50	50	0.5	50
76-85	60	80	0.4	64
> 85	75	100	0.4	80
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### **PegIntron Monotherapy Dose Reduction Guidelines**

Dose modification guidelines for patients who use PegIntron monotherapy are shown in **Table 3a**.

<b>Table 3a Dose modification guidelines for PegIntron monotherapy</b>		
<b>Laboratory values</b>	<b>Reduce PegIntron to one-half dose if:</b>	<b>Discontinue PegIntron if:</b>
Neutrophils	< 0.75 x 10 <sup>9</sup> /l	< 0.5 x 10 <sup>9</sup> /l
Platelets	< 50 x 10 <sup>9</sup> /l	< 25 x 10 <sup>9</sup> /l

Dose reduction for patients who use 0.5 µg/kg PegIntron monotherapy must be accomplished by reducing the prescribed volume by one-half. The 50 µg/0.5 ml vial must be used if necessary since the pen can only deliver a minimum volume of 0.3 ml.

For patients who use 1.0 µg/kg PegIntron monotherapy, dose reduction may be accomplished by reducing the prescribed volume by one-half or by utilizing a lower dose strength as shown in **Table 3b**.

<b>Table 3b – Reduced PegIntron Dose for the 1.0 µg/kg Monotherapy Regimen</b>				
<b>Body Weight (kg)</b>	<b>Target Reduced Dose (µg)</b>	<b>Vial/Pen Strength (µg/0.5ml)</b>	<b>Administer Once Weekly (ml)</b>	<b>Amount Delivered (µg)</b>
30-35	15	50*	0.15	15

36-45	20	50*	0.20	20
46-56	25	50*	0.25	25
57-72	32	50	0.3	30
73-89	40	50	0.4	40
90-106	50	50	0.5	50
> 106	60	80	0.4	64
*Must use vial. Minimum delivery for pen is 0.3 ml.				

### Special populations

**Use in renal impairment:** *Monotherapy:* PegIntron should be used with caution in patients with moderate to severe renal impairment. In patients with moderate renal dysfunction (creatinine clearance 30-50 ml/minute), the starting dose of PegIntron should be reduced by 25 %. Patients with severe renal dysfunction (creatinine clearance 15-29 ml/minute) should have the starting dose of PegIntron reduced by 50 %. Data are not available for the use of PegIntron in patients with creatinine clearance < 15 ml/minute (see section 5.2). Patients with severe renal impairment, including those on hemodialysis, should be closely monitored. If renal function decreases during treatment, PegIntron therapy should be discontinued.

*Combination therapy:* Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see ribavirin SPC). When administered in combination with ribavirin, subjects with impaired renal function should be more carefully monitored with respect to the development of anaemia.

**Use in hepatic impairment:** The safety and efficacy of PegIntron therapy has not been evaluated in patients with severe hepatic dysfunction, therefore PegIntron must not be used for these patients.

**Use in the elderly (≥ 65 years of age):** There are no apparent age-related effects on the pharmacokinetics of PegIntron. Data from elderly patients treated with a single dose of PegIntron suggest no alteration in PegIntron dose is necessary based on age (see section 5.2).

**Use in patients under the age of 18 years:** There is no experience in children (see section 5.2).

### 4.3 Contraindications

- Hypersensitivity to the active substance or to any interferon or to any of the excipients;
- A history of severe pre-existing cardiac disease, including unstable or uncontrolled cardiac disease in the previous six months (see section 4.4);
- Severe, debilitating medical conditions;
- Autoimmune hepatitis or a history of autoimmune disease;
- Severe hepatic dysfunction or decompensated cirrhosis of the liver;
- Pre-existing thyroid disease unless it can be controlled with conventional treatment;
- Epilepsy and/or compromised central nervous system (CNS) function;
- Initiation of PegIntron is contraindicated in HCV/HIV patients with cirrhosis and a Child-Pugh score ≥ 6.

*Combination therapy with ribavirin:* Also see ribavirin Summary of the Product Characteristics (SPC) if PegIntron is to be administered in combination with ribavirin in patients with chronic hepatitis C.

### 4.4 Special warnings and precautions for use

#### **Psychiatric and Central Nervous System (CNS):**

Severe CNS effects, particularly depression, suicidal ideation and attempted suicide have been observed in some patients during PegIntron therapy, and even after treatment discontinuation mainly during the 6-

month follow-up period. Other CNS effects including aggressive behaviour (sometimes directed against others), confusion and alterations of mental status have been observed with alpha interferons. Patients should be closely monitored for any signs or symptoms of psychiatric disorders. If such symptoms appear, the potential seriousness of these undesirable effects must be borne in mind by the prescribing physician and the need for adequate therapeutic management should be considered. If psychiatric symptoms persist or worsen, or suicidal ideation is identified, it is recommended that treatment with PegIntron be discontinued, and the patient followed, with psychiatric intervention as appropriate.

*Patients with existence of, or history of severe psychiatric conditions:* If treatment with peginterferon alfa-2b is judged necessary in patients with existence or history of severe psychiatric conditions, this should only be initiated after having ensured appropriate individualised diagnostic and therapeutic management of the psychiatric condition.

There is no experience with PegIntron in combination with ribavirin in patients who have relapsed after interferon alpha + ribavirin therapy.

All patients in the chronic hepatitis C studies had a liver biopsy before inclusion, but in certain cases (i.e. patients with genotype 2 and 3), treatment may be possible without histological confirmation. Current treatment guidelines should be consulted as to whether a liver biopsy is needed prior to commencing treatment.

**Acute hypersensitivity:** Acute hypersensitivity reactions (e.g., urticaria, angioedema, bronchoconstriction, anaphylaxis) have been observed rarely during interferon alfa-2b therapy. If such a reaction develops during treatment with PegIntron, discontinue treatment and institute appropriate medical therapy immediately. Transient rashes do not necessitate interruption of treatment.

**Cardiovascular system:** As with interferon alfa-2b, patients with a history of congestive heart failure, myocardial infarction and/or previous or current arrhythmic disorders, receiving PegIntron therapy require close monitoring. It is recommended that patients who have pre-existing cardiac abnormalities have electrocardiograms taken prior to and during the course of treatment. Cardiac arrhythmias (primarily supraventricular) usually respond to conventional therapy but may require discontinuation of PegIntron therapy.

More significant obtundation and coma, including cases of encephalopathy, have been observed in some patients, usually elderly, treated at higher doses for oncology indications. While these effects are generally reversible, in a few patients full resolution took up to three weeks. Very rarely, seizures have occurred with high doses of interferon alpha.

**Liver function:** As with all interferons, discontinue treatment with PegIntron in patients who develop prolongation of coagulation markers which might indicate liver decompensation.

**Fever:** While fever may be associated with the flu-like syndrome reported commonly during interferon therapy, other causes of persistent fever must be ruled out.

**Hydration:** Adequate hydration must be maintained in patients undergoing PegIntron therapy since hypotension related to fluid depletion has been seen in some patients treated with alpha interferons. Fluid replacement may be necessary.

**Pulmonary changes:** Pulmonary infiltrates, pneumonitis, and pneumonia, occasionally resulting in fatality, have been observed rarely in interferon alpha treated patients. Any patient developing fever, cough, dyspnea or other respiratory symptoms must have a chest X-ray taken. If the chest X-ray shows pulmonary infiltrates or there is evidence of pulmonary function impairment, the patient is to be monitored closely, and, if appropriate, discontinue interferon alpha. Prompt discontinuation of interferon alpha administration and treatment with corticosteroids appear to be associated with resolution of pulmonary adverse events.

**Autoimmune disease:** The development of auto-antibodies and autoimmune disorders has been reported during treatment with alpha interferons. Patients predisposed to the development of

autoimmune disorders may be at increased risk. Patients with signs or symptoms compatible with autoimmune disorders should be evaluated carefully, and the benefit-risk of continued interferon therapy should be reassessed (see also section 4.4 Thyroid changes and 4.8).

**Ocular changes:** Ophthalmologic disorders, including retinal haemorrhages, cotton wool spots, and retinal artery or vein obstruction have been reported in rare instances after treatment with alpha interferons (see section 4.8). All patients should have a baseline eye examination. Any patient complaining of ocular symptoms, including loss of visual acuity or visual field must have a prompt and complete eye examination. Periodic visual examinations are recommended during PegIntron therapy, particularly in patients with disorders that may be associated with retinopathy, such as diabetes mellitus or hypertension. Discontinuation of PegIntron should be considered in patients who develop new or worsening ophthalmological disorders.

**Thyroid changes:** Infrequently, patients treated for chronic hepatitis C with interferon alpha have developed thyroid abnormalities, either hypothyroidism or hyperthyroidism. Determine thyroid stimulating hormone (TSH) levels if, during the course of therapy, a patient develops symptoms consistent with possible thyroid dysfunction. In the presence of thyroid dysfunction, PegIntron treatment may be continued if TSH levels can be maintained in the normal range by medication.

**Metabolic disturbances:** Hypertriglyceridemia and aggravation of hypertriglyceridemia, sometimes severe, have been observed. Monitoring of lipid levels is, therefore, recommended.

### **HCV/HIV Co-infection**

Mitochondrial toxicity and lactic acidosis:

Patients co-infected with HIV and receiving Highly Active Anti-Retroviral Therapy (HAART) may be at increased risk of developing lactic acidosis. Caution should be used when adding PegIntron and ribavirin to HAART therapy (see ribavirin SPC).

Moreover, patients treated with PegIntron and ribavirin combination therapy and zidovudine are at increased risk of developing anaemia.

Hepatic decompensation in HCV/HIV co-infected patients with advanced cirrhosis:

Co-infected patients with advanced cirrhosis receiving HAART may be at increased risk of hepatic decompensation and death. Adding treatment with alfa interferons alone or in combination with ribavirin may increase the risk in this patient subset. Other baseline factors in co-infected patients that may be associated with a higher risk of hepatic decompensation include treatment with didanosine and elevated bilirubin serum concentration.

Co-infected patients receiving both antiretroviral (ARV) and anti-hepatitis treatment should be closely monitored, assessing their Child-Pugh score during treatment. Patients progressing to hepatic decompensation should have their anti-hepatitis treatment immediately discontinued and the ARV treatment reassessed.

Haematological abnormalities in HCV/HIV co-infected patients:

HCV/HIV co-infected patients receiving peginterferon alfa-2b/ribavirin treatment and HAART may be at increased risk to develop haematological abnormalities (as neutropenia, thrombocytopenia and anaemia) compared to HCV mono-infected patients. Although, the majority of them could be managed by dose reduction, close monitoring of haematological parameters should be undertaken in this population of patients (see section 4.2 and below “Laboratory tests” and section 4.8).

Patients with low CD4 counts:

In patients co-infected with HCV/HIV, limited efficacy and safety data (N = 25) are available in subjects with CD4 counts less than 200 cells/ $\mu$ l. Caution is therefore warranted in the treatment of patients with low CD4 counts.

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron and ribavirin.

**Dental and periodontal disorders:** Dental and periodontal disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and mucous membranes of the mouth during long-term treatment with the combination of PegIntron and ribavirin. Patients should brush their teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If this reaction occurs, they should be advised to rinse out their mouth thoroughly afterwards.

**Organ transplant recipients:** The safety and efficacy of PegIntron alone or in combination with ribavirin for the treatment of hepatitis C in liver or other organ transplant recipients have not been studied. Preliminary data indicates that interferon alpha therapy may be associated with an increased rate of kidney graft rejection. Liver graft rejection has also been reported.

**Other:** Due to reports of interferon alpha exacerbating pre-existing psoriatic disease and sarcoidosis, use of PegIntron in patients with psoriasis or sarcoidosis is recommended only if the potential benefit justifies the potential risk.

**Laboratory tests:** Standard haematologic tests, blood chemistry and a test of thyroid function must be conducted in all patients prior to initiating therapy. Acceptable baseline values that may be considered as a guideline prior to initiation of PegIntron therapy are:

- Platelets  $\geq 100,000/\text{mm}^3$
- Neutrophil count  $\geq 1,500/\text{mm}^3$
- TSH level must be within normal limits

Laboratory evaluations are to be conducted at weeks 2 and 4 of therapy, and periodically thereafter as clinically appropriate.

**Important information about some of the ingredients of PegIntron:**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

Patients with rare hereditary problems of fructose intolerance, glucose galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Results from a multiple-dose probe study assessing P450 substrates in chronic hepatitis C patients receiving once weekly PegIntron (1.5  $\mu\text{g}/\text{kg}$ ) for 4 weeks demonstrated an increase in activity of CYP2D6 and CYP2C8/9. No change in activity of CYP1A2, CYP3A4, or N-acetyltransferase was observed.

Caution should be used when administering peginterferon alfa-2b with medicines metabolised by CYP2D6 and CYP2C8/9, especially those with narrow therapeutic window, such as warfarin and phenytoin (CYP2C9) and flecainide (CYP2D6).

These findings may partly relate to improved metabolic capacity due to reduced hepatic inflammation in patients undergoing treatment with PegIntron. Caution is therefore advised when PegIntron treatment is initiated for chronic hepatitis in patients treated with medication with a narrow therapeutic window and sensitive to mild metabolic impairment of the liver.

No pharmacokinetic interactions were noted between PegIntron and ribavirin in a multiple-dose pharmacokinetic study.

**Methadone:** In patients with chronic hepatitis C that were on stable methadone maintenance therapy and naïve to peginterferon alfa-2b, addition of 1.5 microgram/kg/week of PegIntron subcutaneously for 4 weeks increased R-methadone AUC by approximately 15 % (95 % CI for AUC ratio estimate 103 – 128 %). The clinical significance of this finding is unknown; however, patients should be

monitored for signs and symptoms of increased sedative effect, as well as respiratory depression. Especially in patients on a high dose of methadone, the risk for QTc prolongation should be considered.

#### *HCV/HIV Co-infection*

Nucleoside analogs: ribavirin has been shown *in vitro* to inhibit phosphorylation of zidovudine and stavudine. The clinical significance of these findings is unknown. However, these *in vitro* findings raise the possibility that concurrent use of ribavirin with either zidovudine or stavudine might lead to increased HIV plasma viraemia. Therefore, it is recommended that plasma HIV RNA levels be closely monitored in patients treated with ribavirin concurrently with either of these two agents. If HIV RNA levels increase, the use of ribavirin concomitantly with reverse transcriptase inhibitors must be reviewed (see ribavirin SPC).

Use of nucleoside analogs, alone or in combination with other nucleosides, has resulted in lactic acidosis. Pharmacologically, ribavirin increases phosphorylated metabolites of purine nucleosides *in vitro*. This activity could potentiate the risk of lactic acidosis induced by purine nucleoside analogs (e.g. didanosine or abacavir). Co-administration of ribavirin and didanosine is not recommended. Reports of mitochondrial toxicity, in particular lactic acidosis and pancreatitis, of which some fatal, have been reported (see ribavirin SPC).

## **4.6 Pregnancy and lactation**

PegIntron is recommended for use in fertile women only when they are using effective contraception during the treatment.

There are no adequate data from the use of interferon alfa-2b in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3). Interferon alfa-2b has been shown to be abortifacient in primates. PegIntron is likely to also cause this effect.

The potential risk in humans is unknown. PegIntron is to be used during pregnancy only if the potential benefit justifies the potential risk to the foetus

**Lactation:** It is not known whether the components of this medicinal product are excreted in human milk. Because of the potential for adverse reactions in breast-fed infants, Page: 199  
[0]breast-feeding should be discontinued prior to initiation of treatment.

#### *Combination therapy with ribavirin:*

Ribavirin causes serious birth defects when administered during pregnancy. Ribavirin therapy is contraindicated in women who are pregnant. Extreme care must be taken to avoid pregnancy in female patients or in partners of male patients taking PegIntron in combination with ribavirin. Females of childbearing potential and their partners must each use an effective contraceptive during treatment and for 4 months after treatment has been concluded. Male patients and their female partners must each use an effective contraceptive during treatment and for 7 months after treatment has been concluded (see Ribavirin SPC).

## **4.7 Effects on ability to drive and use machines**

Patients who develop fatigue, somnolence or confusion during treatment with PegIntron are cautioned to avoid driving or operating machinery.

## **4.8 Undesirable effects**

The safety of PegIntron is evaluated from data from two clinical trials: one with PegIntron monotherapy, one with PegIntron in combination with ribavirin. In both cases, patients were treated for one year.

**Table 4** describes the regimens and patient exposure for one year of treatment in patients with no previous exposure to interferon (interferon-naïve patients). Because of a significant overlap in the

pattern of undesirable effects with PegIntron monotherapy, groups of patients have been brought together in **Table 5** to show the pattern of reported effects for all monotherapy groups.

<b>Table 4</b> Regimens and patient exposure		
<b>Treatment</b>	<b>Regimen</b>	<b>Number of patients treated for one year</b>
PegIntron + ribavirin	PegIntron (1.5 micrograms/kg/week) + ribavirin (> 10.6 mg/kg/day)	188
Interferon alfa-2b + ribavirin	Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day)	505
PegIntron monotherapy	PegIntron (0.5 microgram/kg/week)	315
	PegIntron (1.0 microgram/kg/week)	297
	PegIntron (1.5 micrograms/kg/week)	304



<b>Table 5</b> Undesirable effects very commonly reported in clinical trials > 10 % of patients in PegIntron + ribavirin group Very common (> 1/10)			
	<b>PegIntron + ribavirin</b>	<b>Interferon alfa- 2b + ribavirin</b>	<b>PegIntron monotherapy</b>
<b>Infections and infestations</b>			
Infection viral	10 %	5 %	4-5 %
<b>Metabolism and nutrition disorders</b>			
Weight decrease	30 %	19 %	8-18 %
<b>Psychiatric disorders</b>			
Depression	34 %	32 %	26 %
Irritability	32 %	34 %	19 %
Insomnia	37 %	41 %	16-19 %
Anxiety	14 %	14 %	8 %
Concentration impaired	18 %	21 %	9-10 %
Emotional lability	11 %	10 %	5 %
<b>Nervous system disorders</b>			
Headache	58 %	57 %	57-63 %
Mouth dry	10 %	8 %	4-8 %
<b>Respiratory, thoracic and mediastinal disorders</b>			
Pharyngitis	10 %	7 %	3 %
Coughing	14 %	11 %	4 %
Dyspnea	26 %	22 %	5 %
<b>Gastrointestinal disorders</b>			
Nausea	43 %	31 %	20-23 %
Anorexia	35 %	26 %	10-25 %
Diarrhoea	20 %	13 %	14-17 %
Abdominal pain	12 %	9 %	11 %
Vomiting	16 %	10 %	4-7 %
<b>Skin and subcutaneous tissue disorders</b>			
Alopecia	45 %	32 %	20-34 %
Pruritus	27 %	27 %	7-9 %
Skin dry	23 %	21 %	6-9 %
Rash	21 %	21 %	5-7 %
<b>Musculoskeletal and connective tissue disorders</b>			
Myalgia	49 %	49 %	46-60 %
Arthralgia	31 %	26 %	23-28 %
Musculoskeletal pain	15 %	11 %	11-13 %
<b>General disorders and administration site conditions</b>			
Injection site inflammation	20 %	17 %	39-44 %
Injection site reaction	54 %	36 %	7-9 %
Fatigue	56 %	59 %	43 %
Rigors	42 %	40 %	33-43 %
Fever	39 %	32 %	29-43 %
Flu-like symptoms	21 %	23 %	18-25 %
Asthenia	28 %	17 %	12-14 %
Dizziness	17 %	16 %	7-12 %

<b>Table 6</b> Undesirable effects commonly reported in clinical trials in patients treated with PegIntron + ribavirin or PegIntron monotherapy Common (> 1/100, < 1/10)		
<b>Body system</b>	<b>5-10%</b>	<b>1-&lt;5%</b>
Infections and infestations		otitis media, fungal infection, bacterial infection, herpes simplex
Blood and lymphatic system disorders	Anaemia, leukopaenia	Thrombocytopenia, lymphadenopathy
Endocrine disorders	hypothyroidism	hyperthyroidism
Metabolism and nutrition disorders		hyperuricemia, hypocalcemia, thirst
Psychiatric disorders	agitation, nervousness	aggressive behaviour, somnolence, behavior disorder, apathy, appetite increased, sleep disorder, dreaming abnormal, decreased libido
Nervous system disorders	paresthesia, increased sweating	hypoesthesia, hyperaesthesia, hypertonia, confusion, tremor, vertigo, migraine, ataxia, neuralgia
Eye disorders		blurred vision, conjunctivitis, lacrimal gland disorder, eye pain
Cardiac disorders		tachycardia, palpitation
Vascular disorders		hypotension, hypertension, syncope, flushing
Ear and labyrinth disorders		tinnitus, hearing impairment/loss
Respiratory, thoracic and mediastinal disorders		nonproductive cough, rhinitis, sinusitis, bronchitis, respiratory disorder, nasal congestion, rhinorrhea, dysphonia, epistaxis
Gastrointestinal disorders	dyspepsia	constipation, taste perversion, loose stools, stomatitis, ulcerative stomatitis, gingival bleeding, glossitis, flatulence, hemorrhoids, gastroesophageal reflux, gingivitis, dehydration
Hepatobiliary disorders		hepatomegaly, hyperbilirubinemia
Skin and subcutaneous tissue disorders		erythematous rash, eczema, photosensitivity reaction, maculopapular rash, abnormal hair texture, acne, dermatitis, furunculosis, nail disorder, psoriasis, urticaria, erythema, face or peripheral oedema
Musculoskeletal and connective tissue disorders		arthritis
Renal and urinary disorders		micturition frequency, urine abnormal
Reproductive system and breast disorders	menstrual disorder, menorrhagia	ovarian disorder, vaginal disorder, sexual dysfunction (not specified), impotence, breast pain, amenorrhoea, prostatitis
General disorders and administration site conditions	RUQ pain, malaise, chest pain	injection site pain

Most cases of neutropaenia and thrombocytopaenia were mild (WHO grades 1 or 2). There were some cases of more severe neutropenia in patients treated with the recommended doses of PegIntron in combination with ribavirin (WHO grade 3: 39 of 186 [21 %]; and WHO grade 4: 13 of 186 [7 %]).

In a clinical trial, approximately 1.2 % of patients treated with PegIntron or interferon alfa-2b in combination with ribavirin reported life-threatening psychiatric events during treatment. These events included suicidal ideation and attempted suicide (see section 4.4). Following marketing, psychosis and hallucination have been reported rarely.

Rarely ( $> 1/10,000$ ,  $< 1/1,000$ ) or very rarely ( $< 1/10,000$ ) reported events with interferon alfa-2b, including PegIntron, include:

**Blood and lymphatic system disorders:**

Very rarely interferon alfa-2b or PegIntron used alone or in combination with ribavirin may be associated with aplastic anaemia. Pure red cell aplasia has been reported.

**Immune system disorders:**

very rarely: sarcoidosis or exacerbation of sarcoidosis

**Endocrine disorders:**

rarely: diabetes

**Nervous system disorders:**

rarely: seizure, peripheral neuropathy

very rarely: cerebrovascular ischaemia, cerebrovascular haemorrhage, encephalopathy

**Cardiac disorders:**

rarely: arrhythmia

very rarely: cardiac ischaemia, myocardial infarction

**Respiratory, thoracic and mediastinal disorders:**

very rarely: interstitial lung disease

**Gastrointestinal disorders:**

rarely: pancreatitis

very rarely: ulcerative and ischaemic colitis

**Skin and subcutaneous tissue disorders:**

very rarely: erythema multiforme, Stevens Johnson syndrome, toxic epidermal necrolysis, injection site necrosis

**Musculoskeletal and connective tissue disorders:**

rarely: rhabdomyolysis, myositis

**Renal and urinary disorders:**

rarely: renal insufficiency and renal failure

Cardiovascular (CVS) adverse events, particularly arrhythmia, appeared to be correlated mostly with pre-existing CVS disease and prior therapy with cardiotoxic agents (see section 4.4). Cardiomyopathy, that may be reversible upon discontinuation of interferon alpha, has been reported rarely in patients without prior evidence of cardiac disease.

Ophthalmological disorders that have been reported rarely with alpha interferons include retinopathies (including macular oedema), retinal haemorrhages, retinal artery or vein obstruction, cotton wool spots, loss of visual acuity or visual field, optic neuritis, and papilloedema (see section 4.4).

A wide variety of autoimmune and immune-mediated disorders have been reported with alpha interferons including thyroid disorders, systemic lupus erythematosus, rheumatoid arthritis (new or aggravated), idiopathic and thrombotic thrombocytopenic purpura, vasculitis, neuropathies including mononeuropathies (see also section 4.4, **Autoimmune disorders**).

**HCV/HIV co-infected patients**

For HCV/HIV co-infected patients receiving PegIntron in combination with ribavirin, other undesirable effects (that were not reported in mono-infected patients) which have been reported in the larger studies with a frequency  $> 5\%$  were: oral candidiasis (14 %), lipodystrophy acquired (13 %), CD4 lymphocytes decreased (8 %), appetite decreased (8 %), gamma-glutamyltransferase increased (9 %), back pain (5 %), blood amylase increased (6 %), blood lactic acid increased (5 %), cytolytic hepatitis (6 %), lipase increased (6 %) and pain in limb (6 %).

*Mitochondrial toxicity:*

Mitochondrial toxicity and lactic acidosis have been reported in HIV-positive patients receiving NRTI

regimen and associated ribavirin for co-HCV infection (see section 4.4).

*Laboratory values for HCV/HIV co-infected patients:*

Although haematological toxicities of neutropenia, thrombocytopenia and anaemia occurred more frequently in HCV/HIV co-infected patients, the majority could be managed by dose modification and rarely required premature discontinuation of treatment (see section 4.4). Haematological abnormalities were more frequently reported in patients receiving PegIntron in combination with ribavirin when compared to patients receiving interferon alfa-2b in combination with ribavirin. In Study 1 (see section 5.1), decrease in absolute neutrophil count levels below 500 cells/mm<sup>3</sup> was observed in 4 % (8/194) of patients and decrease in platelets below 50,000/mm<sup>3</sup> was observed in 4 % (8/194) of patients receiving PegIntron in combination with ribavirin. Anaemia (hemoglobin < 9.4g/dl) was reported in 12% (23/194) of patients treated with PegIntron in combination with ribavirin.

*CD4 lymphocytes decrease:* Treatment with PegIntron in combination with ribavirin was associated with decreases in absolute CD4+ cell counts within the first 4 weeks without a reduction in CD4+ cell percentage. The decrease in CD4+ cell counts was reversible upon dose reduction or cessation of therapy. The use of PegIntron in combination with ribavirin had no observable negative impact on the control of HIV viraemia during therapy or follow-up. Limited safety data (N= 25) are available in co-infected patients with CD4+ cell counts < 200/μl (see section 4.4).

Please refer to the respective Summary of Product Characteristics of the antiretroviral medicinal products that are to be taken concurrently with HCV therapy for awareness and management of toxicities specific for each product and the potential for overlapping toxicities with PegIntron in combination with ribavirin.

## **4.9 Overdose**

In clinical trials, cases of accidental overdose, at never more than twice the prescribed dose, were reported. There were no serious reactions. Undesirable effects resolved during continued administration of PegIntron.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Immunostimulants, Cytokines and immunomodulators, Interferons, Peginterferon alfa-2b, ATC code: L03A B10.

Recombinant interferon alfa-2b is covalently conjugated with monomethoxy polyethylene glycol at an average degree of substitution of 1 mole of polymer/mole of protein. The average molecular mass is approximately 31,300 daltons of which the protein moiety constitutes approximately 19,300.

#### **Interferon alfa-2b**

*In vitro* and *in vivo* studies suggest that the biological activity of PegIntron is derived from its interferon alfa-2b moiety.

Interferons exert their cellular activities by binding to specific membrane receptors on the cell surface. Studies with other interferons have demonstrated species specificity. However, certain monkey species, e.g., Rhesus monkeys, are susceptible to pharmacodynamic stimulation upon exposure to human type 1 interferons.

Once bound to the cell membrane, interferon initiates a complex sequence of intracellular events that include the induction of certain enzymes. It is thought that this process, at least in part, is responsible for the various cellular responses to interferon, including inhibition of virus replication in virus-infected cells, suppression of cell proliferation and such immunomodulating activities as enhancement of the phagocytic activity of macrophages and augmentation of the specific cytotoxicity of

lymphocytes for target cells. Any or all of these activities may contribute to interferon's therapeutic effects.

Recombinant interferon alfa-2b also inhibits viral replication *in vitro* and *in vivo*. Although the exact antiviral mode of action of recombinant interferon alfa-2b is unknown, it appears to alter the host cell metabolism. This action inhibits viral replication or if replication occurs, the progeny virions are unable to leave the cell.

### **PegIntron**

PegIntron pharmacodynamics were assessed in a rising single-dose trial in healthy subjects by examining changes in oral temperature, concentrations of effector proteins such as serum neopterin and 2'5'-oligoadenylate synthetase (2'5'-OAS), as well as white cell and neutrophil counts. Subjects treated with PegIntron showed mild dose-related elevations in body temperature. Following single doses of PegIntron between 0.25 and 2.0 micrograms/kg/week, serum neopterin concentration was increased in a dose-related manner. Neutrophil and white cell count reductions at the end of week 4 correlated with the dose of PegIntron.

### **PegIntron clinical trials**

Two pivotal trials have been conducted, one (C/I97-010) with PegIntron monotherapy; the other (C/I98-580) with PegIntron in combination with ribavirin. Eligible patients for these trials had chronic hepatitis C confirmed by a positive HCV-RNA polymerase chain reaction (PCR) assay (> 30 IU/ml), a liver biopsy consistent with a histological diagnosis of chronic hepatitis with no other cause for the chronic hepatitis, and abnormal serum ALT.

In the PegIntron monotherapy trial, a total of 916 naïve chronic hepatitis C patients were treated with PegIntron (0.5, 1.0 or 1.5 micrograms/kg/week) for one year with a follow-up period of six months. In addition, 303 patients received interferon alfa-2b (3 million International Units [MIU] three times a week) as a comparator. This study showed that PegIntron was superior to interferon alfa-2b (**Table 8**).

In the PegIntron combination trial, 1,530 naïve patients were treated for one year with one of the following combination regimens:

- PegIntron (1.5 micrograms/kg/week) + ribavirin (800 mg/day), (n = 511).
- PegIntron (1.5 micrograms/kg/week for one month followed by 0.5 microgram/kg/week for 11 months) + ribavirin (1,000/1,200 mg/day), (n = 514).
- Interferon alfa-2b (3 MIU three times a week) + ribavirin (1,000/1,200 mg/day) (n = 505).

In this trial, the combination of PegIntron (1.5 micrograms/kg/week) and ribavirin was significantly more effective than the combination of interferon alfa-2b and ribavirin (**Table 7**), particularly in patients infected with Genotype 1 (**Table 8**). Sustained response was assessed by the response rate six months after the cessation of treatment.

HCV genotype and baseline virus load are prognostic factors which are known to affect response rates. However, response rates in this trial were shown to be dependent also on the dose of ribavirin administered in combination with PegIntron or interferon alfa-2b. In those patients that received > 10.6 mg/kg ribavirin (800 mg dose in typical 75 kg patient), regardless of genotype or viral load, response rates were significantly higher than in those patients that received ≤ 10.6 mg/kg ribavirin (**Table 8**), while response rates in patients that received > 13.2 mg/kg ribavirin were even higher.

<b>Table 7</b> Sustained virological response (% patients HCV negative)							
	<b>PegIntron monotherapy</b>				<b>PegIntron + ribavirin</b>		
<b>Treatment regimen</b>	<b>P 1.5</b>	<b>P 1.0</b>	<b>P 0.5</b>	<b>I</b>	<b>P 1.5/R</b>	<b>P 0.5/R</b>	<b>I/R</b>
<b>Number of patients</b>	304	297	315	303	511	514	505
<b>Response at end of treatment</b>	49 %	41 %	33 %	24 %	65 %	56 %	54 %

Sustained response	23 %*	25 %	18 %	12 %	54 %**	47 %	47 %
P 1.5	PegIntron 1.5 micrograms/kg						
P 1.0	PegIntron 1.0 microgram/kg						
P 0.5	PegIntron 0.5 microgram/kg						
I	Interferon alfa-2b 3 MIU						
P 1.5/R	PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)						
P 0.5/R	PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)						
I/R	Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)						
*	p < 0.001 P 1.5 vs. I						
**	p = 0.0143 P 1.5/R vs. I/R						

<b>Table 8</b> Sustained response rates with PegIntron + ribavirin (by ribavirin dose, genotype and viral load)				
HCV Genotype	Ribavirin dose (mg/kg)	P 1.5/R	P 0.5/R	I/R
<b>All Genotypes</b>	<b>All</b>	<b>54 %</b>	<b>47 %</b>	<b>47 %</b>
	≤ 10.6	50 %	41 %	27 %
	> 10.6	61 %	48 %	47 %
<b>Genotype 1</b>	<b>All</b>	<b>42 %</b>	<b>34 %</b>	<b>33 %</b>
	≤ 10.6	38 %	25 %	20 %
	> 10.6	48 %	34 %	34 %
Genotype 1 ≤ 600,000 IU/ml	All	73 %	51 %	45 %
	≤ 10.6	74 %	25 %	33 %
	> 10.6	71 %	52 %	45 %
Genotype 1 > 600,000 IU/ml	All	30 %	27 %	29 %
	≤ 10.6	27 %	25 %	17 %
	> 10.6	37 %	27 %	29 %
<b>Genotype 2/3</b>	<b>All</b>	<b>82 %</b>	<b>80 %</b>	<b>79 %</b>
	≤ 10.6	79 %	73 %	50 %
	> 10.6	88 %	80 %	80 %

P 1.5/R	PegIntron (1.5 micrograms/kg) + ribavirin (800 mg)
P 0.5/R	PegIntron (1.5 to 0.5 microgram/kg) + ribavirin (1,000/1,200 mg)
I/R	Interferon alfa-2b (3 MIU) + ribavirin (1,000/1,200 mg)

In the PegIntron monotherapy study, the Quality of Life was generally less affected by 0.5 microgram/kg of PegIntron than by either 1.0 microgram/kg of PegIntron once weekly or 3 MIU of interferon alfa-2b three times a week.

In a separate trial, 224 patients with genotype 2 or 3 received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with ribavirin 800 mg –1,400 mg p.o. for 6 months (based on body weight, only three patients weighing > 105 kg, received the 1,400 mg dose, which has not yet been validated) (**Table 9**). Twenty-four % had bridging fibrosis or cirrhosis (Knodell 3/4).

**Table 9. Virologic Response at End of Treatment, Sustained Virologic Response and Relapse by HCV Genotype and Viral Load\***

	PegIntron 1.5 µg/kg once weekly plus Ribavirin 800-1,400 mg/day		
	End of Treatment Response	Sustained Virologic Response	Relapse
All Subjects	94 % (211/224)	81 % (182/224)	12 % (27/224)
HCV 2 ≤ 600,000 IU/ml > 600,000 IU/ml	100 % (42/42)	93 % (39/42)	7 % (3/42)
	100 % (20/20)	95 % (19/20)	5 % (1/20)
	100 % (22/22)	91 % (20/22)	9 % (2/22)
HCV 3	93 % (169/182)	79 % (143/182)	14 %

≤ 600,000 IU/ml	93 % (92/99)	86 % (85/99)	(24/166) 8 % (7/91)
> 600,000 IU/ml	93 % (77/83)	70 % (58/83)	23 % (17/75)

\* Any subject with an undetectable HCV-RNA level at the Follow-Up Week 12 visit and missing data at the Follow-Up Week 24 visit was considered a sustained responder. Any subject with missing data in and after the Follow-Up Week 12 window was considered to be a non-responder at Week 24 of follow-up.

The 6 month treatment duration in this trial was better tolerated than one year of treatment in the pivotal combination trial; for discontinuation 5 % vs. 14 %, for dose modification 18 % vs. 49 %.

In a non-comparative trial, 235 patients with genotype 1 and low viral load (< 600,000 IU/ml) received PegIntron, 1.5 microgram/kg subcutaneously, once weekly, in combination with weight adjusted ribavirin. The overall sustained response rate after a 24-week treatment duration was 50 %. Forty-one percent of subjects (97/235) had nondetectable plasma HCV-RNA levels at Week 4 and Week 24 of therapy. In this subgroup, there was a 92 % (89/97) sustained virological response rate. The high sustained response rate in this subgroup of patients was identified in an interim analysis (n=49) and prospectively confirmed (n=48).

Limited historical data indicate that treatment for 48 weeks might be associated with a higher sustained response rate (11/11) and with a lower risk of relapse (0/11 as compared to 7/96 following 24 weeks of treatment).

#### Predictability of sustained virological response

Virological response by week 12, defined as a 2-log viral load decrease or undetectable levels of HCV-RNA has been shown to be predictive for sustained response (**Table 10**).

<b>Table 10</b> Predictability of sustained response by viral response at week 12 and genotype*				
Treatment	Genotype	Viral response at week 12	Sustained response	Negative predictive value
PegIntron 1.5 + ribavirin (> 10.6 mg/kg) 48-week treatment	1	Yes 75 % (82/110)	71 % (58/82)	----
		No 25 % (28/110)	0 % (0/28)	100 %
PegIntron 1.5 + ribavirin 800-1,400 mg 24-week treatment	2 and 3	Yes 99 % (213/215)	83 % (177/213)	----
		No 1 % (2/215)	50 % (1/2)	50 %

\* reflects patients with 12 week data available

The negative predictive value for sustained response in patients treated with PegIntron in monotherapy was 98 %.

#### HCV/HIV Co-infected patients

Two trials have been conducted in patients co-infected with HIV and HCV. The response to treatment in both of these trials is presented in **Table 11**. Study 1 (RIBAVIC; P01017) was a randomized, multicentre study which enrolled 412 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (1.5 µg/kg/week) plus ribavirin (800 mg/day) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800 mg/day) for 48 weeks with a follow-up period of 6 months. Study 2 (P02080) was a randomized, single centre study that enrolled 95 previously untreated adult patients with chronic hepatitis C who were co-infected with HIV. Patients were randomized to receive either PegIntron (100 or 150 µg /week based

on weight) plus ribavirin (800-1,200 mg/day based on weight) or interferon alfa-2b (3 MIU TIW) plus ribavirin (800-1,200 mg/day based on weight). The duration of therapy was 48 weeks with a follow-up period of 6 months except for patients infected with genotypes 2 or 3 and viral load < 800,000 IU/ml (Amplicor) who were treated for 24 weeks with a 6-month follow-up period.

<b>Table 11</b> Sustained virological response based on genotype after PegIntron in combination with Ribavirin in HCV/HIV Co-infected patients						
	<b>Study 1<sup>1</sup></b>			<b>Study 2<sup>2</sup></b>		
	PegIntron (1.5 µg/kg/week) + ribavirin (800 mg)	Interferon alfa-2b (3 MIU TIW) + ribavirin (800 mg)	p value <sup>a</sup>	PegIntron (100 or 150 <sup>c</sup> µg/week) + ribavirin (800-1,200 mg) <sup>d</sup>	Interferon alfa-2b (3 MIU TIW) + ribavirin (800-1,200 mg) <sup>d</sup>	p value <sup>b</sup>
All	27 % (56/205)	20 % (41/205)	0.047	44 % (23/52)	21 % (9/43)	0.017
Genotype 1, 4	17 % (21/125)	6 % (8/129)	0.006	38 % (12/32)	7 % (2/27)	0.007
Genotype 2, 3	44 % (35/80)	43 % (33/76)	0.88	53 % (10/19)	47 % (7/15)	0.730

MIU = million international units; TIW = three times a week.

a: p value based on Cochran-Mantel Haenszel Chi square test.

b: p value based on chi-square test.

c: subjects < 75 kg received 100 µg/week PegIntron and subjects ≥ 75 kg received 150 µg/week PegIntron.

d: ribavirin dosing was 800 mg for patients < 60 kg, 1,000 mg for patients 60-75 kg, and 1,200 mg for patients > 75 kg.

<sup>1</sup>Carrato F, Bani-Sadir F, Pol S et al. JAMA 2004; 292(23): 2839-2848.

<sup>2</sup>Laguno M, Murillas J, Blanco J et al. AIDS 2004; 18(13): F27-F36.

### *Histological response*

Liver biopsies were obtained before and after treatment in Study 1 and were available for 210 of the 412 subjects (51 %). Both the Metavir score and Ishak grade decreased among subjects treated with PegIntron in combination with ribavirin. This decline was significant among responders (-0.3 for Metavir and -1.2 for Ishak) and stable (-0.1 for Metavir and -0.2 for Ishak) among non-responders. In terms of activity, about one-third of sustained responders showed improvement and none showed worsening. There was no improvement in terms of fibrosis observed in this study. Steatosis was significantly improved in patients infected with HCV Genotype 3.

## **5.2 Pharmacokinetic properties**

PegIntron is a well characterized polyethylene glycol-modified (“pegylated”) derivative of interferon alfa-2b and is predominantly composed of monopegylated species. The plasma half-life of PegIntron is prolonged compared with non-pegylated interferon alfa-2b. PegIntron has a potential to depegylate to free interferon alfa-2b. The biologic activity of the pegylated isomers is qualitatively similar, but weaker than free interferon alfa-2b.

Following subcutaneous administration, maximal serum concentrations occur between 15-44 hours post-dose, and are sustained for up to 48-72 hours post-dose.

PegIntron C<sub>max</sub> and AUC measurements increase in a dose-related manner. Mean apparent volume of distribution is 0.99 l/kg.

Upon multiple dosing, there is an accumulation of immunoreactive interferons. There is, however, only a modest increase in biologic activity as measured by a bioassay.

Mean (SD) PegIntron elimination half-life is approximately 40 hours (13.3 hours), with apparent clearance of 22.0 ml/hr·kg. The mechanisms involved in clearance of interferons in man have not yet been



fully elucidated. However, renal elimination may account for a minority (approximately 30 %) of PegIntron apparent clearance.

**Renal function:** Renal clearance appears to account for 30 % of total clearance of PegIntron. In a single dose study (1.0 microgram/kg) in patients with impaired renal function,  $C_{max}$ , AUC, and half-life increased in relation to the degree of renal impairment.

Following multiple dosing of PegIntron (1.0 microgram/kg subcutaneously administered every week for four weeks) the clearance of PegIntron is reduced by a mean of 17 % in patients with moderate renal impairment (creatinine clearance 30-49 ml/minute) and by a mean of 44 % in patients with severe renal impairment (creatinine clearance 15-29 ml/minute) compared to subjects with normal renal function. Based on single dose data, clearance was similar in patients with severe renal impairment not on dialysis and in patients who were receiving hemodialysis. The dose of PegIntron for monotherapy should be reduced in patients with moderate or severe renal impairment (see sections 4.2 and 4.4). Patients with creatinine clearance < 50 ml/minute must not be treated with PegIntron in combination with ribavirin (see section 4.3).

Because of marked inter-subject variability in interferon pharmacokinetics, it is recommended that patients with severe renal impairment be closely monitored during treatment with PegIntron (see section 4.2)

**Hepatic function:** The pharmacokinetics of PegIntron have not been evaluated in patients with severe hepatic dysfunction.

**Elderly patients  $\geq 65$  years of age:** The pharmacokinetics of PegIntron following a single subcutaneous dose of 1.0 microgram/kg were not affected by age. The data suggest that no alteration in PegIntron dosage is necessary based on advancing age.

**Patients under the age of 18 years:** Specific pharmacokinetic evaluations have not been performed on these patients. PegIntron is indicated for the treatment of chronic hepatitis C only in patients 18 years of age or older.

**Interferon neutralising factors:** Interferon neutralising factor assays were performed on serum samples of patients who received PegIntron in the clinical trial. Interferon neutralising factors are antibodies which neutralise the antiviral activity of interferon. The clinical incidence of neutralising factors in patients who received PegIntron 0.5 micrograms/kg is 1.1 %.

### 5.3 Preclinical safety data

**PegIntron:** Adverse events not observed in clinical trials were not seen in toxicity studies in monkeys. These studies were limited to four weeks due to the appearance of anti-interferon antibodies in most monkeys.

Reproduction studies of PegIntron have not been performed. Interferon alfa-2b has been shown to be an abortifacient in primates. PegIntron is likely to also cause this effect. Effects on fertility have not been determined. It is not known whether the components of this medicinal product are excreted into experimental animal or human milk (see section 4.6 for relevant human data on pregnancy and lactation). PegIntron showed no genotoxic potential.

The relative non-toxicity of monomethoxy-polyethylene glycol (mPEG), which is liberated from PegIntron by metabolism *in vivo* has been demonstrated in preclinical acute and subchronic toxicity studies in rodents and monkeys, standard embryo-foetal development studies and in *in vitro* mutagenicity assays.

**PegIntron plus ribavirin:** When used in combination with ribavirin, PegIntron did not cause any effects not previously seen with either active substance alone. The major treatment-related change was

a reversible, mild to moderate anaemia, the severity of which was greater than that produced by either active substance alone.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Powder for solution for injection:

Disodium phosphate, anhydrous,  
Sodium dihydrogen phosphate dihydrate,  
Sucrose,  
Polysorbate 80.

Solvent for parenteral use:

Water for injections.

Deliverable volume from pen = 0.5 ml.

### **6.2 Incompatibilities**

This medicinal product should only be reconstituted with the solvent provided (see section 6.6). In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

3 years

After reconstitution:

- Chemical and physical in-use stability has been demonstrated for 24 hours at 2°C - 8°C.
- From a microbiological point of view, the product is to be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C - 8°C.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C - 8°C). Do not freeze.

### **6.5 Nature and contents of container**

The powder and solvent are both contained in a two-chamber cartridge, Type I flint glass, separated by a bromobutyl rubber plunger. The cartridge is sealed at one end with a polypropylene cap containing a bromobutyl rubber liner and at the other end by a bromobutyl rubber plunger.

PegIntron 150 micrograms is supplied as:

- 1 pen containing powder and solvent for solution for injection, 1 injection needle and 2 cleansing swabs;
- 4 pens containing powder and solvent for solution for injection, 4 injection needles and 8 cleansing swabs;
- 6 pens containing powder and solvent for solution for injection, 6 injection needles and 12 cleansing swabs;
- 12 pens containing powder and solvent for solution for injection, 12 injection needles and 24 cleansing swabs.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

PegIntron pre-filled pen contains a powder of peginterferon alfa-2b and a solvent for solution at a strength of 150 micrograms for single use. Each pen is reconstituted with the solvent provided in the two-chamber cartridge (water for injections) for administration of up to 0.5 ml of solution. A small volume is lost during preparation of PegIntron for injection when the dose is measured and injected. Therefore, each pen contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection. The reconstituted solution has a concentration of 150 micrograms in 0.5 ml.

PegIntron is injected subcutaneously after reconstituting the powder as instructed, attaching an injection needle and setting the prescribed dose. A complete and illustrated set of instructions is provided in the Annex to the Package Leaflet.

Remove PegIntron pre-filled pen from the refrigerator before administration to allow the solvent to reach room temperature (not more than 25°C).

As for all parenteral medicinal products, inspect visually the reconstituted solution prior to administration. The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. After administering the dose, discard the PegIntron pre-filled pen and any unused solution contained in it.

## **7. MARKETING AUTHORISATION HOLDER**

SP Europe  
73, rue de Stalle  
B-1180 Bruxelles  
Belgium

## **8. MARKETING AUTHORISATION NUMBERS**

EU/1/00/131/047  
EU/1/00/131/048  
EU/1/00/131/049  
EU/1/00/131/050

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 6 February 2002  
Date of last renewal: 25 May 2005

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.emea.europa.eu/>

## **ANNEX II**

- A. MANUFACTURER OF THE BIOLOGICAL ACTIVE  
SUBSTANCE AND MANUFACTURING AUTHORISATION  
HOLDER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OF THE MARKETING AUTHORISATION**

**A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND  
MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH  
RELEASE**

Name and address of the manufacturer of the biological active substance

SP (Brinny) Company  
Innishannon - County Cork  
Ireland

Name and address of the manufacturer responsible for batch release

SP (Brinny) Company  
Innishannon - County Cork  
Ireland

**B. CONDITIONS OF THE MARKETING AUTHORISATION**

• **CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IMPOSED ON  
THE MARKETING AUTHORISATION HOLDER**

Medicinal product subject to restricted medical prescription (See Annex I: Summary of Product Characteristics, section 4.2).

• **CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE  
USE OF THE MEDICINAL PRODUCT**

Not applicable.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****Carton 50 micrograms****1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 50 micrograms powder and solvent for solution for injection  
peginterferon alfa-2b

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

One vial of powder contains 50 micrograms of peginterferon alfa-2b and provides 50 micrograms/0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

**3. LIST OF EXCIPIENTS**

Excipients: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80. One ampoule of solvent contains 0.7 ml of water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

1 vial of powder, 1 ampoule of solvent  
1 vial of powder, 1 ampoule of solvent, 1 injection syringe, 2 injection needles and 1 cleansing swab  
4 vials of powder, 4 ampoules of solvent  
4 vials of powder, 4 ampoules of solvent, 4 injection syringes, 8 injection needles and 4 cleansing swabs  
6 vials of powder, 6 ampoules of solvent  
12 vials of powder, 12 ampoules of solvent, 12 injection syringes, 24 injection needles and 12 cleansing swabs  
50 micrograms/0.5 ml

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Subcutaneous use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP  
After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).



**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

After withdrawal of the dose, any remaining solution must be discarded.

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Marketing authorisation holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/131/001 (1 vial of powder, 1 ampoule of solvent)

EU/1/00/131/002 (1 vial of powder, 1 ampoule of solvent, 1 injection syringe, 2 injection needles and 1 cleansing swab)

EU/1/00/131/003 (4 vials of powder, 4 ampoules of solvent)

EU/1/00/131/004 (4 vials of powder, 4 ampoules of solvent, 4 injection syringes, 8 injection needles and 4 cleansing swabs)

EU/1/00/131/005 (6 vials of powder, 6 ampoules of solvent)

EU/1/00/131/026 (12 vials of powder, 12 ampoules of solvent, 12 injection syringes, 24 injection needles and 12 cleansing swabs)

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

PegIntron 50 mcg

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS****PegIntron 50 micrograms – vial of powder****1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

PegIntron 50 micrograms powder for injection  
SC

**2. METHOD OF ADMINISTRATION**

Read the package leaflet before use.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

50 micrograms/0.5 ml

**6. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**Carton 80 micrograms**

**1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 80 micrograms powder and solvent for solution for injection  
peginterferon alfa-2b

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

One vial of powder contains 80 micrograms of peginterferon alfa-2b and provides 80 micrograms/0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

**3. LIST OF EXCIPIENTS**

Excipients: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80. One ampoule of solvent contains 0.7 ml of water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

1 vial of powder, 1 ampoule of solvent  
1 vial of powder, 1 ampoule of solvent, 1 injection syringe, 2 injection needles and 1 cleansing swab  
4 vials of powder, 4 ampoules of solvent  
4 vials of powder, 4 ampoules of solvent, 4 injection syringes, 8 injection needles and 4 cleansing swabs  
6 vials of powder, 6 ampoules of solvent  
12 vials of powder, 12 ampoules of solvent, 12 injection syringes, 24 injection needles and 12 cleansing swabs  
80 micrograms/0.5 ml

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Subcutaneous use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

#### **9. SPECIAL STORAGE CONDITIONS**

**Store in a refrigerator.**

#### **10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

After withdrawal of the dose, any remaining solution must be discarded.

#### **11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Marketing authorisation holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

#### **12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/131/006 (1 vial of powder, 1 ampoule of solvent)

EU/1/00/131/007 (1 vial of powder, 1 ampoule of solvent, 1 injection syringe, 2 injection needles and 1 cleansing swab)

EU/1/00/131/008 (4 vials of powder, 4 ampoules of solvent)

EU/1/00/131/009 (4 vials of powder, 4 ampoules of solvent, 4 injection syringes, 8 injection needles and 4 cleansing swabs)

EU/1/00/131/010 (6 vials of powder, 6 ampoules of solvent)

EU/1/00/131/027 (12 vials of powder, 12 ampoules of solvent, 12 injection syringes, 24 injection needles and 12 cleansing swabs)

#### **13. BATCH NUMBER**

Lot

#### **14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

#### **15. INSTRUCTIONS ON USE**

#### **16. INFORMATION IN BRAILLE**

PegIntron 80 mcg

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS****PegIntron 80 micrograms - vial of powder****1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

PegIntron 80 micrograms powder for injection  
SC

**2. METHOD OF ADMINISTRATION**

Read the package leaflet before use.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

80 micrograms/0.5 ml

**6. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****Carton 100 micrograms****1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 100 micrograms powder and solvent for solution for injection  
peginterferon alfa-2b

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

One vial of powder contains 100 micrograms of peginterferon alfa-2b and  
provides 100 micrograms/0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

**3. LIST OF EXCIPIENTS**

Excipients: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and  
polysorbate 80. One ampoule of solvent contains 0.7 ml of water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

1 vial of powder, 1 ampoule of solvent  
1 vial of powder, 1 ampoule of solvent, 1 injection syringe, 2 injection needles and 1 cleansing swab  
4 vials of powder, 4 ampoules of solvent  
4 vials of powder, 4 ampoules of solvent, 4 injection syringes, 8 injection needles and 4 cleansing  
swabs  
6 vials of powder, 6 ampoules of solvent  
12 vials of powder, 12 ampoules of solvent, 12 injection syringes, 24 injection needles  
and 12 cleansing swabs  
100 micrograms/0.5 ml

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Subcutaneous use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP  
After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a  
refrigerator (2°C - 8°C).

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

After withdrawal of the dose, any remaining solution must be discarded.

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Marketing authorisation holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/131/011 (1 vial of powder, 1 ampoule of solvent)

EU/1/00/131/012 (1 vial of powder, 1 ampoule of solvent, 1 injection syringe, 2 injection needles and 1 cleansing swab)

EU/1/00/131/013 (4 vials of powder, 4 ampoules of solvent)

EU/1/00/131/014 (4 vials of powder, 4 ampoules of solvent, 4 injection syringes, 8 injection needles and 4 cleansing swabs)

EU/1/00/131/015 (6 vials of powder, 6 ampoules of solvent)

EU/1/00/131/028 (12 vials of powder, 12 ampoules of solvent, 12 injection syringes, 24 injection needles and 12 cleansing swabs)

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

PegIntron 100 mg

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS****PegIntron 100 micrograms - vial of powder****1. NAME OF THE MEDICINAL PRODUCT AND IF NECESSARY ROUTE(S) OF ADMINISTRATION**

PegIntron 100 micrograms powder for injection  
SC

**2. METHOD OF ADMINISTRATION**

Read the package leaflet before use.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

100 micrograms/0.5 ml

**6. OTHER**



**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**Carton 120 micrograms**

**1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 120 micrograms powder and solvent for solution for injection  
peginterferon alfa-2b

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

One vial of powder contains 120 micrograms of peginterferon alfa-2b and provides 120 micrograms/0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

**3. LIST OF EXCIPIENTS**

Excipients: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80. One ampoule of solvent contains 0.7 ml of water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

1 vial of powder, 1 ampoule of solvent  
1 vial of powder, 1 ampoule of solvent, 1 injection syringe, 2 injection needles and 1 cleansing swab  
4 vials of powder, 4 ampoules of solvent  
4 vials of powder, 4 ampoules of solvent, 4 injection syringes, 8 injection needles and 4 cleansing swabs  
6 vials of powder, 6 ampoules of solvent  
12 vials of powder, 12 ampoules of solvent, 12 injection syringes, 24 injection needles and 12 cleansing swabs  
120 micrograms/0.5 ml

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Subcutaneous use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

#### **9. SPECIAL STORAGE CONDITIONS**

**Store in a refrigerator.**

#### **10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

After withdrawal of the dose, any remaining solution must be discarded.

#### **11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Marketing authorisation holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

#### **12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/131/016 (1 vial of powder, 1 ampoule of solvent)

EU/1/00/131/017 (1 vial of powder, 1 ampoule of solvent, 1 injection syringe, 2 injection needles and 1 cleansing swab)

EU/1/00/131/018 (4 vials of powder, 4 ampoules of solvent)

EU/1/00/131/019 (4 vials of powder, 4 ampoules of solvent, 4 injection syringes, 8 injection needles and 4 cleansing swabs)

EU/1/00/131/020 (6 vials of powder, 6 ampoules of solvent)

EU/1/00/131/029 (12 vials of powder, 12 ampoules of solvent, 12 injection syringes, 24 injection needles and 12 cleansing swabs)

#### **13. BATCH NUMBER**

Lot

#### **14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

#### **15. INSTRUCTIONS ON USE**

#### **16. INFORMATION IN BRAILLE**

PegIntron 120 mcg

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS****PegIntron 120 micrograms - vial of powder****1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

PegIntron 120 micrograms powder for injection  
SC

**2. METHOD OF ADMINISTRATION**

Read the package leaflet before use.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

120 micrograms/0.5 ml

**6. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**Carton 150 micrograms**

**1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 150 micrograms powder and solvent for solution for injection  
peginterferon alfa-2b

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

One vial of powder contains 150 micrograms of peginterferon alfa-2b and  
provides 150 micrograms/0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

**3. LIST OF EXCIPIENTS**

Excipients: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and  
polysorbate 80. One ampoule of solvent contains 0.7 ml of water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

1 vial of powder, 1 ampoule of solvent  
1 vial of powder, 1 ampoule of solvent, 1 injection syringe, 2 injection needles and 1 cleansing swab  
4 vials of powder, 4 ampoules of solvent  
4 vials of powder, 4 ampoules of solvent, 4 injection syringes, 8 injection needles and 4 cleansing  
swabs  
6 vials of powder, 6 ampoules of solvent  
12 vials of powder, 12 ampoules of solvent, 12 injection syringes, 24 injection needles  
and 12 cleansing swabs  
150 micrograms/0.5 ml

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Subcutaneous use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

#### **9. SPECIAL STORAGE CONDITIONS**

**Store in a refrigerator.**

#### **10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

After withdrawal of the dose, any remaining solution must be discarded.

#### **11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Marketing authorisation holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

#### **12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/131/021 (1 vial of powder, 1 ampoule of solvent)

EU/1/00/131/022 (1 vial of powder, 1 ampoule of solvent, 1 injection syringe, 2 injection needles and 1 cleansing swab)

EU/1/00/131/023 (4 vials of powder, 4 ampoules of solvent)

EU/1/00/131/024 (4 vials of powder, 4 ampoules of solvent, 4 injection syringes, 8 injection needles and 4 cleansing swabs)

EU/1/00/131/025 (6 vials of powder, 6 ampoules of solvent)

EU/1/00/131/030 (12 vials of powder, 12 ampoules of solvent, 12 injection syringes, 24 injection needles and 12 cleansing swabs)

#### **13. BATCH NUMBER**

Lot

#### **14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

#### **15. INSTRUCTIONS ON USE**

#### **16. INFORMATION IN BRAILLE**

PegIntron 150 mcg

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS****PegIntron 150 micrograms - vial of powder****1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

PegIntron 150 micrograms powder for injection  
SC

**2. METHOD OF ADMINISTRATION**

Read the package leaflet before use.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

150 micrograms/0.5 ml

**6. OTHER**

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS****PegIntron - ampoule of solvent****1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

Solvent for PegIntron

Water for injections

**2. METHOD OF ADMINISTRATION****3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

0.7 ml

**6. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**Carton 50 micrograms powder and solvent for solution for injection**

**1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 50 micrograms powder and solvent for solution for injection pre-filled pen  
peginterferon alfa-2b

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

One PegIntron pre-filled pen contains a sufficient amount of peginterferon alfa-2b to provide 50 micrograms in 0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

**3. LIST OF EXCIPIENTS**

Excipients: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80. Solvent: water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

1 pen, 1 injection needle and 2 cleansing swabs  
4 pens, 4 injection needles and 8 cleansing swabs  
6 pens, 6 injection needles and 12 cleansing swabs  
12 pens, 12 injection needles and 24 cleansing swabs  
50 micrograms/0.5 ml

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Subcutaneous use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).



**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator. Do not freeze.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

After injection of the dose, discard the pen in an appropriate container.

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Marketing authorisation holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/131/031 (1 pen, 1 injection needle and 2 cleansing swabs)

EU/1/00/131/032 (4 pens, 4 injection needles and 8 cleansing swabs)

EU/1/00/131/033 (6 pens, 6 injection needles and 12 cleansing swabs)

EU/1/00/131/034 (12 pens, 12 injection needles and 24 cleansing swabs)

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

PegIntron 50 mcg

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS****PegIntron 50 micrograms powder and solvent for solution for injection****1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

PegIntron 50 micrograms powder and solvent for injection  
SC

**2. METHOD OF ADMINISTRATION**

Read the package leaflet before use.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

50 micrograms/0.5 ml

**6. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****Carton 80 micrograms powder and solvent for solution for injection****1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 80 micrograms powder and solvent for solution for injection pre-filled pen  
peginterferon alfa-2b

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

One PegIntron pre-filled pen contains a sufficient amount of peginterferon alfa-2b to provide 80 micrograms in 0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

**3. LIST OF EXCIPIENTS**

Excipients: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80. Solvent: water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

1 pen, 1 injection needle and 2 cleansing swabs  
4 pens, 4 injection needles and 8 cleansing swabs  
6 pens, 6 injection needles and 12 cleansing swabs  
12 pens, 12 injection needles and 24 cleansing swabs  
80 micrograms/0.5 ml

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Subcutaneous use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator. Do not freeze.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

After injection of the dose, discard the pen in an appropriate container.

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Marketing authorisation holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/131/035 (1 pen, 1 injection needle and 2 cleansing swabs)

EU/1/00/131/036 (4 pens, 4 injection needles and 8 cleansing swabs)

EU/1/00/131/037 (6 pens, 6 injection needles and 12 cleansing swabs)

EU/1/00/131/038 (12 pens, 12 injection needles and 24 cleansing swabs)

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

PegIntron 80 mcg

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS****PegIntron 80 micrograms powder and solvent for solution for injection****1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

PegIntron 80 micrograms powder and solvent for injection  
SC

**2. METHOD OF ADMINISTRATION**

Read the package leaflet before use.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

80 micrograms/0.5 ml

**6. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**Carton 100 micrograms powder and solvent for solution for injection**

**1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 100 micrograms powder and solvent for solution for injection pre-filled pen  
peginterferon alfa-2b

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

One PegIntron pre-filled pen contains a sufficient amount of peginterferon alfa-2b to provide 100 micrograms in 0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

**3. LIST OF EXCIPIENTS**

Excipients: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80. Solvent: water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

1 pen, 1 injection needle and 2 cleansing swabs  
4 pens, 4 injection needles and 8 cleansing swabs  
6 pens, 6 injection needles and 12 cleansing swabs  
12 pens, 12 injection needles and 24 cleansing swabs  
100 micrograms/0.5 ml

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Subcutaneous use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator. Do not freeze.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

After injection of the dose, discard the pen in an appropriate container.

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Marketing authorisation holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/131/039 (1 pen, 1 injection needle and 2 cleansing swabs)

EU/1/00/131/040 (4 pens, 4 injection needles and 8 cleansing swabs)

EU/1/00/131/041 (6 pens, 6 injection needles and 12 cleansing swabs)

EU/1/00/131/042 (12 pens, 12 injection needles and 24 cleansing swabs)

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

PegIntron 100 mcg

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS****PegIntron 100 micrograms powder and solvent for solution for injection****1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

PegIntron 100 micrograms powder and solvent for injection  
SC

**2. METHOD OF ADMINISTRATION**

Read the package leaflet before use.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

100 micrograms/0.5 ml

**6. OTHER**



**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**Carton 120 micrograms powder and solvent for solution for injection**

**1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 120 micrograms powder and solvent for solution for injection pre-filled pen  
peginterferon alfa-2b

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

One PegIntron pre-filled pen contains a sufficient amount of peginterferon alfa-2b to provide 120 micrograms in 0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

**3. LIST OF EXCIPIENTS**

Excipients: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80. Solvent: water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

1 pen, 1 injection needle and 2 cleansing swabs  
4 pens, 4 injection needles and 8 cleansing swabs  
6 pens, 6 injection needles and 12 cleansing swabs  
12 pens, 12 injection needles and 24 cleansing swabs  
120 micrograms/0.5 ml

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Subcutaneous use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator. Do not freeze.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

After injection of the dose, discard the pen in an appropriate container.

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Marketing authorisation holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/131/043 (1 pen, 1 injection needle and 2 cleansing swabs)

EU/1/00/131/044 (4 pens, 4 injection needles and 8 cleansing swabs)

EU/1/00/131/045 (6 pens, 6 injection needles and 12 cleansing swabs)

EU/1/00/131/046 (12 pens, 12 injection needles and 24 cleansing swabs)

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

PegIntron 120 mcg

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS****PegIntron 120 micrograms powder and solvent for solution for injection****1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

PegIntron 120 micrograms powder and solvent for injection  
SC

**2. METHOD OF ADMINISTRATION**

Read the package leaflet before use.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

120 micrograms/0.5 ml

**6. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**Carton 150 micrograms powder and solvent for solution for injection**

**1. NAME OF THE MEDICINAL PRODUCT**

PegIntron 150 micrograms powder and solvent for solution for injection pre-filled pen  
peginterferon alfa-2b

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

One PegIntron pre-filled pen contains a sufficient amount of peginterferon alfa-2b to provide 150 micrograms in 0.5 ml of peginterferon alfa-2b when reconstituted as recommended.

**3. LIST OF EXCIPIENTS**

Excipients: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80. Solvent: water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

1 pen, 1 injection needle and 2 cleansing swabs  
4 pens, 4 injection needles and 8 cleansing swabs  
6 pens, 6 injection needles and 12 cleansing swabs  
12 pens, 12 injection needles and 24 cleansing swabs  
150 micrograms/0.5 ml

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Subcutaneous use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator. Do not freeze.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

After injection of the dose, discard the pen in an appropriate container.

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Marketing authorisation holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/131/047 (1 pen, 1 injection needle and 2 cleansing swabs)

EU/1/00/131/048 (4 pens, 4 injection needles and 8 cleansing swabs)

EU/1/00/131/049 (6 pens, 6 injection needles and 12 cleansing swabs)

EU/1/00/131/050 (12 pens, 12 injection needles and 24 cleansing swabs)

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

PegIntron 150 mcg

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS****PegIntron 150 micrograms powder and solvent for solution for injection****1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

PegIntron 150 micrograms powder and solvent for injection  
SC

**2. METHOD OF ADMINISTRATION**

Read the package leaflet before use.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

150 micrograms/0.5 ml

**6. OTHER**

## **B. PACKAGE LEAFLET**

## **PACKAGE LEAFLET : INFORMATION FOR THE USER**

### **PegIntron 50 micrograms powder and solvent for solution for injection**

peginterferon alfa-2b (conjugation of recombinant interferon alfa-2b with monomethoxy polyethylene glycol)

**Read all of this leaflet carefully before you start using this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What PegIntron is and what it is used for
2. Before you use PegIntron
3. How to use PegIntron
4. Possible side effects
5. How to store PegIntron
6. Further information

## **1. WHAT PEGINTRON IS AND WHAT IT IS USED FOR**

Interferons modify the response of the body's immune system to help fight infections and severe diseases. PegIntron, which contains an interferon, is used for the treatment of chronic hepatitis C, a viral infection of the liver, including in first time users co-infected with clinically stable HIV.

PegIntron is best used for this treatment in combination with ribavirin.

PegIntron is used alone in case of intolerance or contraindication to ribavirin.

## **2. BEFORE YOU USE PEGINTRON**

PegIntron is not recommended for use in patients under the age of 18 years.

**Do not use PegIntron**

- If you are allergic (hypersensitive) to peginterferon alfa-2b or any of the other ingredients of PegIntron.
- If you are allergic (hypersensitive) to any interferon.
- If you have had severe heart problems, or if you have heart disease that has not been well controlled during the past 6 months.
- If you have severe medical conditions that leave you very weak.
- If you have autoimmune hepatitis or any other problem with your immune system; if you are taking medicine that suppresses your immune system (your immune system protects you against infection and some diseases).
- If you have advanced, uncontrolled liver disease (other than hepatitis C).
- If you have thyroid disease that is not well controlled with medicines.
- If you have a condition that causes convulsions (seizures, or "fits").

**Take special care with PegIntron**

- If you develop symptoms of a severe allergic reaction (such as difficulty in breathing, wheezing, or hives) while on this medication, seek medical help immediately.
- If you ever had a heart attack or a heart problem.
- If you have ever been treated for depression or any other nervous or mental disorder.



- If you have kidney disease, your doctor may prescribe a lower than usual dose and monitor your kidney blood values regularly during treatment. If you have a kidney disease and you are using PegIntron in combination with medicinal products containing ribavirin, your doctor should monitor you more carefully for a decrease in red blood cell count.
- If you have had a problem with your liver (other than hepatitis C).
- If you develop symptoms associated with a cold or other respiratory infection, such as fever, cough, or any difficulty in breathing, tell your doctor.
- If you are diabetic, your doctor may ask you to have an eye examination.
- If you have had any serious illness affecting your breathing or your blood.
- If you have psoriasis, it may become worse while you are using PegIntron.
- If you are planning to become pregnant, discuss this with your doctor before starting to use PegIntron.
- If you are also being treated for HIV, please see **Using other medicines**.
- If you have had a severe nervous or mental disorder.
- If you ever had depression or develop symptoms associated with depression (e.g. feelings of sadness, dejection, etc) while on treatment with PegIntron (see section 4).
- If you have received an organ transplant, either kidney or liver, interferon treatment may increase the risk of rejection. Be sure to discuss this with your doctor.

Dental and gum disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and membranes of the mouth during long-term treatment with the combination of PegIntron with ribavirin. You should brush your teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If you have this reaction, be sure to rinse your mouth thoroughly afterwards.

#### **While being treated with PegIntron**

Your doctor may want you to drink extra fluids to help prevent low blood pressure.

#### **Using other medicines**

Please inform your doctor or pharmacist if you are taking or have recently taken any other medicines, even those not prescribed.

#### **Pregnancy and breast-feeding**

Ask your doctor or pharmacist for advice before taking any medicine. In studies in pregnant animals, interferons have sometimes caused miscarriage. The effect on human pregnancy is not known. In combination therapy with ribavirin, ribavirin can be very damaging to an unborn baby, thus both female and male patients must take special precautions in their sexual activity if there is any chance for pregnancy to occur:

- if you are a **girl** or a **woman** of childbearing age, you must have a negative pregnancy test before treatment, each month during treatment, and for the 4 months after treatment is stopped. You and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 4 months after stopping treatment. This can be discussed with your doctor.
- if you are a **man** who is taking ribavirin, do not have sex with a pregnant woman unless you use a condom. This will lessen the chance for ribavirin to be left in the woman's body. If your female partner is not pregnant now but is of childbearing age, she must be tested for pregnancy each month during treatment and for the 7 months after treatment has stopped. This can be discussed with your doctor. If you are a male patient, you and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 7 months after stopping treatment. This can be discussed with your doctor.

It is not known whether this product is present in human milk. Therefore, do not breast-feed an infant if you are taking PegIntron. In combination therapy with ribavirin, take notice of the respective informing texts of ribavirin containing medicinal products.

#### **Driving and using machines**

Do not drive or operate any tools or machines if you feel tired, sleepy or confused while taking PegIntron.

### **Important information about some of the ingredients of PegIntron**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

This medicinal product contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### **Patients who also have HIV infection**

Lactic acidosis and worsening liver function are side effects associated with Highly Active Anti-Retroviral Therapy (HAART), an HIV treatment. If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis and of liver failure. Your doctor will monitor you for signs and symptoms of these conditions (Please be sure to read the ribavirin Patient Leaflet also). Additionally, patients treated with PegIntron and ribavirin combination therapy and zidovudine could be at increased risk of developing anaemia (low number of red blood cells). If you are also being treated for Human Immunodeficiency Virus (HIV) infection (AIDS) with zidovudine or stavudine, it is not certain if ribavirin will change the way these medicines work. Therefore, your blood will be checked regularly to be sure that the HIV infection is not getting worse. If it gets worse, your doctor will decide whether or not your ribavirin treatment needs to be changed.

Co-administration of ribavirin and didanosine and/or stavudine is not recommended due to the risk of lactic acidosis (a build-up of lactic acid in the body) and pancreatitis.

## **3. HOW TO USE PEGINTRON**

Your doctor has prescribed PegIntron specifically for you and your current condition; do not share this medicine with anyone else.

Your doctor has determined your dose of PegIntron based on your weight. If necessary, the dose may be changed during treatment.

### **Combination treatment**

PegIntron, when given with ribavirin capsules, is usually given at a dose of 1.5 microgram/kg once a week.

Ribavirin capsules are taken every day, morning and evening. The number of ribavirin capsules you take depends on your weight.

- If you weigh less than 65 kg, take 2 capsules in the morning and 2 in the evening (total of 800 mg each day).
- If you weigh between 65 and 85 kg, take 2 capsules in the morning and 3 in the evening (total of 1,000 mg each day).
- If you weigh more than 85 kg, take 3 capsules in the morning and 3 in the evening (total 1,200 mg each day).

The combination treatment is continued for 3 to 6 months, and sometimes for one year depending on your physicians judgement. Take notice of the respective informing texts of ribavirin containing medicinal products.

For HCV/HIV co-infected patients, the duration of treatment is 48 weeks.

### **PegIntron alone:**

PegIntron, when given alone, is usually given at a dose of 0.5 or 1.0 microgram/kg once a week, for at least 6 months, and possibly for 1 year. If you have kidney disease, your dose may be lower, depending upon your kidney function.

PegIntron alone was not studied in HCV/HIV co-infected patients.

**All patients:**

If you are injecting PegIntron yourself, please be sure that the dose that has been prescribed for you is clearly provided on the package of medicine you receive.

If you have the impression that the effect of PegIntron is too strong or too weak, talk to your doctor or pharmacist.

PegIntron is intended for subcutaneous use. This means that it is injected through a short injection needle into the fatty tissue just under your skin. If you are injecting this medicine yourself, you will be instructed how to prepare and give the injection. Detailed instructions for subcutaneous administration are provided with this leaflet (see How to self-inject PegIntron at the end of the package leaflet).

Prepare the dose just before you intend to inject it and use it immediately. Look carefully at the reconstituted solution prior to administration. Do not use if there is discolouration of the reconstituted solution or if particulate matter is present. Discard any solution that is left in the vial after you give yourself the injection.

Inject PegIntron once each week on the same day. Injecting it at the same time of day each week will help you not to forget to take it.

Always use PegIntron exactly as your doctor has told you. Do not exceed the recommended dosage, and take it for as long as prescribed.

**If you use more PegIntron than you should**

Tell your doctor or healthcare professional as soon as possible.

**If you forget to take PegIntron**

Take the dose as soon as you remember, then continue your treatment as usual.

Do not take a double dose to make up for a forgotten dose. Contact your doctor or pharmacist if needed.

**4. POSSIBLE SIDE EFFECTS**

Like all medicines, PegIntron can cause side effects, although not everybody gets them. Although not all of these side effects may occur, they may need medical attention if they do.

Some people get depressed when taking PegIntron alone or in combination treatment with ribavirin, and in some cases people had suicidal thoughts or aggressive behaviour (sometimes directed against others). Some patients have actually committed suicide. Be sure to seek emergency care if you notice that you are becoming depressed or have suicidal thoughts or change in your behaviour. You may want to consider asking a family member or close friend to help you stay alert to signs of depression or changes in your behaviour.

Check with your doctor immediately if any of the following side effects occur: chest pain; changes in the way your heart beats; breathing problems (including shortness of breath), confusion; feeling depressed, wanting to harm yourself, hallucinations, numbness or tingling feeling; dizziness, convulsion ("fit"); trouble sleeping, thinking or concentrating; difficulty remaining alert, severe stomach pain or cramps; blood or clots in stool (or black, tarry stool); fever or chills beginning after a few weeks of treatment, pain in your lower back or side; difficulty or inability to pass urine, painful or inflamed muscles (sometimes severe); problems with your eyes or your eyesight or hearing; severe or painful reddening of your skin or mucous membrane, severe bleeding from your nose. Your doctor will test your blood to ensure that your white blood cell (cells that fight infection) and red blood cell (cells that carry oxygen) counts, platelets (blood clotting cells) and other laboratory values are at acceptable levels.

Other side effects that may occur with the combination of PegIntron and ribavirin capsules include:

Very commonly reported side effects (greater than 1 in every 10 patients):

Irritation or redness (and rarely, skin damage) at the site of injection, headache, tired feeling, shaking chills, fever, flu-like symptoms, weakness, loss of weight, nausea, loss of appetite, diarrhoea or loose stools, stomach pain, vomiting, muscle aches, pain in joints and muscles, feeling depressed, irritability, trouble falling asleep or staying asleep, feeling anxious or nervous, difficulty concentrating, mood swings, hair loss, itching, dry skin, sore throat, coughing, difficult breathing, dizziness, virus infection, rash, and dry mouth.

Commonly reported side effects (at least 1 in every 100 patients, but not greater than 1 in every 10 patients):

Increased sweating, chest pain, pain on the right side around your ribs, numbness, pain or tingling feeling, change in thyroid gland activity (which may make you feel tired or, less commonly, energetic), stomach upset, rapid heart rate, agitation, nervousness, difficult or irregular menstrual period

Less common are pain at the place of injection, flushing, low or high blood pressure, dry or teary eyes, redness of skin or skin disorders, psoriasis, hives, nail disorder, feeling unwell, feeling faint, poor coordination, confusion, increased or decreased sensitivity to touch, tense muscles, arthritis, bruising, loss of interest in activities including sex, sexual problem, unusual dreams, shaky hands, vertigo (spinning feeling), increased appetite, heartburn, intestinal gas (flatulence), constipation, hemorrhoids, red or bleeding gums, redness or sores in mouth, change in taste, changes in hearing or ringing in ears, thirst, changed behaviour or aggressive behaviour (sometimes directed against others), feeling sleepy, cold sores, fungal or bacterial infections, irritation of prostate gland, increased need to pass urine, ear or respiratory infections, sinusitis, stuffy or runny nose, abnormal hair texture, sensitivity to sunlight, migraine headache, eye pain or infection, blurred vision, puffy face, puffy hands or feet, enlarged liver, problem affecting ovary or vagina, pain in breast, difficulty in speaking and swollen glands.

The following rare and very rare events have been reported with PegIntron:

Rarely reported side effects (greater than 1 in every 10,000 patients, but less than 1 in every 1,000 patients):

Diabetes, abnormal heart rhythm, inflammation of pancreas, inflammation and degeneration of muscle tissue and peripheral nerves, kidney problems and seizures.

Very rarely reported side effects (less than 1 in every 10,000 patients):

Very rarely sarcoidosis (a disease characterised by persistent fever, weight loss, joint pain and swelling, skin lesions and swollen glands) has been reported. Loss of consciousness has occurred very rarely with alpha interferons, mostly in elderly patients treated at high doses. Cases of stroke (cerebrovascular events) have been reported. Check with your doctor immediately if you have any of these symptoms or any other symptoms that are troubling.

Very rarely, PegIntron alone or in combination with ribavirin may cause aplastic anaemia. Pure red cell aplasia, a condition where the body stopped or reduced the production of red blood cells, has been reported. This causes severe anaemia, symptoms of which would include unusual tiredness and a lack of energy.

When PegIntron is used alone, some of these effects are less likely to occur, and some have not occurred at all.

If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis, liver failure, and development of blood abnormalities (reduction in number of red blood cells which carry oxygen, certain white blood cells that fight infection, and blood clotting cells called platelets).

The following other side effects (not listed above) have occurred with the combination of PegIntron and ribavirin capsules (adults) in HCV/HIV co-infected patients receiving HAART:

oral candidiasis (oral thrush), defective metabolism of fat, CD4 lymphocytes decreased, appetite decreased, back pain, hepatitis, limb pain, and various laboratory blood values abnormalities.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE PEGINTRON**

Keep out of the reach and sight of children.

Store in a refrigerator (2°C - 8°C).

Do not use PegIntron after the expiry date which is stated on the carton.

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

Do not use PegIntron if you notice discolouration of the powder.

The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. Discard any unused material.

## **6. FURTHER INFORMATION**

### **What PegIntron contains**

- The active substance is peginterferon alfa-2b, 50 micrograms/0.5 ml.
- The other ingredients are:  
Powder: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80;  
Solvent: water for injections 0.7 ml/ampoule.

### **What PegIntron looks like and contents of the pack**

The pharmaceutical form is: powder and solvent for solution for injection.

The white powder is contained in a 2 ml glass vial and the clear and colourless solvent is presented in a 2 ml glass ampoule.

PegIntron 50 micrograms is available in different pack sizes:

- 1 vial of powder for solution for injection and 1 ampoule of solvent for parenteral use;
- 1 vial of powder for solution for injection, 1 ampoule of solvent for parenteral use, 1 injection syringe, 2 injection needles and 1 cleansing swab;
- 4 vials of powder for solution for injection and 4 ampoules of solvent for parenteral use;
- 4 vials of powder for solution for injection, 4 ampoules of solvent for parenteral use, 4 injection syringes, 8 injection needles and 4 cleansing swabs;
- 6 vials of powder for solution for injection and 6 ampoules of solvent for parenteral use;
- 12 vials of powder for solution for injection, 12 ampoules of solvent for parenteral use, 12 injection syringes, 24 injection needles and 12 cleansing swabs.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

Marketing Authorisation Holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

Manufacturer: SP (Brinny) Company, Innishannon, County Cork, Ireland

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Rue de Stalle/Stallestraat 73  
B-1180 Bruxelles/Brussel/Brüssel  
Tél/Tel: + 32-(0)2 370 92 11

**Luxembourg/Luxemburg**

Rue de Stalle 73  
B-1180 Bruxelles/Brüssel  
Belgique/Belgien  
Tél/Tel: + 32-(0)2 370 92 11

**България**

ул. "Проф. Милко Бичев" № 7, ет.3  
BG-София 1504  
Тел.: +359 2 944 1073

**Magyarország**

Alkotás u. 53.  
H-1123 Budapest  
Tel.: +36 1 457-8500

**Česká republika**

Na Příkopě 25  
CZ-110 00 Praha 1  
Tel: +420 221771250

**Malta**

168 Christopher Street  
MT-VLT02 Valletta  
Tel: + 356-21 23 21 75

**Danmark**

Lautrupbjerg 2  
DK-2750 Ballerup  
Tlf: + 45-44 39 50 00

**Nederland**

Maarssenbroeksedijk 4  
NL-3542 DN Utrecht  
Tel: + 31-(0)800 778 78 78

**Deutschland**

Thomas-Dehler-Straße 27  
D-81737 München  
Tel: + 49-(0)89 627 31-0

**Norge**

Pb. 398  
N-1326 Lysaker  
Tlf: + 47 67 16 64 50

**Eesti**

Järvevana tee 9  
EE-11314 Tallinn  
Tel: + 372 654 96 86

**Österreich**

Badener Strasse 23  
A-2514 Traiskirchen  
Tel: + 43-(0)2252 502-0

**Ελλάδα**

Αγίου Δημητρίου 63  
GR-174 55 Αλιμος  
Τηλ.: + 30-210 98 97 300

**Polska**

Al. Jerozolimskie 195a  
PL-02-222 Warszawa  
Tel.: + 48-(0)22 478 41 50

**España**

Km. 36, Ctra. Nacional I  
E-28750 San Agustín de Guadalix – Madrid  
Tel: + 34-91 848 85 00

**Portugal**

Rua Agualva dos Açores 16  
P-2735-557 Agualva-Cacém  
Tel: +351-21 433 93 00

**France**

92 rue Baudin  
F-92300 Levallois-Perret  
Tél: + 33-(0)1 41 06 35 00

**România**

Șos. București-Ploiești, nr. 17-21, Băneasa  
Center, et. 7, sector 1  
RO-013682 București  
Tel: + 40 21 233 35 30

**Ireland**

Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW  
Tel: +44-(0)1 707 363 636

**Slovenija**

Dunajska 22  
SI-1000 Ljubljana  
Tel: + 386 01 3001070

**Ísland**

Hörgatún 2  
IS-210 Garðabær  
Sími: + 354 535 70 00

**Slovenská republika**

Strakova 5  
SK-811 01 Bratislava  
Tel: + 421 (2) 5920 2712

**Italia**

Via fratelli Cervi snc,  
Centro Direzionale Milano Due  
Palazzo Borromini  
I-20090 Segrate (Milano)  
Tel: + 39-02 21018.1

**Κύπρος**

Οδός Αγίου Νικολάου, 8  
CY-1055 Λευκωσία  
Τηλ: +357-22 757188

**Latvija**

Bauskas 58a -401  
Rīga, LV-1004  
Tel: + 371-7 21 38 25

**Lietuva**

Kęstučio g. 65/40  
LT-08124 Vilnius  
Tel: + 370 52 101868

**Suomi/Finland**

PL 3/PB 3  
FIN-02201 Espoo/Esbo  
Puh/Tel: + 358-(0)20-7570 300

**Sverige**

Box 27190  
S-102 52 Stockholm  
Tel: + 46-(0)8 522 21 500

**United Kingdom**

Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW - UK  
Tel: + 44-(0)1 707 363 636

**This leaflet was last approved on**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.ema.europa.eu>

## HOW TO SELF-INJECT PEGINTRON?

The following instructions explain how to inject PegIntron yourself. Please read the instructions carefully and follow them step by step. Your doctor or his/her assistant will instruct you how to self-inject PegIntron. Do not attempt to inject yourself unless you are sure you understand the procedure and requirement of self-injection.

### Preparation

Collect necessary items before you begin:

- a vial of PegIntron powder for injection;
- an ampoule of solvent for PegIntron (water for injections);
- a 1 ml syringe;
- a long needle (for example  $0.8 \times 40$  mm [21 gauge 1.5 inch]) to be used to add water for injections to the PegIntron powder vial;
- a short needle (for example  $0.3 \times 13$  mm [30 gauge 0.5 inch]) for the subcutaneous injection;
- a cleansing swab.

Wash your hands carefully.

### Reconstituting PegIntron powder for injection

Before reconstitution, PegIntron may appear either as a white, tablet-shaped solid that is whole or in pieces, or as a white powder.

Remove the protective cap from the PegIntron vial. Clean the rubber top of the vial with a cleansing swab. You can save the swab to clean the skin area where you will inject the dose. Remove the syringe from the wrapping. Do not touch the tip of the syringe. Take the long needle and place it firmly on to the tip of the syringe. Remove the needle guard without touching the needle and keep the syringe with the needle in your hand. Tap the top of the ampoule of solvent gently to make sure that all the liquid is at the bottom of the ampoule. Break off the top of the ampoule of solvent. Insert the needle in the ampoule of solvent and withdraw the total amount of solvent.

When the total amount of solvent is combined with the full amount of PegIntron powder, the solution will be at the correct concentration to measure your dose (i.e., the labelled amount is contained in 0.5 ml).

A small volume is lost during preparation of PegIntron for injection and when the dose is measured and injected. Therefore, each vial contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection.

To prepare the PegIntron solution, insert the needle through the rubber top of the PegIntron vial and gently place the needle tip against the glass wall of the vial without touching the cleaned top of the vial with your hands.

Inject the solvent **SLOWLY**, aiming the stream of liquid at the glass wall of the vial. It is best not to aim the stream directly at the white solid or powder, or to inject the liquid quickly, as this causes a greater amount of bubbles. The solution may appear cloudy or bubbly for a few minutes. This is to be expected and is not cause for concern.

To dissolve the entire contents, swirl the PegIntron vial with a gentle rotary motion leaving the needle and attached syringe in the vial. **Do not shake**, but gently turn the vial upside down until any powder at the top of the vial is dissolved. The contents should now be completely dissolved. Stand the vial upright and let any bubbles present in the solution rise to the top of the solution. Once the solution has settled and all bubbles have risen to the top of the solution, you should have a clear solution with a small ring of tiny bubbles around the top. Now you can withdraw your dose from the vial. Use this solution immediately. If it cannot be used immediately, the solution may be refrigerated for up to 24 hours.

### Measuring the dose of PegIntron from the reconstituted powder for injection

Turn the vial and the syringe upside down in one hand. Be sure the tip of needle is in the PegIntron reconstituted solution. Your other hand will be free to move the plunger. Pull back on the plunger



slowly to draw just more than the dose prescribed by your doctor into the syringe. Hold the syringe with the needle in the vial pointing up, remove the syringe from the long needle leaving the needle in the vial and without touching the tip of the syringe. Take the short needle and place it firmly on to the tip of the syringe. Remove the needle guard from the syringe needle and check for air bubbles in the syringe. If you see any bubbles, pull the plunger slightly back; tap the syringe gently, with the needle pointing upwards, until the bubbles disappear. Push up the plunger slowly back to the correct dose. Replace the needle guard and place the syringe with the needle on a flat surface.

Be sure the solution is at room temperature up to 25°C. If the solution is cold, warm the syringe between your palms. Inspect visually the reconstituted solution prior to administration: do not use if discolouration or particulate matter is present. You are now ready to inject the dose.

#### Injecting the solution

Select the injection site. The best sites for injection are tissues with a layer of fat between skin and muscle: thigh, outer surface of the upper arm (you may need the assistance of another person to use this site), abdomen (except the navel or waistline). If you are exceptionally thin, use only the thigh or outer surface of the arm for injection.

#### **Change your injection site each time.**

Cleanse and disinfect the skin where the injection is to be made. Wait for the area to dry. Remove the needle guard. With one hand, pinch a fold of loose skin. With your other hand, hold the syringe as you would a pencil. Insert the needle into the pinched skin at an angle of approximately 45°. After the needle is in, remove the hand used to pinch the skin and use it to hold the syringe barrel. Pull back the plunger very slightly with one hand. If blood comes into the syringe, the needle has entered a blood vessel. Do not inject into this site; withdraw the needle and repeat the procedure. Inject the solution by pushing the plunger all the way down gently.

Pull the needle straight out of the skin. Press the injection site with a small bandage or sterile gauze if necessary for several seconds. Do not massage the injection site. If there is bleeding, cover with an adhesive bandage.

The vial, ampoule and injection materials intended for single use must be discarded. Dispose of the syringe and needles safely in a closed container.

## **PACKAGE LEAFLET: INFORMATION FOR THE USER**

### **PegIntron 80 micrograms powder and solvent for solution for injection**

peginterferon alfa-2b (conjugation of recombinant interferon alfa-2b with monomethoxy polyethylene glycol)

**Read all of this leaflet carefully before you start using this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

#### **In this leaflet:**

1. What PegIntron is and what it is used for
2. Before you use PegIntron
3. How to use PegIntron
4. Possible side effects
5. Storing PegIntron
6. Further information

## **1. WHAT PEGINTRON IS AND WHAT IT IS USED FOR**

Interferons modify the response of the body's immune system to help fight infections and severe diseases. PegIntron, which contains an interferon, is used for the treatment of chronic hepatitis C, a viral infection of the liver, including in first time users co-infected with clinically stable HIV.

PegIntron is best used for this treatment in combination with ribavirin.

PegIntron is used alone in case of intolerance or contraindication to ribavirin.

## **2. BEFORE YOU USE PEGINTRON**

PegIntron is not recommended for use in patients under the age of 18 years.

#### **Do not use PegIntron**

- If you are allergic (hypersensitive) to peginterferon alfa-2b or any of the other ingredients of PegIntron.
- If you are allergic (hypersensitive) to any interferon.
- If you have had severe heart problems, or if you have heart disease that has not been well controlled during the past 6 months.
- If you have severe medical conditions that leave you very weak.
- If you have autoimmune hepatitis or any other problem with your immune system; if you are taking medicine that suppresses your immune system (your immune system protects you against infection and some diseases).
- If you have advanced, uncontrolled liver disease (other than hepatitis C).
- If you have thyroid disease that is not well controlled with medicines.
- If you have a condition that causes convulsions (seizures, or "fits").

#### **Take special care with PegIntron**

- If you develop symptoms of a severe allergic reaction (such as difficulty in breathing, wheezing, or hives) while on this medication, seek medical help immediately.
- If you ever had a heart attack or a heart problem.
- If you have ever been treated for depression or any other nervous or mental disorder.

- If you have kidney disease, your doctor may prescribe a lower than usual dose and monitor your kidney blood values regularly during treatment. If you have a kidney disease and you are using PegIntron in combination with medicinal products containing ribavirin, your doctor should monitor you more carefully for a decrease in red blood cell count.
- If you have had a problem with your liver (other than hepatitis C).
- If you develop symptoms associated with a cold or other respiratory infection, such as fever, cough, or any difficulty in breathing, tell your doctor.
- If you are diabetic, your doctor may ask you to have an eye examination.
- If you have had any serious illness affecting your breathing or your blood.
- If you have psoriasis, it may become worse while you are using PegIntron.
- If you are planning to become pregnant, discuss this with your doctor before starting to use PegIntron.
- If you are also being treated for HIV, please see **Using other medicines**.
- If you have had a severe nervous or mental disorder.
- If you ever had depression or develop symptoms associated with depression (e.g. feelings of sadness, dejection, etc) while on treatment with PegIntron (see section 4).
- If you have received an organ transplant, either kidney or liver, interferon treatment may increase the risk of rejection. Be sure to discuss this with your doctor.

Dental and gum disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and membranes of the mouth during long-term treatment with the combination of PegIntron with ribavirin. You should brush your teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If you have this reaction, be sure to rinse your mouth thoroughly afterwards.

#### **While being treated with PegIntron**

Your doctor may want you to drink extra fluids to help prevent low blood pressure.

#### **Using other medicines**

Please inform your doctor or pharmacist if you are taking or have recently taken any other medicines, even those not prescribed.

#### **Pregnancy and breast-feeding**

Ask your doctor or pharmacist for advice before taking any medicine. In studies in pregnant animals, interferons have sometimes caused miscarriage. The effect on human pregnancy is not known. In combination therapy with ribavirin, ribavirin can be very damaging to an unborn baby, thus both female and male patients must take special precautions in their sexual activity if there is any chance for pregnancy to occur:

- if you are a **girl** or a **woman** of childbearing age, you must have a negative pregnancy test before treatment, each month during treatment, and for the 4 months after treatment is stopped. You and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 4 months after stopping treatment. This can be discussed with your doctor.
- if you are a **man** who is taking ribavirin, do not have sex with a pregnant woman unless you use a condom. This will lessen the chance for ribavirin to be left in the woman's body. If your female partner is not pregnant now but is of childbearing age, she must be tested for pregnancy each month during treatment and for the 7 months after treatment has stopped. This can be discussed with your doctor. If you are a male patient, you and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 7 months after stopping treatment. This can be discussed with your doctor.

It is not known whether this product is present in human milk. Therefore, do not breast-feed an infant if you are taking PegIntron. In combination therapy with ribavirin, take notice of the respective informing texts of ribavirin containing medicinal products.

#### **Driving and using machines**

Do not drive or operate any tools or machines if you feel tired, sleepy or confused while taking PegIntron.

### **Important information about some of the ingredients of PegIntron**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

This medicinal product contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### **Patients who also have HIV infection**

Lactic acidosis and worsening liver function are side effects associated with Highly Active Anti-Retroviral Therapy (HAART), an HIV treatment. If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis and of liver failure. Your doctor will monitor you for signs and symptoms of these conditions (Please be sure to read the ribavirin Patient Leaflet also). Additionally, patients treated with PegIntron and ribavirin combination therapy and zidovudine could be at increased risk of developing anaemia (low number of red blood cells). If you are also being treated for Human Immunodeficiency Virus (HIV) infection (AIDS) with zidovudine or stavudine, it is not certain if ribavirin will change the way these medicines work. Therefore, your blood will be checked regularly to be sure that the HIV infection is not getting worse. If it gets worse, your doctor will decide whether or not your ribavirin treatment needs to be changed.

Co-administration of ribavirin and didanosine and/or stavudine is not recommended due to the risk of lactic acidosis (a build-up of lactic acid in the body) and pancreatitis.

## **3. HOW TO USE PEGINTRON**

Your doctor has prescribed PegIntron specifically for you and your current condition; do not share this medicine with anyone else.

Your doctor has determined your dose of PegIntron based on your weight. If necessary, the dose may be changed during treatment.

### **Combination treatment**

PegIntron, when given with ribavirin capsules, is usually given at a dose of 1.5 microgram/kg once a week.

Ribavirin capsules are taken every day, morning and evening. The number of ribavirin capsules you take depends on your weight.

- If you weigh less than 65 kg, take 2 capsules in the morning and 2 in the evening (total of 800 mg each day).
- If you weigh between 65 and 85 kg, take 2 capsules in the morning and 3 in the evening (total of 1,000 mg each day).
- If you weigh more than 85 kg, take 3 capsules in the morning and 3 in the evening (total 1,200 mg each day).

The combination treatment is continued for 3 to 6 months, and sometimes for one year depending on your physicians judgement. Take notice of the respective informing texts of ribavirin containing medicinal products.

For HCV/HIV co-infected patients, the duration of treatment is 48 weeks.

### **PegIntron alone:**

PegIntron, when given alone, is usually given at a dose of 0.5 or 1.0 microgram/kg once a week, for at least 6 months, and possibly for 1 year. If you have kidney disease, your dose may be lower, depending upon your kidney function.

PegIntron alone was not studied in HCV/HIV co-infected patients.

**All patients:**

If you are injecting PegIntron yourself, please be sure that the dose that has been prescribed for you is clearly provided on the package of medicine you receive.

If you have the impression that the effect of PegIntron is too strong or too weak, talk to your doctor or pharmacist.

PegIntron is intended for subcutaneous use. This means that it is injected through a short injection needle into the fatty tissue just under your skin. If you are injecting this medicine yourself, you will be instructed how to prepare and give the injection. Detailed instructions for subcutaneous administration are provided with this leaflet (see How to self-inject PegIntron at the end of the package leaflet).

Prepare the dose just before you intend to inject it and use it immediately. Look carefully at the reconstituted solution prior to administration. Do not use if there is discolouration of the reconstituted solution or if particulate matter is present. Discard any solution that is left in the vial after you give yourself the injection.

Inject PegIntron once each week on the same day. Injecting it at the same time of day each week will help you not to forget to take it.

Always use PegIntron exactly as your doctor has told you. Do not exceed the recommended dosage, and take it for as long as prescribed.

**If you use more PegIntron than you should**

Tell your doctor or healthcare professional as soon as possible.

**If you forget to take PegIntron**

Take the dose as soon as you remember, then continue your treatment as usual.

Do not take a double dose to make up for a forgotten dose. Contact your doctor or pharmacist if needed.

**4. POSSIBLE SIDE EFFECTS**

Like all medicines, PegIntron can cause side effects, although not everybody gets them. Although not all of these side effects may occur, they may need medical attention if they do.

Some people get depressed when taking PegIntron alone or in combination treatment with ribavirin, and in some cases people had suicidal thoughts or aggressive behaviour (sometimes directed against others). Some patients have actually committed suicide. Be sure to seek emergency care if you notice that you are becoming depressed or have suicidal thoughts or change in your behaviour. You may want to consider asking a family member or close friend to help you stay alert to signs of depression or changes in your behaviour.

Check with your doctor immediately if any of the following side effects occur: chest pain; changes in the way your heart beats; breathing problems (including shortness of breath), confusion; feeling depressed, wanting to harm yourself, hallucinations, numbness or tingling feeling; dizziness, convulsion ("fit"); trouble sleeping, thinking or concentrating; difficulty remaining alert, severe stomach pain or cramps; blood or clots in stool (or black, tarry stool); fever or chills beginning after a few weeks of treatment, pain in your lower back or side; difficulty or inability to pass urine, painful or inflamed muscles (sometimes severe); problems with your eyes or your eyesight or hearing; severe or painful reddening of your skin or mucous membrane, severe bleeding from your nose. Your doctor will test your blood to ensure that your white blood cell (cells that fight infection) and red blood cell (cells that carry oxygen) counts, platelets (blood clotting cells) and other laboratory values are at acceptable levels.

Other side effects that may occur with the combination of PegIntron and ribavirin capsules include:

Very commonly reported side effects (greater than 1 in every 10 patients):

Irritation or redness (and rarely, skin damage) at the site of injection, headache, tired feeling, shaking chills, fever, flu-like symptoms, weakness, loss of weight, nausea, loss of appetite, diarrhoea or loose stools, stomach pain, vomiting, muscle aches, pain in joints and muscles, feeling depressed, irritability, trouble falling asleep or staying asleep, feeling anxious or nervous, difficulty concentrating, mood swings, hair loss, itching, dry skin, sore throat, coughing, difficult breathing, dizziness, virus infection, rash, and dry mouth.

Commonly reported side effects (at least 1 in every 100 patients, but not greater than 1 in every 10 patients):

Increased sweating, chest pain, pain on the right side around your ribs, numbness, pain or tingling feeling, change in thyroid gland activity (which may make you feel tired or, less commonly, energetic), stomach upset, rapid heart rate, agitation, nervousness, difficult or irregular menstrual period

Less common are pain at the place of injection, flushing, low or high blood pressure, dry or teary eyes, redness of skin or skin disorders, psoriasis, hives, nail disorder, feeling unwell, feeling faint, poor coordination, confusion, increased or decreased sensitivity to touch, tense muscles, arthritis, bruising, loss of interest in activities including sex, sexual problem, unusual dreams, shaky hands, vertigo (spinning feeling), increased appetite, heartburn, intestinal gas (flatulence), constipation, hemorrhoids, red or bleeding gums, redness or sores in mouth, change in taste, changes in hearing or ringing in ears, thirst, changed behaviour or aggressive behaviour (sometimes directed against others), feeling sleepy, cold sores, fungal or bacterial infections, irritation of prostate gland, increased need to pass urine, ear or respiratory infections, sinusitis, stuffy or runny nose, abnormal hair texture, sensitivity to sunlight, migraine headache, eye pain or infection, blurred vision, puffy face, puffy hands or feet, enlarged liver, problem affecting ovary or vagina, pain in breast, difficulty in speaking and swollen glands.

The following rare and very rare events have been reported with PegIntron:

Rarely reported side effects (greater than 1 in every 10,000 patients, but less than 1 in every 1,000 patients):

Diabetes, abnormal heart rhythm, inflammation of pancreas, inflammation and degeneration of muscle tissue and peripheral nerves, kidney problems and seizures.

Very rarely reported side effects (less than 1 in every 10,000 patients):

Very rarely sarcoidosis (a disease characterised by persistent fever, weight loss, joint pain and swelling, skin lesions and swollen glands) has been reported. Loss of consciousness has occurred very rarely with alpha interferons, mostly in elderly patients treated at high doses. Cases of stroke (cerebrovascular events) have been reported. Check with your doctor immediately if you have any of these symptoms or any other symptoms that are troubling.

Very rarely, PegIntron alone or in combination with ribavirin may cause aplastic anaemia. Pure red cell aplasia, a condition where the body stopped or reduced the production of red blood cells, has been reported. This causes severe anaemia, symptoms of which would include unusual tiredness and a lack of energy.

When PegIntron is used alone, some of these effects are less likely to occur, and some have not occurred at all.

If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis, liver failure, and development of blood abnormalities (reduction in number of red blood cells which carry oxygen, certain white blood cells that fight infection, and blood clotting cells called platelets).

The following other side effects (not listed above) have occurred with the combination of PegIntron and ribavirin capsules (adults) in HCV/HIV co-infected patients receiving HAART: oral candidiasis (oral thrush), defective metabolism of fat, CD4 lymphocytes decreased, appetite decreased, back pain, hepatitis, limb pain, and various laboratory blood values abnormalities.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE PEGINTRON**

Keep out of the reach and sight of children.

Store in a refrigerator (2°C - 8°C).

Do not use PegIntron after the expiry date which is stated on the carton.

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

Do not use PegIntron if you notice discolouration of the powder.

The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. Discard any unused material.

## **6. FURTHER INFORMATION**

### **What PegIntron contains**

- The active substance is peginterferon alfa-2b, 80 micrograms/0.5 ml.
- The other ingredients are:  
Powder: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80;  
Solvent: water for injections 0.7 ml/ampoule.

### **What PegIntron looks like and contents of the pack**

The pharmaceutical form is: powder and solvent for solution for injection.

The white powder is contained in a 2 ml glass vial and the clear and colourless solvent is presented in a 2 ml glass ampoule.

PegIntron 80 micrograms is available in different pack sizes:

- 1 vial of powder for solution for injection and 1 ampoule of solvent for parenteral use;
- 1 vial of powder for solution for injection, 1 ampoule of solvent for parenteral use, 1 injection syringe, 2 injection needles and 1 cleansing swab;
- 4 vials of powder for solution for injection and 4 ampoules of solvent for parenteral use;
- 4 vials of powder for solution for injection, 4 ampoules of solvent for parenteral use, 4 injection syringes, 8 injection needles and 4 cleansing swabs;
- 6 vials of powder for solution for injection and 6 ampoules of solvent for parenteral use;
- 12 vials of powder for solution for injection, 12 ampoules of solvent for parenteral use, 12 injection syringes, 24 injection needles and 12 cleansing swabs.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

Marketing Authorisation Holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

Manufacturer: SP (Brinny) Company, Innishannon, County Cork, Ireland

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Rue de Stalle/Stallestraat 73  
B-1180 Bruxelles/Brussel/Brüssel  
Tél/Tel: + 32-(0)2 370 92 11

**Luxembourg/Luxemburg**

Rue de Stalle 73  
B-1180 Bruxelles/Brüssel  
Belgique/Belgien  
Tél/Tel: + 32-(0)2 370 92 11

**България**

ул. "Проф. Милко Бичев" № 7, ет.3  
BG-София 1504  
Тел.: +359 2 944 1073

**Magyarország**

Alkotás u. 53.  
H-1123 Budapest  
Tel.: +36 1 457-8500

**Česká republika**

Na Příkopě 25  
CZ-110 00 Praha 1  
Tel: +420 221771250

**Malta**

168 Christopher Street  
MT-VLT02 Valletta  
Tel: + 356-21 23 21 75

**Danmark**

Lautrupbjerg 2  
DK-2750 Ballerup  
Tlf: + 45-44 39 50 00

**Nederland**

Maarssenbroeksedijk 4  
NL-3542 DN Utrecht  
Tel: + 31-(0)800 778 78 78

**Deutschland**

Thomas-Dehler-Straße 27  
D-81737 München  
Tel: + 49-(0)89 627 31-0

**Norge**

Pb. 398  
N-1326 Lysaker  
Tlf: + 47 67 16 64 50

**Eesti**

Järvevana tee 9  
EE-11314 Tallinn  
Tel: + 372 654 96 86

**Österreich**

Badener Strasse 23  
A-2514 Traiskirchen  
Tel: + 43-(0)2252 502-0

**Ελλάδα**

Αγίου Δημητρίου 63  
GR-174 55 Αλιμος  
Τηλ.: + 30-210 98 97 300

**Polska**

Al. Jerozolimskie 195a  
PL-02-222 Warszawa  
Tel.: + 48-(0)22 478 41 50

**España**

Km. 36, Ctra. Nacional I  
E-28750 San Agustín de Guadalix – Madrid  
Tel: + 34-91 848 85 00

**Portugal**

Rua Agualva dos Açores 16  
P-2735-557 Agualva-Cacém  
Tel: +351-21 433 93 00

**France**

92 rue Baudin  
F-92300 Levallois-Perret  
Tél: + 33-(0)1 41 06 35 00

**România**

Șos. București-Ploiești, nr. 17-21, Băneasa  
Center, et. 7, sector 1  
RO-013682 București  
Tel: + 40 21 233 35 30

**Ireland**

Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW  
Tel: +44-(0)1 707 363 636

**Slovenija**

Dunajska 22  
SI-1000 Ljubljana  
Tel: + 386 01 3001070

**Ísland**

Hörgatún 2  
IS-210 Garðabær  
Sími: + 354 535 70 00

**Slovenská republika**

Strakova 5  
SK-811 01 Bratislava  
Tel: + 421 (2) 5920 2712



**Italia**

Via fratelli Cervi snc,  
Centro Direzionale Milano Due  
Palazzo Borromini  
I-20090 Segrate (Milano)  
Tel: + 39-02 21018.1

**Κύπρος**

Οδός Αγίου Νικολάου, 8  
CY-1055 Λευκωσία  
Τηλ: +357-22 757188

**Latvija**

Bauskas 58a -401  
Rīga, LV-1004  
Tel: + 371-7 21 38 25

**Lietuva**

Kęstučio g. 65/40  
LT-08124 Vilnius  
Tel: + 370 52 101868

**Suomi/Finland**

PL 3/PB 3  
FIN-02201 Espoo/Esbo  
Puh/Tel: + 358-(0)20-7570 300

**Sverige**

Box 27190  
S-102 52 Stockholm  
Tel: + 46-(0)8 522 21 500

**United Kingdom**

Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW - UK  
Tel: + 44-(0)1 707 363 636

**This leaflet was last approved on**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>

## HOW TO SELF-INJECT PEGINTRON?

The following instructions explain how to inject PegIntron yourself. Please read the instructions carefully and follow them step by step. Your doctor or his/her assistant will instruct you how to self-inject PegIntron. Do not attempt to inject yourself unless you are sure you understand the procedure and requirement of self-injection.

### Preparation

Collect necessary items before you begin:

- a vial of PegIntron powder for injection;
- an ampoule of solvent for PegIntron (water for injections);
- a 1 ml syringe;
- a long needle (for example  $0.8 \times 40$  mm [21 gauge 1.5 inch]) to be used to add water for injections to the PegIntron powder vial;
- a short needle (for example  $0.3 \times 13$  mm [30 gauge 0.5 inch]) for the subcutaneous injection;
- a cleansing swab.

Wash your hands carefully.

### Reconstituting PegIntron powder for injection

Before reconstitution, PegIntron may appear either as a white, tablet-shaped solid that is whole or in pieces, or as a white powder.

Remove the protective cap from the PegIntron vial. Clean the rubber top of the vial with a cleansing swab. You can save the swab to clean the skin area where you will inject the dose. Remove the syringe from the wrapping. Do not touch the tip of the syringe. Take the long needle and place it firmly on to the tip of the syringe. Remove the needle guard without touching the needle and keep the syringe with the needle in your hand. Tap the top of the ampoule of solvent gently to make sure that all the liquid is at the bottom of the ampoule. Break off the top of the ampoule of solvent. Insert the needle in the ampoule of solvent and withdraw the total amount of solvent.

When the total amount of solvent is combined with the full amount of PegIntron powder, the solution will be at the correct concentration to measure your dose (i.e., the labelled amount is contained in 0.5 ml).

A small volume is lost during preparation of PegIntron for injection and when the dose is measured and injected. Therefore, each vial contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection.

To prepare the PegIntron solution, insert the needle through the rubber top of the PegIntron vial and gently place the needle tip against the glass wall of the vial without touching the cleaned top of the vial with your hands.

Inject the solvent **SLOWLY**, aiming the stream of liquid at the glass wall of the vial. It is best not to aim the stream directly at the white solid or powder, or to inject the liquid quickly, as this causes a greater amount of bubbles. The solution may appear cloudy or bubbly for a few minutes. This is to be expected and is not cause for concern.

To dissolve the entire contents, swirl the PegIntron vial with a gentle rotary motion leaving the needle and attached syringe in the vial. **Do not shake**, but gently turn the vial upside down until any powder at the top of the vial is dissolved. The contents should now be completely dissolved. Stand the vial upright and let any bubbles present in the solution rise to the top of the solution. Once the solution has settled and all bubbles have risen to the top of the solution, you should have a clear solution with a small ring of tiny bubbles around the top. Now you can withdraw your dose from the vial. Use this solution immediately. If it cannot be used immediately, the solution may be refrigerated for up to 24 hours.

### Measuring the dose of PegIntron from the reconstituted powder for injection

Turn the vial and the syringe upside down in one hand. Be sure the tip of needle is in the PegIntron reconstituted solution. Your other hand will be free to move the plunger. Pull back on the plunger

slowly to draw just more than the dose prescribed by your doctor into the syringe. Hold the syringe with the needle in the vial pointing up, remove the syringe from the long needle leaving the needle in the vial and without touching the tip of the syringe. Take the short needle and place it firmly on to the tip of the syringe. Remove the needle guard from the syringe needle and check for air bubbles in the syringe. If you see any bubbles, pull the plunger slightly back; tap the syringe gently, with the needle pointing upwards, until the bubbles disappear. Push up the plunger slowly back to the correct dose. Replace the needle guard and place the syringe with the needle on a flat surface.

Be sure the solution is at room temperature up to 25°C. If the solution is cold, warm the syringe between your palms. Inspect visually the reconstituted solution prior to administration: do not use if discolouration or particulate matter is present. You are now ready to inject the dose.

#### Injecting the solution

Select the injection site. The best sites for injection are tissues with a layer of fat between skin and muscle: thigh, outer surface of the upper arm (you may need the assistance of another person to use this site), abdomen (except the navel or waistline). If you are exceptionally thin, use only the thigh or outer surface of the arm for injection.

#### **Change your injection site each time.**

Cleanse and disinfect the skin where the injection is to be made. Wait for the area to dry. Remove the needle guard. With one hand, pinch a fold of loose skin. With your other hand, hold the syringe as you would a pencil. Insert the needle into the pinched skin at an angle of approximately 45°. After the needle is in, remove the hand used to pinch the skin and use it to hold the syringe barrel. Pull back the plunger very slightly with one hand. If blood comes into the syringe, the needle has entered a blood vessel. Do not inject into this site; withdraw the needle and repeat the procedure. Inject the solution by pushing the plunger all the way down gently.

Pull the needle straight out of the skin. Press the injection site with a small bandage or sterile gauze if necessary for several seconds. Do not massage the injection site. If there is bleeding, cover with an adhesive bandage.

The vial, ampoule and injection materials intended for single use must be discarded. Dispose of the syringe and needles safely in a closed container.

## **PACKAGE LEAFLET: INFORMATION FOR THE USER**

### **PegIntron 100 micrograms powder and solvent for solution for injection**

peginterferon alfa-2b (conjugation of recombinant interferon alfa-2b with monomethoxy polyethylene glycol)

**Read all of this leaflet carefully before you start using this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What PegIntron is and what it is used for
2. Before you use PegIntron
3. How to use PegIntron
4. Possible side effects
5. Storing PegIntron
6. Further information

## **1. WHAT PEGINTRON IS AND WHAT IT IS USED FOR**

Interferons modify the response of the body's immune system to help fight infections and severe diseases. PegIntron, which contains an interferon, is used for the treatment of chronic hepatitis C, a viral infection of the liver, including in first time users co-infected with clinically stable HIV.

PegIntron is best used for this treatment in combination with ribavirin.

PegIntron is used alone in case of intolerance or contraindication to ribavirin.

## **2. BEFORE YOU USE PEGINTRON**

PegIntron is not recommended for use in patients under the age of 18 years.

**Do not use PegIntron**

- If you are allergic (hypersensitive) to peginterferon alfa-2b or any of the other ingredients of PegIntron.
- If you are allergic (hypersensitive) to any interferon.
- If you have had severe heart problems, or if you have heart disease that has not been well controlled during the past 6 months.
- If you have severe medical conditions that leave you very weak.
- If you have autoimmune hepatitis or any other problem with your immune system; if you are taking medicine that suppresses your immune system (your immune system protects you against infection and some diseases).
- If you have advanced, uncontrolled liver disease (other than hepatitis C).
- If you have thyroid disease that is not well controlled with medicines.
- If you have a condition that causes convulsions (seizures, or "fits").

**Take special care with PegIntron**

- If you develop symptoms of a severe allergic reaction (such as difficulty in breathing, wheezing, or hives) while on this medication, seek medical help immediately.
- If you ever had a heart attack or a heart problem.
- If you have ever been treated for depression or any other nervous or mental disorder.

- If you have kidney disease, your doctor may prescribe a lower than usual dose and monitor your kidney blood values regularly during treatment. If you have a kidney disease and you are using PegIntron in combination with medicinal products containing ribavirin, your doctor should monitor you more carefully for a decrease in red blood cell count.
- If you have had a problem with your liver (other than hepatitis C).
- If you develop symptoms associated with a cold or other respiratory infection, such as fever, cough, or any difficulty in breathing, tell your doctor.
- If you are diabetic, your doctor may ask you to have an eye examination.
- If you have had any serious illness affecting your breathing or your blood.
- If you have psoriasis, it may become worse while you are using PegIntron.
- If you are planning to become pregnant, discuss this with your doctor before starting to use PegIntron.
- If you are also being treated for HIV, please see **Using other medicines**.
- If you have had a severe nervous or mental disorder.
- If you ever had depression or develop symptoms associated with depression (e.g. feelings of sadness, dejection, etc) while on treatment with PegIntron (see section 4).
- If you have received an organ transplant, either kidney or liver, interferon treatment may increase the risk of rejection. Be sure to discuss this with your doctor.

Dental and gum disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and membranes of the mouth during long-term treatment with the combination of PegIntron with ribavirin. You should brush your teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If you have this reaction, be sure to rinse your mouth thoroughly afterwards.

#### **While being treated with PegIntron**

Your doctor may want you to drink extra fluids to help prevent low blood pressure.

#### **Using other medicines**

Please inform your doctor or pharmacist if you are taking or have recently taken any other medicines, even those not prescribed.

#### **Pregnancy and breast-feeding**

Ask your doctor or pharmacist for advice before taking any medicine. In studies in pregnant animals, interferons have sometimes caused miscarriage. The effect on human pregnancy is not known. In combination therapy with ribavirin, ribavirin can be very damaging to an unborn baby, thus both female and male patients must take special precautions in their sexual activity if there is any chance for pregnancy to occur:

- if you are a **girl** or a **woman** of childbearing age, you must have a negative pregnancy test before treatment, each month during treatment, and for the 4 months after treatment is stopped. You and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 4 months after stopping treatment. This can be discussed with your doctor.
- if you are a **man** who is taking ribavirin, do not have sex with a pregnant woman unless you use a condom. This will lessen the chance for ribavirin to be left in the woman's body. If your female partner is not pregnant now but is of childbearing age, she must be tested for pregnancy each month during treatment and for the 7 months after treatment has stopped. This can be discussed with your doctor. If you are a male patient, you and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 7 months after stopping treatment. This can be discussed with your doctor.

It is not known whether this product is present in human milk. Therefore, do not breast-feed an infant if you are taking PegIntron. In combination therapy with ribavirin, take notice of the respective informing texts of ribavirin containing medicinal products.

#### **Driving and using machines**

Do not drive or operate any tools or machines if you feel tired, sleepy or confused while taking PegIntron.

### **Important information about some of the ingredients of PegIntron**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

This medicinal product contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### **Patients who also have HIV infection**

Lactic acidosis and worsening liver function are side effects associated with Highly Active Anti-Retroviral Therapy (HAART), an HIV treatment. If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis and of liver failure. Your doctor will monitor you for signs and symptoms of these conditions. (Please be sure to read the ribavirin Patient Leaflet also). Additionally, patients treated with PegIntron and ribavirin combination therapy and zidovudine could be at increased risk of developing anaemia (low number of red blood cells). If you are also being treated for Human Immunodeficiency Virus (HIV) infection (AIDS) with zidovudine or stavudine, it is not certain if ribavirin will change the way these medicines work. Therefore, your blood will be checked regularly to be sure that the HIV infection is not getting worse. If it gets worse, your doctor will decide whether or not your ribavirin treatment needs to be changed.

Co-administration of ribavirin and didanosine and/or stavudine is not recommended due to the risk of lactic acidosis (a build-up of lactic acid in the body) and pancreatitis.

## **3. HOW TO USE PEGINTRON**

Your doctor has prescribed PegIntron specifically for you and your current condition; do not share this medicine with anyone else.

Your doctor has determined your dose of PegIntron based on your weight. If necessary, the dose may be changed during treatment.

### **Combination treatment**

PegIntron, when given with ribavirin capsules, is usually given at a dose of 1.5 microgram/kg once a week.

Ribavirin capsules are taken every day, morning and evening. The number of ribavirin capsules you take depends on your weight.

- If you weigh less than 65 kg, take 2 capsules in the morning and 2 in the evening (total of 800 mg each day).
- If you weigh between 65 and 85 kg, take 2 capsules in the morning and 3 in the evening (total of 1,000 mg each day).
- If you weigh more than 85 kg, take 3 capsules in the morning and 3 in the evening (total 1,200 mg each day).

The combination treatment is continued for 3 to 6 months, and sometimes for one year depending on your physicians judgement. Take notice of the respective informing texts of ribavirin containing medicinal products.

For HCV/HIV co-infected patients, the duration of treatment is 48 weeks.

### **PegIntron alone:**

PegIntron, when given alone, is usually given at a dose of 0.5 or 1.0 microgram/kg once a week, for at least 6 months, and possibly for 1 year. If you have kidney disease, your dose may be lower, depending upon your kidney function.

PegIntron alone was not studied in HCV/HIV co-infected patients.

### **All patients:**

If you are injecting PegIntron yourself, please be sure that the dose that has been prescribed for you is clearly provided on the package of medicine you receive.

If you have the impression that the effect of PegIntron is too strong or too weak, talk to your doctor or pharmacist.

PegIntron is intended for subcutaneous use. This means that it is injected through a short injection needle into the fatty tissue just under your skin. If you are injecting this medicine yourself, you will be instructed how to prepare and give the injection. Detailed instructions for subcutaneous administration are provided with this leaflet (see How to self-inject PegIntron at the end of the package leaflet).

Prepare the dose just before you intend to inject it and use it immediately. Look carefully at the reconstituted solution prior to administration. Do not use if there is discolouration of the reconstituted solution or if particulate matter is present. Discard any solution that is left in the vial after you give yourself the injection.

Inject PegIntron once each week on the same day. Injecting it at the same time of day each week will help you not to forget to take it.

Always use PegIntron exactly as your doctor told you. Do not exceed the recommended dosage, and take it for as long as prescribed.

**If you use more PegIntron than you should**

Tell your doctor or healthcare professional as soon as possible.

**If you forget to take PegIntron**

Take the dose as soon as you remember, then continue your treatment as usual.

Do not take a double dose to make up for a forgotten dose. Contact your doctor or pharmacist if needed.

#### **4. POSSIBLE SIDE EFFECTS**

Like all medicines, PegIntron can cause side effects, although not everybody gets them. Although not all of these side effects may occur, they may need medical attention if they do.

Some people get depressed when taking PegIntron alone or in combination treatment with ribavirin, and in some cases people had suicidal thoughts or aggressive behaviour (sometimes directed against others). Some patients have actually committed suicide. Be sure to seek emergency care if you notice that you are becoming depressed or have suicidal thoughts or change in your behaviour. You may want to consider asking a family member or close friend to help you stay alert to signs of depression or changes in your behaviour.

Check with your doctor immediately if any of the following side effects occur: chest pain; changes in the way your heart beats; breathing problems (including shortness of breath), confusion; feeling depressed, wanting to harm yourself, hallucinations, numbness or tingling feeling; dizziness, convulsion ("fit"); trouble sleeping, thinking or concentrating; difficulty remaining alert, severe stomach pain or cramps; blood or clots in stool (or black, tarry stool); fever or chills beginning after a few weeks of treatment, pain in your lower back or side; difficulty or inability to pass urine, painful or inflamed muscles (sometimes severe); problems with your eyes or your eyesight or hearing; severe or painful reddening of your skin or mucous membrane, severe bleeding from your nose. Your doctor will test your blood to ensure that your white blood cell (cells that fight infection) and red blood cell (cells that carry oxygen) counts, platelets (blood clotting cells) and other laboratory values are at acceptable levels.

Other side effects that may occur with the combination of PegIntron and ribavirin capsules include:

Very commonly reported side effects (greater than 1 in every 10 patients):

Irritation or redness (and rarely, skin damage) at the site of injection, headache, tired feeling, shaking chills, fever, flu-like symptoms, weakness, loss of weight, nausea, loss of appetite, diarrhoea or loose stools, stomach pain, vomiting, muscle aches, pain in joints and muscles, feeling depressed, irritability, trouble falling asleep or staying asleep, feeling anxious or nervous, difficulty concentrating, mood swings, hair loss, itching, dry skin, sore throat, coughing, difficult breathing, dizziness, virus infection, rash, and dry mouth.

Commonly reported side effects (at least 1 in every 100 patients, but not greater than 1 in every 10 patients):

Increased sweating, chest pain, pain on the right side around your ribs, numbness, pain or tingling feeling, change in thyroid gland activity (which may make you feel tired or, less commonly, energetic), stomach upset, rapid heart rate, agitation, nervousness, difficult or irregular menstrual period

Less common are pain at the place of injection, flushing, low or high blood pressure, dry or teary eyes, redness of skin or skin disorders, psoriasis, hives, nail disorder, feeling unwell, feeling faint, poor coordination, confusion, increased or decreased sensitivity to touch, tense muscles, arthritis, bruising, loss of interest in activities including sex, sexual problem, unusual dreams, shaky hands, vertigo (spinning feeling), increased appetite, heartburn, intestinal gas (flatus), constipation, hemorrhoids, red or bleeding gums, redness or sores in mouth, change in taste, changes in hearing or ringing in ears, thirst, changed behaviour or aggressive behaviour (sometimes directed against others), feeling sleepy, cold sores, fungal or bacterial infections, irritation of prostate gland, increased need to pass urine, ear or respiratory infections, sinusitis, stuffy or runny nose, abnormal hair texture, sensitivity to sunlight, migraine headache, eye pain or infection, blurred vision, puffy face, puffy hands or feet, enlarged liver, problem affecting ovary or vagina, pain in breast, difficulty in speaking and swollen glands.

The following rare and very rare events have been reported with PegIntron:

Rarely reported side effects (greater than 1 in every 10,000 patients, but less than 1 in every 1,000 patients):

Diabetes, abnormal heart rhythm, inflammation of pancreas, inflammation and degeneration of muscle tissue and peripheral nerves, kidney problems and seizures.

Very rarely reported side effects (less than 1 in every 10,000 patients):

Very rarely sarcoidosis (a disease characterised by persistent fever, weight loss, joint pain and swelling, skin lesions and swollen glands) has been reported. Loss of consciousness has occurred very rarely with alpha interferons, mostly in elderly patients treated at high doses. Cases of stroke (cerebrovascular events) have been reported. Check with your doctor immediately if you have any of these symptoms or any other symptoms that are troubling.

Very rarely, PegIntron alone or in combination with ribavirin may cause aplastic anaemia. Pure red cell aplasia, a condition where the body stopped or reduced the production of red blood cells, has been reported. This causes severe anaemia, symptoms of which would include unusual tiredness and a lack of energy.

When PegIntron is used alone, some of these effects are less likely to occur, and some have not occurred at all.

If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis, liver failure, and development of blood abnormalities (reduction in number of red blood cells which carry oxygen, certain white blood cells that fight infection, and blood clotting cells called platelets).

The following other side effects (not listed above) have occurred with the combination of PegIntron and ribavirin capsules (adults) in HCV/HIV co-infected patients receiving HAART:

oral candidiasis (oral thrush), defective metabolism of fat, CD4 lymphocytes decreased, appetite decreased,



back pain, hepatitis, limb pain, and various laboratory blood values abnormalities.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE PEGINTRON**

Keep out of the reach and sight of children.

Store in a refrigerator (2°C - 8°C).

Do not use PegIntron after the expiry date which is stated on the carton.

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

Do not use PegIntron if you notice discolouration of the powder.

The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. Discard any unused material.

## **6. FURTHER INFORMATION**

### **What PegIntron contains**

- The active substance is peginterferon alfa-2b, 100 micrograms/0.5 ml.
- The other ingredients are:  
Powder: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80;  
Solvent: water for injections 0.7 ml/ampoule.

### **What PegIntron looks like and contents of the pack**

The pharmaceutical form is: powder and solvent for solution for injection.

The white powder is contained in a 2 ml glass vial and the clear and colourless solvent is presented in a 2 ml glass ampoule.

PegIntron 100 micrograms is available in different pack sizes:

- 1 vial of powder for solution for injection and 1 ampoule of solvent for parenteral use;
- 1 vial of powder for solution for injection, 1 ampoule of solvent for parenteral use, 1 injection syringe, 2 injection needles and 1 cleansing swab;
- 4 vials of powder for solution for injection and 4 ampoules of solvent for parenteral use;
- 4 vials of powder for solution for injection, 4 ampoules of solvent for parenteral use, 4 injection syringes, 8 injection needles and 4 cleansing swabs;
- 6 vials of powder for solution for injection and 6 ampoules of solvent for parenteral use;
- 12 vials of powder for solution for injection, 12 ampoules of solvent for parenteral use, 12 injection syringes, 24 injection needles and 12 cleansing swabs.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

Marketing Authorisation Holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

Manufacturer: SP (Brinny) Company, Innishannon, County Cork, Ireland

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Rue de Stalle/Stallestraat 73  
B-1180 Bruxelles/Brussel/Brüssel  
Tél/Tel: + 32-(0)2 370 92 11

**Luxembourg/Luxemburg**

Rue de Stalle 73  
B-1180 Bruxelles/Brüssel  
Belgique/Belgien  
Tél/Tel: + 32-(0)2 370 92 11

**България**

ул. "Проф. Милко Бичев" № 7, ет.3  
BG-София 1504  
Тел.: +359 2 944 1073

**Magyarország**

Alkotás u. 53.  
H-1123 Budapest  
Tel.: +36 1 457-8500

**Česká republika**

Na Příkopě 25  
CZ-110 00 Praha 1  
Tel: +420 221771250

**Malta**

168 Christopher Street  
MT-VLT02 Valletta  
Tel: + 356-21 23 21 75

**Danmark**

Lautrupbjerg 2  
DK-2750 Ballerup  
Tlf: + 45-44 39 50 00

**Nederland**

Maarssebroeksedijk 4  
NL-3542 DN Utrecht  
Tel: + 31-(0)800 778 78 78

**Deutschland**

Thomas-Dehler-Straße 27  
D-81737 München  
Tel: + 49-(0)89 627 31-0

**Norge**

Pb. 398  
N-1326 Lysaker  
Tlf: + 47 67 16 64 50

**Eesti**

Järvevana tee 9  
EE-11314 Tallinn  
Tel: + 372 654 96 86

**Österreich**

Badener Strasse 23  
A-2514 Traiskirchen  
Tel: + 43-(0)2252 502-0

**Ελλάδα**

Αγίου Δημητρίου 63  
GR-174 55 Άλιμος  
Τηλ.: + 30-210 98 97 300

**Polska**

Al. Jerozolimskie 195a  
PL-02-222 Warszawa  
Tel.: + 48-(0)22 478 41 50

**España**

Km. 36, Ctra. Nacional I  
E-28750 San Agustín de Guadalix – Madrid  
Tel: + 34-91 848 85 00

**Portugal**

Rua Agualva dos Açores 16  
P-2735-557 Agualva-Cacém  
Tel: +351-21 433 93 00

**France**

92 rue Baudin  
F-92300 Levallois-Perret  
Tél: + 33-(0)1 41 06 35 00

**România**

Șos. București-Ploiești, nr. 17-21, Băneasa  
Center, et. 7, sector 1  
RO-013682 București  
Tel: + 40 21 233 35 30

**Ireland**

Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW  
Tel: +44-(0)1 707 363 636

**Slovenija**

Dunajska 22  
SI-1000 Ljubljana  
Tel: + 386 01 3001070

**Ísland**

Hörgatún 2  
IS-210 Garðabær  
Sími: + 354 535 70 00

**Italia**

Via fratelli Cervi snc,  
Centro Direzionale Milano Due  
Palazzo Borromini  
I-20090 Segrate (Milano)  
Tel: + 39-02 21018.1

**Κύπρος**

Οδός Αγίου Νικολάου, 8  
CY-1055 Λευκωσία  
Τηλ: +357-22 757188

**Latvija**

Bauskas 58a -401  
Rīga, LV-1004  
Tel: + 371-7 21 38 25

**Lietuva**

Kęstučio g. 65/40  
LT-08124 Vilnius  
Tel: + 370 52 101868

**Slovenská republika**

Strakova 5  
SK-811 01 Bratislava  
Tel: + 421 (2) 5920 2712

**Suomi/Finland**

PL 3/PB 3  
FIN-02201 Espoo/Esbo  
Puh/Tel: + 358-(0)20-7570 300

**Sverige**

Box 27190  
S-102 52 Stockholm  
Tel: + 46-(0)8 522 21 500

**United Kingdom**

Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW - UK  
Tel: + 44-(0)1 707 363 636

**This leaflet was last approved on**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>

## HOW TO SELF-INJECT PEGINTRON?

The following instructions explain how to inject PegIntron yourself. Please read the instructions carefully and follow them step by step. Your doctor or his/her assistant will instruct you how to self-inject PegIntron. Do not attempt to inject yourself unless you are sure you understand the procedure and requirement of self-injection.

### Preparation

Collect necessary items before you begin:

- a vial of PegIntron powder for injection;
- an ampoule of solvent for PegIntron (water for injections);
- a 1 ml syringe;
- a long needle (for example  $0.8 \times 40$  mm [21 gauge 1.5 inch]) to be used to add water for injections to the PegIntron powder vial;
- a short needle (for example  $0.3 \times 13$  mm [30 gauge 0.5 inch]) for the subcutaneous injection;
- a cleansing swab.

Wash your hands carefully.

### Reconstituting PegIntron powder for injection

Before reconstitution, PegIntron may appear either as a white, tablet-shaped solid that is whole or in pieces, or as a white powder.

Remove the protective cap from the PegIntron vial. Clean the rubber top of the vial with a cleansing swab. You can save the swab to clean the skin area where you will inject the dose. Remove the syringe from the wrapping. Do not touch the tip of the syringe. Take the long needle and place it firmly on to the tip of the syringe. Remove the needle guard without touching the needle and keep the syringe with the needle in your hand. Tap the top of the ampoule of solvent gently to make sure that all the liquid is at the bottom of the ampoule. Break off the top of the ampoule of solvent. Insert the needle in the ampoule of solvent and withdraw the total amount of solvent.

When the total amount of solvent is combined with the full amount of PegIntron powder, the solution will be at the correct concentration to measure your dose (i.e., the labelled amount is contained in 0.5 ml).

A small volume is lost during preparation of PegIntron for injection and when the dose is measured and injected. Therefore, each vial contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection.

To prepare the PegIntron solution, insert the needle through the rubber top of the PegIntron vial and gently place the needle tip against the glass wall of the vial without touching the cleaned top of the vial with your hands.

Inject the solvent **SLOWLY**, aiming the stream of liquid at the glass wall of the vial. It is best not to aim the stream directly at the white solid or powder, or to inject the liquid quickly, as this causes a greater amount of bubbles. The solution may appear cloudy or bubbly for a few minutes. This is to be expected and is not cause for concern.

To dissolve the entire contents, swirl the PegIntron vial with a gentle rotary motion leaving the needle and attached syringe in the vial. **Do not shake**, but gently turn the vial upside down until any powder at the top of the vial is dissolved. The contents should now be completely dissolved. Stand the vial upright and let any bubbles present in the solution rise to the top of the solution. Once the solution has settled and all bubbles have risen to the top of the solution, you should have a clear solution with a small ring of tiny bubbles around the top. Now you can withdraw your dose from the vial. Use this solution immediately. If it cannot be used immediately, the solution may be refrigerated for up to 24 hours.

### Measuring the dose of PegIntron from the reconstituted powder for injection

Turn the vial and the syringe upside down in one hand. Be sure the tip of needle is in the PegIntron reconstituted solution. Your other hand will be free to move the plunger. Pull back on the plunger

slowly to draw just more than the dose prescribed by your doctor into the syringe. Hold the syringe with the needle in the vial pointing up, remove the syringe from the long needle leaving the needle in the vial and without touching the tip of the syringe. Take the short needle and place it firmly on to the tip of the syringe. Remove the needle guard from the syringe needle and check for air bubbles in the syringe. If you see any bubbles, pull the plunger slightly back; tap the syringe gently, with the needle pointing upwards, until the bubbles disappear. Push up the plunger slowly back to the correct dose. Replace the needle guard and place the syringe with the needle on a flat surface.

Be sure the solution is at room temperature up to 25°C. If the solution is cold, warm the syringe between your palms. Inspect visually the reconstituted solution prior to administration: do not use if discolouration or particulate matter is present. You are now ready to inject the dose.

#### Injecting the solution

Select the injection site. The best sites for injection are tissues with a layer of fat between skin and muscle: thigh, outer surface of the upper arm (you may need the assistance of another person to use this site), abdomen (except the navel or waistline). If you are exceptionally thin, use only the thigh or outer surface of the arm for injection.

#### **Change your injection site each time.**

Cleanse and disinfect the skin where the injection is to be made. Wait for the area to dry. Remove the needle guard. With one hand, pinch a fold of loose skin. With your other hand, hold the syringe as you would a pencil. Insert the needle into the pinched skin at an angle of approximately 45°. After the needle is in, remove the hand used to pinch the skin and use it to hold the syringe barrel. Pull back the plunger very slightly with one hand. If blood comes into the syringe, the needle has entered a blood vessel. Do not inject into this site; withdraw the needle and repeat the procedure. Inject the solution by pushing the plunger all the way down gently.

Pull the needle straight out of the skin. Press the injection site with a small bandage or sterile gauze if necessary for several seconds. Do not massage the injection site. If there is bleeding, cover with an adhesive bandage.

The vial, ampoule and injection materials intended for single use must be discarded. Dispose of the syringe and needles safely in a closed container.

## **PACKAGE LEAFLET: INFORMATION FOR THE USER**

### **PegIntron 120 micrograms powder and solvent for solution for injection**

peginterferon alfa-2b (conjugation of recombinant interferon alfa-2b with monomethoxy polyethylene glycol)

#### **Read all of this leaflet carefully before you start using this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

#### **In this leaflet:**

1. What PegIntron is and what it is used for
2. Before you use PegIntron
3. How to use PegIntron
4. Possible side effects
5. Storing PegIntron
6. Further information

### **1. WHAT PEGINTRON IS AND WHAT IT IS USED FOR**

Interferons modify the response of the body's immune system to help fight infections and severe diseases. PegIntron, which contains an interferon, is used for the treatment of chronic hepatitis C, a viral infection of the liver, including in first time users co-infected with clinically stable HIV.

PegIntron is best used for this treatment in combination with ribavirin.

PegIntron is used alone in case of intolerance or contraindication to ribavirin.

### **2. BEFORE YOU USE PEGINTRON**

PegIntron is not recommended for use in patients under the age of 18 years.

#### **Do not use PegIntron**

- If you are allergic (hypersensitive) to peginterferon alfa-2b or any of the other ingredients of PegIntron.
- If you are allergic (hypersensitive) to any interferon.
- If you have had severe heart problems, or if you have heart disease that has not been well controlled during the past 6 months.
- If you have severe medical conditions that leave you very weak.
- If you have autoimmune hepatitis or any other problem with your immune system; if you are taking medicine that suppresses your immune system (your immune system protects you against infection and some diseases).
- If you have advanced, uncontrolled liver disease (other than hepatitis C).
- If you have thyroid disease that is not well controlled with medicines.
- If you have a condition that causes convulsions (seizures, or "fits").

#### **Take special care with PegIntron**

- If you develop symptoms of a severe allergic reaction (such as difficulty in breathing, wheezing, or hives) while on this medication, seek medical help immediately.
- If you ever had a heart attack or a heart problem.
- If you have ever been treated for depression or any other nervous or mental disorder.

- If you have kidney disease, your doctor may prescribe a lower than usual dose and monitor your kidney blood values regularly during treatment. If you have a kidney disease and you are using PegIntron in combination with medicinal products containing ribavirin, your doctor should monitor you more carefully for a decrease in red blood cell count.
- If you have had a problem with your liver (other than hepatitis C).
- If you develop symptoms associated with a cold or other respiratory infection, such as fever, cough, or any difficulty in breathing, tell your doctor.
- If you are diabetic, your doctor may ask you to have an eye examination.
- If you have had any serious illness affecting your breathing or your blood.
- If you have psoriasis, it may become worse while you are using PegIntron.
- If you are planning to become pregnant, discuss this with your doctor before starting to use PegIntron.
- If you are also being treated for HIV, please see **Using other medicines**.
- If you have had a severe nervous or mental disorder.
- If you ever had depression or develop symptoms associated with depression (e.g. feelings of sadness, dejection, etc) while on treatment with PegIntron (see section 4).
- If you have received an organ transplant, either kidney or liver, interferon treatment may increase the risk of rejection. Be sure to discuss this with your doctor.

Dental and gum disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and membranes of the mouth during long-term treatment with the combination of PegIntron with ribavirin. You should brush your teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If you have this reaction, be sure to rinse your mouth thoroughly afterwards.

#### **While being treated with PegIntron**

Your doctor may want you to drink extra fluids to help prevent low blood pressure.

#### **Using other medicines**

Please inform your doctor or pharmacist if you are taking or have recently taken any other medicines, even those not prescribed.

#### **Pregnancy and breast-feeding**

Ask your doctor or pharmacist for advice before taking any medicine. In studies in pregnant animals, interferons have sometimes caused miscarriage. The effect on human pregnancy is not known. In combination therapy with ribavirin, ribavirin can be very damaging to an unborn baby, thus both female and male patients must take special precautions in their sexual activity if there is any chance for pregnancy to occur:

- if you are a **girl** or a **woman** of childbearing age, you must have a negative pregnancy test before treatment, each month during treatment, and for the 4 months after treatment is stopped. You and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 4 months after stopping treatment. This can be discussed with your doctor.
- if you are a **man** who is taking ribavirin, do not have sex with a pregnant woman unless you use a condom. This will lessen the chance for ribavirin to be left in the woman's body. If your female partner is not pregnant now but is of childbearing age, she must be tested for pregnancy each month during treatment and for the 7 months after treatment has stopped. This can be discussed with your doctor. If you are a male patient, you and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 7 months after stopping treatment. This can be discussed with your doctor.

It is not known whether this product is present in human milk. Therefore, do not breast-feed an infant if you are taking PegIntron. In combination therapy with ribavirin, take notice of the respective informing texts of ribavirin containing medicinal products.

#### **Driving and using machines**

Do not drive or operate any tools or machines if you feel tired, sleepy or confused while taking PegIntron.

### **Important information about some of the ingredients of PegIntron**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

This medicinal product contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### **Patients who also have HIV infection**

Lactic acidosis and worsening liver function are side effects associated with Highly Active Anti-Retroviral Therapy (HAART), an HIV treatment. If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis and of liver failure. Your doctor will monitor you for signs and symptoms of these conditions. (Please be sure to read the ribavirin Patient Leaflet also). Additionally, patients treated with PegIntron and ribavirin combination therapy and zidovudine could be at increased risk of developing anaemia (low number of red blood cells). If you are also being treated for Human Immunodeficiency Virus (HIV) infection (AIDS) with zidovudine or stavudine, it is not certain if ribavirin will change the way these medicines work. Therefore, your blood will be checked regularly to be sure that the HIV infection is not getting worse. If it gets worse, your doctor will decide whether or not your ribavirin treatment needs to be changed.

Co-administration of ribavirin and didanosine and/or stavudine is not recommended due to the risk of lactic acidosis (a build-up of lactic acid in the body) and pancreatitis.

## **3. HOW TO USE PEGINTRON**

Your doctor has prescribed PegIntron specifically for you and your current condition; do not share this medicine with anyone else.

Your doctor has determined your dose of PegIntron based on your weight. If necessary, the dose may be changed during treatment.

### **Combination treatment**

PegIntron, when given with ribavirin capsules, is usually given at a dose of 1.5 microgram/kg once a week.

Ribavirin capsules are taken every day, morning and evening. The number of ribavirin capsules you take depends on your weight.

- If you weigh less than 65 kg, take 2 capsules in the morning and 2 in the evening (total of 800 mg each day).
- If you weigh between 65 and 85 kg, take 2 capsules in the morning and 3 in the evening (total of 1,000 mg each day).
- If you weigh more than 85 kg, take 3 capsules in the morning and 3 in the evening (total 1,200 mg each day).

The combination treatment is continued for 3 to 6 months, and sometimes for one year depending on your physicians judgement. Take notice of the respective informing texts of ribavirin containing medicinal products.

For HCV/HIV co-infected patients, the duration of treatment is 48 weeks.

### **PegIntron alone:**

PegIntron, when given alone, is usually given at a dose of 0.5 or 1.0 microgram/kg once a week, for at least 6 months, and possibly for 1 year. If you have kidney disease, your dose may be lower, depending upon your kidney function.

PegIntron alone was not studied in HCV/HIV co-infected patients.



**All patients:**

If you are injecting PegIntron yourself, please be sure that the dose that has been prescribed for you is clearly provided on the package of medicine you receive.

If you have the impression that the effect of PegIntron is too strong or too weak, talk to your doctor or pharmacist.

PegIntron is intended for subcutaneous use. This means that it is injected through a short injection needle into the fatty tissue just under your skin. If you are injecting this medicine yourself, you will be instructed how to prepare and give the injection. Detailed instructions for subcutaneous administration are provided with this leaflet (see How to self-inject PegIntron at the end of the package leaflet).

Prepare the dose just before you intend to inject it and use it immediately. Look carefully at the reconstituted solution prior to administration. Do not use if there is discolouration of the reconstituted solution or if particulate matter is present. Discard any solution that is left in the vial after you give yourself the injection.

Inject PegIntron once each week on the same day. Injecting it at the same time of day each week will help you not to forget to take it.

Always use PegIntron exactly as your doctor told you. Do not exceed the recommended dosage, and take it for as long as prescribed.

**If you use more PegIntron than you should**

Tell your doctor or healthcare professional as soon as possible.

**If you forget to take PegIntron**

Take the dose as soon as you remember, then continue your treatment as usual.

Do not take a double dose to make up for a forgotten dose. Contact your doctor or pharmacist if needed.

**4. POSSIBLE SIDE EFFECTS**

Like all medicines, PegIntron can cause side effects, although not everybody gets them. Although not all of these side effects may occur, they may need medical attention if they do.

Some people get depressed when taking PegIntron alone or in combination treatment with ribavirin, and in some cases people had suicidal thoughts or aggressive behaviour (sometimes directed against others). Some patients have actually committed suicide. Be sure to seek emergency care if you notice that you are becoming depressed or have suicidal thoughts or change in your behaviour. You may want to consider asking a family member or close friend to help you stay alert to signs of depression or changes in your behaviour.

Check with your doctor immediately if any of the following side effects occur: chest pain; changes in the way your heart beats; breathing problems (including shortness of breath), confusion; feeling depressed, wanting to harm yourself, hallucinations, numbness or tingling feeling; dizziness, convulsion ("fit"); trouble sleeping, thinking or concentrating; difficulty remaining alert, severe stomach pain or cramps; blood or clots in stool (or black, tarry stool); fever or chills beginning after a few weeks of treatment, pain in your lower back or side; difficulty or inability to pass urine, painful or inflamed muscles (sometimes severe); problems with your eyes or your eyesight or hearing; severe or painful reddening of your skin or mucous membrane, severe bleeding from your nose. Your doctor will test your blood to ensure that your white blood cell (cells that fight infection) and red blood cell (cells that carry oxygen) counts, platelets (blood clotting cells) and other laboratory values are at acceptable levels.

Other side effects that may occur with the combination of PegIntron and ribavirin capsules include:

Very commonly reported side effects (greater than 1 in every 10 patients):

Irritation or redness (and rarely, skin damage) at the site of injection, headache, tired feeling, shaking chills, fever, flu-like symptoms, weakness, loss of weight, nausea, loss of appetite, diarrhoea or loose stools, stomach pain, vomiting, muscle aches, pain in joints and muscles, feeling depressed, irritability, trouble falling asleep or staying asleep, feeling anxious or nervous, difficulty concentrating, mood swings, hair loss, itching, dry skin, sore throat, coughing, difficult breathing, dizziness, virus infection, rash, and dry mouth.

Commonly reported side effects (at least 1 in every 100 patients, but not greater than 1 in every 10 patients):

Increased sweating, chest pain, pain on the right side around your ribs, numbness, pain or tingling feeling, change in thyroid gland activity (which may make you feel tired or, less commonly, energetic), stomach upset, rapid heart rate, agitation, nervousness, difficult or irregular menstrual period

Less common are pain at the place of injection, flushing, low or high blood pressure, dry or teary eyes, redness of skin or skin disorders, psoriasis, hives, nail disorder, feeling unwell, feeling faint, poor coordination, confusion, increased or decreased sensitivity to touch, tense muscles, arthritis, bruising, loss of interest in activities including sex, sexual problem, unusual dreams, shaky hands, vertigo (spinning feeling), increased appetite, heartburn, intestinal gas (flatulence), constipation, hemorrhoids, red or bleeding gums, redness or sores in mouth, change in taste, changes in hearing or ringing in ears, thirst, changed behaviour or aggressive behaviour (sometimes directed against others), feeling sleepy, cold sores, fungal or bacterial infections, irritation of prostate gland, increased need to pass urine, ear or respiratory infections, sinusitis, stuffy or runny nose, abnormal hair texture, sensitivity to sunlight, migraine headache, eye pain or infection, blurred vision, puffy face, puffy hands or feet, enlarged liver, problem affecting ovary or vagina, pain in breast, difficulty in speaking and swollen glands.

The following rare and very rare events have been reported with PegIntron:

Rarely reported side effects (greater than 1 in every 10,000 patients, but less than 1 in every 1,000 patients):

Diabetes, abnormal heart rhythm, inflammation of pancreas, inflammation and degeneration of muscle tissue and peripheral nerves, kidney problems and seizures.

Very rarely reported side effects (less than 1 in every 10,000 patients):

Very rarely sarcoidosis (a disease characterised by persistent fever, weight loss, joint pain and swelling, skin lesions and swollen glands) has been reported. Loss of consciousness has occurred very rarely with alpha interferons, mostly in elderly patients treated at high doses. Cases of stroke (cerebrovascular events) have been reported. Check with your doctor immediately if you have any of these symptoms or any other symptoms that are troubling.

Very rarely, PegIntron alone or in combination with ribavirin may cause aplastic anaemia. Pure red cell aplasia, a condition where the body stopped or reduced the production of red blood cells, has been reported. This causes severe anaemia, symptoms of which would include unusual tiredness and a lack of energy.

When PegIntron is used alone, some of these effects are less likely to occur, and some have not occurred at all.

If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis, liver failure, and development of blood abnormalities (reduction in number of red blood cells which carry oxygen, certain white blood cells that fight infection, and blood clotting cells called platelets).

The following other side effects (not listed above) have occurred with the combination of PegIntron and ribavirin capsules (adults) in HCV/HIV co-infected patients receiving HAART: oral candidiasis (oral thrush), defective metabolism of fat, CD4 lymphocytes decreased, appetite decreased, back pain, hepatitis, limb pain, and various laboratory blood values abnormalities.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE PEGINTRON**

Keep out of the reach and sight of children.

Store in a refrigerator (2°C - 8°C).

Do not use PegIntron after the expiry date which is stated on the carton.

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

Do not use PegIntron if you notice discolouration of the powder.

The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. Discard any unused material.

## **6. FURTHER INFORMATION**

### **What PegIntron contains**

- The active substance is peginterferon alfa-2b, 120 micrograms/0.5 ml.
- The other ingredients are:  
Powder: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80;  
Solvent: water for injections 0.7 ml/ampoule.

### **What PegIntron looks like and contents of the pack**

The pharmaceutical form is: powder and solvent for solution for injection.

The white powder is contained in a 2 ml glass vial and the clear and colourless solvent is presented in a 2 ml glass ampoule.

PegIntron 120 micrograms is available in different pack sizes:

- 1 vial of powder for solution for injection and 1 ampoule of solvent for parenteral use;
- 1 vial of powder for solution for injection, 1 ampoule of solvent for parenteral use, 1 injection syringe, 2 injection needles and 1 cleansing swab;
- 4 vials of powder for solution for injection and 4 ampoules of solvent for parenteral use;
- 4 vials of powder for solution for injection, 4 ampoules of solvent for parenteral use, 4 injection syringes, 8 injection needles and 4 cleansing swabs;
- 6 vials of powder for solution for injection and 6 ampoules of solvent for parenteral use;
- 12 vials of powder for solution for injection, 12 ampoules of solvent for parenteral use, 12 injection syringes, 24 injection needles and 12 cleansing swabs.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

Marketing Authorisation Holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

Manufacturer: SP (Brinny) Company, Innishannon, County Cork, Ireland

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Rue de Stalle/Stallestraat 73  
B-1180 Bruxelles/Brussel/Brüssel  
Tél/Tel: + 32-(0)2 370 92 11

**Luxembourg/Luxemburg**

Rue de Stalle 73  
B-1180 Bruxelles/Brüssel  
Belgique/Belgien  
Tél/Tel: + 32-(0)2 370 92 11

**България**

ул. "Проф. Милко Бичев" № 7, ет.3  
BG-София 1504  
Тел.: +359 2 944 1073

**Magyarország**

Alkotás u. 53.  
H-1123 Budapest  
Tel.: +36 1 457-8500

**Česká republika**

Na Příkopě 25  
CZ-110 00 Praha 1  
Tel: +420 221771250

**Malta**

168 Christopher Street  
MT-VLT02 Valletta  
Tel: + 356-21 23 21 75

**Danmark**

Lautrupbjerg 2  
DK-2750 Ballerup  
Tlf: + 45-44 39 50 00

**Nederland**

Maarssenbroeksedijk 4  
NL-3542 DN Utrecht  
Tel: + 31-(0)800 778 78 78

**Deutschland**

Thomas-Dehler-Straße 27  
D-81737 München  
Tel: + 49-(0)89 627 31-0

**Norge**

Pb. 398  
N-1326 Lysaker  
Tlf: + 47 67 16 64 50

**Eesti**

Järvevana tee 9  
EE-11314 Tallinn  
Tel: + 372 654 96 86

**Österreich**

Badener Strasse 23  
A-2514 Traiskirchen  
Tel: + 43-(0)2252 502-0

**Ελλάδα**

Αγίου Δημητρίου 63  
GR-174 55 Αλιμος  
Τηλ.: + 30-210 98 97 300

**Polska**

Al. Jerozolimskie 195a  
PL-02-222 Warszawa  
Tel.: + 48-(0)22 478 41 50

**España**

Km. 36, Ctra. Nacional I  
E-28750 San Agustín de Guadalix – Madrid  
Tel: + 34-91 848 85 00

**Portugal**

Rua Agualva dos Açores 16  
P-2735-557 Agualva-Cacém  
Tel: +351-21 433 93 00

**France**

92 rue Baudin  
F-92300 Levallois-Perret  
Tél: + 33-(0)1 41 06 35 00

**România**

Șos. București-Ploiești, nr. 17-21, Băneasa  
Center, et. 7, sector 1  
RO-013682 București  
Tel: + 40 21 233 35 30

**Ireland**

Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW  
Tel: +44-(0)1 707 363 636

**Slovenija**

Dunajska 22  
SI-1000 Ljubljana  
Tel: + 386 01 3001070

**Ísland**

Hörgatún 2  
IS-210 Garðabær  
Sími: + 354 535 70 00

**Slovenská republika**

Strakova 5  
SK-811 01 Bratislava  
Tel: + 421 (2) 5920 2712

**Italia**

Via fratelli Cervi snc,  
Centro Direzionale Milano Due  
Palazzo Borromini  
I-20090 Segrate (Milano)  
Tel: + 39-02 21018.1

**Κύπρος**

Οδός Αγίου Νικολάου, 8  
CY-1055 Λευκωσία  
Τηλ: +357-22 757188

**Latvija**

Bauskas 58a -401  
Rīga, LV-1004  
Tel: + 371-7 21 38 25

**Lietuva**

Kęstučio g. 65/40  
LT-08124 Vilnius  
Tel: + 370 52 101868

**Suomi/Finland**

PL 3/PB 3  
FIN-02201 Espoo/Esbo  
Puh/Tel: + 358-(0)20-7570 300

**Sverige**

Box 27190  
S-102 52 Stockholm  
Tel: + 46-(0)8 522 21 500

**United Kingdom**

Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW - UK  
Tel: + 44-(0)1 707 363 636

**This leaflet was last approved on**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>

## HOW TO SELF-INJECT PEGINTRON?

The following instructions explain how to inject PegIntron yourself. Please read the instructions carefully and follow them step by step. Your doctor or his/her assistant will instruct you how to self-inject PegIntron. Do not attempt to inject yourself unless you are sure you understand the procedure and requirement of self-injection.

### Preparation

Collect necessary items before you begin:

- a vial of PegIntron powder for injection;
- an ampoule of solvent for PegIntron (water for injections);
- a 1 ml syringe;
- a long needle (for example  $0.8 \times 40$  mm [21 gauge 1.5 inch]) to be used to add water for injections to the PegIntron powder vial;
- a short needle (for example  $0.3 \times 13$  mm [30 gauge 0.5 inch]) for the subcutaneous injection;
- a cleansing swab.

Wash your hands carefully.

### Reconstituting PegIntron powder for injection

Before reconstitution, PegIntron may appear either as a white, tablet-shaped solid that is whole or in pieces, or as a white powder.

Remove the protective cap from the PegIntron vial. Clean the rubber top of the vial with a cleansing swab. You can save the swab to clean the skin area where you will inject the dose. Remove the syringe from the wrapping. Do not touch the tip of the syringe. Take the long needle and place it firmly on to the tip of the syringe. Remove the needle guard without touching the needle and keep the syringe with the needle in your hand. Tap the top of the ampoule of solvent gently to make sure that all the liquid is at the bottom of the ampoule. Break off the top of the ampoule of solvent. Insert the needle in the ampoule of solvent and withdraw the total amount of solvent.

When the total amount of solvent is combined with the full amount of PegIntron powder, the solution will be at the correct concentration to measure your dose (i.e., the labelled amount is contained in 0.5 ml).

A small volume is lost during preparation of PegIntron for injection and when the dose is measured and injected. Therefore, each vial contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection.

To prepare the PegIntron solution, insert the needle through the rubber top of the PegIntron vial and gently place the needle tip against the glass wall of the vial without touching the cleaned top of the vial with your hands.

Inject the solvent **SLOWLY**, aiming the stream of liquid at the glass wall of the vial. It is best not to aim the stream directly at the white solid or powder, or to inject the liquid quickly, as this causes a greater amount of bubbles. The solution may appear cloudy or bubbly for a few minutes. This is to be expected and is not cause for concern.

To dissolve the entire contents, swirl the PegIntron vial with a gentle rotary motion leaving the needle and attached syringe in the vial. **Do not shake**, but gently turn the vial upside down until any powder at the top of the vial is dissolved. The contents should now be completely dissolved. Stand the vial upright and let any bubbles present in the solution rise to the top of the solution. Once the solution has settled and all bubbles have risen to the top of the solution, you should have a clear solution with a small ring of tiny bubbles around the top. Now you can withdraw your dose from the vial. Use this solution immediately. If it cannot be used immediately, the solution may be refrigerated for up to 24 hours.

### Measuring the dose of PegIntron from the reconstituted powder for injection

Turn the vial and the syringe upside down in one hand. Be sure the tip of needle is in the PegIntron reconstituted solution. Your other hand will be free to move the plunger. Pull back on the plunger

slowly to draw just more than the dose prescribed by your doctor into the syringe. Hold the syringe with the needle in the vial pointing up, remove the syringe from the long needle leaving the needle in the vial and without touching the tip of the syringe. Take the short needle and place it firmly on to the tip of the syringe. Remove the needle guard from the syringe needle and check for air bubbles in the syringe. If you see any bubbles, pull the plunger slightly back; tap the syringe gently, with the needle pointing upwards, until the bubbles disappear. Push up the plunger slowly back to the correct dose. Replace the needle guard and place the syringe with the needle on a flat surface.

Be sure the solution is at room temperature up to 25°C. If the solution is cold, warm the syringe between your palms. Inspect visually the reconstituted solution prior to administration: do not use if discolouration or particulate matter is present. You are now ready to inject the dose.

#### Injecting the solution

Select the injection site. The best sites for injection are tissues with a layer of fat between skin and muscle: thigh, outer surface of the upper arm (you may need the assistance of another person to use this site), abdomen (except the navel or waistline). If you are exceptionally thin, use only the thigh or outer surface of the arm for injection.

#### **Change your injection site each time.**

Cleanse and disinfect the skin where the injection is to be made. Wait for the area to dry. Remove the needle guard. With one hand, pinch a fold of loose skin. With your other hand, hold the syringe as you would a pencil. Insert the needle into the pinched skin at an angle of approximately 45°. After the needle is in, remove the hand used to pinch the skin and use it to hold the syringe barrel. Pull back the plunger very slightly with one hand. If blood comes into the syringe, the needle has entered a blood vessel. Do not inject into this site; withdraw the needle and repeat the procedure. Inject the solution by pushing the plunger all the way down gently.

Pull the needle straight out of the skin. Press the injection site with a small bandage or sterile gauze if necessary for several seconds. Do not massage the injection site. If there is bleeding, cover with an adhesive bandage.

The vial, ampoule and injection materials intended for single use must be discarded. Dispose of the syringe and needles safely in a closed container.

## **PACKAGE LEAFLET: INFORMATION FOR THE USER**

### **PegIntron 150 micrograms powder and solvent for solution for injection**

peginterferon alfa-2b (conjugation of recombinant interferon alfa-2b with monomethoxy polyethylene glycol)

**Read all of this leaflet carefully before you start using this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What PegIntron is and what it is used for
2. Before you use PegIntron
3. How to use PegIntron
4. Possible side effects
5. Storing PegIntron
6. Further information

## **1. WHAT PEGINTRON IS AND WHAT IT IS USED FOR**

Interferons modify the response of the body's immune system to help fight infections and severe diseases. PegIntron, which contains an interferon, is used for the treatment of chronic hepatitis C, a viral infection of the liver, including in first time users co-infected with clinically stable HIV.

PegIntron is best used for this treatment in combination with ribavirin.

PegIntron is used alone in case of intolerance or contraindication to ribavirin.

## **2. BEFORE YOU USE PEGINTRON**

PegIntron is not recommended for use in patients under the age of 18 years.

**Do not use PegIntron**

- If you are allergic (hypersensitive) to peginterferon alfa-2b or any of the other ingredients of PegIntron.
- If you are allergic (hypersensitive) to any interferon.
- If you have had severe heart problems, or if you have heart disease that has not been well controlled during the past 6 months.
- If you have severe medical conditions that leave you very weak.
- If you have autoimmune hepatitis or any other problem with your immune system; if you are taking medicine that suppresses your immune system (your immune system protects you against infection and some diseases).
- If you have advanced, uncontrolled liver disease (other than hepatitis C).
- If you have thyroid disease that is not well controlled with medicines.
- If you have a condition that causes convulsions (seizures, or "fits").

**Take special care with PegIntron**

- If you develop symptoms of a severe allergic reaction (such as difficulty in breathing, wheezing, or hives) while on this medication, seek medical help immediately.
- If you ever had a heart attack or a heart problem.
- If you have ever been treated for depression or any other nervous or mental disorder.



- If you have kidney disease, your doctor may prescribe a lower than usual dose and monitor your kidney blood values regularly during treatment. If you have a kidney disease and you are using PegIntron in combination with medicinal products containing ribavirin, your doctor should monitor you more carefully for a decrease in red blood cell count.
- If you have had a problem with your liver (other than hepatitis C).
- If you develop symptoms associated with a cold or other respiratory infection, such as fever, cough, or any difficulty in breathing, tell your doctor.
- If you are diabetic, your doctor may ask you to have an eye examination.
- If you have had any serious illness affecting your breathing or your blood.
- If you have psoriasis, it may become worse while you are using PegIntron.
- If you are planning to become pregnant, discuss this with your doctor before starting to use PegIntron.
- If you are also being treated for HIV, please see **Using other medicines**.
- If you have had a severe nervous or mental disorder.
- If you ever had depression or develop symptoms associated with depression (e.g. feelings of sadness, dejection, etc) while on treatment with PegIntron (see section 4).
- If you have received an organ transplant, either kidney or liver, interferon treatment may increase the risk of rejection. Be sure to discuss this with your doctor.

Dental and gum disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and membranes of the mouth during long-term treatment with the combination of PegIntron with ribavirin. You should brush your teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If you have this reaction, be sure to rinse your mouth thoroughly afterwards.

#### **While being treated with PegIntron**

Your doctor may want you to drink extra fluids to help prevent low blood pressure.

#### **Using other medicines:**

Please inform your doctor or pharmacist if you are taking or have recently taken any other medicines, even those not prescribed.

#### **Pregnancy and breast-feeding**

Ask your doctor or pharmacist for advice before taking any medicine. In studies in pregnant animals, interferons have sometimes caused miscarriage. The effect on human pregnancy is not known. In combination therapy with ribavirin, ribavirin can be very damaging to an unborn baby, thus both female and male patients must take special precautions in their sexual activity if there is any chance for pregnancy to occur:

- if you are a **girl** or a **woman** of childbearing age, you must have a negative pregnancy test before treatment, each month during treatment, and for the 4 months after treatment is stopped. You and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 4 months after stopping treatment. This can be discussed with your doctor.
- if you are a **man** who is taking ribavirin, do not have sex with a pregnant woman unless you use a condom. This will lessen the chance for ribavirin to be left in the woman's body. If your female partner is not pregnant now but is of childbearing age, she must be tested for pregnancy each month during treatment and for the 7 months after treatment has stopped. This can be discussed with your doctor. If you are a male patient, you and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 7 months after stopping treatment. This can be discussed with your doctor.

It is not known whether this product is present in human milk. Therefore, do not breast-feed an infant if you are taking PegIntron. In combination therapy with ribavirin, take notice of the respective informing texts of ribavirin containing medicinal products.

#### **Driving and using machines**

Do not drive or operate any tools or machines if you feel tired, sleepy or confused while taking PegIntron.

### **Important information about some of the ingredients of PegIntron**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

This medicinal product contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### **Patients who also have HIV infection**

Lactic acidosis and worsening liver function are side effects associated with Highly Active Anti-Retroviral Therapy (HAART), an HIV treatment. If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis and of liver failure. Your doctor will monitor you for signs and symptoms of these conditions. (Please be sure to read the ribavirin Patient Leaflet also). Additionally, patients treated with PegIntron and ribavirin combination therapy and zidovudine could be at increased risk of developing anaemia (low number of red blood cells). If you are also being treated for Human Immunodeficiency Virus (HIV) infection (AIDS) with zidovudine or stavudine, it is not certain if ribavirin will change the way these medicines work. Therefore, your blood will be checked regularly to be sure that the HIV infection is not getting worse. If it gets worse, your doctor will decide whether or not your ribavirin treatment needs to be changed.

Co-administration of ribavirin and didanosine and/or stavudine is not recommended due to the risk of lactic acidosis (a build-up of lactic acid in the body) and pancreatitis.

## **3. HOW TO USE PEGINTRON**

Your doctor has prescribed PegIntron specifically for you and your current condition; do not share this medicine with anyone else.

Your doctor has determined your dose of PegIntron based on your weight. If necessary, the dose may be changed during treatment.

### **Combination treatment**

PegIntron, when given with ribavirin capsules, is usually given at a dose of 1.5 microgram/kg once a week.

Ribavirin capsules are taken every day, morning and evening. The number of ribavirin capsules you take depends on your weight.

- If you weigh less than 65 kg, take 2 capsules in the morning and 2 in the evening (total of 800 mg each day).
- If you weigh between 65 and 85 kg, take 2 capsules in the morning and 3 in the evening (total of 1,000 mg each day).
- If you weigh more than 85 kg, take 3 capsules in the morning and 3 in the evening (total 1,200 mg each day).

The combination treatment is continued for 3 to 6 months, and sometimes for one year depending on your physicians judgement. Take notice of the respective informing texts of ribavirin containing medicinal products.

For HCV/HIV co-infected patients, the duration of treatment is 48 weeks.

### **PegIntron alone:**

PegIntron, when given alone, is usually given at a dose of 0.5 or 1.0 microgram/kg once a week, for at least 6 months, and possibly for 1 year. If you have kidney disease, your dose may be lower, depending upon your kidney function.

PegIntron alone was not studied in HCV/HIV co-infected patients.

**All patients:**

If you are injecting PegIntron yourself, please be sure that the dose that has been prescribed for you is clearly provided on the package of medicine you receive.

If you have the impression that the effect of PegIntron is too strong or too weak, talk to your doctor or pharmacist.

PegIntron is intended for subcutaneous use. This means that it is injected through a short injection needle into the fatty tissue just under your skin. If you are injecting this medicine yourself, you will be instructed how to prepare and give the injection. Detailed instructions for subcutaneous administration are provided with this leaflet (see How to self-inject PegIntron at the end of the package leaflet).

Prepare the dose just before you intend to inject it and use it immediately. Look carefully at the reconstituted solution prior to administration. Do not use if there is discolouration of the reconstituted solution or if particulate matter is present. Discard any solution that is left in the vial after you give yourself the injection.

Inject PegIntron once each week on the same day. Injecting it at the same time of day each week will help you not to forget to take it.

Always use PegIntron exactly as your doctor told you. Do not exceed the recommended dosage, and take it for as long as prescribed.

**If you use more PegIntron than you should**

Tell your doctor or healthcare professional as soon as possible.

**If you forget to take PegIntron**

Take the dose as soon as you remember, then continue your treatment as usual.

Do not take a double dose to make up for a forgotten dose. Contact your doctor or pharmacist if needed.

**4. POSSIBLE SIDE EFFECTS**

Like all medicines, PegIntron can cause side effects, although not everybody gets them. Although not all of these side effects may occur, they may need medical attention if they do.

Some people get depressed when taking PegIntron alone or in combination treatment with ribavirin, and in some cases people had suicidal thoughts or aggressive behaviour (sometimes directed against others). Some patients have actually committed suicide. Be sure to seek emergency care if you notice that you are becoming depressed or have suicidal thoughts or change in your behaviour. You may want to consider asking a family member or close friend to help you stay alert to signs of depression or changes in your behaviour.

Check with your doctor immediately if any of the following side effects occur: chest pain; changes in the way your heart beats; breathing problems (including shortness of breath), confusion; feeling depressed, wanting to harm yourself, hallucinations, numbness or tingling feeling; dizziness, convulsion ("fit"); trouble sleeping, thinking or concentrating; difficulty remaining alert, severe stomach pain or cramps; blood or clots in stool (or black, tarry stool); fever or chills beginning after a few weeks of treatment, pain in your lower back or side; difficulty or inability to pass urine, painful or inflamed muscles (sometimes severe); problems with your eyes or your eyesight or hearing; severe or painful reddening of your skin or mucous membrane, severe bleeding from your nose. Your doctor will test your blood to ensure that your white blood cell (cells that fight infection) and red blood cell (cells that carry oxygen) counts, platelets (blood clotting cells) and other laboratory values are at acceptable levels.

Other side effects that may occur with the combination of PegIntron and ribavirin capsules include:

Very commonly reported side effects (greater than 1 in every 10 patients):

Irritation or redness (and rarely, skin damage) at the site of injection, headache, tired feeling, shaking chills, fever, flu-like symptoms, weakness, loss of weight, nausea, loss of appetite, diarrhoea or loose stools, stomach pain, vomiting, muscle aches, pain in joints and muscles, feeling depressed, irritability, trouble falling asleep or staying asleep, feeling anxious or nervous, difficulty concentrating, mood swings, hair loss, itching, dry skin, sore throat, coughing, difficult breathing, dizziness, virus infection, rash, and dry mouth.

Commonly reported side effects (at least 1 in every 100 patients, but not greater than 1 in every 10 patients):

Increased sweating, chest pain, pain on the right side around your ribs, numbness, pain or tingling feeling, change in thyroid gland activity (which may make you feel tired or, less commonly, energetic), stomach upset, rapid heart rate, agitation, nervousness, difficult or irregular menstrual period

Less common are pain at the place of injection, flushing, low or high blood pressure, dry or teary eyes, redness of skin or skin disorders, psoriasis, hives, nail disorder, feeling unwell, feeling faint, poor coordination, confusion, increased or decreased sensitivity to touch, tense muscles, arthritis, bruising, loss of interest in activities including sex, sexual problem, unusual dreams, shaky hands, vertigo (spinning feeling), increased appetite, heartburn, intestinal gas (flatulence), constipation, hemorrhoids, red or bleeding gums, redness or sores in mouth, change in taste, changes in hearing or ringing in ears, thirst, changed behaviour or aggressive behaviour (sometimes directed against others), feeling sleepy, cold sores, fungal or bacterial infections, irritation of prostate gland, increased need to pass urine, ear or respiratory infections, sinusitis, stuffy or runny nose, abnormal hair texture, sensitivity to sunlight, migraine headache, eye pain or infection, blurred vision, puffy face, puffy hands or feet, enlarged liver, problem affecting ovary or vagina, pain in breast, difficulty in speaking and swollen glands.

The following rare and very rare events have been reported with PegIntron:

Rarely reported side effects (greater than 1 in every 10,000 patients, but less than 1 in every 1,000 patients):

Diabetes, abnormal heart rhythm, inflammation of pancreas, inflammation and degeneration of muscle tissue and peripheral nerves, kidney problems and seizures.

Very rarely reported side effects (less than 1 in every 10,000 patients):

Very rarely sarcoidosis (a disease characterised by persistent fever, weight loss, joint pain and swelling, skin lesions and swollen glands) has been reported. Loss of consciousness has occurred very rarely with alpha interferons, mostly in elderly patients treated at high doses. Cases of stroke (cerebrovascular events) have been reported. Check with your doctor immediately if you have any of these symptoms or any other symptoms that are troubling.

Very rarely, PegIntron alone or in combination with ribavirin may cause aplastic anaemia. Pure red cell aplasia, a condition where the body stopped or reduced the production of red blood cells, has been reported. This causes severe anaemia, symptoms of which would include unusual tiredness and a lack of energy.

When PegIntron is used alone, some of these effects are less likely to occur, and some have not occurred at all.

If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis, liver failure, and development of blood abnormalities (reduction in number of red blood cells which carry oxygen, certain white blood cells that fight infection, and blood clotting cells called platelets).

The following other side effects (not listed above) have occurred with the combination of PegIntron and ribavirin capsules (adults) in HCV/HIV co-infected patients receiving HAART: oral candidiasis (oral thrush), defective metabolism of fat, CD4 lymphocytes decreased, appetite decreased,

back pain, hepatitis, limb pain, and various laboratory blood values abnormalities.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE PEGINTRON**

Keep out of the reach and sight of children.

Store in a refrigerator (2°C - 8°C).

Do not use PegIntron after the expiry date which is stated on the carton.

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

Do not use PegIntron if you notice discolouration of the powder.

The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. Discard any unused material.

## **6. FURTHER INFORMATION**

### **What PegIntron contains**

- The active substance is peginterferon alfa-2b, 150 micrograms/0.5 ml.
- The other ingredients are:  
Powder: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80;  
Solvent: water for injections 0.7 ml/ampoule.

### **What PegIntron looks like and contents of the pack**

The pharmaceutical form is: powder and solvent for solution for injection.

The white powder is contained in a 2 ml glass vial and the clear and colourless solvent is presented in a 2 ml glass ampoule.

PegIntron 150 micrograms is available in different pack sizes:

- 1 vial of powder for solution for injection and 1 ampoule of solvent for parenteral use;
- 1 vial of powder for solution for injection, 1 ampoule of solvent for parenteral use, 1 injection syringe, 2 injection needles and 1 cleansing swab;
- 4 vials of powder for solution for injection and 4 ampoules of solvent for parenteral use;
- 4 vials of powder for solution for injection, 4 ampoules of solvent for parenteral use, 4 injection syringes, 8 injection needles and 4 cleansing swabs;
- 6 vials of powder for solution for injection and 6 ampoules of solvent for parenteral use;
- 12 vials of powder for solution for injection, 12 ampoules of solvent for parenteral use, 12 injection syringes, 24 injection needles and 12 cleansing swabs.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

Marketing Authorisation Holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

Manufacturer: SP (Brinny) Company, Innishannon, County Cork, Ireland

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**  
Rue de Stalle/Stallestraat 73  
B-1180 Bruxelles/Brussel/Brüssel  
Tél/Tel: + 32-(0)2 370 92 11

**България**  
ул. "Проф. Милко Бичев" № 7, ет.3  
BG-София 1504  
Тел.: +359 2 944 1073

**Česká republika**  
Na Příkopě 25  
CZ-110 00 Praha 1  
Tel: +420 221771250

**Danmark**  
Lautrupbjerg 2  
DK-2750 Ballerup  
Tlf: + 45-44 39 50 00

**Deutschland**  
Thomas-Dehler-Straße 27  
D-81737 München  
Tel: + 49-(0)89 627 31-0

**Eesti**  
Järvevana tee 9  
EE-11314 Tallinn  
Tel: + 372 654 96 86

**Ελλάδα**  
Αγίου Δημητρίου 63  
GR-174 55 Αλιμος  
Τηλ.: + 30-210 98 97 300

**España**  
Km. 36, Ctra. Nacional I  
E-28750 San Agustín de Guadalix – Madrid  
Tel: + 34-91 848 85 00

**France**  
92 rue Baudin  
F-92300 Levallois-Perret  
Tél: + 33-(0)1 41 06 35 00

**Ireland**  
Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW  
Tel: +44-(0)1 707 363 636

**Luxembourg/Luxemburg**  
Rue de Stalle 73  
B-1180 Bruxelles/Brüssel  
Belgique/Belgien  
Tél/Tel: + 32-(0)2 370 92 11

**Magyarország**  
Alkotás u. 53.  
H-1123 Budapest  
Tel.: +36 1 457-8500

**Malta**  
168 Christopher Street  
MT-VLT02 Valletta  
Tel: + 356-21 23 21 75

**Nederland**  
Maarssenbroeksedijk 4  
NL-3542 DN Utrecht  
Tel: + 31-(0)800 778 78 78

**Norge**  
Pb. 398  
N-1326 Lysaker  
Tlf: + 47 67 16 64 50

**Österreich**  
Badener Strasse 23  
A-2514 Traiskirchen  
Tel: + 43-(0)2252 502-0

**Polska**  
Al. Jerozolimskie 195a  
PL-02-222 Warszawa  
Tel.: + 48-(0)22 478 41 50

**Portugal**  
Rua Agualva dos Açores 16  
P-2735-557 Agualva-Cacém  
Tel: +351-21 433 93 00

**România**  
Șos. București-Ploiești, nr. 17-21, Băneasa  
Center, et. 7, sector 1  
RO-013682 București  
Tel: + 40 21 233 35 30

**Slovenija**  
Dunajska 22  
SI-1000 Ljubljana  
Tel: + 386 01 3001070

**Ísland**

Hörgatún 2  
IS-210 Garðabær  
Sími: + 354 535 70 00

**Italia**

Via fratelli Cervi snc,  
Centro Direzionale Milano Due  
Palazzo Borromini  
I-20090 Segrate (Milano)  
Tel: + 39-02 21018.1

**Κύπρος**

Οδός Αγίου Νικολάου, 8  
CY-1055 Λευκωσία  
Τηλ: +357-22 757188

**Latvija**

Bauskas 58a -401  
Rīga, LV-1004  
Tel: + 371-7 21 38 25

**Lietuva**

Kęstučio g. 65/40  
LT-08124 Vilnius  
Tel: + 370 52 101868

**Slovenská republika**

Strakova 5  
SK-811 01 Bratislava  
Tel: + 421 (2) 5920 2712

**Suomi/Finland**

PL 3/PB 3  
FIN-02201 Espoo/Esbo  
Puh/Tel: + 358-(0)20-7570 300

**Sverige**

Box 27190  
S-102 52 Stockholm  
Tel: + 46-(0)8 522 21 500

**United Kingdom**

Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW - UK  
Tel: + 44-(0)1 707 363 636

**This leaflet was last approved on**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>

## HOW TO SELF-INJECT PEGINTRON?

The following instructions explain how to inject PegIntron yourself. Please read the instructions carefully and follow them step by step. Your doctor or his/her assistant will instruct you how to self-inject PegIntron. Do not attempt to inject yourself unless you are sure you understand the procedure and requirement of self-injection.

### Preparation

Collect necessary items before you begin:

- a vial of PegIntron powder for injection;
- an ampoule of solvent for PegIntron (water for injections);
- a 1 ml syringe;
- a long needle (for example  $0.8 \times 40$  mm [21 gauge 1.5 inch]) to be used to add water for injections to the PegIntron powder vial;
- a short needle (for example  $0.3 \times 13$  mm [30 gauge 0.5 inch]) for the subcutaneous injection;
- a cleansing swab.

Wash your hands carefully.

### Reconstituting PegIntron powder for injection

Before reconstitution, PegIntron may appear either as a white, tablet-shaped solid that is whole or in pieces, or as a white powder.

Remove the protective cap from the PegIntron vial. Clean the rubber top of the vial with a cleansing swab. You can save the swab to clean the skin area where you will inject the dose. Remove the syringe from the wrapping. Do not touch the tip of the syringe. Take the long needle and place it firmly on to the tip of the syringe. Remove the needle guard without touching the needle and keep the syringe with the needle in your hand. Tap the top of the ampoule of solvent gently to make sure that all the liquid is at the bottom of the ampoule. Break off the top of the ampoule of solvent. Insert the needle in the ampoule of solvent and withdraw the total amount of solvent.

When the total amount of solvent is combined with the full amount of PegIntron powder, the solution will be at the correct concentration to measure your dose (i.e., the labelled amount is contained in 0.5 ml).

A small volume is lost during preparation of PegIntron for injection and when the dose is measured and injected. Therefore, each vial contains an excess amount of solvent and PegIntron powder to ensure delivery of the labelled dose in 0.5 ml of PegIntron, solution for injection.

To prepare the PegIntron solution, insert the needle through the rubber top of the PegIntron vial and gently place the needle tip against the glass wall of the vial without touching the cleaned top of the vial with your hands.

Inject the solvent **SLOWLY**, aiming the stream of liquid at the glass wall of the vial. It is best not to aim the stream directly at the white solid or powder, or to inject the liquid quickly, as this causes a greater amount of bubbles. The solution may appear cloudy or bubbly for a few minutes. This is to be expected and is not cause for concern.

To dissolve the entire contents, swirl the PegIntron vial with a gentle rotary motion leaving the needle and attached syringe in the vial. **Do not shake**, but gently turn the vial upside down until any powder at the top of the vial is dissolved. The contents should now be completely dissolved. Stand the vial upright and let any bubbles present in the solution rise to the top of the solution. Once the solution has settled and all bubbles have risen to the top of the solution, you should have a clear solution with a small ring of tiny bubbles around the top. Now you can withdraw your dose from the vial. Use this solution immediately. If it cannot be used immediately, the solution may be refrigerated for up to 24 hours.

### Measuring the dose of PegIntron from the reconstituted powder for injection

Turn the vial and the syringe upside down in one hand. Be sure the tip of needle is in the PegIntron reconstituted solution. Your other hand will be free to move the plunger. Pull back on the plunger



slowly to draw just more than the dose prescribed by your doctor into the syringe. Hold the syringe with the needle in the vial pointing up, remove the syringe from the long needle leaving the needle in the vial and without touching the tip of the syringe. Take the short needle and place it firmly on to the tip of the syringe. Remove the needle guard from the syringe needle and check for air bubbles in the syringe. If you see any bubbles, pull the plunger slightly back; tap the syringe gently, with the needle pointing upwards, until the bubbles disappear. Push up the plunger slowly back to the correct dose. Replace the needle guard and place the syringe with the needle on a flat surface.

Be sure the solution is at room temperature up to 25°C. If the solution is cold, warm the syringe between your palms. Inspect visually the reconstituted solution prior to administration: do not use if discolouration or particulate matter is present. You are now ready to inject the dose.

#### Injecting the solution

Select the injection site. The best sites for injection are tissues with a layer of fat between skin and muscle: thigh, outer surface of the upper arm (you may need the assistance of another person to use this site), abdomen (except the navel or waistline). If you are exceptionally thin, use only the thigh or outer surface of the arm for injection.

#### **Change your injection site each time.**

Cleanse and disinfect the skin where the injection is to be made. Wait for the area to dry. Remove the needle guard. With one hand, pinch a fold of loose skin. With your other hand, hold the syringe as you would a pencil. Insert the needle into the pinched skin at an angle of approximately 45°. After the needle is in, remove the hand used to pinch the skin and use it to hold the syringe barrel. Pull back the plunger very slightly with one hand. If blood comes into the syringe, the needle has entered a blood vessel. Do not inject into this site; withdraw the needle and repeat the procedure. Inject the solution by pushing the plunger all the way down gently.

Pull the needle straight out of the skin. Press the injection site with a small bandage or sterile gauze if necessary for several seconds. Do not massage the injection site. If there is bleeding, cover with an adhesive bandage.

The vial, ampoule and injection materials intended for single use must be discarded. Dispose of the syringe and needles safely in a closed container.

## PACKAGE LEAFLET: INFORMATION FOR THE USER

**PegIntron 50 micrograms powder and solvent for solution for injection in pre-filled pen**  
Peginterferon alfa-2b (conjugation of recombinant interferon alfa-2b with monomethoxy polyethylene glycol)

**Read all of this leaflet carefully before you start using this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What PegIntron is and what it is used for
2. Before you use PegIntron
3. How to use PegIntron
4. Possible side effects
5. Storing PegIntron
6. Further information

### **1. WHAT PEGINTRON IS AND WHAT IT IS USED FOR**

Interferons modify the response of the body's immune system to help fight infections and severe diseases. PegIntron, which contains an interferon, is used for the treatment of chronic hepatitis C, a viral infection of the liver, including in first time users co-infected with clinically stable HIV.

PegIntron is best used for this treatment in combination with ribavirin.

PegIntron is used alone in case of intolerance or contraindication to ribavirin.

### **2. BEFORE YOU USE PEGINTRON**

PegIntron is not recommended for use in patients under the age of 18 years.

**Do not use PegIntron**

- If you are allergic (hypersensitive) to peginterferon alfa-2b or any of the other ingredients of PegIntron.
- If you are allergic (hypersensitive) to any interferon.
- If you have had severe heart problems, or if you have heart disease that has not been well controlled during the past 6 months.
- If you have severe medical conditions that leave you very weak.
- If you have autoimmune hepatitis or any other problem with your immune system; if you are taking medicine that suppresses your immune system (your immune system protects you against infection and some diseases).
- If you have advanced, uncontrolled liver disease (other than hepatitis C).
- If you have thyroid disease that is not well controlled with medicines.
- If you have a condition that causes convulsions (seizures, or "fits").

**Take special care with PegIntron**

- If you develop symptoms of a severe allergic reaction (such as difficulty in breathing, wheezing, or hives) while on this medication, seek medical help immediately.
- If you ever had a heart attack or a heart problem.
- If you have ever been treated for depression or any other nervous or mental disorder.

- If you have kidney disease, your doctor may prescribe a lower than usual dose and monitor your kidney blood values regularly during treatment. If you have a kidney disease and you are using PegIntron in combination with medicinal products containing ribavirin, your doctor should monitor you more carefully for a decrease in red blood cell count.
- If you have had a problem with your liver (other than hepatitis C).
- If you develop symptoms associated with a cold or other respiratory infection, such as fever, cough, or any difficulty in breathing, tell your doctor.
- If you are diabetic, your doctor may ask you to have an eye examination.
- If you have had any serious illness affecting your breathing or your blood.
- If you have psoriasis, it may become worse while you are using PegIntron.
- If you are planning to become pregnant, discuss this with your doctor before starting to use PegIntron.
- If you are also being treated for HIV, please see **Using other medicines**.
- If you have had a severe nervous or mental disorder.
- If you ever had depression or develop symptoms associated with depression (e.g. feelings of sadness, dejection, etc) while on treatment with PegIntron (see section 4).
- If you have received an organ transplant, either kidney or liver, interferon treatment may increase the risk of rejection. Be sure to discuss this with your doctor.

Dental and gum disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and membranes of the mouth during long-term treatment with the combination of PegIntron with ribavirin. You should brush your teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If you have this reaction, be sure to rinse your mouth thoroughly afterwards.

#### **While being treated with PegIntron**

Your doctor may want you to drink extra fluids to help prevent low blood pressure.

#### **Using other medicines**

Please inform your doctor or pharmacist if you are taking or have recently taken any other medicines, even those not prescribed.

#### **Pregnancy and breast-feeding**

Ask your doctor or pharmacist for advice before taking any medicine. In studies in pregnant animals, interferons have sometimes caused miscarriage. The effect on human pregnancy is not known. In combination therapy with ribavirin, ribavirin can be very damaging to an unborn baby, thus both female and male patients must take special precautions in their sexual activity if there is any chance for pregnancy to occur:

- if you are a **girl** or a **woman** of childbearing age, you must have a negative pregnancy test before treatment, each month during treatment, and for the 4 months after treatment is stopped. You and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 4 months after stopping treatment. This can be discussed with your doctor.
- if you are a **man** who is taking ribavirin, do not have sex with a pregnant woman unless you use a condom. This will lessen the chance for ribavirin to be left in the woman's body. If your female partner is not pregnant now but is of childbearing age, she must be tested for pregnancy each month during treatment and for the 7 months after treatment has stopped. This can be discussed with your doctor. If you are a male patient, you and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 7 months after stopping treatment. This can be discussed with your doctor.

It is not known whether this product is present in human milk. Therefore, do not breast-feed an infant if you are taking PegIntron. In combination therapy with ribavirin, take notice of the respective informing texts of ribavirin containing medicinal products.

#### **Driving and using machines**

Do not drive or operate any tools or machines if you feel tired, sleepy or confused while taking PegIntron.

### **Important information about some of the ingredients of PegIntron**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

This medicinal product contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### **Patients who also have HIV infection**

Lactic acidosis and worsening liver function are side effects associated with Highly Active Anti-Retroviral Therapy (HAART), an HIV treatment. If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis and of liver failure. Your doctor will monitor you for signs and symptoms of these conditions. (Please be sure to read the ribavirin Patient Leaflet also). Additionally, patients treated with PegIntron and ribavirin combination therapy and zidovudine could be at increased risk of developing anaemia (low number of red blood cells). If you are also being treated for Human Immunodeficiency Virus (HIV) infection (AIDS) with zidovudine or stavudine, it is not certain if ribavirin will change the way these medicines work. Therefore, your blood will be checked regularly to be sure that the HIV infection is not getting worse. If it gets worse, your doctor will decide whether or not your ribavirin treatment needs to be changed.

Co-administration of ribavirin and didanosine and/or stavudine is not recommended due to the risk of lactic acidosis (a build-up of lactic acid in the body) and pancreatitis.

## **3. HOW TO USE PEGINTRON**

Your doctor has prescribed PegIntron specifically for you and your current condition; do not share this medicine with anyone else.

Your doctor has determined your dose of PegIntron based on your weight. If necessary, the dose may be changed during treatment.

### **Combination treatment**

PegIntron, when given with ribavirin capsules, is usually given at a dose of 1.5 microgram/kg once a week.

Ribavirin capsules are taken every day, morning and evening. The number of ribavirin capsules you take depends on your weight.

- If you weigh less than 65 kg, take 2 capsules in the morning and 2 in the evening (total of 800 mg each day).
- If you weigh between 65 and 85 kg, take 2 capsules in the morning and 3 in the evening (total of 1,000 mg each day).
- If you weigh more than 85 kg, take 3 capsules in the morning and 3 in the evening (total 1,200 mg each day).

The combination treatment is continued for 3 to 6 months, and sometimes for one year depending on your physicians judgement. Take notice of the respective informing texts of ribavirin containing medicinal products.

For HCV/HIV co-infected patients, the duration of treatment is 48 weeks.

### **PegIntron alone:**

PegIntron, when given alone, is usually given at a dose of 0.5 or 1.0 microgram/kg once a week, for at least 6 months, and possibly for 1 year. If you have kidney disease, your dose may be lower, depending upon your kidney function.

PegIntron alone was not studied in HCV/HIV co-infected patients.

**All patients:**

If you are injecting PegIntron yourself, please be sure that the dose that has been prescribed for you is clearly provided on the package of medicine you receive.

If you have the impression that the effect of PegIntron is too strong or too weak, talk to your doctor or pharmacist.

PegIntron is intended for subcutaneous use. This means that it is injected through a short injection needle into the fatty tissue just under your skin. If you are injecting this medicine yourself, you will be instructed how to prepare and give the injection. Detailed instructions for subcutaneous administration are provided with this leaflet (see How to use the PegIntron pre-filled pen).

Prepare the dose just before you intend to inject it and use it immediately. Look carefully at the reconstituted solution prior to administration. Do not use if there is discolouration of the reconstituted solution or if particulate matter is present. PegIntron is intended for single use only. Thus, after you give yourself the injection, discard the PegIntron pre-filled pen with any solution that is left in it.

Inject PegIntron once each week on the same day. Injecting it at the same time of day each week will help you not to forget to take it.

Always use PegIntron exactly as your doctor told you. Do not exceed the recommended dosage, and take it for as long as prescribed.

**If you use more PegIntron than you should**

Tell your doctor or healthcare professional as soon as possible.

**If you forget to take PegIntron**

Take the dose as soon as you remember, then continue your treatment as usual.

Do not take a double dose to make up for a forgotten dose. Contact your doctor or pharmacist if needed.

**4. POSSIBLE SIDE EFFECTS**

Like all medicines, PegIntron can cause side effects, although not everybody gets them. Although not all of these side effects may occur, they may need medical attention if they do.

Some people get depressed when taking PegIntron alone or in combination treatment with ribavirin, and in some cases people had suicidal thoughts or aggressive behaviour (sometimes directed against others). Some patients have actually committed suicide. Be sure to seek emergency care if you notice that you are becoming depressed or have suicidal thoughts or change in your behaviour. You may want to consider asking a family member or close friend to help you stay alert to signs of depression or changes in your behaviour.

Check with your doctor immediately if any of the following side effects occur: chest pain; changes in the way your heart beats; breathing problems (including shortness of breath), confusion; feeling depressed, wanting to harm yourself, hallucinations, numbness or tingling feeling; dizziness, convulsion ("fit"); trouble sleeping, thinking or concentrating; difficulty remaining alert, severe stomach pain or cramps; blood or clots in stool (or black, tarry stool); fever or chills beginning after a few weeks of treatment, pain in your lower back or side; difficulty or inability to pass urine, painful or inflamed muscles (sometimes severe); problems with your eyes or your eyesight or hearing; severe or painful reddening of your skin or mucous membrane, severe bleeding from your nose. Your doctor will test your blood to ensure that your white blood cell (cells that fight infection) and red blood cell (cells that carry oxygen) counts, platelets (blood clotting cells) and other laboratory values are at acceptable levels.

Other side effects that may occur with the combination of PegIntron and ribavirin capsules include:

Very commonly reported side effects (greater than 1 in every 10 patients): Irritation or redness (and rarely, skin damage) at the site of injection, headache, tired feeling, shaking chills, fever, flu-like symptoms, weakness, loss of weight, nausea, loss of appetite, diarrhoea or loose stools, stomach pain, vomiting, muscle aches, pain in joints and muscles, feeling depressed, irritability, trouble falling asleep or staying asleep, feeling anxious or nervous, difficulty concentrating, mood swings, hair loss, itching, dry skin, sore throat, coughing, difficult breathing, dizziness, virus infection, rash, and dry mouth.

Commonly reported side effects (at least 1 in every 100 patients, but not greater than 1 in every 10 patients):

Increased sweating, chest pain, pain on the right side around your ribs, numbness, pain or tingling feeling, change in thyroid gland activity (which may make you feel tired or, less commonly, energetic), stomach upset, rapid heart rate, agitation, nervousness, difficult or irregular menstrual period

Less common are pain at the place of injection, flushing, low or high blood pressure, dry or teary eyes, redness of skin or skin disorder, psoriasis, hives, nail disorder, feeling unwell, feeling faint, poor coordination, confusion, increased or decreased sensitivity to touch, tense muscles, arthritis, bruising, loss of interest in activities including sex, sexual problem, unusual dreams, shaky hands, vertigo (spinning feeling), increased appetite, heartburn, intestinal gas (flatulence), constipation, hemorrhoids, red or bleeding gums, redness or sores in mouth, change in taste, changes in hearing or ringing in ears, thirst, changed behaviour or aggressive behaviour (sometimes directed against others), feeling sleepy, cold sores, fungal or bacterial infections, irritation of prostate gland, increased need to pass urine, ear or respiratory infections, sinusitis, stuffy or runny nose, abnormal hair texture, sensitivity to sunlight, migraine headache, eye pain or infection, blurred vision, puffy face, puffy hands or feet, enlarged liver, problem affecting ovary or vagina, pain in breast, difficulty in speaking and swollen glands.

The following rare and very rare events have been reported with PegIntron:

Rarely reported side effects (greater than 1 in every 10,000 patients, but less than 1 in every 1,000 patients):

Diabetes, abnormal heart rhythm, inflammation of pancreas, inflammation and degeneration of muscle tissue and peripheral nerves, kidney problems and seizures.

Very rarely reported side effects (less than 1 in every 10,000 patients):

Very rarely sarcoidosis (a disease characterised by persistent fever, weight loss, joint pain and swelling, skin lesions and swollen glands) has been reported. Loss of consciousness has occurred very rarely with alpha interferons, mostly in elderly patients treated at high doses. Cases of stroke (cerebrovascular events) have been reported. Check with your doctor immediately if you have any of these symptoms or any other symptoms that are troubling.

Very rarely, PegIntron alone or in combination with ribavirin may cause aplastic anaemia. Pure red cell aplasia, a condition where the body stopped or reduced the production of red blood cells, has been reported. This causes severe anaemia, symptoms of which would include unusual tiredness and a lack of energy.

When PegIntron is used alone, some of these effects are less likely to occur, and some have not occurred at all.

If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis, liver failure, and development of blood abnormalities (reduction in number of red blood cells which carry oxygen, certain white blood cells that fight infection, and blood clotting cells called platelets).

The following other side effects (not listed above) have occurred with the combination of PegIntron and ribavirin capsules (adults) in HCV/HIV co-infected patients receiving HAART:

oral candidiasis (oral thrush), defective metabolism of fat, , CD4 lymphocytes decreased, appetite decreased,

back pain, hepatitis, limb pain, and various laboratory blood values abnormalities.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE PEGINTRON**

Keep out of the reach and sight of children.

Store in a refrigerator (2°C - 8°C). Do not freeze.

Do not use PegIntron after the expiry date which is stated on the carton.

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

Do not use PegIntron if you notice discolouration of the powder.

The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. After administering the dose, discard the PegIntron pre-filled pen and any unused solution contained in it.

## **6. FURTHER INFORMATION**

### **What PegIntron contains**

- The active substance is peginterferon alfa-2b, 50 micrograms/0.5 ml.
- The other ingredients are:  
Powder: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80;  
Solvent: water for injections

### **What PegIntron looks like and contents of the pack**

The pharmaceutical form is: powder and solvent for solution for injection in a pre-filled pen. The white powder and the clear and colourless solvent are both contained in a two-chamber glass cartridge assembled into a single use pre-filled pen.

PegIntron 50 micrograms is available in different pack sizes:

- 1 pen containing powder and solvent for solution for injection, 1 injection needle and 2 cleansing swabs;
- 4 pens containing powder and solvent for solution for injection, 4 injection needles and 8 cleansing swabs;
- 6 pens containing powder and solvent for solution for injection, 6 injection needles and 12 cleansing swabs;
- 12 pens containing powder and solvent for solution for injection, 12 injection needles and 24 cleansing swabs.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

Marketing Authorisation Holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

Manufacturer: SP (Brinny) Company, Innishannon, County Cork, Ireland

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**  
Rue de Stalle/Stallestraat 73  
B-1180 Bruxelles/Brussel/Brüssel  
Tél/Tel: + 32-(0)2 370 92 11

**България**  
ул. "Проф. Милко Бичев" № 7, ет.3  
BG-София 1504  
Тел.: +359 2 944 1073

**Česká republika**  
Na Příkopě 25  
CZ-110 00 Praha 1  
Tel: +420 221771250

**Danmark**  
Lautrupbjerg 2  
DK-2750 Ballerup  
Tlf: + 45-44 39 50 00

**Deutschland**  
Thomas-Dehler-Straße 27  
D-81737 München  
Tel: + 49-(0)89 627 31-0

**Eesti**  
Järvevana tee 9  
EE-11314 Tallinn  
Tel: + 372 654 96 86

**Ελλάδα**  
Αγίου Δημητρίου 63  
GR-174 55 Αλιμος  
Τηλ.: + 30-210 98 97 300

**España**  
Km. 36, Ctra. Nacional I  
E-28750 San Agustín de Guadalix – Madrid  
Tel: + 34-91 848 85 00

**France**  
92 rue Baudin  
F-92300 Levallois-Perret  
Tél: + 33-(0)1 41 06 35 00

**Ireland**  
Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW  
Tel: +44-(0)1 707 363 636

**Luxembourg/Luxemburg**  
Rue de Stalle 73  
B-1180 Bruxelles/Brüssel  
Belgique/Belgien  
Tél/Tel: + 32-(0)2 370 92 11

**Magyarország**  
Alkotás u. 53.  
H-1123 Budapest  
Tel.: +36 1 457-8500

**Malta**  
168 Christopher Street  
MT-VLT02 Valletta  
Tel: + 356-21 23 21 75

**Nederland**  
Maarssenbroeksedijk 4  
NL-3542 DN Utrecht  
Tel: + 31-(0)800 778 78 78

**Norge**  
Pb. 398  
N-1326 Lysaker  
Tlf: + 47 67 16 64 50

**Österreich**  
Badener Strasse 23  
A-2514 Traiskirchen  
Tel: + 43-(0)2252 502-0

**Polska**  
Al. Jerozolimskie 195a  
PL-02-222 Warszawa  
Tel.: + 48-(0)22 478 41 50

**Portugal**  
Rua Agualva dos Açores 16  
P-2735-557 Agualva-Cacém  
Tel: +351-21 433 93 00

**România**  
Șos. București-Ploiești, nr. 17-21, Băneasa  
Center, et. 7, sector 1  
RO-013682 București  
Tel: + 40 21 233 35 30

**Slovenija**  
Dunajska 22  
SI-1000 Ljubljana  
Tel: + 386 01 3001070



**Ísland**

Hörgatún 2  
IS-210 Garðabær  
Sími: + 354 535 70 00

**Italia**

Via fratelli Cervi snc,  
Centro Direzionale Milano Due  
Palazzo Borromini  
I-20090 Segrate (Milano)  
Tel: + 39-02 21018.1

**Κύπρος**

Οδός Αγίου Νικολάου, 8  
CY-1055 Λευκωσία  
Τηλ: +357-22 757188

**Latvija**

Bauskas 58a -401  
Rīga, LV-1004  
Tel: + 371-7 21 38 25

**Lietuva**

Kęstučio g. 65/40  
LT-08124 Vilnius  
Tel: + 370 52 101868

**Slovenská republika**

Strakova 5  
SK-811 01 Bratislava  
Tel: + 421 (2) 5920 2712

**Suomi/Finland**

PL 3/PB 3  
FIN-02201 Espoo/Esbo  
Puh/Tel: + 358-(0)20-7570 300

**Sverige**

Box 27190  
S-102 52 Stockholm  
Tel: + 46-(0)8 522 21 500

**United Kingdom**



Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW - UK  
Tel: + 44-(0)1 707 363 636

**This leaflet was last approved on**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>

## ANNEX TO THE PACKAGE LEAFLET

### How to use the PegIntron pre-filled pen



The following instructions explain how to inject yourself with the single use PegIntron pre-filled pen. **Please read the instructions carefully and completely before attempting to use the pen and follow them step by step.** Your doctor or his/her assistant will instruct you on how to self-inject with the PegIntron pre-filled pen. Do not attempt to inject yourself unless you are sure you understand the procedure and requirements for self-injection.


The PegIntron pre-filled pen is intended for use by one person only and must not be shared. Use the injection needle and cleansing swabs provided in the packaging only for the PegIntron pre-filled pen. Be sure the solution is at room temperature at the time of injection. Your doctor will have told you what dose you require for your therapy.

**Note:** The colour of the dosing button is different for each strength of the PegIntron pre-filled pen.

#### Step 1: Mixing

**It is important that you keep the PegIntron pre-filled pen upright (as shown in figure 1) during mixing, unless otherwise instructed.**

- Take the PegIntron pre-filled pen out of the refrigerator. Allow the medicine to come to room temperature.



- Wash your hands with soap and water.
- **Place the PegIntron pre-filled pen upright** in the holder of the tray provided in the pack (the dosing button will be on the **bottom**).
- You may want to hold the pre-filled pen using the grip. To mix the powder and the liquid, press the two halves together firmly by pressing down until you hear the pre-filled pen click. The two stoppers should come together.
- Wait for several seconds until the powder is completely dissolved.
- **Gently turn the PegIntron pre-filled pen upside down twice. To avoid excessive foaming, do not shake.**
- Maintaining the PegIntron pre-filled pen upright, check the mixed PegIntron solution through the window. If there is still foam, wait until it settles. The solution should be clear and colourless. **Do not use the pen if the solution is discoloured or contains any particles.**



Figure 2

- Keeping the PegIntron pre-filled pen upright in the holder provided in the packaging, disinfect the rubber membrane of the PegIntron pre-filled pen with one cleansing swab.
- Take the injection needle provided in the tray and remove its protective paper tab.
- **Maintaining the PegIntron pre-filled pen upright in the holder**, gently push the injection needle onto the pre-filled pen and **screw it securely in place**.

- **Keep the PegIntron pre-filled pen pointed up.**
- **Do not take off the outer needle cap at this point.**
- You may see some liquid trickle out from under the cap, as the air has been expelled out of the pen.
- Wait about 5 seconds for this process to finish.




Figure 3

- **Check through the window to be sure that the two stoppers are together. If they are not together, do not use this pen because you may not be able to dial your dose.**

### Step 2: Setting the dose



Figure 4

- Remove the PegIntron pre-filled pen from the holder.
- Holding the PegIntron pre-filled pen firmly, pull the dosing button out as far as it will go, until you see a **dark ring** on the pen. The dosing button should be easy to pull out without excessive force being required.

**Note: Do not push the dosing button back in at this time. You will push it in when you are ready to self-inject the PegIntron.**



Figure 5

- Turn the dosing button until your prescribed dose is aligned with the dosing tab. The button should turn easily without excessive force being required.

**Note:** **If you cannot easily pull out the dosing button or dial the dose, do not use excessive force and do not use this pen because it may not deliver the correct dose.**

### **Step 3: Injecting the solution**

- Select the injection site. Your doctor will have told you which sites to use (e.g., thigh or abdomen).

**Note:** change your injection site each time.

- Clean the injection site skin with the second cleansing swab.




Figure 6

- Pull off the **outer needle cap**.
- There may be some liquid around the inner needle cap. This liquid is not part of your dose, this is extra. This is normal, as the air has been expelled out of the needle.
- Once the injection site is dry, pull off the **yellow inner needle cap** carefully exposing the injection needle.



Figure 7

- Hold the PegIntron pre-filled pen with your fingers wrapped around the barrel and your thumb on the dosing button.
- With your other hand, pinch a fold of loose skin.
- Insert the needle into the pinched skin at an angle of 45° to 90°.
- Press the dosing button down **slowly** and **firmly** until the button can no longer move.
- **Maintain pressure on the dosing button for an additional 5 seconds** to ensure that you get the complete dose.

- Remove the needle from your skin.
- Press the injection site with a small bandage or sterile gauze if necessary for a few seconds.
- Do not massage the injection site. If there is bleeding, cover with an adhesive bandage.
- Discard the PegIntron pre-filled pen with the needle safely in a closed rigid container.

## PACKAGE LEAFLET: INFORMATION FOR THE USER

**PegIntron 80 micrograms powder and solvent for solution for injection in pre-filled pen**  
Peginterferon alfa-2b (conjugation of recombinant interferon alfa-2b with monomethoxy polyethylene glycol)

**Read all of this leaflet carefully before you start using this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What PegIntron is and what it is used for
2. Before you use PegIntron
3. How to use PegIntron
4. Possible side effects
5. Storing PegIntron
6. Further information

### **1. WHAT PEGINTRON IS AND WHAT IT IS USED FOR**

Interferons modify the response of the body's immune system to help fight infections and severe diseases. PegIntron, which contains an interferon, is used for the treatment of chronic hepatitis C, a viral infection of the liver, including in first time users co-infected with clinically stable HIV.

PegIntron is best used for this treatment in combination with ribavirin.

PegIntron is used alone in case of intolerance or contraindication to ribavirin.

### **2. BEFORE YOU USE PEGINTRON**

PegIntron is not recommended for use in patients under the age of 18 years.

**Do not use PegIntron**

- If you are allergic (hypersensitive) to peginterferon alfa-2b or any of the other ingredients of PegIntron.
- If you are allergic (hypersensitive) to any interferon.
- If you have had severe heart problems, or if you have heart disease that has not been well controlled during the past 6 months.
- If you have severe medical conditions that leave you very weak.
- If you have autoimmune hepatitis or any other problem with your immune system; if you are taking medicine that suppresses your immune system (your immune system protects you against infection and some diseases).
- If you have advanced, uncontrolled liver disease (other than hepatitis C).
- If you have thyroid disease that is not well controlled with medicines.
- If you have a condition that causes convulsions (seizures, or "fits").

**Take special care with PegIntron**

- If you develop symptoms of a severe allergic reaction (such as difficulty in breathing, wheezing, or hives) while on this medication, seek medical help immediately.
- If you ever had a heart attack or a heart problem.
- If you have ever been treated for depression or any other nervous or mental disorder.

- If you have kidney disease, your doctor may prescribe a lower than usual dose and monitor your kidney blood values regularly during treatment. If you have a kidney disease and you are using PegIntron in combination with medicinal products containing ribavirin, your doctor should monitor you more carefully for a decrease in red blood cell count.
- If you have had a problem with your liver (other than hepatitis C).
- If you develop symptoms associated with a cold or other respiratory infection, such as fever, cough, or any difficulty in breathing, tell your doctor.
- If you are diabetic, your doctor may ask you to have an eye examination.
- If you have had any serious illness affecting your breathing or your blood.
- If you have psoriasis, it may become worse while you are using PegIntron.
- If you are planning to become pregnant, discuss this with your doctor before starting to use PegIntron.
- If you are also being treated for HIV, please see **Using other medicines**.
- If you have had a severe nervous or mental disorder.
- If you ever had depression or develop symptoms associated with depression (e.g. feelings of sadness, dejection, etc) while on treatment with PegIntron (see section 4).
- If you have received an organ transplant, either kidney or liver, interferon treatment may increase the risk of rejection. Be sure to discuss this with your doctor.

Dental and gum disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and membranes of the mouth during long-term treatment with the combination of PegIntron with ribavirin. You should brush your teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If you have this reaction, be sure to rinse your mouth thoroughly afterwards.

### **While being treated with PegIntron**

Your doctor may want you to drink extra fluids to help prevent low blood pressure.

### **Using other medicines**

Please inform your doctor or pharmacist if you are taking or have recently taken any other medicines, even those not prescribed.

### **Pregnancy and breast-feeding**

Ask your doctor or pharmacist for advice before taking any medicine. In studies in pregnant animals, interferons have sometimes caused miscarriage. The effect on human pregnancy is not known. In combination therapy with ribavirin, ribavirin can be very damaging to an unborn baby, thus both female and male patients must take special precautions in their sexual activity if there is any chance for pregnancy to occur:

- if you are a **girl** or a **woman** of childbearing age, you must have a negative pregnancy test before treatment, each month during treatment, and for the 4 months after treatment is stopped. You and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 4 months after stopping treatment. This can be discussed with your doctor.
- if you are a **man** who is taking ribavirin, do not have sex with a pregnant woman unless you use a condom. This will lessen the chance for ribavirin to be left in the woman's body. If your female partner is not pregnant now but is of childbearing age, she must be tested for pregnancy each month during treatment and for the 7 months after treatment has stopped. This can be discussed with your doctor. If you are a male patient, you and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 7 months after stopping treatment. This can be discussed with your doctor.

It is not known whether this product is present in human milk. Therefore, do not breast-feed an infant if you are taking PegIntron. In combination therapy with ribavirin, take notice of the respective informing texts of ribavirin containing medicinal products.

### **Driving and using machines**

Do not drive or operate any tools or machines if you feel tired, sleepy or confused while taking PegIntron.

### **Important information about some of the ingredients of PegIntron**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

This medicinal product contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### **Patients who also have HIV infection**

Lactic acidosis and worsening liver function are side effects associated with Highly Active Anti-Retroviral Therapy (HAART), an HIV treatment. If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis and of liver failure. Your doctor will monitor you for signs and symptoms of these conditions. (Please be sure to read the ribavirin Patient Leaflet also). Additionally, patients treated with PegIntron and ribavirin combination therapy and zidovudine could be at increased risk of developing anaemia (low number of red blood cells). If you are also being treated for Human Immunodeficiency Virus (HIV) infection (AIDS) with zidovudine or stavudine, it is not certain if ribavirin will change the way these medicines work. Therefore, your blood will be checked regularly to be sure that the HIV infection is not getting worse. If it gets worse, your doctor will decide whether or not your ribavirin treatment needs to be changed.

Co-administration of ribavirin and didanosine and/or stavudine is not recommended due to the risk of lactic acidosis (a build-up of lactic acid in the body) and pancreatitis.

## **3. HOW TO USE PEGINTRON**

Your doctor has prescribed PegIntron specifically for you and your current condition; do not share this medicine with anyone else.

Your doctor has determined your dose of PegIntron based on your weight. If necessary, the dose may be changed during treatment.

### **Combination treatment**

PegIntron, when given with ribavirin capsules, is usually given at a dose of 1.5 microgram/kg once a week.

Ribavirin capsules are taken every day, morning and evening. The number of ribavirin capsules you take depends on your weight.

- If you weigh less than 65 kg, take 2 capsules in the morning and 2 in the evening (total of 800 mg each day).
- If you weigh between 65 and 85 kg, take 2 capsules in the morning and 3 in the evening (total of 1,000 mg each day).
- If you weigh more than 85 kg, take 3 capsules in the morning and 3 in the evening (total 1,200 mg each day).

The combination treatment is continued for 3 to 6 months, and sometimes for one year depending on your physicians judgement. Take notice of the respective informing texts of ribavirin containing medicinal products.

For HCV/HIV co-infected patients, the duration of treatment is 48 weeks.

### **PegIntron alone:**

PegIntron, when given alone, is usually given at a dose of 0.5 or 1.0 microgram/kg once a week, for at least 6 months, and possibly for 1 year. If you have kidney disease, your dose may be lower, depending upon your kidney function.

PegIntron alone was not studied in HCV/HIV co-infected patients.



**All patients:**

If you are injecting PegIntron yourself, please be sure that the dose that has been prescribed for you is clearly provided on the package of medicine you receive.

If you have the impression that the effect of PegIntron is too strong or too weak, talk to your doctor or pharmacist.

PegIntron is intended for subcutaneous use. This means that it is injected through a short injection needle into the fatty tissue just under your skin. If you are injecting this medicine yourself, you will be instructed how to prepare and give the injection. Detailed instructions for subcutaneous administration are provided with this leaflet (see How to use the PegIntron pre-filled pen).

Prepare the dose just before you intend to inject it and use it immediately. Look carefully at the reconstituted solution prior to administration. Do not use if there is discolouration of the reconstituted solution or if particulate matter is present. PegIntron is intended for single use only. Thus, after you give yourself the injection, discard the PegIntron pre-filled pen with any solution that is left in it.

Inject PegIntron once each week on the same day. Injecting it at the same time of day each week will help you not to forget to take it.

Always use PegIntron exactly as your doctor told you. Do not exceed the recommended dosage, and take it for as long as prescribed.

**If you use more PegIntron than you should**

Tell your doctor or healthcare professional as soon as possible.

**If you forget to take PegIntron**

Take the dose as soon as you remember, then continue your treatment as usual.

Do not take a double dose to make up for a forgotten dose. Contact your doctor or pharmacist if needed.

**4. POSSIBLE SIDE EFFECTS**

Like all medicines, PegIntron can cause side effects, although not everybody gets them. Although not all of these side effects may occur, they may need medical attention if they do.

Some people get depressed when taking PegIntron alone or in combination treatment with ribavirin, and in some cases people had suicidal thoughts or aggressive behaviour (sometimes directed against others). Some patients have actually committed suicide. Be sure to seek emergency care if you notice that you are becoming depressed or have suicidal thoughts or change in your behaviour. You may want to consider asking a family member or close friend to help you stay alert to signs of depression or changes in your behaviour.

Check with your doctor immediately if any of the following side effects occur: chest pain; changes in the way your heart beats; breathing problems (including shortness of breath), confusion; feeling depressed, wanting to harm yourself, hallucinations, numbness or tingling feeling; dizziness, convulsion ("fit"); trouble sleeping, thinking or concentrating; difficulty remaining alert, severe stomach pain or cramps; blood or clots in stool (or black, tarry stool); fever or chills beginning after a few weeks of treatment, pain in your lower back or side; difficulty or inability to pass urine, painful or inflamed muscles (sometimes severe); problems with your eyes or your eyesight or hearing; severe or painful reddening of your skin or mucous membrane, severe bleeding from your nose. Your doctor will test your blood to ensure that your white blood cell (cells that fight infection) and red blood cell (cells that carry oxygen) counts, platelets (blood clotting cells) and other laboratory values are at acceptable levels.

Other side effects that may occur with the combination of PegIntron and ribavirin capsules include:

Very commonly reported side effects (greater than 1 in every 10 patients):

Irritation or redness (and rarely, skin damage) at the site of injection, headache, tired feeling, shaking chills, fever, flu-like symptoms, weakness, loss of weight, nausea, loss of appetite, diarrhoea or loose stools, stomach pain, vomiting, muscle aches, pain in joints and muscles, feeling depressed, irritability, trouble falling asleep or staying asleep, feeling anxious or nervous, difficulty concentrating, mood swings, hair loss, itching, dry skin, sore throat, coughing, difficult breathing, dizziness, virus infection, rash, and dry mouth.

Commonly reported side effects (at least 1 in every 100 patients, but not greater than 1 in every 10 patients):

Increased sweating, chest pain, pain on the right side around your ribs, numbness, pain or tingling feeling, change in thyroid gland activity (which may make you feel tired or, less commonly, energetic), stomach upset, rapid heart rate, agitation, nervousness, difficult or irregular menstrual period

Less common are pain at the place of injection, flushing, low or high blood pressure, dry or teary eyes, redness of skin or skin disorder, psoriasis, hives, nail disorder, feeling unwell, feeling faint, poor coordination, confusion, increased or decreased sensitivity to touch, tense muscles, arthritis, bruising, loss of interest in activities including sex, sexual problem, unusual dreams, shaky hands, vertigo (spinning feeling), increased appetite, heartburn, intestinal gas (flatus), constipation, hemorrhoids, red or bleeding gums, redness or sores in mouth, change in taste, changes in hearing or ringing in ears, thirst, changed behaviour or aggressive behaviour (sometimes directed against others), feeling sleepy, cold sores, fungal or bacterial infections, irritation of prostate gland, increased need to pass urine, ear or respiratory infections, sinusitis, stuffy or runny nose, abnormal hair texture, sensitivity to sunlight, migraine headache, eye pain or infection, blurred vision, puffy face, puffy hands or feet, enlarged liver, problem affecting ovary or vagina, pain in breast, difficulty in speaking and swollen glands.

The following rare and very rare events have been reported with PegIntron:

Rarely reported side effects (greater than 1 in every 10,000 patients, but less than 1 in every 1,000 patients):

Diabetes, abnormal heart rhythm, inflammation of pancreas, inflammation and degeneration of muscle tissue and peripheral nerves, kidney problems and seizures.

Very rarely reported side effects (less than 1 in every 10,000 patients):

Very rarely sarcoidosis (a disease characterised by persistent fever, weight loss, joint pain and swelling, skin lesions and swollen glands) has been reported. Loss of consciousness has occurred very rarely with alpha interferons, mostly in elderly patients treated at high doses. Cases of stroke (cerebrovascular events) have been reported. Check with your doctor immediately if you have any of these symptoms or any other symptoms that are troubling.

Very rarely, PegIntron alone or in combination with ribavirin may cause aplastic anaemia. Pure red cell aplasia, a condition where the body stopped or reduced the production of red blood cells, has been reported. This causes severe anaemia, symptoms of which would include unusual tiredness and a lack of energy.

When PegIntron is used alone, some of these effects are less likely to occur, and some have not occurred at all.

If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis, liver failure, and development of blood abnormalities (reduction in number of red blood cells which carry oxygen, certain white blood cells that fight infection, and blood clotting cells called platelets).

The following other side effects (not listed above) have occurred with the combination of PegIntron and ribavirin capsules (adults) in HCV/HIV co-infected patients receiving HAART: oral candidiasis (oral thrush), defective metabolism of fat, CD4 lymphocytes decreased, appetite decreased, back pain, hepatitis, limb pain, and various laboratory blood values abnormalities.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE PEGINTRON**

Keep out of the reach and sight of children.

Store in a refrigerator (2°C - 8°C). Do not freeze.

Do not use PegIntron after the expiry date which is stated on the carton.

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

Do not use PegIntron if you notice discolouration of the powder.

The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. After administering the dose, discard the PegIntron pre-filled pen and any unused solution contained in it.

## **6. FURTHER INFORMATION**

### **What PegIntron contains**

- The active substance is peginterferon alfa-2b, 80 micrograms/0.5 ml.
- The other ingredients are:  
Powder: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80;  
Solvent: water for injections

### **What PegIntron looks like and contents of the pack**

The pharmaceutical form is: powder and solvent for solution for injection in a pre-filled pen. The white powder and the clear and colourless solvent are both contained in a two-chamber glass cartridge assembled into a single use pre-filled pen.

PegIntron 80 micrograms is available in different pack sizes:

- 1 pen containing powder and solvent for solution for injection, 1 injection needle and 2 cleansing swabs;
- 4 pens containing powder and solvent for solution for injection, 4 injection needles and 8 cleansing swabs;
- 6 pens containing powder and solvent for solution for injection, 6 injection needles and 12 cleansing swabs;
- 12 pens containing powder and solvent for solution for injection, 12 injection needles and 24 cleansing swabs.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

Marketing Authorisation Holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

Manufacturer: SP (Brinny) Company, Innishannon, County Cork, Ireland

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Rue de Stalle/Stallestraat 73  
B-1180 Bruxelles/Brussel/Brüssel  
Tél/Tel: + 32-(0)2 370 92 11

**Luxembourg/Luxemburg**

Rue de Stalle 73  
B-1180 Bruxelles/Brüssel  
Belgique/Belgien  
Tél/Tel: + 32-(0)2 370 92 11

**България**

ул. "Проф. Милко Бичев" № 7, ет.3  
BG-София 1504  
Тел.: +359 2 944 1073

**Magyarország**

Alkotás u. 53.  
H-1123 Budapest  
Tel.: +36 1 457-8500

**Česká republika**

Na Příkopě 25  
CZ-110 00 Praha 1  
Tel: +420 221771250

**Malta**

168 Christopher Street  
MT-VLT02 Valletta  
Tel: + 356-21 23 21 75

**Danmark**

Lautrupbjerg 2  
DK-2750 Ballerup  
Tlf: + 45-44 39 50 00

**Nederland**

Maarssenbroeksedijk 4  
NL-3542 DN Utrecht  
Tel: + 31-(0)800 778 78 78

**Deutschland**

Thomas-Dehler-Straße 27  
D-81737 München  
Tel: + 49-(0)89 627 31-0

**Norge**

Pb. 398  
N-1326 Lysaker  
Tlf: + 47 67 16 64 50

**Eesti**

Järvevana tee 9  
EE-11314 Tallinn  
Tel: + 372 654 96 86

**Österreich**

Badener Strasse 23  
A-2514 Traiskirchen  
Tel: + 43-(0)2252 502-0

**Ελλάδα**

Αγίου Δημητρίου 63  
GR-174 55 Αλιμος  
Τηλ.: + 30-210 98 97 300

**Polska**

Al. Jerozolimskie 195a  
PL-02-222 Warszawa  
Tel.: + 48-(0)22 478 41 50

**España**

Km. 36, Ctra. Nacional I  
E-28750 San Agustín de Guadalix – Madrid  
Tel: + 34-91 848 85 00

**Portugal**

Rua Agualva dos Açores 16  
P-2735-557 Agualva-Cacém  
Tel: +351-21 433 93 00

**France**

92 rue Baudin  
F-92300 Levallois-Perret  
Tél: + 33-(0)1 41 06 35 00

**România**

Șos. București-Ploiești, nr. 17-21, Băneasa  
Center, et. 7, sector 1  
RO-013682 București  
Tel: + 40 21 233 35 30

**Ireland**

Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW  
Tel: +44-(0)1 707 363 636

**Slovenija**

Dunajska 22  
SI-1000 Ljubljana  
Tel: + 386 01 3001070

**Ísland**

Hörgatún 2  
IS-210 Garðabær  
Sími: + 354 535 70 00

**Slovenská republika**

Strakova 5  
SK-811 01 Bratislava  
Tel: + 421 (2) 5920 2712

**Italia**

Via fratelli Cervi snc,  
Centro Direzionale Milano Due  
Palazzo Borromini  
I-20090 Segrate (Milano)  
Tel: + 39-02 21018.1

**Κύπρος**

Οδός Αγίου Νικολάου, 8  
CY-1055 Λευκωσία  
Τηλ: +357-22 757188

**Latvija**

Bauskas 58a -401  
Rīga, LV-1004  
Tel: + 371-7 21 38 25

**Lietuva**

Kęstučio g. 65/40  
LT-08124 Vilnius  
Tel: + 370 52 101868

**Suomi/Finland**

PL 3/PB 3  
FIN-02201 Espoo/Esbo  
Puh/Tel: + 358-(0)20-7570 300

**Sverige**

Box 27190  
S-102 52 Stockholm  
Tel: + 46-(0)8 522 21 500

**United Kingdom**



Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW - UK  
Tel: + 44-(0)1 707 363 636

**This leaflet was last approved on**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>

## ANNEX TO THE PACKAGE LEAFLET

### How to use the PegIntron pre-filled pen



The following instructions explain how to inject yourself with the single use PegIntron pre-filled pen. **Please read the instructions carefully and completely before attempting to use the pen and follow them step by step.** Your doctor or his/her assistant will instruct you on how to self-inject with the PegIntron pre-filled pen. Do not attempt to inject yourself unless you are sure you understand the procedure and requirements for self-injection.


The PegIntron pre-filled pen is intended for use by one person only and must not be shared. Use the injection needle and cleansing swabs provided in the packaging only for the PegIntron pre-filled pen. Be sure the solution is at room temperature at the time of injection. Your doctor will have told you what dose you require for your therapy.

**Note:** The colour of the dosing button is different for each strength of the PegIntron pre-filled pen.

#### Step 1: Mixing

**It is important that you keep the PegIntron pre-filled pen upright (as shown in figure 1) during mixing, unless otherwise instructed.**

- Take the PegIntron pre-filled pen out of the refrigerator. Allow the medicine to come to room temperature.



- Wash your hands with soap and water.
- **Place the PegIntron pre-filled pen upright** in the holder of the tray provided in the pack (the dosing button will be on the **bottom**).
- You may want to hold the pre-filled pen using the grip. To mix the powder and the liquid, press the two halves together firmly by pressing down until you hear the pre-filled pen click. The two stoppers should come together.
- Wait for several seconds until the powder is completely dissolved.
- **Gently turn the PegIntron pre-filled pen upside down twice. To avoid excessive foaming, do not shake.**
- Maintaining the PegIntron pre-filled pen upright, check the mixed PegIntron solution through the window. If there is still foam, wait until it settles. The solution should be clear and colourless. **Do not use the pen if the solution is discoloured or contains any particles.**



Figure 2

- Keeping the PegIntron pre-filled pen upright in the holder provided in the packaging, disinfect the rubber membrane of the PegIntron pre-filled pen with one cleansing swab.
- Take the injection needle provided in the tray and remove its protective paper tab.
- **Maintaining the PegIntron pre-filled pen upright in the holder**, gently push the injection needle onto the pre-filled pen and **screw it securely in place**.

- **Keep the PegIntron pre-filled pen pointed up.**
- **Do not take off the outer needle cap at this point.**
- You may see some liquid trickle out from under the cap, as the air has been expelled out of the pen.
- Wait about 5 seconds for this process to finish.




Figure 3

- **Check through the window to be sure that the two stoppers are together. If they are not together, do not use this pen because you may not be able to dial your dose.**

### Step 2: Setting the dose



Figure 4

- Remove the PegIntron pre-filled pen from the holder.
- Holding the PegIntron pre-filled pen firmly, pull the dosing button out as far as it will go, until you see a **dark ring** on the pen. The dosing button should be easy to pull out without excessive force being required.

**Note: Do not push the dosing button back in at this time. You will push it in when you are ready to self-inject the PegIntron.**



Figure 5

- Turn the dosing button until your prescribed dose is aligned with the dosing tab. The button should turn easily without excessive force being required.

**Note:** **If you cannot easily pull out the dosing button or dial the dose, do not use excessive force and do not use this pen because it may not deliver the correct dose.**

### **Step 3: Injecting the solution**

- Select the injection site. Your doctor will have told you which sites to use (e.g., thigh or abdomen).

**Note:** change your injection site each time.

- Clean the injection site skin with the second cleansing swab.




Figure 6

- Pull off the **outer needle cap**.
- There may be some liquid around the inner needle cap. This liquid is not part of your dose, this is extra. This is normal, as the air has been expelled out of the needle.
- Once the injection site is dry, pull off the **yellow inner needle cap** carefully exposing the injection needle.



Figure 7

- Hold the PegIntron pre-filled pen with your fingers wrapped around the barrel and your thumb on the dosing button.
- With your other hand, pinch a fold of loose skin.
- Insert the needle into the pinched skin at an angle of 45° to 90°.
- Press the dosing button down **slowly** and **firmly** until the button can no longer move.
- **Maintain pressure on the dosing button for an additional 5 seconds** to ensure that you get the complete dose.



- Remove the needle from your skin.
- Press the injection site with a small bandage or sterile gauze if necessary for a few seconds.
- Do not massage the injection site. If there is bleeding, cover with an adhesive bandage.
- Discard the PegIntron pre-filled pen with the needle safely in a closed rigid container.

## **PACKAGE LEAFLET: INFORMATION FOR THE USER**

**PegIntron 100 micrograms powder and solvent for solution for injection in pre-filled pen**  
Peginterferon alfa-2b (conjugation of recombinant interferon alfa-2b with monomethoxy polyethylene glycol)

**Read all of this leaflet carefully before you start using this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What PegIntron is and what it is used for
2. Before you use PegIntron
3. How to use PegIntron
4. Possible side effects
5. Storing PegIntron
6. Further information

### **1. WHAT PEGINTRON IS AND WHAT IT IS USED FOR**

Interferons modify the response of the body's immune system to help fight infections and severe diseases. PegIntron, which contains an interferon, is used for the treatment of chronic hepatitis C, a viral infection of the liver, including in first time users co-infected with clinically stable HIV.

PegIntron is best used for this treatment in combination with ribavirin.

PegIntron is used alone in case of intolerance or contraindication to ribavirin.

### **2. BEFORE YOU USE PEGINTRON**

PegIntron is not recommended for use in patients under the age of 18 years.

**Do not use PegIntron**

- If you are allergic (hypersensitive) to peginterferon alfa-2b or any of the other ingredients of PegIntron.
- If you are allergic (hypersensitive) to any interferon.
- If you have had severe heart problems, or if you have heart disease that has not been well controlled during the past 6 months.
- If you have severe medical conditions that leave you very weak.
- If you have autoimmune hepatitis or any other problem with your immune system; if you are taking medicine that suppresses your immune system (your immune system protects you against infection and some diseases).
- If you have advanced, uncontrolled liver disease (other than hepatitis C).
- If you have thyroid disease that is not well controlled with medicines.
- If you have a condition that causes convulsions (seizures, or "fits").

**Take special care with PegIntron**

- If you develop symptoms of a severe allergic reaction (such as difficulty in breathing, wheezing, or hives) while on this medication, seek medical help immediately.
- If you ever had a heart attack or a heart problem.
- If you have ever been treated for depression or any other nervous or mental disorder.

- If you have kidney disease, your doctor may prescribe a lower than usual dose and monitor your kidney blood values regularly during treatment. If you have a kidney disease and you are using PegIntron in combination with medicinal products containing ribavirin, your doctor should monitor you more carefully for a decrease in red blood cell count.
- If you have had a problem with your liver (other than hepatitis C).
- If you develop symptoms associated with a cold or other respiratory infection, such as fever, cough, or any difficulty in breathing, tell your doctor.
- If you are diabetic, your doctor may ask you to have an eye examination.
- If you have had any serious illness affecting your breathing or your blood.
- If you have psoriasis, it may become worse while you are using PegIntron.
- If you are planning to become pregnant, discuss this with your doctor before starting to use PegIntron.
- If you are also being treated for HIV, please see **Using other medicines**.
- If you have had a severe nervous or mental disorder.
- If you ever had depression or develop symptoms associated with depression (e.g. feelings of sadness, dejection, etc) while on treatment with PegIntron (see section 4).
- If you have received an organ transplant, either kidney or liver, interferon treatment may increase the risk of rejection. Be sure to discuss this with your doctor.

Dental and gum disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and membranes of the mouth during long-term treatment with the combination of PegIntron with ribavirin. You should brush your teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If you have this reaction, be sure to rinse your mouth thoroughly afterwards.

### **While being treated with PegIntron**

Your doctor may want you to drink extra fluids to help prevent low blood pressure.

### **Using other medicines**

Please inform your doctor or pharmacist if you are taking or have recently taken any other medicines, even those not prescribed.

### **Pregnancy and breast-feeding**

Ask your doctor or pharmacist for advice before taking any medicine. In studies in pregnant animals, interferons have sometimes caused miscarriage. The effect on human pregnancy is not known. In combination therapy with ribavirin, ribavirin can be very damaging to an unborn baby, thus both female and male patients must take special precautions in their sexual activity if there is any chance for pregnancy to occur:

- if you are a **girl** or a **woman** of childbearing age, you must have a negative pregnancy test before treatment, each month during treatment, and for the 4 months after treatment is stopped. You and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 4 months after stopping treatment. This can be discussed with your doctor.
- if you are a **man** who is taking ribavirin, do not have sex with a pregnant woman unless you use a condom. This will lessen the chance for ribavirin to be left in the woman's body. If your female partner is not pregnant now but is of childbearing age, she must be tested for pregnancy each month during treatment and for the 7 months after treatment has stopped. This can be discussed with your doctor. If you are a male patient, you and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 7 months after stopping treatment. This can be discussed with your doctor.

It is not known whether this product is present in human milk. Therefore, do not breast-feed an infant if you are taking PegIntron. In combination therapy with ribavirin, take notice of the respective informing texts of ribavirin containing medicinal products.

### **Driving and using machines**

Do not drive or operate any tools or machines if you feel tired, sleepy or confused while taking PegIntron.

### **Important information about some of the ingredients of PegIntron**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

This medicinal product contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### **Patients who also have HIV infection**

Lactic acidosis and worsening liver function are side effects associated with Highly Active Anti-Retroviral Therapy (HAART), an HIV treatment. If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis and of liver failure. Your doctor will monitor you for signs and symptoms of these conditions. (Please be sure to read the ribavirin Patient Leaflet also). Additionally, patients treated with PegIntron and ribavirin combination therapy and zidovudine could be at increased risk of developing anaemia (low number of red blood cells). If you are also being treated for Human Immunodeficiency Virus (HIV) infection (AIDS) with zidovudine or stavudine, it is not certain if ribavirin will change the way these medicines work. Therefore, your blood will be checked regularly to be sure that the HIV infection is not getting worse. If it gets worse, your doctor will decide whether or not your ribavirin treatment needs to be changed.

Co-administration of ribavirin and didanosine and/or stavudine is not recommended due to the risk of lactic acidosis (a build-up of lactic acid in the body) and pancreatitis.

## **3. HOW TO USE PEGINTRON**

Your doctor has prescribed PegIntron specifically for you and your current condition; do not share this medicine with anyone else.

Your doctor has determined your dose of PegIntron based on your weight. If necessary, the dose may be changed during treatment.

### **Combination treatment**

PegIntron, when given with ribavirin capsules, is usually given at a dose of 1.5 microgram/kg once a week.

Ribavirin capsules are taken every day, morning and evening. The number of ribavirin capsules you take depends on your weight.

- If you weigh less than 65 kg, take 2 capsules in the morning and 2 in the evening (total of 800 mg each day).
- If you weigh between 65 and 85 kg, take 2 capsules in the morning and 3 in the evening (total of 1,000 mg each day).
- If you weigh more than 85 kg, take 3 capsules in the morning and 3 in the evening (total 1,200 mg each day).

The combination treatment is continued for 3 to 6 months, and sometimes for one year depending on your physicians judgement. Take notice of the respective informing texts of ribavirin containing medicinal products.

For HCV/HIV co-infected patients, the duration of treatment is 48 weeks.

### **PegIntron alone:**

PegIntron, when given alone, is usually given at a dose of 0.5 or 1.0 microgram/kg once a week, for at least 6 months, and possibly for 1 year. If you have kidney disease, your dose may be lower, depending upon your kidney function.

PegIntron alone was not studied in HCV/HIV co-infected patients.

**All patients:**

If you are injecting PegIntron yourself, please be sure that the dose that has been prescribed for you is clearly provided on the package of medicine you receive.

If you have the impression that the effect of PegIntron is too strong or too weak, talk to your doctor or pharmacist.

PegIntron is intended for subcutaneous use. This means that it is injected through a short injection needle into the fatty tissue just under your skin. If you are injecting this medicine yourself, you will be instructed how to prepare and give the injection. Detailed instructions for subcutaneous administration are provided with this leaflet (see How to use the PegIntron pre-filled pen).

Prepare the dose just before you intend to inject it and use it immediately. Look carefully at the reconstituted solution prior to administration. Do not use if there is discolouration of the reconstituted solution or if particulate matter is present. PegIntron is intended for single use only. Thus, after you give yourself the injection, discard the PegIntron pre-filled pen with any solution that is left in it.

Inject PegIntron once each week on the same day. Injecting it at the same time of day each week will help you not to forget to take it.

Always use PegIntron exactly as your doctor told you. Do not exceed the recommended dosage, and take it for as long as prescribed.

**If you use more PegIntron than you should**

Tell your doctor or healthcare professional as soon as possible.

**If you forget to take PegIntron**

Take the dose as soon as you remember, then continue your treatment as usual.

Do not take a double dose to make up for a forgotten dose. Contact your doctor or pharmacist if needed.

**4. POSSIBLE SIDE EFFECTS**

Like all medicines, PegIntron can cause side effects, although not everybody gets them. Although not all of these side effects may occur, they may need medical attention if they do.

Some people get depressed when taking PegIntron alone or in combination treatment with ribavirin, and in some cases people had suicidal thoughts or aggressive behaviour (sometimes directed against others). Some patients have actually committed suicide. Be sure to seek emergency care if you notice that you are becoming depressed or have suicidal thoughts or change in your behaviour. You may want to consider asking a family member or close friend to help you stay alert to signs of depression or changes in your behaviour.

Check with your doctor immediately if any of the following side effects occur: chest pain; changes in the way your heart beats; breathing problems (including shortness of breath), confusion; feeling depressed, wanting to harm yourself, hallucinations, numbness or tingling feeling; dizziness, convulsion ("fit"); trouble sleeping, thinking or concentrating; difficulty remaining alert, severe stomach pain or cramps; blood or clots in stool (or black, tarry stool); fever or chills beginning after a few weeks of treatment, pain in your lower back or side; difficulty or inability to pass urine, painful or inflamed muscles (sometimes severe); problems with your eyes or your eyesight or hearing; severe or painful reddening of your skin or mucous membrane, severe bleeding from your nose. Your doctor will test your blood to ensure that your white blood cell (cells that fight infection) and red blood cell (cells that carry oxygen) counts, platelets (blood clotting cells) and other laboratory values are at acceptable levels.

Other side effects that may occur with the combination of PegIntron and ribavirin capsules include:

Very commonly reported side effects (greater than 1 in every 10 patients):

Irritation or redness (and rarely, skin damage) at the site of injection, headache, tired feeling, shaking chills, fever, flu-like symptoms, weakness, loss of weight, nausea, loss of appetite, diarrhoea or loose stools, stomach pain, vomiting, muscle aches, pain in joints and muscles, feeling depressed, irritability, trouble falling asleep or staying asleep, feeling anxious or nervous, difficulty concentrating, mood swings, hair loss, itching, dry skin, sore throat, coughing, difficult breathing, dizziness, virus infection, rash, and dry mouth.

Commonly reported side effects (at least 1 in every 100 patients, but not greater than 1 in every 10 patients):

Increased sweating, chest pain, pain on the right side around your ribs, numbness, pain or tingling feeling, change in thyroid gland activity (which may make you feel tired or, less commonly, energetic), stomach upset, rapid heart rate, agitation, nervousness, difficult or irregular menstrual period

Less common are pain at the place of injection, flushing, low or high blood pressure, dry or teary eyes, redness of skin or skin disorder, psoriasis, hives, nail disorder, feeling unwell, feeling faint, poor coordination, confusion, increased or decreased sensitivity to touch, tense muscles, arthritis, bruising, loss of interest in activities including sex, sexual problem, unusual dreams, shaky hands, vertigo (spinning feeling), increased appetite, heartburn, intestinal gas (flatus), constipation, hemorrhoids, red or bleeding gums, redness or sores in mouth, change in taste, changes in hearing or ringing in ears, thirst, changed behaviour or aggressive behaviour (sometimes directed against others), feeling sleepy, cold sores, fungal or bacterial infections, irritation of prostate gland, increased need to pass urine, ear or respiratory infections, sinusitis, stuffy or runny nose, abnormal hair texture, sensitivity to sunlight, migraine headache, eye pain or infection, blurred vision, puffy face, puffy hands or feet, enlarged liver, problem affecting ovary or vagina, pain in breast, difficulty in speaking and swollen glands.

The following rare and very rare events have been reported with PegIntron:

Rarely reported side effects (greater than 1 in every 10,000 patients, but less than 1 in every 1,000 patients):

Diabetes, abnormal heart rhythm, inflammation of pancreas, inflammation and degeneration of muscle tissue and peripheral nerves, kidney problems and seizures.

Very rarely reported side effects (less than 1 in every 10,000 patients):

Very rarely sarcoidosis (a disease characterised by persistent fever, weight loss, joint pain and swelling, skin lesions and swollen glands) has been reported. Loss of consciousness has occurred very rarely with alpha interferons, mostly in elderly patients treated at high doses. Cases of stroke (cerebrovascular events) have been reported. Check with your doctor immediately if you have any of these symptoms or any other symptoms that are troubling.

Very rarely, PegIntron alone or in combination with ribavirin may cause aplastic anaemia. Pure red cell aplasia, a condition where the body stopped or reduced the production of red blood cells, has been reported. This causes severe anaemia, symptoms of which would include unusual tiredness and a lack of energy.

When PegIntron is used alone, some of these effects are less likely to occur, and some have not occurred at all.

If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis, liver failure, and development of blood abnormalities (reduction in number of red blood cells which carry oxygen, certain white blood cells that fight infection, and blood clotting cells called platelets).

The following other side effects (not listed above) have occurred with the combination of PegIntron and ribavirin capsules (adults) in HCV/HIV co-infected patients receiving HAART: oral candidiasis (oral thrush), defective metabolism of fat, CD4 lymphocytes decreased, appetite decreased, back pain, hepatitis, limb pain, and various laboratory blood values abnormalities.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE PEGINTRON**

Keep out of the reach and sight of children.

Store in a refrigerator (2°C - 8°C). Do not freeze.

Do not use PegIntron after the expiry date which is stated on the carton.

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

Do not use PegIntron if you notice discolouration of the powder.

The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. After administering the dose, discard the PegIntron pre-filled pen and any unused solution contained in it.

## **6. FURTHER INFORMATION**

### **What PegIntron contains**

- The active substance is peginterferon alfa-2b, 100 micrograms/0.5 ml.
- The other ingredients are:  
Powder: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80;  
Solvent: water for injections

### **What PegIntron looks like and contents of the pack**

The pharmaceutical form is: powder and solvent for solution for injection in a pre-filled pen. The white powder and the clear and colourless solvent are both contained in a two-chamber glass cartridge assembled into a single use pre-filled pen.

PegIntron 100 micrograms is available in different pack sizes:

- 1 pen containing powder and solvent for solution for injection, 1 injection needle and 2 cleansing swabs;
- 4 pens containing powder and solvent for solution for injection, 4 injection needles and 8 cleansing swabs;
- 6 pens containing powder and solvent for solution for injection, 6 injection needles and 12 cleansing swabs;
- 12 pens containing powder and solvent for solution for injection, 12 injection needles and 24 cleansing swabs.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

Marketing Authorisation Holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

Manufacturer: SP (Brinny) Company, Innishannon, County Cork, Ireland

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Rue de Stalle/Stallestraat 73  
B-1180 Bruxelles/Brussel/Brüssel  
Tél/Tel: + 32-(0)2 370 92 11

**Luxembourg/Luxemburg**

Rue de Stalle 73  
B-1180 Bruxelles/Brüssel  
Belgique/Belgien  
Tél/Tel: + 32-(0)2 370 92 11

**България**

ул. "Проф. Милко Бичев" № 7, ет.3  
BG-София 1504  
Тел.: +359 2 944 1073

**Magyarország**

Alkotás u. 53.  
H-1123 Budapest  
Tel.: +36 1 457-8500

**Česká republika**

Na Příkopě 25  
CZ-110 00 Praha 1  
Tel: +420 221771250

**Malta**

168 Christopher Street  
MT-VLT02 Valletta  
Tel: + 356-21 23 21 75

**Danmark**

Lautrupbjerg 2  
DK-2750 Ballerup  
Tlf: + 45-44 39 50 00

**Nederland**

Maarssenbroeksedijk 4  
NL-3542 DN Utrecht  
Tel: + 31-(0)800 778 78 78

**Deutschland**

Thomas-Dehler-Straße 27  
D-81737 München  
Tel: + 49-(0)89 627 31-0

**Norge**

Pb. 398  
N-1326 Lysaker  
Tlf: + 47 67 16 64 50

**Eesti**

Järvevana tee 9  
EE-11314 Tallinn  
Tel: + 372 654 96 86

**Österreich**

Badener Strasse 23  
A-2514 Traiskirchen  
Tel: + 43-(0)2252 502-0

**Ελλάδα**

Αγίου Δημητρίου 63  
GR-174 55 Αλιμος  
Τηλ.: + 30-210 98 97 300

**Polska**

Al. Jerozolimskie 195a  
PL-02-222 Warszawa  
Tel.: + 48-(0)22 478 41 50

**España**

Km. 36, Ctra. Nacional I  
E-28750 San Agustín de Guadalix – Madrid  
Tel: + 34-91 848 85 00

**Portugal**

Rua Agualva dos Açores 16  
P-2735-557 Agualva-Cacém  
Tel: +351-21 433 93 00

**France**

92 rue Baudin  
F-92300 Levallois-Perret  
Tél: + 33-(0)1 41 06 35 00

**România**

Șos. București-Ploiești, nr. 17-21, Băneasa  
Center, et. 7, sector 1  
RO-013682 București  
Tel: + 40 21 233 35 30

**Ireland**

Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW  
Tel: +44-(0)1 707 363 636

**Slovenija**

Dunajska 22  
SI-1000 Ljubljana  
Tel: + 386 01 3001070

**Ísland**

Hörgatún 2  
IS-210 Garðabær  
Sími: + 354 535 70 00

**Slovenská republika**

Strakova 5  
SK-811 01 Bratislava  
Tel: + 421 (2) 5920 2712



**Italia**

Via fratelli Cervi snc,  
Centro Direzionale Milano Due  
Palazzo Borromini  
I-20090 Segrate (Milano)  
Tel: + 39-02 21018.1

**Κύπρος**

Οδός Αγίου Νικολάου, 8  
CY-1055 Λευκωσία  
Τηλ: +357-22 757188

**Latvija**

Bauskas 58a -401  
Rīga, LV-1004  
Tel: + 371-7 21 38 25

**Lietuva**

Kęstučio g. 65/40  
LT-08124 Vilnius  
Tel: + 370 52 101868

**Suomi/Finland**

PL 3/PB 3  
FIN-02201 Espoo/Esbo  
Puh/Tel: + 358-(0)20-7570 300

**Sverige**

Box 27190  
S-102 52 Stockholm  
Tel: + 46-(0)8 522 21 500

**United Kingdom**



Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW - UK  
Tel: + 44-(0)1 707 363 636

**This leaflet was last approved on**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>

## ANNEX TO THE PACKAGE LEAFLET

### How to use the PegIntron pre-filled pen



The following instructions explain how to inject yourself with the single use PegIntron pre-filled pen. **Please read the instructions carefully and completely before attempting to use the pen and follow them step by step.** Your doctor or his/her assistant will instruct you on how to self-inject with the PegIntron pre-filled pen. Do not attempt to inject yourself unless you are sure you understand the procedure and requirements for self-injection.


The PegIntron pre-filled pen is intended for use by one person only and must not be shared. Use the injection needle and cleansing swabs provided in the packaging only for the PegIntron pre-filled pen. Be sure the solution is at room temperature at the time of injection. Your doctor will have told you what dose you require for your therapy.

**Note:** The colour of the dosing button is different for each strength of the PegIntron pre-filled pen.

#### Step 1: Mixing

**It is important that you keep the PegIntron pre-filled pen upright (as shown in figure 1) during mixing, unless otherwise instructed.**

- Take the PegIntron pre-filled pen out of the refrigerator. Allow the medicine to come to room temperature.



- Wash your hands with soap and water.
- **Place the PegIntron pre-filled pen upright** in the holder of the tray provided in the pack (the dosing button will be on the **bottom**).
- You may want to hold the pre-filled pen using the grip. To mix the powder and the liquid, press the two halves together firmly by pressing down until you hear the pre-filled pen click. The two stoppers should come together.
- Wait for several seconds until the powder is completely dissolved.
- **Gently turn the PegIntron pre-filled pen upside down twice. To avoid excessive foaming, do not shake.**
- Maintaining the PegIntron pre-filled pen upright, check the mixed PegIntron solution through the window. If there is still foam, wait until it settles. The solution should be clear and colourless. **Do not use the pen if the solution is discoloured or contains any particles.**



Figure 2

- Keeping the PegIntron pre-filled pen upright in the holder provided in the packaging, disinfect the rubber membrane of the PegIntron pre-filled pen with one cleansing swab.
- Take the injection needle provided in the tray and remove its protective paper tab.
- **Maintaining the PegIntron pre-filled pen upright in the holder**, gently push the injection needle onto the pre-filled pen and **screw it securely in place**.

- **Keep the PegIntron pre-filled pen pointed up.**
- **Do not take off the outer needle cap at this point.**
- You may see some liquid trickle out from under the cap, as the air has been expelled out of the pen.
- Wait about 5 seconds for this process to finish.




Figure 3

- **Check through the window to be sure that the two stoppers are together. If they are not together, do not use this pen because you may not be able to dial your dose.**

### Step 2: Setting the dose



Figure 4

- Remove the PegIntron pre-filled pen from the holder.
- Holding the PegIntron pre-filled pen firmly, pull the dosing button out as far as it will go, until you see a **dark ring** on the pen. The dosing button should be easy to pull out without excessive force being required.

**Note: Do not push the dosing button back in at this time. You will push it in when you are ready to self-inject the PegIntron.**



Figure 5

- Turn the dosing button until your prescribed dose is aligned with the dosing tab. The button should turn easily without excessive force being required.

**Note:** **If you cannot easily pull out the dosing button or dial the dose, do not use excessive force and do not use this pen because it may not deliver the correct dose.**

### **Step 3: Injecting the solution**

- Select the injection site. Your doctor will have told you which sites to use (e.g., thigh or abdomen).

**Note:** change your injection site each time.

- Clean the injection site skin with the second cleansing swab.




Figure 6

- Pull off the **outer needle cap**.
- There may be some liquid around the inner needle cap. This liquid is not part of your dose, this is extra. This is normal, as the air has been expelled out of the needle.
- Once the injection site is dry, pull off the **yellow inner needle cap** carefully exposing the injection needle.



Figure 7

- Hold the PegIntron pre-filled pen with your fingers wrapped around the barrel and your thumb on the dosing button.
- With your other hand, pinch a fold of loose skin.
- Insert the needle into the pinched skin at an angle of 45° to 90°.
- Press the dosing button down **slowly** and **firmly** until the button can no longer move.
- **Maintain pressure on the dosing button for an additional 5 seconds** to ensure that you get the complete dose.

- Remove the needle from your skin.
- Press the injection site with a small bandage or sterile gauze if necessary for a few seconds.
- Do not massage the injection site. If there is bleeding, cover with an adhesive bandage.
- Discard the PegIntron pre-filled pen with the needle safely in a closed rigid container.

## **PACKAGE LEAFLET: INFORMATION FOR THE USER**

**PegIntron 120 micrograms powder and solvent for solution for injection in pre-filled pen**  
Peginterferon alfa-2b (conjugation of recombinant interferon alfa-2b with monomethoxy polyethylene glycol)

**Read all of this leaflet carefully before you start using this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What PegIntron is and what it is used for
2. Before you use PegIntron
3. How to use PegIntron
4. Possible side effects
5. Storing PegIntron
6. Further information

### **1. WHAT PEGINTRON IS AND WHAT IT IS USED FOR**

Interferons modify the response of the body's immune system to help fight infections and severe diseases. PegIntron, which contains an interferon, is used for the treatment of chronic hepatitis C, a viral infection of the liver, including in first time users co-infected with clinically stable HIV.

PegIntron is best used for this treatment in combination with ribavirin.

PegIntron is used alone in case of intolerance or contraindication to ribavirin.

### **2. BEFORE YOU USE PEGINTRON**

PegIntron is not recommended for use in patients under the age of 18 years.

**Do not use PegIntron**

- If you are allergic (hypersensitive) to peginterferon alfa-2b or any of the other ingredients of PegIntron.
- If you are allergic (hypersensitive) to any interferon.
- If you have had severe heart problems, or if you have heart disease that has not been well controlled during the past 6 months.
- If you have severe medical conditions that leave you very weak.
- If you have autoimmune hepatitis or any other problem with your immune system; if you are taking medicine that suppresses your immune system (your immune system protects you against infection and some diseases).
- If you have advanced, uncontrolled liver disease (other than hepatitis C).
- If you have thyroid disease that is not well controlled with medicines.
- If you have a condition that causes convulsions (seizures, or "fits").

**Take special care with PegIntron**

- If you develop symptoms of a severe allergic reaction (such as difficulty in breathing, wheezing, or hives) while on this medication, seek medical help immediately.
- If you ever had a heart attack or a heart problem.
- If you have ever been treated for depression or any other nervous or mental disorder.

- If you have kidney disease, your doctor may prescribe a lower than usual dose and monitor your kidney blood values regularly during treatment. If you have a kidney disease and you are using PegIntron in combination with medicinal products containing ribavirin, your doctor should monitor you more carefully for a decrease in red blood cell count.
- If you have had a problem with your liver (other than hepatitis C).
- If you develop symptoms associated with a cold or other respiratory infection, such as fever, cough, or any difficulty in breathing, tell your doctor.
- If you are diabetic, your doctor may ask you to have an eye examination.
- If you have had any serious illness affecting your breathing or your blood.
- If you have psoriasis, it may become worse while you are using PegIntron.
- If you are planning to become pregnant, discuss this with your doctor before starting to use PegIntron.
- If you are also being treated for HIV, please see **Using other medicines**.
- If you have had a severe nervous or mental disorder.
- If you ever had depression or develop symptoms associated with depression (e.g. feelings of sadness, dejection, etc) while on treatment with PegIntron (see section 4).
- If you have received an organ transplant, either kidney or liver, interferon treatment may increase the risk of rejection. Be sure to discuss this with your doctor.

Dental and gum disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and membranes of the mouth during long-term treatment with the combination of PegIntron with ribavirin. You should brush your teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If you have this reaction, be sure to rinse your mouth thoroughly afterwards.

### **While being treated with PegIntron**

Your doctor may want you to drink extra fluids to help prevent low blood pressure.

### **Using other medicines**

Please inform your doctor or pharmacist if you are taking or have recently taken any other medicines, even those not prescribed.

### **Pregnancy and breast-feeding**

Ask your doctor or pharmacist for advice before taking any medicine. In studies in pregnant animals, interferons have sometimes caused miscarriage. The effect on human pregnancy is not known. In combination therapy with ribavirin, ribavirin can be very damaging to an unborn baby, thus both female and male patients must take special precautions in their sexual activity if there is any chance for pregnancy to occur:

- if you are a **girl** or a **woman** of childbearing age, you must have a negative pregnancy test before treatment, each month during treatment, and for the 4 months after treatment is stopped. You and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 4 months after stopping treatment. This can be discussed with your doctor.
- if you are a **man** who is taking ribavirin, do not have sex with a pregnant woman unless you use a condom. This will lessen the chance for ribavirin to be left in the woman's body. If your female partner is not pregnant now but is of childbearing age, she must be tested for pregnancy each month during treatment and for the 7 months after treatment has stopped. This can be discussed with your doctor. If you are a male patient, you and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 7 months after stopping treatment. This can be discussed with your doctor.

It is not known whether this product is present in human milk. Therefore, do not breast-feed an infant if you are taking PegIntron. In combination therapy with ribavirin, take notice of the respective informing texts of ribavirin containing medicinal products.

### **Driving and using machines**

Do not drive or operate any tools or machines if you feel tired, sleepy or confused while taking PegIntron.

### **Important information about some of the ingredients of PegIntron**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

This medicinal product contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### **Patients who also have HIV infection**

Lactic acidosis and worsening liver function are side effects associated with Highly Active Anti-Retroviral Therapy (HAART), an HIV treatment. If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis and of liver failure. Your doctor will monitor you for signs and symptoms of these conditions. (Please be sure to read the ribavirin Patient Leaflet also). Additionally, patients treated with PegIntron and ribavirin combination therapy and zidovudine could be at increased risk of developing anaemia (low number of red blood cells). If you are also being treated for Human Immunodeficiency Virus (HIV) infection (AIDS) with zidovudine or stavudine, it is not certain if ribavirin will change the way these medicines work. Therefore, your blood will be checked regularly to be sure that the HIV infection is not getting worse. If it gets worse, your doctor will decide whether or not your ribavirin treatment needs to be changed.

Co-administration of ribavirin and didanosine and/or stavudine is not recommended due to the risk of lactic acidosis (a build-up of lactic acid in the body) and pancreatitis.

## **3. HOW TO USE PEGINTRON**

Your doctor has prescribed PegIntron specifically for you and your current condition; do not share this medicine with anyone else.

Your doctor has determined your dose of PegIntron based on your weight. If necessary, the dose may be changed during treatment.

### **Combination treatment**

PegIntron, when given with ribavirin capsules, is usually given at a dose of 1.5 microgram/kg once a week.

Ribavirin capsules are taken every day, morning and evening. The number of ribavirin capsules you take depends on your weight.

- If you weigh less than 65 kg, take 2 capsules in the morning and 2 in the evening (total of 800 mg each day).
- If you weigh between 65 and 85 kg, take 2 capsules in the morning and 3 in the evening (total of 1,000 mg each day).
- If you weigh more than 85 kg, take 3 capsules in the morning and 3 in the evening (total 1,200 mg each day).

The combination treatment is continued for 3 to 6 months, and sometimes for one year depending on your physicians judgement. Take notice of the respective informing texts of ribavirin containing medicinal products.

For HCV/HIV co-infected patients, the duration of treatment is 48 weeks.

### **PegIntron alone:**

PegIntron, when given alone, is usually given at a dose of 0.5 or 1.0 microgram/kg once a week, for at least 6 months, and possibly for 1 year. If you have kidney disease, your dose may be lower, depending upon your kidney function.

PegIntron alone was not studied in HCV/HIV co-infected patients.



**All patients:**

If you are injecting PegIntron yourself, please be sure that the dose that has been prescribed for you is clearly provided on the package of medicine you receive.

If you have the impression that the effect of PegIntron is too strong or too weak, talk to your doctor or pharmacist.

PegIntron is intended for subcutaneous use. This means that it is injected through a short injection needle into the fatty tissue just under your skin. If you are injecting this medicine yourself, you will be instructed how to prepare and give the injection. Detailed instructions for subcutaneous administration are provided with this leaflet (see How to use the PegIntron pre-filled pen).

Prepare the dose just before you intend to inject it and use it immediately. Look carefully at the reconstituted solution prior to administration. Do not use if there is discolouration of the reconstituted solution or if particulate matter is present. PegIntron is intended for single use only. Thus, after you give yourself the injection, discard the PegIntron pre-filled pen with any solution that is left in it.

Inject PegIntron once each week on the same day. Injecting it at the same time of day each week will help you not to forget to take it.

Always use PegIntron exactly as your doctor told you. Do not exceed the recommended dosage, and take it for as long as prescribed.

**If you use more PegIntron than you should**

Tell your doctor or healthcare professional as soon as possible.

**If you forget to take PegIntron**

Take the dose as soon as you remember, then continue your treatment as usual.

Do not take a double dose to make up for a forgotten dose. Contact your doctor or pharmacist if needed.

**4. POSSIBLE SIDE EFFECTS**

Like all medicines, PegIntron can cause side effects, although not everybody gets them. Although not all of these side effects may occur, they may need medical attention if they do.

Some people get depressed when taking PegIntron alone or in combination treatment with ribavirin, and in some cases people had suicidal thoughts or aggressive behaviour (sometimes directed against others). Some patients have actually committed suicide. Be sure to seek emergency care if you notice that you are becoming depressed or have suicidal thoughts or change in your behaviour. You may want to consider asking a family member or close friend to help you stay alert to signs of depression or changes in your behaviour.

Check with your doctor immediately if any of the following side effects occur: chest pain; changes in the way your heart beats; breathing problems (including shortness of breath), confusion; feeling depressed, wanting to harm yourself, hallucinations, numbness or tingling feeling; dizziness, convulsion ("fit"); trouble sleeping, thinking or concentrating; difficulty remaining alert, severe stomach pain or cramps; blood or clots in stool (or black, tarry stool); fever or chills beginning after a few weeks of treatment, pain in your lower back or side; difficulty or inability to pass urine, painful or inflamed muscles (sometimes severe); problems with your eyes or your eyesight or hearing; severe or painful reddening of your skin or mucous membrane, severe bleeding from your nose. Your doctor will test your blood to ensure that your white blood cell (cells that fight infection) and red blood cell (cells that carry oxygen) counts, platelets (blood clotting cells) and other laboratory values are at acceptable levels.

Other side effects that may occur with the combination of PegIntron and ribavirin capsules include:

Very commonly reported side effects (greater than 1 in every 10 patients):

Irritation or redness (and rarely, skin damage) at the site of injection, headache, tired feeling, shaking chills, fever, flu-like symptoms, weakness, loss of weight, nausea, loss of appetite, diarrhoea or loose stools, stomach pain, vomiting, muscle aches, pain in joints and muscles, feeling depressed, irritability, trouble falling asleep or staying asleep, feeling anxious or nervous, difficulty concentrating, mood swings, hair loss, itching, dry skin, sore throat, coughing, difficult breathing, dizziness, virus infection, rash, and dry mouth.

Commonly reported side effects (at least 1 in every 100 patients, but not greater than 1 in every 10 patients):

Increased sweating, chest pain, pain on the right side around your ribs, numbness, pain or tingling feeling, change in thyroid gland activity (which may make you feel tired or, less commonly, energetic), stomach upset, rapid heart rate, agitation, nervousness, difficult or irregular menstrual period

Less common are pain at the place of injection, flushing, low or high blood pressure, dry or teary eyes, redness of skin or skin disorder, psoriasis, hives, nail disorder, feeling unwell, feeling faint, poor coordination, confusion, increased or decreased sensitivity to touch, tense muscles, arthritis, bruising, loss of interest in activities including sex, sexual problem, unusual dreams, shaky hands, vertigo (spinning feeling), increased appetite, heartburn, intestinal gas (flatus), constipation, hemorrhoids, red or bleeding gums, redness or sores in mouth, change in taste, changes in hearing or ringing in ears, thirst, changed behaviour or aggressive behaviour (sometimes directed against others), feeling sleepy, cold sores, fungal or bacterial infections, irritation of prostate gland, increased need to pass urine, ear or respiratory infections, sinusitis, stuffy or runny nose, abnormal hair texture, sensitivity to sunlight, migraine headache, eye pain or infection, blurred vision, puffy face, puffy hands or feet, enlarged liver, problem affecting ovary or vagina, pain in breast, difficulty in speaking and swollen glands.

The following rare and very rare events have been reported with PegIntron:

Rarely reported side effects (greater than 1 in every 10,000 patients, but less than 1 in every 1,000 patients):

Diabetes, abnormal heart rhythm, inflammation of pancreas, inflammation and degeneration of muscle tissue and peripheral nerves, kidney problems and seizures.

Very rarely reported side effects (less than 1 in every 10,000 patients):

Very rarely sarcoidosis (a disease characterised by persistent fever, weight loss, joint pain and swelling, skin lesions and swollen glands) has been reported. Loss of consciousness has occurred very rarely with alpha interferons, mostly in elderly patients treated at high doses. Cases of stroke (cerebrovascular events) have been reported. Check with your doctor immediately if you have any of these symptoms or any other symptoms that are troubling.

Very rarely, PegIntron alone or in combination with ribavirin may cause aplastic anaemia. Pure red cell aplasia, a condition where the body stopped or reduced the production of red blood cells, has been reported. This causes severe anaemia, symptoms of which would include unusual tiredness and a lack of energy.

When PegIntron is used alone, some of these effects are less likely to occur, and some have not occurred at all.

If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis, liver failure, and development of blood abnormalities (reduction in number of red blood cells which carry oxygen, certain white blood cells that fight infection, and blood clotting cells called platelets).

The following other side effects (not listed above) have occurred with the combination of PegIntron and ribavirin capsules (adults) in HCV/HIV co-infected patients receiving HAART:

oral candidiasis (oral thrush), defective metabolism of fat, CD4 lymphocytes decreased, appetite decreased,

back pain, hepatitis, limb pain, and various laboratory blood values abnormalities.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE PEGINTRON**

Keep out of the reach and sight of children.

Store in a refrigerator (2°C - 8°C). Do not freeze.

Do not use PegIntron after the expiry date which is stated on the carton.

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

Do not use PegIntron if you notice discolouration of the powder.

The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. After administering the dose, discard the PegIntron pre-filled pen and any unused solution contained in it.

## **6. FURTHER INFORMATION**

### **What PegIntron contains**

- The active substance is peginterferon alfa-2b, 120 micrograms/0.5 ml.
- The other ingredients are:  
Powder: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80;  
Solvent: water for injections

### **What PegIntron looks like and contents of the pack**

The pharmaceutical form is: powder and solvent for solution for injection in a pre-filled pen. The white powder and the clear and colourless solvent are both contained in a two-chamber glass cartridge assembled into a single use pre-filled pen.

PegIntron 120 micrograms is available in different pack sizes:

- 1 pen containing powder and solvent for solution for injection, 1 injection needle and 2 cleansing swabs;
- 4 pens containing powder and solvent for solution for injection, 4 injection needles and 8 cleansing swabs;
- 6 pens containing powder and solvent for solution for injection, 6 injection needles and 12 cleansing swabs;
- 12 pens containing powder and solvent for solution for injection, 12 injection needles and 24 cleansing swabs.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

Marketing Authorisation Holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

Manufacturer: SP (Brinny) Company, Innishannon, County Cork, Ireland

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Rue de Stalle/Stallestraat 73  
B-1180 Bruxelles/Brussel/Brüssel  
Tél/Tel: + 32-(0)2 370 92 11

**Luxembourg/Luxemburg**

Rue de Stalle 73  
B-1180 Bruxelles/Brüssel  
Belgique/Belgien  
Tél/Tel: + 32-(0)2 370 92 11

**България**

ул. "Проф. Милко Бичев" № 7, ет.3  
BG-София 1504  
Тел.: +359 2 944 1073

**Magyarország**

Alkotás u. 53.  
H-1123 Budapest  
Tel.: +36 1 457-8500

**Česká republika**

Na Příkopě 25  
CZ-110 00 Praha 1  
Tel: +420 221771250

**Malta**

168 Christopher Street  
MT-VLT02 Valletta  
Tel: + 356-21 23 21 75

**Danmark**

Lautrupbjerg 2  
DK-2750 Ballerup  
Tlf: + 45-44 39 50 00

**Nederland**

Maarssenbroeksedijk 4  
NL-3542 DN Utrecht  
Tel: + 31-(0)800 778 78 78

**Deutschland**

Thomas-Dehler-Straße 27  
D-81737 München  
Tel: + 49-(0)89 627 31-0

**Norge**

Pb. 398  
N-1326 Lysaker  
Tlf: + 47 67 16 64 50

**Eesti**

Järvevana tee 9  
EE-11314 Tallinn  
Tel: + 372 654 96 86

**Österreich**

Badener Strasse 23  
A-2514 Traiskirchen  
Tel: + 43-(0)2252 502-0

**Ελλάδα**

Αγίου Δημητρίου 63  
GR-174 55 Αλιμος  
Τηλ.: + 30-210 98 97 300

**Polska**

Al. Jerozolimskie 195a  
PL-02-222 Warszawa  
Tel.: + 48-(0)22 478 41 50

**España**

Km. 36, Ctra. Nacional I  
E-28750 San Agustín de Guadalix – Madrid  
Tel: + 34-91 848 85 00

**Portugal**

Rua Agualva dos Açores 16  
P-2735-557 Agualva-Cacém  
Tel: +351-21 433 93 00

**France**

92 rue Baudin  
F-92300 Levallois-Perret  
Tél: + 33-(0)1 41 06 35 00

**România**

Șos. București-Ploiești, nr. 17-21, Băneasa  
Center, et. 7, sector 1  
RO-013682 București  
Tel: + 40 21 233 35 30

**Ireland**

Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW  
Tel: +44-(0)1 707 363 636

**Slovenija**

Dunajska 22  
SI-1000 Ljubljana  
Tel: + 386 01 3001070

**Ísland**

Hörgatún 2  
IS-210 Garðabær  
Sími: + 354 535 70 00

**Italia**

Via fratelli Cervi snc,  
Centro Direzionale Milano Due  
Palazzo Borromini  
I-20090 Segrate (Milano)  
Tel: + 39-02 21018.1

**Κύπρος**

Οδός Αγίου Νικολάου, 8  
CY-1055 Λευκωσία  
Τηλ: +357-22 757188

**Latvija**

Bauskas 58a -401  
Rīga, LV-1004  
Tel: + 371-7 21 38 25

**Lietuva**

Kęstučio g. 65/40  
LT-08124 Vilnius  
Tel: + 370 52 101868

**Slovenská republika**

Strakova 5  
SK-811 01 Bratislava  
Tel: + 421 (2) 5920 2712

**Suomi/Finland**

PL 3/PB 3  
FIN-02201 Espoo/Esbo  
Puh/Tel: + 358-(0)20-7570 300

**Sverige**

Box 27190  
S-102 52 Stockholm  
Tel: + 46-(0)8 522 21 500

**United Kingdom**



Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW - UK  
Tel: + 44-(0)1 707 363 636

**This leaflet was last approved on**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>

## ANNEX TO THE PACKAGE LEAFLET

### How to use the PegIntron pre-filled pen



The following instructions explain how to inject yourself with the single use PegIntron pre-filled pen. **Please read the instructions carefully and completely before attempting to use the pen and follow them step by step.** Your doctor or his/her assistant will instruct you on how to self-inject with the PegIntron pre-filled pen. Do not attempt to inject yourself unless you are sure you understand the procedure and requirements for self-injection.


The PegIntron pre-filled pen is intended for use by one person only and must not be shared. Use the injection needle and cleansing swabs provided in the packaging only for the PegIntron pre-filled pen. Be sure the solution is at room temperature at the time of injection. Your doctor will have told you what dose you require for your therapy.

**Note:** The colour of the dosing button is different for each strength of the PegIntron pre-filled pen.

#### Step 1: Mixing

**It is important that you keep the PegIntron pre-filled pen upright (as shown in figure 1) during mixing, unless otherwise instructed.**

- Take the PegIntron pre-filled pen out of the refrigerator. Allow the medicine to come to room temperature.



- Wash your hands with soap and water.
- **Place the PegIntron pre-filled pen upright** in the holder of the tray provided in the pack (the dosing button will be on the **bottom**).
- You may want to hold the pre-filled pen using the grip. To mix the powder and the liquid, press the two halves together firmly by pressing down until you hear the pre-filled pen click. The two stoppers should come together.
- Wait for several seconds until the powder is completely dissolved.
- **Gently turn the PegIntron pre-filled pen upside down twice. To avoid excessive foaming, do not shake.**
- Maintaining the PegIntron pre-filled pen upright, check the mixed PegIntron solution through the window. If there is still foam, wait until it settles. The solution should be clear and colourless. **Do not use the pen if the solution is discoloured or contains any particles.**



Figure 2

- Keeping the PegIntron pre-filled pen upright in the holder provided in the packaging, disinfect the rubber membrane of the PegIntron pre-filled pen with one cleansing swab.
- Take the injection needle provided in the tray and remove its protective paper tab.
- **Maintaining the PegIntron pre-filled pen upright in the holder**, gently push the injection needle onto the pre-filled pen and **screw it securely in place**.

- **Keep the PegIntron pre-filled pen pointed up.**
- **Do not take off the outer needle cap at this point.**
- You may see some liquid trickle out from under the cap, as the air has been expelled out of the pen.
- Wait about 5 seconds for this process to finish.




Figure 3

- **Check through the window to be sure that the two stoppers are together. If they are not together, do not use this pen because you may not be able to dial your dose.**

### Step 2: Setting the dose



Figure 4

- Remove the PegIntron pre-filled pen from the holder.
- Holding the PegIntron pre-filled pen firmly, pull the dosing button out as far as it will go, until you see a **dark ring** on the pen. The dosing button should be easy to pull out without excessive force being required.

**Note: Do not push the dosing button back in at this time. You will push it in when you are ready to self-inject the PegIntron.**



Figure 5

- Turn the dosing button until your prescribed dose is aligned with the dosing tab. The button should turn easily without excessive force being required.

**Note:** **If you cannot easily pull out the dosing button or dial the dose, do not use excessive force and do not use this pen because it may not deliver the correct dose.**

### **Step 3: Injecting the solution**

- Select the injection site. Your doctor will have told you which sites to use (e.g., thigh or abdomen).

**Note:** change your injection site each time.

- Clean the injection site skin with the second cleansing swab.




Figure 6

- Pull off the **outer needle cap**.
- There may be some liquid around the inner needle cap. This liquid is not part of your dose, this is extra. This is normal, as the air has been expelled out of the needle.
- Once the injection site is dry, pull off the **yellow inner needle cap** carefully exposing the injection needle.



Figure 7

- Hold the PegIntron pre-filled pen with your fingers wrapped around the barrel and your thumb on the dosing button.
- With your other hand, pinch a fold of loose skin.
- Insert the needle into the pinched skin at an angle of 45° to 90°.
- Press the dosing button down **slowly** and **firmly** until the button can no longer move.
- **Maintain pressure on the dosing button for an additional 5 seconds** to ensure that you get the complete dose.



- Remove the needle from your skin.
- Press the injection site with a small bandage or sterile gauze if necessary for a few seconds.
- Do not massage the injection site. If there is bleeding, cover with an adhesive bandage.
- Discard the PegIntron pre-filled pen with the needle safely in a closed rigid container.

## PACKAGE LEAFLET: INFORMATION FOR THE USER

**PegIntron 150 micrograms powder and solvent for solution for injection in pre-filled pen**  
Peginterferon alfa-2b (conjugation of recombinant interferon alfa-2b with monomethoxy polyethylene glycol)

**Read all of this leaflet carefully before you start using this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What PegIntron is and what it is used for
2. Before you use PegIntron
3. How to use PegIntron
4. Possible side effects
5. Storing PegIntron
6. Further information

### **1. WHAT PEGINTRON IS AND WHAT IT IS USED FOR**

Interferons modify the response of the body's immune system to help fight infections and severe diseases. PegIntron, which contains an interferon, is used for the treatment of chronic hepatitis C, a viral infection of the liver, including in first time users co-infected with clinically stable HIV.

PegIntron is best used for this treatment in combination with ribavirin.

PegIntron is used alone in case of intolerance or contraindication to ribavirin.

### **2. BEFORE YOU USE PEGINTRON**

PegIntron is not recommended for use in patients under the age of 18 years.

**Do not use PegIntron**

- If you are allergic (hypersensitive) to peginterferon alfa-2b or any of the other ingredients of PegIntron.
- If you are allergic (hypersensitive) to any interferon.
- If you have had severe heart problems, or if you have heart disease that has not been well controlled during the past 6 months.
- If you have severe medical conditions that leave you very weak.
- If you have autoimmune hepatitis or any other problem with your immune system; if you are taking medicine that suppresses your immune system (your immune system protects you against infection and some diseases).
- If you have advanced, uncontrolled liver disease (other than hepatitis C).
- If you have thyroid disease that is not well controlled with medicines.
- If you have a condition that causes convulsions (seizures, or "fits").

**Take special care with PegIntron**

- If you develop symptoms of a severe allergic reaction (such as difficulty in breathing, wheezing, or hives) while on this medication, seek medical help immediately.
- If you ever had a heart attack or a heart problem.
- If you have ever been treated for depression or any other nervous or mental disorder.

- If you have kidney disease, your doctor may prescribe a lower than usual dose and monitor your kidney blood values regularly during treatment. If you have a kidney disease and you are using PegIntron in combination with medicinal products containing ribavirin, your doctor should monitor you more carefully for a decrease in red blood cell count.
- If you have had a problem with your liver (other than hepatitis C).
- If you develop symptoms associated with a cold or other respiratory infection, such as fever, cough, or any difficulty in breathing, tell your doctor.
- If you are diabetic, your doctor may ask you to have an eye examination.
- If you have had any serious illness affecting your breathing or your blood.
- If you have psoriasis, it may become worse while you are using PegIntron.
- If you are planning to become pregnant, discuss this with your doctor before starting to use PegIntron.
- If you are also being treated for HIV, please see **Using other medicines**.
- If you have had a severe nervous or mental disorder.
- If you ever had depression or develop symptoms associated with depression (e.g. feelings of sadness, dejection, etc) while on treatment with PegIntron (see section 4).
- If you have received an organ transplant, either kidney or liver, interferon treatment may increase the risk of rejection. Be sure to discuss this with your doctor.

Dental and gum disorders, which may lead to loss of teeth, have been reported in patients receiving PegIntron and ribavirin combination therapy. In addition, dry mouth could have a damaging effect on teeth and membranes of the mouth during long-term treatment with the combination of PegIntron with ribavirin. You should brush your teeth thoroughly twice daily and have regular dental examinations. In addition some patients may experience vomiting. If you have this reaction, be sure to rinse your mouth thoroughly afterwards.

### **While being treated with PegIntron**

Your doctor may want you to drink extra fluids to help prevent low blood pressure.

### **Using other medicines**

Please inform your doctor or pharmacist if you are taking or have recently taken any other medicines, even those not prescribed.

### **Pregnancy and breast-feeding**

Ask your doctor or pharmacist for advice before taking any medicine. In studies in pregnant animals, interferons have sometimes caused miscarriage. The effect on human pregnancy is not known. In combination therapy with ribavirin, ribavirin can be very damaging to an unborn baby, thus both female and male patients must take special precautions in their sexual activity if there is any chance for pregnancy to occur:

- if you are a **girl** or a **woman** of childbearing age, you must have a negative pregnancy test before treatment, each month during treatment, and for the 4 months after treatment is stopped. You and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 4 months after stopping treatment. This can be discussed with your doctor.
- if you are a **man** who is taking ribavirin, do not have sex with a pregnant woman unless you use a condom. This will lessen the chance for ribavirin to be left in the woman's body. If your female partner is not pregnant now but is of childbearing age, she must be tested for pregnancy each month during treatment and for the 7 months after treatment has stopped. This can be discussed with your doctor. If you are a male patient, you and your partner must each use an effective contraceptive during the time you are taking ribavirin and for 7 months after stopping treatment. This can be discussed with your doctor.

It is not known whether this product is present in human milk. Therefore, do not breast-feed an infant if you are taking PegIntron. In combination therapy with ribavirin, take notice of the respective informing texts of ribavirin containing medicinal products.

### **Driving and using machines**

Do not drive or operate any tools or machines if you feel tired, sleepy or confused while taking PegIntron.

### **Important information about some of the ingredients of PegIntron**

This medicinal product contains less than 1 mmol sodium (23 mg) per 0.7 ml, i.e., essentially "sodium-free".

This medicinal product contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### **Patients who also have HIV infection**

Lactic acidosis and worsening liver function are side effects associated with Highly Active Anti-Retroviral Therapy (HAART), an HIV treatment. If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis and of liver failure. Your doctor will monitor you for signs and symptoms of these conditions. (Please be sure to read the ribavirin Patient Leaflet also). Additionally, patients treated with PegIntron and ribavirin combination therapy and zidovudine could be at increased risk of developing anaemia (low number of red blood cells). If you are also being treated for Human Immunodeficiency Virus (HIV) infection (AIDS) with zidovudine or stavudine, it is not certain if ribavirin will change the way these medicines work. Therefore, your blood will be checked regularly to be sure that the HIV infection is not getting worse. If it gets worse, your doctor will decide whether or not your ribavirin treatment needs to be changed.

Co-administration of ribavirin and didanosine and/or stavudine is not recommended due to the risk of lactic acidosis (a build-up of lactic acid in the body) and pancreatitis.

## **3. HOW TO USE PEGINTRON**

Your doctor has prescribed PegIntron specifically for you and your current condition; do not share this medicine with anyone else.

Your doctor has determined your dose of PegIntron based on your weight. If necessary, the dose may be changed during treatment.

### **Combination treatment**

PegIntron, when given with ribavirin capsules, is usually given at a dose of 1.5 microgram/kg once a week.

Ribavirin capsules are taken every day, morning and evening. The number of ribavirin capsules you take depends on your weight.

- If you weigh less than 65 kg, take 2 capsules in the morning and 2 in the evening (total of 800 mg each day).
- If you weigh between 65 and 85 kg, take 2 capsules in the morning and 3 in the evening (total of 1,000 mg each day).
- If you weigh more than 85 kg, take 3 capsules in the morning and 3 in the evening (total 1,200 mg each day).

The combination treatment is continued for 3 to 6 months, and sometimes for one year depending on your physicians judgement. Take notice of the respective informing texts of ribavirin containing medicinal products.

For HCV/HIV co-infected patients, the duration of treatment is 48 weeks.

### **PegIntron alone:**

PegIntron, when given alone, is usually given at a dose of 0.5 or 1.0 microgram/kg once a week, for at least 6 months, and possibly for 1 year. If you have kidney disease, your dose may be lower, depending upon your kidney function.

PegIntron alone was not studied in HCV/HIV co-infected patients.

**All patients:**

If you are injecting PegIntron yourself, please be sure that the dose that has been prescribed for you is clearly provided on the package of medicine you receive.

If you have the impression that the effect of PegIntron is too strong or too weak, talk to your doctor or pharmacist.

PegIntron is intended for subcutaneous use. This means that it is injected through a short injection needle into the fatty tissue just under your skin. If you are injecting this medicine yourself, you will be instructed how to prepare and give the injection. Detailed instructions for subcutaneous administration are provided with this leaflet (see How to use the PegIntron pre-filled pen).

Prepare the dose just before you intend to inject it and use it immediately. Look carefully at the reconstituted solution prior to administration. Do not use if there is discolouration of the reconstituted solution or if particulate matter is present. PegIntron is intended for single use only. Thus, after you give yourself the injection, discard the PegIntron pre-filled pen with any solution that is left in it.

Inject PegIntron once each week on the same day. Injecting it at the same time of day each week will help you not to forget to take it.

Always use PegIntron exactly as your doctor told you. Do not exceed the recommended dosage, and take it for as long as prescribed.

**If you use more PegIntron than you should**

Tell your doctor or healthcare professional as soon as possible.

**If you forget to take PegIntron**

Take the dose as soon as you remember, then continue your treatment as usual.

Do not take a double dose to make up for a forgotten dose. Contact your doctor or pharmacist if needed.

**4. POSSIBLE SIDE EFFECTS**

Like all medicines, PegIntron can cause side effects, although not everybody gets them. Although not all of these side effects may occur, they may need medical attention if they do.

Some people get depressed when taking PegIntron alone or in combination treatment with ribavirin, and in some cases people had suicidal thoughts or aggressive behaviour (sometimes directed against others). Some patients have actually committed suicide. Be sure to seek emergency care if you notice that you are becoming depressed or have suicidal thoughts or change in your behaviour. You may want to consider asking a family member or close friend to help you stay alert to signs of depression or changes in your behaviour.

Check with your doctor immediately if any of the following side effects occur: chest pain; changes in the way your heart beats; breathing problems (including shortness of breath), confusion; feeling depressed, wanting to harm yourself, hallucinations, numbness or tingling feeling; dizziness, convulsion ("fit"); trouble sleeping, thinking or concentrating; difficulty remaining alert, severe stomach pain or cramps; blood or clots in stool (or black, tarry stool); fever or chills beginning after a few weeks of treatment, pain in your lower back or side; difficulty or inability to pass urine, painful or inflamed muscles (sometimes severe); problems with your eyes or your eyesight or hearing; severe or painful reddening of your skin or mucous membrane, severe bleeding from your nose. Your doctor will test your blood to ensure that your white blood cell (cells that fight infection) and red blood cell (cells that carry oxygen) counts, platelets (blood clotting cells) and other laboratory values are at acceptable levels.

Other side effects that may occur with the combination of PegIntron and ribavirin capsules include:

Very commonly reported side effects (greater than 1 in every 10 patients):

Irritation or redness (and rarely, skin damage) at the site of injection, headache, tired feeling, shaking chills, fever, flu-like symptoms, weakness, loss of weight, nausea, loss of appetite, diarrhoea or loose stools, stomach pain, vomiting, muscle aches, pain in joints and muscles, feeling depressed, irritability, trouble falling asleep or staying asleep, feeling anxious or nervous, difficulty concentrating, mood swings, hair loss, itching, dry skin, sore throat, coughing, difficult breathing, dizziness, virus infection, rash, and dry mouth.

Commonly reported side effects (at least 1 in every 100 patients, but not greater than 1 in every 10 patients):

Increased sweating, chest pain, pain on the right side around your ribs, numbness, pain or tingling feeling, change in thyroid gland activity (which may make you feel tired or, less commonly, energetic), stomach upset, rapid heart rate, agitation, nervousness, difficult or irregular menstrual period

Less common are pain at the place of injection, flushing, low or high blood pressure, dry or teary eyes, redness of skin or skin disorder, psoriasis, hives, nail disorder, feeling unwell, feeling faint, poor coordination, confusion, increased or decreased sensitivity to touch, tense muscles, arthritis, bruising, loss of interest in activities including sex, sexual problem, unusual dreams, shaky hands, vertigo (spinning feeling), increased appetite, heartburn, intestinal gas (flatus), constipation, hemorrhoids, red or bleeding gums, redness or sores in mouth, change in taste, changes in hearing or ringing in ears, thirst, changed behaviour or aggressive behaviour (sometimes directed against others), feeling sleepy, cold sores, fungal or bacterial infections, irritation of prostate gland, increased need to pass urine, ear or respiratory infections, sinusitis, stuffy or runny nose, abnormal hair texture, sensitivity to sunlight, migraine headache, eye pain or infection, blurred vision, puffy face, puffy hands or feet, enlarged liver, problem affecting ovary or vagina, pain in breast, difficulty in speaking and swollen glands.

The following rare and very rare events have been reported with PegIntron:

Rarely reported side effects (greater than 1 in every 10,000 patients, but less than 1 in every 1,000 patients):

Diabetes, abnormal heart rhythm, inflammation of pancreas, inflammation and degeneration of muscle tissue and peripheral nerves, kidney problems and seizures.

Very rarely reported side effects (less than 1 in every 10,000 patients):

Very rarely sarcoidosis (a disease characterised by persistent fever, weight loss, joint pain and swelling, skin lesions and swollen glands) has been reported. Loss of consciousness has occurred very rarely with alpha interferons, mostly in elderly patients treated at high doses. Cases of stroke (cerebrovascular events) have been reported. Check with your doctor immediately if you have any of these symptoms or any other symptoms that are troubling.

Very rarely, PegIntron alone or in combination with ribavirin may cause aplastic anaemia. Pure red cell aplasia, a condition where the body stopped or reduced the production of red blood cells, has been reported. This causes severe anaemia, symptoms of which would include unusual tiredness and a lack of energy.

When PegIntron is used alone, some of these effects are less likely to occur, and some have not occurred at all.

If you are receiving HAART, the addition of PegIntron and ribavirin may increase your risk of lactic acidosis, liver failure, and development of blood abnormalities (reduction in number of red blood cells which carry oxygen, certain white blood cells that fight infection, and blood clotting cells called platelets).

The following other side effects (not listed above) have occurred with the combination of PegIntron and ribavirin capsules (adults) in HCV/HIV co-infected patients receiving HAART: oral candidiasis (oral thrush), defective metabolism of fat, CD4 lymphocytes decreased, appetite decreased, back pain, hepatitis, limb pain, and various laboratory blood values abnormalities.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE PEGINTRON**

Keep out of the reach and sight of children.

Store in a refrigerator (2°C - 8°C). Do not freeze.

Do not use PegIntron after the expiry date which is stated on the carton.

After reconstitution, use the reconstituted solution immediately or within 24 hours when stored in a refrigerator (2°C - 8°C).

Do not use PegIntron if you notice discolouration of the powder.

The reconstituted solution should be clear and colourless. Do not use if discolouration or particulate matter is present. After administering the dose, discard the PegIntron pre-filled pen and any unused solution contained in it.

## **6. FURTHER INFORMATION**

### **What PegIntron contains**

- The active substance is peginterferon alfa-2b, 150 micrograms/0.5 ml.
- The other ingredients are:  
Powder: disodium phosphate, anhydrous; sodium dihydrogen phosphate dihydrate, sucrose and polysorbate 80;  
Solvent: water for injections

### **What PegIntron looks like and contents of the pack**

The pharmaceutical form is: powder and solvent for solution for injection in a pre-filled pen. The white powder and the clear and colourless solvent are both contained in a two-chamber glass cartridge assembled into a single use pre-filled pen.

PegIntron 150 micrograms is available in different pack sizes:

- 1 pen containing powder and solvent for solution for injection, 1 injection needle and 2 cleansing swabs;
- 4 pens containing powder and solvent for solution for injection, 4 injection needles and 8 cleansing swabs;
- 6 pens containing powder and solvent for solution for injection, 6 injection needles and 12 cleansing swabs;
- 12 pens containing powder and solvent for solution for injection, 12 injection needles and 24 cleansing swabs.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

Marketing Authorisation Holder: SP Europe, 73, rue de Stalle, B-1180 Bruxelles, Belgium

Manufacturer: SP (Brinny) Company, Innishannon, County Cork, Ireland

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Rue de Stalle/Stallestraat 73  
B-1180 Bruxelles/Brussel/Brüssel  
Tél/Tel: + 32-(0)2 370 92 11

**Luxembourg/Luxemburg**

Rue de Stalle 73  
B-1180 Bruxelles/Brüssel  
Belgique/Belgien  
Tél/Tel: + 32-(0)2 370 92 11

**България**

ул. "Проф. Милко Бичев" № 7, ет.3  
BG-София 1504  
Тел.: +359 2 944 1073

**Magyarország**

Alkotás u. 53.  
H-1123 Budapest  
Tel.: +36 1 457-8500

**Česká republika**

Na Příkopě 25  
CZ-110 00 Praha 1  
Tel: +420 221771250

**Malta**

168 Christopher Street  
MT-VLT02 Valletta  
Tel: + 356-21 23 21 75

**Danmark**

Lautrupbjerg 2  
DK-2750 Ballerup  
Tlf: + 45-44 39 50 00

**Nederland**

Maarssenbroeksedijk 4  
NL-3542 DN Utrecht  
Tel: + 31-(0)800 778 78 78

**Deutschland**

Thomas-Dehler-Straße 27  
D-81737 München  
Tel: + 49-(0)89 627 31-0

**Norge**

Pb. 398  
N-1326 Lysaker  
Tlf: + 47 67 16 64 50

**Eesti**

Järvevana tee 9  
EE-11314 Tallinn  
Tel: + 372 654 96 86

**Österreich**

Badener Strasse 23  
A-2514 Traiskirchen  
Tel: + 43-(0)2252 502-0

**Ελλάδα**

Αγίου Δημητρίου 63  
GR-174 55 Αλιμος  
Τηλ.: + 30-210 98 97 300

**Polska**

Al. Jerozolimskie 195a  
PL-02-222 Warszawa  
Tel.: + 48-(0)22 478 41 50

**España**

Km. 36, Ctra. Nacional I  
E-28750 San Agustín de Guadalix – Madrid  
Tel: + 34-91 848 85 00

**Portugal**

Rua Agualva dos Açores 16  
P-2735-557 Agualva-Cacém  
Tel: +351-21 433 93 00

**France**

92 rue Baudin  
F-92300 Levallois-Perret  
Tél: + 33-(0)1 41 06 35 00

**România**

Șos. București-Ploiești, nr. 17-21, Băneasa  
Center, et. 7, sector 1  
RO-013682 București  
Tel: + 40 21 233 35 30

**Ireland**

Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW  
Tel: +44-(0)1 707 363 636

**Slovenija**

Dunajska 22  
SI-1000 Ljubljana  
Tel: + 386 01 3001070

**Ísland**

Hörgatún 2  
IS-210 Garðabær  
Sími: + 354 535 70 00

**Slovenská republika**

Strakova 5  
SK-811 01 Bratislava  
Tel: + 421 (2) 5920 2712



**Italia**

Via fratelli Cervi snc,  
Centro Direzionale Milano Due  
Palazzo Borromini  
I-20090 Segrate (Milano)  
Tel: + 39-02 21018.1

**Κύπρος**

Οδός Αγίου Νικολάου, 8  
CY-1055 Λευκωσία  
Τηλ: +357-22 757188

**Latvija**

Bauskas 58a -401  
Rīga, LV-1004  
Tel: + 371-7 21 38 25

**Lietuva**

Kęstučio g. 65/40  
LT-08124 Vilnius  
Tel: + 370 52 101868

**Suomi/Finland**

PL 3/PB 3  
FIN-02201 Espoo/Esbo  
Puh/Tel: + 358-(0)20-7570 300

**Sverige**

Box 27190  
S-102 52 Stockholm  
Tel: + 46-(0)8 522 21 500

**United Kingdom**



Shire Park  
Welwyn Garden City  
Hertfordshire AL7 1TW - UK  
Tel: + 44-(0)1 707 363 636

**This leaflet was last approved on**

Detailed information on this product is available on the web-site of the European Medicines Agency (EMA) <http://www.ema.europa.eu/>

## ANNEX TO THE PACKAGE LEAFLET

### How to use the PegIntron pre-filled pen



The following instructions explain how to inject yourself with the single use PegIntron pre-filled pen. **Please read the instructions carefully and completely before attempting to use the pen and follow them step by step.** Your doctor or his/her assistant will instruct you on how to self-inject with the PegIntron pre-filled pen. Do not attempt to inject yourself unless you are sure you understand the procedure and requirements for self-injection.


The PegIntron pre-filled pen is intended for use by one person only and must not be shared. Use the injection needle and cleansing swabs provided in the packaging only for the PegIntron pre-filled pen. Be sure the solution is at room temperature at the time of injection. Your doctor will have told you what dose you require for your therapy.

**Note:** The colour of the dosing button is different for each strength of the PegIntron pre-filled pen.

#### Step 1: Mixing

**It is important that you keep the PegIntron pre-filled pen upright (as shown in figure 1) during mixing, unless otherwise instructed.**

- Take the PegIntron pre-filled pen out of the refrigerator. Allow the medicine to come to room temperature.



- Wash your hands with soap and water.
- **Place the PegIntron pre-filled pen upright** in the holder of the tray provided in the pack (the dosing button will be on the **bottom**).
- You may want to hold the pre-filled pen using the grip. To mix the powder and the liquid, press the two halves together firmly by pressing down until you hear the pre-filled pen click. The two stoppers should come together.
- Wait for several seconds until the powder is completely dissolved.
- **Gently turn the PegIntron pre-filled pen upside down twice. To avoid excessive foaming, do not shake.**
- Maintaining the PegIntron pre-filled pen upright, check the mixed PegIntron solution through the window. If there is still foam, wait until it settles. The solution should be clear and colourless. **Do not use the pen if the solution is discoloured or contains any particles.**



Figure 2

- Keeping the PegIntron pre-filled pen upright in the holder provided in the packaging, disinfect the rubber membrane of the PegIntron pre-filled pen with one cleansing swab.
- Take the injection needle provided in the tray and remove its protective paper tab.
- **Maintaining the PegIntron pre-filled pen upright in the holder**, gently push the injection needle onto the pre-filled pen and **screw it securely in place**.

- **Keep the PegIntron pre-filled pen pointed up.**
- **Do not take off the outer needle cap at this point.**
- You may see some liquid trickle out from under the cap, as the air has been expelled out of the pen.
- Wait about 5 seconds for this process to finish.




Figure 3

- **Check through the window to be sure that the two stoppers are together. If they are not together, do not use this pen because you may not be able to dial your dose.**

### Step 2: Setting the dose



Figure 4

- Remove the PegIntron pre-filled pen from the holder.
- Holding the PegIntron pre-filled pen firmly, pull the dosing button out as far as it will go, until you see a **dark ring** on the pen. The dosing button should be easy to pull out without excessive force being required.

**Note: Do not push the dosing button back in at this time. You will push it in when you are ready to self-inject the PegIntron.**



Figure 5

- Turn the dosing button until your prescribed dose is aligned with the dosing tab. The button should turn easily without excessive force being required.

**Note:** **If you cannot easily pull out the dosing button or dial the dose, do not use excessive force and do not use this pen because it may not deliver the correct dose.**

### **Step 3: Injecting the solution**

- Select the injection site. Your doctor will have told you which sites to use (e.g., thigh or abdomen).

**Note:** change your injection site each time.

- Clean the injection site skin with the second cleansing swab.




Figure 6

- Pull off the **outer needle cap**.
- There may be some liquid around the inner needle cap. This liquid is not part of your dose, this is extra. This is normal, as the air has been expelled out of the needle.
- Once the injection site is dry, pull off the **yellow inner needle cap** carefully exposing the injection needle.



Figure 7

- Hold the PegIntron pre-filled pen with your fingers wrapped around the barrel and your thumb on the dosing button.
- With your other hand, pinch a fold of loose skin.
- Insert the needle into the pinched skin at an angle of 45° to 90°.
- Press the dosing button down **slowly** and **firmly** until the button can no longer move.
- **Maintain pressure on the dosing button for an additional 5 seconds** to ensure that you get the complete dose.

- Remove the needle from your skin.
- Press the injection site with a small bandage or sterile gauze if necessary for a few seconds.
- Do not massage the injection site. If there is bleeding, cover with an adhesive bandage.
- Discard the PegIntron pre-filled pen with the needle safely in a closed rigid container.