# ANNEX I SUMMARY OF PRODUCT CHARACTERISTICS

### 1. NAME OF THE MEDICINAL PRODUCT

Faslodex 250 mg solution for injection.

# 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One pre-filled syringe contains 250 mg fulvestrant in 5 ml solution.

### Excipients with known effect

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Solution for injection.

Clear, colourless to yellow, viscous solution.

### 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Faslodex is indicated for the treatment of postmenopausal women with estrogen receptor positive, locally advanced or metastatic breast cancer for disease relapse on or after adjuvant anti-estrogen therapy, or disease progression on therapy with an anti-estrogen.

# 4.2 Posology and method of administration

### **Posology**

Adult females (including Elderly)

The recommended dose is 500 mg at intervals of one month, with an additional 500 mg dose given two weeks after the initial dose.

# **Special populations**

# Renal impairment

No dose adjustments are recommended for patients with mild to moderate renal impairment (creatinine clearance ≥30 ml/min). Safety and efficacy have not been evaluated in patients with severe renal impairment (creatinine clearance <30 ml/min), and, therefore, caution is recommended in these patients (see section 4.4).

### Hepatic impairment

No dose adjustments are recommended for patients with mild to moderate hepatic impairment. However, as fulvestrant exposure may be increased, Faslodex should be used with caution in these patients. There are no data in patients with severe hepatic impairment (see sections 4.3, 4.4 and 5.2).

### Paediatric population

The safety and efficacy of Faslodex in children from birth to 18 years of age have not been established. Currently available data are described in sections 5.1 and 5.2, but no recommendation on a posology can be made.

# Method of administration

Faslodex should be administered as two consecutive 5 ml injections by slow intramuscular injection (1-2 minutes/injection), one in each buttock (gluteal area).

Caution should be taken if injecting Faslodex at the dorsogluteal site due to the proximity of the

underlying sciatic nerve.

For detailed instructions for administration, see section 6.6.

### 4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Pregnancy and lactation (see section 4.6).

Severe hepatic impairment (see sections 4.4 and 5.2).

# 4.4 Special warnings and precautions for use

Faslodex should be used with caution in patients with mild to moderate hepatic impairment (see sections 4.2, 4.3 and 5.2).

Faslodex should be used with caution in patients with severe renal impairment (creatinine clearance less than 30 ml/min).

Due to the intramuscular route of administration, Faslodex should be used with caution if treating patients with bleeding diatheses, thrombocytopenia or those taking anticoagulant treatment.

Thromboembolic events are commonly observed in women with advanced breast cancer and have been observed in clinical trials with Faslodex (see section 4.8). This should be taken into consideration when prescribing Faslodex to patients at risk.

Injection site related events including sciatica, neuralgia, neuropathic pain, and peripheral neuropathy have been reported with Faslodex injection. Caution should be taken while administering Faslodex at the dorsogluteal injection site due to the proximity of the underlying sciatic nerve (see sections 4.2 and 4.8).

There are no long-term data on the effect of fulvestrant on bone. Due to the mechanism of action of fulvestrant, there is a potential risk of osteoporosis.

# Interference with estradiol antibody assays

Due to the structural similarity of fulvestrant and estradiol, fulvestrant may interfere with antibody based-estradiol assays and may result in falsely increased levels of estradiol.

### Paediatric population

Faslodex is not recommended for use in children and adolescents as safety and efficacy have not been established in this group of patients (see section 5.1).

# 4.5 Interaction with other medicinal products and other forms of interaction

A clinical interaction study with midazolam (substrate of CYP3A4) demonstrated that fulvestrant does not inhibit CYP3A4. Clinical interaction studies with rifampicin (inducer of CYP3A4) and ketoconazole (inhibitor of CYP3A4) showed no clinically relevant change in fulvestrant clearance. Dose adjustment is therefore not necessary in patients who are receiving fulvestrant and CYP3A4 inhibitors or inducers concomitantly.

# 4.6 Fertility, pregnancy and lactation

### Women of childbearing potential

Patients of child-bearing potential should be advised to use effective contraception while on treatment.

# **Pregnancy**

Faslodex is contraindicated in pregnancy (see section 4.3). Fulvestrant has been shown to cross the placenta after single intramuscular doses in rat and rabbit. Studies in animals have shown reproductive

toxicity including an increased incidence of foetal abnormalities and deaths (see section 5.3). If pregnancy occurs while taking Faslodex, the patient must be informed of the potential hazard to the foetus and potential risk for loss of pregnancy.

### **Breast-feeding**

Breast-feeding must be discontinued during treatment with Faslodex. Fulvestrant is excreted in milk in lactating rats. It is not known whether fulvestrant is excreted in human milk. Considering the potential for serious adverse reactions due to fulvestrant in breast-fed infants, use during lactation is contraindicated (see section 4.3).

### Fertility

The effects of Faslodex on fertility in humans has not been studied.

### 4.7 Effects on ability to drive and use machines

Faslodex has no or negligible influence on the ability to drive or use machines. However, since asthenia has been reported very commonly with Faslodex, caution should be observed by those patients who experience this adverse reaction when driving or operating machinery.

### 4.8 Undesirable effects

This section provides information based on all adverse reactions from clinical trials, post-marketing studies or spontaneous reports. The most frequently reported adverse reactions are injection site reactions, asthenia, nausea, and increased hepatic enzymes (ALT, AST, ALP).

The following frequency categories for adverse drug reactions (ADRs) were calculated based on the Faslodex 500 mg treatment group in pooled safety analyses of the CONFIRM (Study D6997C00002), FINDER 1 (Study D6997C00004), FINDER 2 (Study D6997C00006), and NEWEST (Study D6997C00003) studies that compared Faslodex 500 mg with Faslodex 250 mg. The frequencies in the following table were based on all reported adverse drug reactions, regardless of the investigator assessment of causality.

Adverse reactions listed below are classified according to frequency and System Organ Class (SOC). Frequency groupings are defined according to the following convention: Very common ( $\geq 1/10$ ), Common ( $\geq 1/100$ ) to < 1/10), Uncommon ( $\geq 1/100$ ). Within each frequency grouping adverse reactions are reported in order of decreasing seriousness.

**Table 1** Adverse Drug Reactions

Adverse reactions by system organ class and frequency					
Infections and infestations	Common	Urinary tract infections			
Blood and lymphatic system disorders	Uncommon	Reduced platelet count			
Immune system disorders	Common	Hypersensitivity reactions			
Metabolism and nutrition disorders	Common	Anorexia <sup>a</sup>			
Nervous system disorders	Common	Headache			
Vascular disorders	Common	Venous thromboembolism <sup>a</sup> , hot			
		flushes			
Gastrointestinal disorders	Very common	Nausea			
	Common	Vomiting, diarrhoea			
Hepatobiliary disorders	Very common	Increased hepatic enzymes (ALT,			
		AST, ALP) <sup>a</sup>			
	Common	Elevated bilirubin <sup>a</sup>			
	Uncommon	Hepatic failure <sup>c</sup> , hepatitis, elevated			
		gamma-GT			
Skin and subcutaneous tissue disorders	Common	Rash			
Musculoskeletal and connective tissue	Common	Back pain <sup>a</sup>			

disorders			
Reproductive system and breast disorders	Uncommon	Vaginal moniliasis, leukorrhea,	
		vaginal haemorrhage	
General disorders and administration site	Very common	Asthenia <sup>a</sup> , injection site reactions <sup>b</sup>	
conditions	Uncommon	Injection site haemorrhage, injection	
		site haematoma, sciatica, neuralgia <sup>c</sup> ,	
		neuropathy peripheral	

- Includes adverse drug reactions for which the exact contribution of Faslodex cannot be assessed due to the underlying disease.
- The term injection site reactions does not include the terms injection site haemorrhage, injection site haematoma, sciatica, neuralgia and neuropathy peripheral.
- The event was not observed in major clinical studies (CONFIRM, FINDER 1, FINDER 2, NEWEST). The frequency has been calculated using the upper limit of the 95% confidence interval for the point estimate. This is calculated as 3/560 (where 560 is the number of patients in the major clinical studies), which equates to a frequency category of 'uncommon'.

# Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

### 4.9 Overdose

There is no human experience of overdose. Animal studies suggest that no effects other than those related directly or indirectly to anti-estrogenic activity were evident with higher doses of fulvestrant (see section 5.3). If overdose occurs, symptomatic supportive treatment is recommended.

# 5. PHARMACOLOGICAL PROPERTIES

# 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Endocrine therapy, Anti-estrogens, ATC code: L02BA03

# Mechanism of action and pharmacodynamic effects

Fulvestrant is a competitive estrogen receptor (ER) antagonist with an affinity comparable to estradiol. Fulvestrant blocks the trophic actions of estrogens without any partial agonist (estrogen-like) activity. The mechanism of action is associated with down-regulation of estrogen receptor protein levels. Clinical trials in postmenopausal women with primary breast cancer have shown that fulvestrant significantly down-regulates ER protein in ER positive tumours compared with placebo. There was also a significant decrease in progesterone receptor expression consistent with a lack of intrinsic estrogen agonist effects. It has also been shown that fulvestrant 500 mg downregulates ER and the proliferation marker Ki67, to a greater degree than fulvestrant 250 mg in breast tumours in postmenopausal neoadjuvant setting.

### Clinical efficacy and safety in advanced breast cancer

A phase III clinical trial was completed in 736 postmenopausal women with advanced breast cancer who had disease recurrence on or after adjuvant endocrine therapy or progression following endocrine therapy for advanced disease. The study included 423 patients whose disease had recurred or progressed during anti-estrogen therapy (AE subgroup) and 313 patients whose disease had recurred or progressed during aromatase inhibitor therapy (AI subgroup). This trial compared the efficacy and safety of Faslodex 500 mg (n=362) with Faslodex 250 mg (n=374). Progression-free survival (PFS) was the primary endpoint; key secondary efficacy endpoints included objective response rate (ORR), clinical benefit rate (CBR) and overall survival (OS). Efficacy results for the CONFIRM study are summarized in Table 2.

Table 2 Summary of results of the primary efficacy endpoint (PFS) and key secondary efficacy endpoints in the CONFIRM study

Variable	Type of estimate;	Faslodex 500 mg (N=362)	Faslodex 250 mg (N=374)	Comparison between groups (Faslodex 500 mg/Faslodex 250 mg)		
	treatment comparison			Hazard ratio	95% CI	p-value
PFS	K-M median					
	in months;					
	hazard ratio					
All Patients		6.5	5.5	0.80	0.68, 0.94	0.006
-AE subgrou	ıp (n=423)	8.6	5.8	0.76	0.62, 0.94	0.013
-AI subgrou	p (n=313) <sup>a</sup>	5.4	4.1	0.85	0.67, 1.08	0.195
OS <sup>b</sup>	K-M median					
	in months;					
	hazard ratio					
All Patients		26.4	22.3	0.81	0.69, 0.96	$0.016^{c}$
-AE subgro	oup (n=423)	30.6	23.9	0.79	0.63, 0.99	$0.038^{c}$
-AI subgrou	up (n=313) <sup>a</sup>	24.1	20.8	0.86	0.67, 1.11	0.241 <sup>c</sup>
Variable	Type of	Faslodex	Faslodex	Comparis	on between grou	ups
	estimate;	500 mg	250 mg	(Faslodex 500 mg/Faslodex 250 mg)		
	treatment	(N=362)	(N=374)	Absolute	95% CI	
	comparison		· · ·	difference in %		
ORR <sup>d</sup>	% of patients with OR; absolute difference in %					
All Patients		13.8	14.6	-0.8	-5.8, 6.3	
-AE subgro	oup (n=296)	18.1	19.1	-1.0	-8.2, 9.3	
-AI subgrou	up (n=205) <sup>a</sup>	7.3	8.3	-1.0	-5.5, 9.8	
CBR <sup>e</sup>	% of patients					
	with CB;					
	absolute					
	difference in					
	%					
All Patients		45.6	39.6	6.0	-1.1, 13.3	
-AE subgro	oup (n=423)	52.4	45.1	7.3	-2.2, 16.6	
_	up (n=313) <sup>a</sup>	36.2	32.3	3.9	-6.1, 15.2	

Faslodex is indicated in patients whose disease had recurred or progressed on an anti-estrogen therapy. The results in the AI subgroup are inconclusive.

Two phase-III clinical trials were completed in a total of 851 postmenopausal women with advanced breast cancer who had disease recurrence on or after adjuvant endocrine therapy or progression following endocrine therapy for advanced disease. 77% of the study population had estrogen receptor positive breast cancer. These trials compared the safety and efficacy of monthly administration of

OS is presented for the final survival analyses at 75% maturity.

Nominal p-value with no adjustments made for multiplicity between the initial overall survival analyses at 50% maturity and the updated survival analyses at 75% maturity.

ORR was assessed in patients who were evaluable for response at baseline (i.e. those with measurable disease at baseline: 240 patients in the Faslodex 500 mg group and 261 patients in the Faslodex 250 mg group).

Patients with a best objective response of complete response, partial response or stable disease ≥24 weeks. PFS:Progression-free survival; ORR:Objective response rate; OR:Objective response; CBR:Clinical benefit rate; CB:Clinical benefit; OS:Overall survival; K-M:Kaplan-Meier; CI:Confidence interval; AI:Aromatase inhibitor; AE:Anti-estrogen.

Faslodex 250 mg versus the daily administration of 1 mg anastrozole (aromatase inhibitor). Overall, Faslodex at the 250 mg monthly dose was at least as effective as anastrozole in terms of progression-free survival, objective response, and time to death. There were no statistically significant differences in any of these endpoints between the two treatment groups. Progression-free survival was the primary endpoint. Combined analysis of both trials showed that 83% of patients who received Faslodex progressed, compared with 85% of patients who received anastrozole. Combined analysis of both trials showed the hazard ratio of Faslodex 250 mg to anastrozole for progression-free survival was 0.95 (95% CI 0.82 to 1.10). The objective response rate for Faslodex 250 mg was 19.2% compared with 16.5% for anastrozole. The median time to death was 27.4 months for patients treated with Faslodex and 27.6 months for patients treated with anastrozole. The hazard ratio of Faslodex 250 mg to anastrozole for time to death was 1.01 (95% CI 0.86 to 1.19).

### Effects on the postmenopausal endometrium

Preclinical data do not suggest a stimulatory effect of fulvestrant on the postmenopausal endometrium (see section 5.3). A 2-week study in healthy postmenopausal volunteers treated with 20 µg per day ethinylestradiol showed that pre-treatment with Faslodex 250 mg resulted in significantly reduced stimulation of the postmenopausal endometrium, compared to pre-treatment with placebo, as judged by ultrasound measurement of endometrium thickness.

Neoadjuvant treatment for up to 16 weeks in breast cancer patients treated with either Faslodex 500 mg or Faslodex 250 mg did not result in clinically significant changes in endometrial thickness, indicating a lack of agonist effect. There is no evidence of adverse endometrial effects in the breast cancer patients studied. No data are available regarding endometrial morphology.

In two short-term studies (1 and 12 weeks) in premenopausal patients with benign gynaecologic disease, no significant differences in endometrial thickness were observed by ultrasound measurement between fulvestrant and placebo groups.

# Effects on bone

There are no long-term data on the effect of fulvestrant on bone. Neoadjuvant treatment for up to 16 weeks in breast cancer patients with either Faslodex 500 mg or Faslodex 250 mg did not result in clinically significant changes in serum bone-turnover markers.

# Paediatric population

Faslodex is not indicated for use in children. The European Medicines Agency has waived the obligation to submit the results of studies with Faslodex in all subsets of the paediatric population in breast cancer (see section 4.2 for information on paediatric use).

An open-label phase II study investigated the safety, efficacy and pharmacokinetics of fulvestrant in 30 girls aged 1 to 8 years with Progressive Precocious Puberty associated with McCune Albright Syndrome (MAS). The paediatric patients received 4 mg/kg monthly intramuscular dose of fulvestrant. This 12-month study investigated a range of MAS endpoints and showed a reduction in the frequency of vaginal bleeding and a reduction in the rate of bone age advancement. The steady-state trough concentrations of fulvestrant in children in this study were consistent with that in adults (see section 5.2). There were no new safety concerns arising from this small study, but 5-year data are yet not available.

# 5.2 Pharmacokinetic properties

# **Absorption**

After administration of Faslodex long-acting intramuscular injection, fulvestrant is slowly absorbed and maximum plasma concentrations ( $C_{max}$ ) are reached after about 5 days. Administration of Faslodex 500 mg regimen achieves exposure levels at, or close to, steady state within the first month of dosing (mean [CV]: AUC 475 [33.4%] ng.days/ml,  $C_{max}$  25.1 [35.3%] ng/ml,  $C_{min}$  16.3 [25.9%] ng/ml, respectively). At steady state, fulvestrant plasma concentrations are maintained within a relatively narrow range with up to an approximately 3-fold difference between maximum and

trough concentrations. After intramuscular administration, the exposure is approximately dose-proportional in the dose range 50 to 500 mg.

### Distribution

Fulvestrant is subject to extensive and rapid distribution. The large apparent volume of distribution at steady state ( $Vd_{ss}$ ) of approximately 3 to 5 l/kg suggests that distribution is largely extravascular. Fulvestrant is highly (99%) bound to plasma proteins. Very low density lipoprotein (VLDL), low density lipoprotein (LDL), and high density lipoprotein (HDL) fractions are the major binding components. No interaction studies were conducted on competitive protein binding. The role of sex hormone-binding globulin (SHBG) has not been determined.

### Biotransformation

The metabolism of fulvestrant has not been fully evaluated, but involves combinations of a number of possible biotransformation pathways analogous to those of endogenous steroids. Identified metabolites (includes 17-ketone, sulphone, 3-sulphate, 3- and 17-glucuronide metabolites) are either less active or exhibit similar activity to fulvestrant in anti-estrogen models. Studies using human liver preparations and recombinant human enzymes indicate that CYP3A4 is the only P450 isoenzyme involved in the oxidation of fulvestrant; however, non-P450 routes appear to be more predominant *in vivo*. *In vitro* data suggest that fulvestrant does not inhibit CYP450 isoenzymes.

### Elimination

Fulvestrant is eliminated mainly in metabolised form. The major route of excretion is via the faeces, with less than 1% being excreted in the urine. Fulvestrant has a high clearance,  $11\pm1.7$  ml/min/kg, suggesting a high hepatic extraction ratio. The terminal half-life ( $t_{1/2}$ ) after intramuscular administration is governed by the absorption rate and was estimated to be 50 days.

# Special populations

In a population pharmacokinetic analysis of data from phase III studies, no difference in fulvestrant's pharmacokinetic profile was detected with regard to age (range 33 to 89 years), weight (40-127 kg) or race.

### Renal impairment

Mild to moderate impairment of renal function did not influence the pharmacokinetics of fulvestrant to any clinically relevant extent.

### Hepatic impairment

The pharmacokinetics of fulvestrant has been evaluated in a single-dose clinical trial conducted in subjects with mild to moderate hepatic impairment (Child-Pugh class A and B). A high dose of a shorter duration intramuscular injection formulation was used. There was up to about 2.5-fold increase in AUC in subjects with hepatic impairment compared to healthy subjects. In patients administered Faslodex, an increase in exposure of this magnitude is expected to be well tolerated. Subjects with severe hepatic impairment (Child-Pugh class C) were not evaluated.

# Paediatric population

The pharmacokinetics of fulvestrant has been evaluated in a clinical trial conducted in 30 girls with Progressive Precocious Puberty associated with McCune Albright Syndrome (see section 5.1). The paediatric patients were aged 1 to 8 years and received 4 mg/kg monthly intramuscular dose of fulvestrant. The geometric mean (standard deviation) steady state trough concentration (Cmin,ss) and AUCss was 4.2 (0.9) ng/mL and 3680 (1020) ng\*hr/mL, respectively. Although the data collected were limited, the steady-state trough concentrations of fulvestrant in children appear to be consistent with those in adults.

### 5.3 Preclinical safety data

The acute toxicity of fulvestrant is low.

Faslodex and other formulations of fulvestrant were well tolerated in animal species used in multiple dose studies. Local reactions, including myositis and granulomata at the injection site were attributed to the vehicle but the severity of myositis in rabbits increased with fulvestrant, compared to the saline control. In toxicity studies with multiple intramuscular doses of fulvestrant in rats and dogs, the antiestrogenic activity of fulvestrant was responsible for most of the effects seen, particularly in the female reproductive system, but also in other organs sensitive to hormones in both sexes. Arteritis involving a range of different tissues was seen in some dogs after chronic (12 months) dosing.

In dog studies following oral and intravenous administration, effects on the cardiovascular system (slight elevations of the S-T segment of the ECG [oral], and sinus arrest in one dog [intravenous]) were seen. These occurred at exposure levels higher than in patients ( $C_{max} > 15$  times) and are likely to be of limited significance for human safety at the clinical dose.

Fulvestrant showed no genotoxic potential.

Fulvestrant showed effects upon reproduction and embryo/foetal development consistent with its anti-estrogenic activity, at doses similar to the clinical dose. In rats, a reversible reduction in female fertility and embryonic survival, dystocia and an increased incidence of foetal abnormalities including tarsal flexure were observed. Rabbits given fulvestrant failed to maintain pregnancy. Increases in placental weight and post-implantation loss of foetuses were seen. There was an increased incidence of foetal variations in rabbits (backwards displacement of the pelvic girdle and 27 pre-sacral vertebrae).

A two-year oncogenicity study in rats (intramuscular administration of Faslodex) showed increased incidence of ovarian benign granulosa cell tumours in female rats at the high dose, 10 mg/rat/15 days and an increased incidence of testicular Leydig cell tumours in males. In a two-year mouse oncogenicity study (daily oral administration) there was an increased incidence of ovarian sex cord stromal tumours (both benign and malignant) at doses of 150 and 500 mg/kg/day. At the no-effect level for these findings, systemic exposure levels (AUC) were, in rats, approximately 1.5–fold the expected human exposure levels in females and 0.8-fold in males, and in mice, approximately 0.8-fold the expected human exposure levels in both males and females. Induction of such tumours is consistent with pharmacology-related endocrine feedback alterations in gonadotropin levels caused by anti-estrogens in cycling animals. Therefore these findings are not considered to be relevant to the use of fulvestrant in postmenopausal women with advanced breast cancer.

# 6. PHARMACEUTICAL PARTICULARS

# 6.1 List of excipients

Ethanol (96 per cent) Benzyl alcohol Benzyl benzoate Castor oil

### 6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### 6.3 Shelf life

4 years

### 6.4 Special precautions for storage

Store and transport in a refrigerator (2°C - 8°C).

Temperature excursions outside  $2^{\circ}\text{C}$  -  $8^{\circ}\text{C}$  should be limited. This includes avoiding storage at temperatures exceeding  $30^{\circ}\text{C}$ , and not exceeding a 28 day period where the average storage temperature for the product is below  $25^{\circ}\text{C}$  (but above  $2^{\circ}\text{C}$  -  $8^{\circ}\text{C}$ ). After temperature excursions, the product should be returned immediately to the recommended storage conditions (store and transport in a refrigerator  $2^{\circ}\text{C}$  -  $8^{\circ}\text{C}$ ). Temperature excursions have a cumulative effect on the product quality and the 28 day time period must not be exceeded over the duration of the 4-year shelf life of Faslodex (see section 6.3). Exposure to temperatures below  $2^{\circ}\text{C}$  will not damage the product providing it is not stored below  $-20^{\circ}\text{C}$ .

Store the pre-filled syringe in the original package in order to protect from light.

### 6.5 Nature and contents of container

BD SafetyGlide is a trademark of Becton Dickinson and Company and is CE-marked: CE 0050.

The pre-filled syringe presentation consists of:

One clear type 1 glass pre-filled syringe with polystyrene plunger rod, fitted with a tamper-evident closure, containing 5 ml Faslodex solution for injection.

A safety needle (BD SafetyGlide<sup>TM</sup>) for connection to the barrel is also provided.

Or

Two clear type 1 glass pre-filled syringes with polystyrene plunger rod, fitted with a tamper-evident closure, each containing 5 ml Faslodex solution for injection. Safety needles (BD SafetyGlide<sup>TM</sup>) for connection to each barrel are also provided.

Not all pack sizes may be marketed.

# 6.6 Special precautions for disposal and other handling

# Instructions for administration

Administer the injection according to the local guidelines for performing large volume intramuscular injections.


NOTE: Due to the proximity of the underlying sciatic nerve, caution should be taken if administering Faslodex at the dorsogluteal injection site (see section 4.4).

Warning - Do not autoclave safety needle (BD SafetyGlide $^{\text{\tiny TM}}$  Shielding Hypodermic Needle) before use. Hands must remain behind the needle at all times during use and disposal.

For each of the two syringes:


- Remove glass syringe barrel from tray and check that it is not damaged.
- Peel open the safety needle (SafetyGlide<sup>™</sup>) outer packaging.
- Parenteral solutions must be inspected visually for particulate matter and discolouration prior to administration.
- Hold the syringe upright on the ribbed part (C). With the other hand, take hold of the cap (A) and carefully tilt back and forth until the cap disconnects and can be pulled off, do not twist (see Figure 1).

Figure 1




• Remove the cap (A) in a straight upward direction. To maintain sterility do not touch the syringe tip (B) (see Figure 2).

Figure 2




- Attach the safety needle to the Luer-Lok and twist until firmly seated (see Figure 3).
- Check that the needle is locked to the Luer connector before moving out of the vertical plane.
- Pull shield straight off needle to avoid damaging needle point.
- Transport filled syringe to point of administration.
- Remove needle sheath.
- Expel excess gas from the syringe.

Figure 3



• Administer intramuscularly slowly (1-2 minutes/injection) into the buttock (gluteal area). For user convenience, the needle bevel-up position is oriented to the lever arm (see Figure 4).

Figure 4



 After injection, immediately apply a single-finger stroke to the activation assisted lever arm to activate the shielding mechanism (see Figure 5).

NOTE: Activate away from self and others. Listen for click and visually confirm needle tip is fully covered.

Figure 5



### Disposal

Pre-filled syringes are for single use **only**.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

### 7. MARKETING AUTHORISATION HOLDER

AstraZeneca UK Limited Charter Way, Macclesfield, Cheshire SK10 2NA United Kingdom

# 8. MARKETING AUTHORISATION NUMBER(S)

EU/1/03/269/001 EU/1/03/269/002

### 9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 10 March 2004

Date of latest renewal: 10 March 2009

# 10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency  $\underline{\text{http://www.ema.europa.eu}}$ 

# **ANNEX II**

- A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

# A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer responsible for batch release AstraZeneca UK Limited Silk Road Business Park, Macclesfield, SK10 2NA United Kingdom

### B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to medical prescription.

# C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

# • Periodic Safety Update Reports

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

# D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

# • Risk Management Plan (RMP)

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

# ANNEX III LABELLING AND PACKAGE LEAFLET

A. LABELLING

# PARTICULARS TO APPEAR ON THE OUTER PACKAGING AND THE IMMEDIATE PACKAGING

### **CARTON**

# 1. NAME OF THE MEDICINAL PRODUCT

Faslodex 250 mg solution for injection. fulvestrant

# 2. STATEMENT OF ACTIVE SUBSTANCE(S)

One pre-filled syringe contains 250 mg fulvestrant in 5 ml solution

# 3. LIST OF EXCIPIENTS

Ethanol (96 per cent), benzyl alcohol, benzyl benzoate and castor oil. See the package leaflet for further information.

# 4. PHARMACEUTICAL FORM AND CONTENTS

Solution for injection in a pre-filled syringe.

1 pre-filled syringe (5 ml)

1 safety needle

2 pre-filled syringes (5 ml each)

2 safety needles

# 5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.

Intramuscular use.

For single use only.

For full instructions on the administration of Faslodex and the use of the safety needle see enclosed, Instructions for administration.

Two syringes must be administered to receive the 500 mg recommended monthly dose.

# 6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

# 7. OTHER SPECIAL WARNING(S), IF NECESSARY

### 8. EXPIRY DATE

# 9. SPECIAL STORAGE CONDITIONS

Store and transport in a refrigerator.

Store the pre-filled syringe in the original package in order to protect from light. See package leaflet for information on temperature excursions.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

### 11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

AstraZeneca UK Limited Charter Way, Macclesfield, Cheshire SK10 2NA United Kingdom

# 12. MARKETING AUTHORISATION NUMBER(S)

EU/1/03/269/001 EU/1/03/269/002

# 13. BATCH NUMBER

Lot

# 14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

# 15. INSTRUCTIONS ON USE

# 16. INFORMATION IN BRAILLE

Justification for not including Braille accepted

# 17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

# 18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

PC:

SN: NN:

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS			
PRE-FILLED SYRINGE LABEL			
1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION			
Faslodex 250 mg solution for injection fulvestrant IM use			
2. METHOD OF ADMINISTRATION			
3. EXPIRY DATE			
EXP			
4. BATCH NUMBER			
Lot			
5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT			
5 ml			
6. OTHER			

**B. PACKAGE LEAFLET** 

### Package leaflet: Information for the user

# Faslodex 250 mg solution for injection

**Fulvestrant** 

# Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist, or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist, or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet

- 1. What Faslodex is and what it is used for
- 2. What you need to know before you use Faslodex
- 3. How to use Faslodex
- 4. Possible side effects
- 5. How to store Faslodex
- 6. Contents of the pack and other information

### 1. What Faslodex is and what it is used for

Faslodex contains the active substance fulvestrant, which belongs to the group of estrogen blockers. Estrogens, a type of female sex hormones, can in some cases be involved in the growth of breast cancer.

Faslodex is used to treat advanced or metastatic breast cancer in postmenopausal women.

# 2. What you need to know before you use Faslodex

### Do not use Faslodex:

- if you are allergic to fulvestrant or to any of the other ingredients of this medicine (listed in section 6)
- if you are pregnant or breast-feeding
- if you have severe liver problems

### Warnings and precautions

Talk to your doctor or pharmacist or nurse before using Faslodex if any of these apply to you:

- kidney or liver problems
- low numbers of platelets (which help blood clotting) or bleeding disorders
- previous problems with blood clots
- osteoporosis (loss of bone density)
- alcoholism

### Children and adolescents

Faslodex is not indicated in children and adolescents under 18 years.

### Other medicines and Faslodex

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

In particular, you should tell your doctor if you are using anticoagulants (medicines to prevent blood clots).

### **Pregnancy and breast-feeding**

You must not use Faslodex if you are pregnant. If you can become pregnant, you should use effective contraception while being treated with Faslodex.

You must not breast-feed while on treatment with Faslodex.

### **Driving and using machines**

Faslodex is not expected to affect your ability to drive or use machines. However, if you feel tired after treatment do not drive or use machines.

**Faslodex contains 10% w/v ethanol (alcohol)**, i.e. up to 1000 mg per dose, equivalent to 20 ml beer or 8 ml wine per dose.

Harmful for those suffering from alcoholism.

To be taken into account in pregnant or breast-feeding women, children and high-risk groups such as patients with liver disease, or epilepsy.

### 3. How to use Faslodex

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is 500 mg fulvestrant (two 250 mg/5 ml injections) given once a month, with an additional 500 mg dose given 2 weeks after the initial dose.

Your doctor or nurse will give you Faslodex as a slow intramuscular injection, one into each of your buttocks.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

# You may need immediate medical treatment if you experience any of the following side effects:

- Allergic (hypersensitivity) reactions, including swelling of the face, lips, tongue and/or throat
- Thromboembolism (increased risk of blood clots)\*
- Inflammation of the liver (hepatitis)
- Liver failure

# Tell your doctor, pharmacist, or nurse if you notice any of the following side effects:

Very common side effects (may affect more than 1 in 10 people)

- Injection site reactions, such as pain and/or inflammation
- Abnormal levels of liver enzymes (in blood tests)\*
- Nausea (feeling sick)
- Weakness, tiredness\*

### All other side effects:

**Common side effects** (may affect up to 1 in 10 people)

- Headache
- Hot flushes
- Vomiting, diarrhoea, or loss of appetite\*
- Rash
- Urinary tract infections
- Back pain\*
- Increase of bilirubin (bile pigment produced by the liver)
- Thromboembolism (increased risk of blood clots)\*
- Allergic (hypersensitivity) reactions, including swelling of the face, lips, tongue and/or throat

# **Uncommon side effects** (may affect up to 1 in 100 people)

- Decreased levels of platelets (thrombocytopenia)
- Vaginal bleeding, thick, whitish discharge and candidiasis (infection)
- Bruising and bleeding at the site of injection
- Increase of gamma-GT, a liver enzyme seen in a blood test
- Inflammation of the liver (hepatitis)
- Liver failure
- Lower back pain irradiating to leg on one side (sciatica)
- Numbness, tingling and pain
- Sudden weakness, numbness, tingling, or loss of movement in your leg, especially on only one side of your body, sudden problems with walking or balance (peripheral neuropathy)

# **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in <a href="Appendix V">Appendix V</a>. By reporting side effects you can help provide more information on the safety of this medicine.

# 5. How to store Faslodex

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton or syringe labels after EXP. The expiry date refers to the last day of that month.

Store and transport in a refrigerator (2°C - 8°C).

Temperature excursions outside 2°C - 8°C should be limited. This includes avoiding storage at temperatures exceeding 30°C, and not exceeding a 28 day period where the average storage temperature for the product is below 25°C (but above 2°C - 8°C). After temperature excursions, the product should be returned immediately to the recommended storage conditions (store and transport in a refrigerator 2°C - 8°C). Temperature excursions have a cumulative effect on the product quality and the 28 day time period must not be exceeded over the duration of the 4-year shelf life of Faslodex. Exposure to temperatures below 2°C will not damage the product providing it is not stored below -20°C.

Keep the pre-filled syringe in the original package, in order to protect from light.

Your healthcare professional will be responsible for the correct storage, use and disposal of Faslodex.

<sup>\*</sup> Includes side effects for which the exact role of Faslodex cannot be assessed due to the underlying disease.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

# 6. Contents of the pack and other information

### What Faslodex contains

- The active substance is fulvestrant. Each pre-filled syringe (5 ml) contains 250 mg fulvestrant.
- The other ingredients (excipients) are ethanol (96 per cent), benzyl alcohol, benzyl benzoate and castor oil.

# What Faslodex looks like and contents of the pack

Faslodex is a clear, colourless to yellow, viscous solution in a pre-filled syringe fitted with a tamper-evident closure, containing 5 ml solution for injection. Two syringes must be administered to receive the 500 mg recommended monthly dose.

Faslodex has 2 pack presentations, either a pack containing 1 glass pre-filled syringe or a pack containing 2 glass pre-filled syringes. Safety needles (BD SafetyGlide<sup>TM</sup>) for connection to each barrel are also provided.

Not all pack sizes may be marketed.

# **Marketing Authorisation Holder**

AstraZeneca UK Limited Charter Way, Macclesfield, Cheshire SK10 2NA United Kingdom

### Manufacturer

AstraZeneca UK Limited Silk Road Business Park Macclesfield Cheshire SK10 2NA United Kingdom

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

# België/Belgique/Belgien

AstraZeneca S.A./N.V. Tel: +32 2 370 48 11

### България

АстраЗенека България ЕООД Тел.: +359 24455000

### Česká republika

AstraZeneca Czech Republic s.r.o.

Tel: +420 222 807 111

#### Danmark

AstraZeneca A/S Tlf: +45 43 66 64 62

### Lietuva

UAB AstraZeneca Lietuva Tel: +370 5 2660550

### Luxembourg/Luxemburg

AstraZeneca S.A./N.V. Tél/Tel: +32 2 370 48 11

# Magyarország

AstraZeneca Kft Tel: +36 1 883 6500

#### Malta

Associated Drug Co. Ltd Tel: +356 2277 8000

**Deutschland** 

AstraZeneca GmbH Tel: +49 41 03 7080

**Eesti** 

AstraZeneca

Tel: +372 6549 600

Ελλάδα

AstraZeneca A.E.

 $T\eta\lambda$ : + 30 210 6871500

España

AstraZeneca Farmacéutica Spain, S.A.

Tel: +34 91 301 91 00

France

AstraZeneca

Tél: +33 1 41 29 40 00

Hrvatska

AstraZeneca d.o.o.

Tel: +385 1 4628 000

**Ireland** 

AstraZeneca Pharmaceuticals (Ireland) Ltd

Tel: +353 1609 7100

Ísland

Vistor hf.

Sími: +354 535 7000

Italia

AstraZeneca S.p.A.

Tel: +39 02 9801 1

Κύπρος

Αλέκτωρ Φαρμακευτική Λτδ

Τηλ: +357 22490305

Latvija

SIA AstraZeneca Latvija

Tel: +371 67377100

Nederland

AstraZeneca BV

Tel: +31 79 363 2222

Norge

AstraZeneca AS

Tlf: +47 21 00 64 00

Österreich

AstraZeneca Österreich GmbH

Tel: +43 1 711 31 0

Polska

AstraZeneca Pharma Poland Sp. z o.o.

Tel.: +48 22 245 73 00

**Portugal** 

AstraZeneca Produtos Farmacêuticos, Lda.

Tel: +351 21 434 61 00

România

AstraZeneca Pharma SRL

Tel: +40 21 317 60 41

Slovenija

AstraZeneca UK Limited

Tel: +386 1 51 35 600

Slovenská republika

AstraZeneca AB, o.z.

Tel: +421 2 5737 7777

Suomi/Finland

AstraZeneca Oy

Puh/Tel: +358 10 23 010

Sverige

AstraZeneca AB

Tel: +46 8 553 26 000

**United Kingdom** 

AstraZeneca UK Ltd

Tel: +44 1582 836 836

This leaflet was last revised in

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site: http://www.ema.europa.eu

This leaflet is available in all EU/EEA languages on the European Medicines Agency website.

### The following information is intended for healthcare professionals only:

Faslodex 500 mg (2 x 250 mg/5 ml solution for injection) should be administered using two pre-filled syringes, see section 3.

BD SafetyGlide is a trademark of Becton Dickinson and Company and is CE-marked: CE 0050.


# Instructions for administration

Warning - Do not autoclave safety needle (BD SafetyGlide<sup>™</sup> Shielding Hypodermic Needle) before use. Hands must remain behind the needle at all times during use and disposal.

# For each of the two syringes:


- Remove glass syringe barrel from tray and check that it is not damaged.
- Peel open the safety needle (SafetyGlide<sup>™</sup>) outer packaging.
- Parenteral solutions must be inspected visually for particulate matter and discolouration prior to administration.
- Hold the syringe upright on the ribbed part (C). With the
  other hand, take hold of the cap (A) and carefully tilt back
  and forth until the cap disconnects and can be pulled off,
  do not twist (see Figure 1).

Figure 1




• Remove the cap (A) in a straight upward direction. To maintain sterility do not touch the syringe tip (B) (see Figure 2).

Figure 2




- Attach the safety needle to the Luer-Lok and twist until firmly seated (see Figure 3).
- Check that the needle is locked to the Luer connector before moving out of the vertical plane.
- Pull shield straight off needle to avoid damaging needle point.
- Transport filled syringe to point of administration.
- Remove needle sheath.
- Expel excess gas from the syringe.

Figure 3



• Administer intramuscularly slowly (1-2 minutes/injection) into the buttock (gluteal area). For user convenience, the needle bevel-up position is oriented to the lever arm (see Figure 4).

Figure 4



• After injection, immediately apply a single-finger stroke to the activation assisted lever arm to activate the shielding mechanism (see Figure 5).

NOTE: Activate away from self and others. Listen for click and visually confirm needle tip is fully covered.

Figure 5



# **Disposal**

Pre-filled syringes are for single use **only**.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.