

**ANNEX I**

**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Faslodex 250 mg solution for injection.

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One pre-filled syringe contains 250 mg fulvestrant in 5 ml solution.

### Excipients with known effect

Ethanol

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Solution for injection.

Clear, colourless to yellow, viscous solution.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Faslodex is indicated:

- as monotherapy for the treatment of estrogen receptor positive, locally advanced or metastatic breast cancer in postmenopausal women:
  - not previously treated with endocrine therapy, or
  - with disease relapse on or after adjuvant antiestrogen therapy, or disease progression on antiestrogen therapy.
- in combination with palbociclib for the treatment of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative locally advanced or metastatic breast cancer in women who have received prior endocrine therapy (see section 5.1).

In pre- or perimenopausal women, the combination treatment with palbociclib should be combined with a luteinizing hormone releasing hormone (LHRH) agonist.

### 4.2 Posology and method of administration

#### Posology

##### *Adult females (including Elderly)*

The recommended dose is 500 mg at intervals of one month, with an additional 500 mg dose given two weeks after the initial dose.

When Faslodex is used in combination with palbociclib, please also refer to the Summary of Product Characteristics of palbociclib.

Prior to the start of treatment with the combination of Faslodex plus palbociclib, and throughout its duration, pre/perimenopausal women should be treated with LHRH agonists according to local clinical practice.

#### Special populations

##### *Renal impairment*

No dose adjustments are recommended for patients with mild to moderate renal impairment (creatinine clearance  $\geq 30$  ml/min). Safety and efficacy have not been evaluated in patients with severe renal

impairment (creatinine clearance <30 ml/min), and, therefore, caution is recommended in these patients (see section 4.4).

#### *Hepatic impairment*

No dose adjustments are recommended for patients with mild to moderate hepatic impairment. However, as fulvestrant exposure may be increased, Faslodex should be used with caution in these patients. There are no data in patients with severe hepatic impairment (see sections 4.3, 4.4 and 5.2).

#### *Paediatric population*

The safety and efficacy of Faslodex in children from birth to 18 years of age have not been established. Currently available data are described in sections 5.1 and 5.2, but no recommendation on a posology can be made.

#### Method of administration

Faslodex should be administered as two consecutive 5 ml injections by slow intramuscular injection (1-2 minutes/injection), one in each buttock (gluteal area).

Caution should be taken if injecting Faslodex at the dorsogluteal site due to the proximity of the underlying sciatic nerve.

For detailed instructions for administration, see section 6.6.

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Pregnancy and lactation (see section 4.6).

Severe hepatic impairment (see sections 4.4 and 5.2).

### **4.4 Special warnings and precautions for use**

Faslodex should be used with caution in patients with mild to moderate hepatic impairment (see sections 4.2, 4.3 and 5.2).

Faslodex should be used with caution in patients with severe renal impairment (creatinine clearance less than 30 ml/min).

Due to the intramuscular route of administration, Faslodex should be used with caution if treating patients with bleeding diatheses, thrombocytopenia or those taking anticoagulant treatment.

Thromboembolic events are commonly observed in women with advanced breast cancer and have been observed in clinical studies with Faslodex (see section 4.8). This should be taken into consideration when prescribing Faslodex to patients at risk.

Injection site related events including sciatica, neuralgia, neuropathic pain, and peripheral neuropathy have been reported with Faslodex injection. Caution should be taken while administering Faslodex at the dorsogluteal injection site due to the proximity of the underlying sciatic nerve (see sections 4.2 and 4.8).

There are no long-term data on the effect of fulvestrant on bone. Due to the mechanism of action of fulvestrant, there is a potential risk of osteoporosis.

The efficacy and safety of Faslodex (either as monotherapy or in combination with palbociclib) have not been studied in patients with critical visceral disease.

When Faslodex is combined with palbociclib, please also refer to the Summary of Product Characteristics of palbociclib.

#### *Interference with estradiol antibody assays*

Due to the structural similarity of fulvestrant and estradiol, fulvestrant may interfere with antibody based-estradiol assays and may result in falsely increased levels of estradiol.

#### Paediatric population

Faslodex is not recommended for use in children and adolescents as safety and efficacy have not been established in this group of patients (see section 5.1).

### **4.5 Interaction with other medicinal products and other forms of interaction**

A clinical interaction study with midazolam (substrate of CYP3A4) demonstrated that fulvestrant does not inhibit CYP3A4. Clinical interaction studies with rifampicin (inducer of CYP3A4) and ketoconazole (inhibitor of CYP3A4) showed no clinically relevant change in fulvestrant clearance. Dose adjustment is therefore not necessary in patients who are receiving fulvestrant and CYP3A4 inhibitors or inducers concomitantly.

### **4.6 Fertility, pregnancy and lactation**

#### Women of childbearing potential

Patients of childbearing potential should be advised to use effective contraception while on treatment.

#### Pregnancy

Faslodex is contraindicated in pregnancy (see section 4.3). Fulvestrant has been shown to cross the placenta after single intramuscular doses in rat and rabbit. Studies in animals have shown reproductive toxicity including an increased incidence of foetal abnormalities and deaths (see section 5.3). If pregnancy occurs while taking Faslodex, the patient must be informed of the potential hazard to the foetus and potential risk for loss of pregnancy.

#### Breast-feeding

Breast-feeding must be discontinued during treatment with Faslodex. Fulvestrant is excreted in milk in lactating rats. It is not known whether fulvestrant is excreted in human milk. Considering the potential for serious adverse reactions due to fulvestrant in breast-fed infants, use during lactation is contraindicated (see section 4.3).

#### Fertility

The effects of Faslodex on fertility in humans has not been studied.

### **4.7 Effects on ability to drive and use machines**

Faslodex has no or negligible influence on the ability to drive or use machines. However, since asthenia has been reported very commonly with Faslodex, caution should be observed by those patients who experience this adverse reaction when driving or operating machinery.

### **4.8 Undesirable effects**

#### Summary of the safety profile

##### *Monotherapy*

This section provides information based on all adverse reactions from clinical studies, post-marketing studies or spontaneous reports. In the pooled dataset of fulvestrant monotherapy, the most frequently reported adverse reactions were injection site reactions, asthenia, nausea, and increased hepatic enzymes (ALT, AST, ALP).

In Table 1, the following frequency categories for adverse drug reactions (ADRs) were calculated based on the Faslodex 500 mg treatment group in pooled safety analyses of studies that compared Faslodex 500 mg with Faslodex 250 mg [CONFIRM (Study D6997C00002), FINDER 1 (Study D6997C00004), FINDER 2 (Study D6997C00006), and NEWEST (Study D6997C00003) studies], or from FALCON (Study D699BC00001) alone that compared Faslodex 500 mg with anastrozole 1 mg.

Where frequencies differ between the pooled safety analysis and FALCON, the highest frequency is presented. The frequencies in Table 1 were based on all reported adverse drug reactions, regardless of the investigator assessment of causality. The median duration of fulvestrant 500 mg treatment across the pooled dataset (including the studies mentioned above plus FALCON) was 6.5 months.

#### Tabulated list of adverse reactions

Adverse reactions listed below are classified according to frequency and System Organ Class (SOC). Frequency groupings are defined according to the following convention: Very common ( $\geq 1/10$ ), Common ( $\geq 1/100$  to  $< 1/10$ ), Uncommon ( $\geq 1/1,000$  to  $< 1/100$ ). Within each frequency grouping adverse reactions are reported in order of decreasing seriousness.

**Table 1 Adverse Drug Reactions reported in patients treated with Faslodex monotherapy**

Adverse reactions by system organ class and frequency		
Infections and infestations	Common	Urinary tract infections
Blood and lymphatic system disorders	Common	Reduced platelet count <sup>c</sup>
Immune system disorders	Very common	Hypersensitivity reactions <sup>c</sup>
	Uncommon	Anaphylactic reactions
Metabolism and nutrition disorders	Common	Anorexia <sup>a</sup>
Nervous system disorders	Common	Headache
Vascular disorders	Very common	Hot flushes <sup>c</sup>
	Common	Venous thromboembolism <sup>a</sup>
Gastrointestinal disorders	Very common	Nausea
	Common	Vomiting, diarrhoea
Hepatobiliary disorders	Very common	Elevated hepatic enzymes (ALT, AST, ALP) <sup>a</sup>
	Common	Elevated bilirubin <sup>a</sup>
	Uncommon	Hepatic failure <sup>c, f</sup> , hepatitis <sup>f</sup> , elevated gamma-GT <sup>f</sup>
Skin and subcutaneous tissue disorders	Very common	Rash <sup>c</sup>
Musculoskeletal and connective tissue disorders	Very common	Joint and musculoskeletal pain <sup>d</sup>
	Common	Back pain <sup>a</sup>
Reproductive system and breast disorders	Common	Vaginal haemorrhage <sup>c</sup>
	Uncommon	Vaginal moniliasis <sup>f</sup> , leukorrhea <sup>f</sup>
General disorders and administration site conditions	Very common	Asthenia <sup>a</sup> , injection site reactions <sup>b</sup>
	Common	Neuropathy peripheral <sup>e</sup> , sciatica <sup>c</sup>
	Uncommon	Injection site haemorrhage <sup>f</sup> , injection site haematoma <sup>f</sup> , neuralgia <sup>c, f</sup>

<sup>a</sup> Includes adverse drug reactions for which the exact contribution of Faslodex cannot be assessed due to the underlying disease.

<sup>b</sup> The term injection site reactions does not include the terms injection site haemorrhage, injection site haematoma, sciatica, neuralgia and neuropathy peripheral.

<sup>c</sup> The event was not observed in major clinical studies (CONFIRM, FINDER 1, FINDER 2, NEWEST). The frequency has been calculated using the upper limit of the 95% confidence interval for the point estimate. This is calculated as 3/560 (where 560 is the number of patients in the major clinical studies), which equates to a frequency category of 'uncommon'.

<sup>d</sup> Includes: arthralgia, and less frequently musculoskeletal pain, myalgia and pain in extremity.

<sup>e</sup> Frequency category differs between pooled safety dataset and FALCON.

<sup>f</sup> ADR was not observed in FALCON.

#### Description of selected adverse reactions

The descriptions included below are based on the safety analysis set of 228 patients who received at least one (1) dose of fulvestrant and 232 patients who received at least one (1) dose of anastrozole, respectively in the Phase 3 FALCON study.

### *Joint and musculoskeletal pain*

In the FALCON study, the number of patients who reported an adverse reaction of joint and musculoskeletal pain was 65 (31.2%) and 48 (24.1%) for fulvestrant and anastrozole arms, respectively. Of the 65 patients in the Faslodex arm, 40% (26/65) of patients reported joint and musculoskeletal pain within the first month of treatment, and 66.2% (43/65) of patients within the first 3 months of treatment. No patients reported events that were CTCAE Grade  $\geq 3$  or that required a dose reduction, dose interruption, or discontinued treatment due to these adverse reactions.

### *Combination therapy with palbociclib*

The overall safety profile of fulvestrant when used in combination with palbociclib is based on data from 517 patients with HR-positive, HER2-negative advanced or metastatic breast cancer in the randomised PALOMA3 study (see section 5.1). The most common ( $\geq 20\%$ ) adverse reactions of any grade reported in patients receiving fulvestrant in combination with palbociclib were neutropenia, leukopenia, infections, fatigue, nausea, anaemia, stomatitis, diarrhoea, and thrombocytopenia. The most common ( $\geq 2\%$ ) Grade  $\geq 3$  adverse reactions were neutropenia, leukopenia, anaemia, infections, AST increased, thrombocytopenia, and fatigue.

Table 2 reports the adverse reactions from PALOMA3.

Median duration of exposure to fulvestrant was 11.2 months in the fulvestrant + palbociclib arm and 4.9 months in the fulvestrant + placebo arm. Median duration of exposure to palbociclib in the fulvestrant + palbociclib arm was 10.8 months.

**Table 2 Adverse reactions based on PALOMA3 Study (N=517)**

System Organ Class Frequency Preferred Term <sup>a</sup>	Faslodex + Palbociclib (N=345)		Faslodex + placebo (N=172)	
	All Grades n (%)	Grade $\geq 3$ n (%)	All Grades n (%)	Grade $\geq 3$ n (%)
<b>Infections and infestations</b>				
<i>Very common</i>				
Infections <sup>b</sup>	163 (47.2)	11 (3.2)	54 (31.4)	5 (2.9)
<b>Blood and lymphatic system disorders</b>				
<i>Very common</i>				
Neutropenia <sup>c</sup>	287 (83.2)	228 (66.1)	7 (4.1)	1 (0.6)
Leukopenia <sup>d</sup>	183 (53.0)	105 (30.4)	9 (5.2)	2 (1.2)
Anaemia <sup>e</sup>	102 (29.6)	12 (3.5)	22 (12.8)	3 (1.7)
Thrombocytopenia <sup>f</sup>	78 (22.6)	8 (2.3)	0 (0.0)	0
<i>Uncommon</i>				
Febrile neutropenia	3 (0.9)	3 (0.9)	1 (0.6)	1 (0.6)
<b>Metabolism and nutrition disorders</b>				
<i>Very common</i>				
Decreased appetite	55 (15.9)	3 (0.9)	14 (8.1)	1 (0.6)
<b>Nervous system disorders</b>				
<i>Common</i>				
Dysgeusia	23 (6.7)	0	5 (2.9)	0
<b>Eye disorders</b>				
<i>Common</i>				
Lacrimation increased	22 (6.4)	0	2 (1.2)	0
Vision blurred	20 (5.8)	0	3 (1.7)	0
Dry eye	13 (3.8)	0	3 (1.7)	0

<b>Respiratory, thoracic and mediastinal disorders</b>				
<i>Common</i>				
Epistaxis	23 (6.7)	0	3 (1.7)	0
<b>Gastrointestinal disorders</b>				
<i>Very common</i>				
Nausea	117 (33.9)	0	48 (27.9)	1 (0.6)
Stomatitis <sup>g</sup>	97 (28.1)	2 (0.6)	22 (12.8)	0
Diarrhoea	81 (23.5)	0	33 (19.2)	2 (1.2)
Vomiting	65 (18.8)	2 (0.6)	26 (15.1)	1 (0.6)
<b>Skin and subcutaneous tissue disorders</b>				
<i>Very common</i>				
Alopecia	62 (18.0)	0	11 (6.4)	0
Rash <sup>h</sup>	58 (16.8)	2 (0.6)	11 (6.4)	0
<i>Common</i>				
Dry skin	21 (6.1)	0	2 (1.2)	0
<b>General disorders and administration site conditions</b>				
<i>Very common</i>				
Fatigue	142 (41.2)	8 (2.3)	50 (29.1)	2 (1.2)
Pyrexia	44 (12.8)	1 (0.3)	9 (5.2)	0
<i>Common</i>				
Asthenia	26 (7.5)	0	9 (5.2)	1 (0.6)
<b>Investigations</b>				
<i>Common</i>				
AST increased	26 (7.5)	10 (2.9)	9 (5.2)	3 (1.7)
ALT increased	20 (5.8)	6 (1.7)	6 (3.5)	0

ALT=alanine aminotransferase; AST=aspartate aminotransferase; N/n=number of patients

<sup>a</sup> Preferred Terms (PTs) are listed according to MedDRA 17.1.

<sup>b</sup> Infections includes all PTs that are part of the System Organ Class Infections and infestations.

<sup>c</sup> Neutropenia includes the following PTs: Neutropenia, Neutrophil count decreased.

<sup>d</sup> Leukopenia includes the following PTs: Leukopenia, White blood cell count decreased.

<sup>e</sup> Anaemia includes the following PTs: Anaemia, Haemoglobin decreased, Haematocrit decreased.

<sup>f</sup> Thrombocytopenia includes the following PTs: Thrombocytopenia, Platelet count decreased.

<sup>g</sup> Stomatitis includes the following PTs: Aphthous stomatitis, Cheilitis, Glossitis, Glossodynia, Mouth ulceration, Mucosal inflammation, Oral pain, Oropharyngeal discomfort, Oropharyngeal pain, Stomatitis.

<sup>h</sup> Rash includes the following PTs: Rash, Rash maculo-papular, Rash pruritic, Rash erythematous, Rash papular, Dermatitis, Dermatitis acneiform, Toxic skin eruption.

### Description of selected adverse reactions

#### *Neutropenia*

In patients receiving fulvestrant in combination with palbociclib in the PALOMA3 study, neutropenia of any grade was reported in 287 (83.2%) patients, with Grade 3 neutropenia being reported in 191 (55.4%) patients, and Grade 4 neutropenia being reported in 37 (10.7%) patients. In the fulvestrant + placebo arm (n=172), neutropenia of any grade was reported in 7 (4.1%) patients, with Grade 3 neutropenia reported in 1 (0.6%) patient. There were no reports of Grade 4 neutropenia in the fulvestrant + placebo arm.

In patients receiving fulvestrant in combination with palbociclib, the median time to first episode of any grade neutropenia was 15 days (range: 13-317) and the median duration of Grade  $\geq$ 3 neutropenia was 7 days. Febrile neutropenia has been reported in 0.9% patients receiving fulvestrant in combination with palbociclib.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

### **4.9 Overdose**

There are isolated reports of overdose with Faslodex in humans. If overdose occurs, symptomatic supportive treatment is recommended. Animal studies suggest that no effects other than those related directly or indirectly to antiestrogenic activity were evident with higher doses of fulvestrant (see section 5.3).

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Endocrine therapy, Antiestrogens, ATC code: L02BA03

#### Mechanism of action and pharmacodynamic effects

Fulvestrant is a competitive estrogen receptor (ER) antagonist with an affinity comparable to estradiol. Fulvestrant blocks the trophic actions of estrogens without any partial agonist (estrogen-like) activity. The mechanism of action is associated with downregulation of estrogen receptor protein levels. Clinical studies in postmenopausal women with primary breast cancer have shown that fulvestrant significantly downregulates ER protein in ER positive tumours compared with placebo. There was also a significant decrease in progesterone receptor expression consistent with a lack of intrinsic estrogen agonist effects. It has also been shown that fulvestrant 500 mg downregulates ER and the proliferation marker Ki67, to a greater degree than fulvestrant 250 mg in breast tumours in postmenopausal neoadjuvant setting.

#### Clinical efficacy and safety in advanced breast cancer

##### *Monotherapy*

A Phase 3 clinical study was completed in 736 postmenopausal women with advanced breast cancer who had disease recurrence on or after adjuvant endocrine therapy or progression following endocrine therapy for advanced disease. The study included 423 patients whose disease had recurred or progressed during antiestrogen therapy (AE subgroup) and 313 patients whose disease had recurred or progressed during aromatase inhibitor therapy (AI subgroup). This study compared the efficacy and safety of Faslodex 500 mg (n=362) with Faslodex 250 mg (n=374). Progression-free survival (PFS) was the primary endpoint; key secondary efficacy endpoints included objective response rate (ORR), clinical benefit rate (CBR) and overall survival (OS). Efficacy results for the CONFIRM study are summarized in Table 3.



**Table 3 Summary of results of the primary efficacy endpoint (PFS) and key secondary efficacy endpoints in the CONFIRM study**

Variable	Type of estimate; treatment comparison	Faslodex 500 mg (N=362)	Faslodex 250 mg (N=374)	Comparison between groups (Faslodex 500 mg/Faslodex 250 mg)		
				Hazard ratio	95% CI	p-value
<b>PFS</b>	<b>K-M median in months; hazard ratio</b>					
All Patients		6.5	5.5	0.80	0.68, 0.94	0.006
-AE subgroup (n=423)		8.6	5.8	0.76	0.62, 0.94	0.013
-AI subgroup (n=313) <sup>a</sup>		5.4	4.1	0.85	0.67, 1.08	0.195
<b>OS<sup>b</sup></b>	<b>K-M median in months; hazard ratio</b>					
All Patients		26.4	22.3	0.81	0.69, 0.96	0.016 <sup>c</sup>
-AE subgroup (n=423)		30.6	23.9	0.79	0.63, 0.99	0.038 <sup>c</sup>
-AI subgroup (n=313) <sup>a</sup>		24.1	20.8	0.86	0.67, 1.11	0.241 <sup>c</sup>
Variable	Type of estimate; treatment comparison	Faslodex 500 mg (N=362)	Faslodex 250 mg (N=374)	Comparison between groups (Faslodex 500 mg/Faslodex 250 mg)		
				Absolute difference in %	95% CI	
<b>ORR<sup>d</sup></b>	<b>% of patients with OR; absolute difference in %</b>					
All Patients		13.8	14.6	-0.8	-5.8, 6.3	
-AE subgroup (n=296)		18.1	19.1	-1.0	-8.2, 9.3	
-AI subgroup (n=205) <sup>a</sup>		7.3	8.3	-1.0	-5.5, 9.8	
<b>CBR<sup>e</sup></b>	<b>% of patients with CB; absolute difference in %</b>					
All Patients		45.6	39.6	6.0	-1.1, 13.3	
-AE subgroup (n=423)		52.4	45.1	7.3	-2.2, 16.6	
-AI subgroup (n=313) <sup>a</sup>		36.2	32.3	3.9	-6.1, 15.2	

<sup>a</sup> Faslodex is indicated in patients whose disease had recurred or progressed on an antiestrogen therapy. The results in the AI subgroup are inconclusive.

<sup>b</sup> OS is presented for the final survival analyses at 75% maturity.

<sup>c</sup> Nominal p-value with no adjustments made for multiplicity between the initial overall survival analyses at 50% maturity and the updated survival analyses at 75% maturity.

<sup>d</sup> ORR was assessed in patients who were evaluable for response at baseline (i.e. those with measurable disease at baseline: 240 patients in the Faslodex 500 mg group and 261 patients in the Faslodex 250 mg group).

<sup>e</sup> Patients with a best objective response of complete response, partial response or stable disease  $\geq 24$  weeks. PFS: Progression-free survival; ORR: Objective response rate; OR: Objective response; CBR: Clinical benefit rate; CB: Clinical benefit; OS: Overall survival; K-M: Kaplan-Meier; CI: Confidence interval; AI: Aromatase inhibitor; AE: Antiestrogen.

A Phase 3, randomised, double-blind, double-dummy, multicentre study of Faslodex 500 mg versus anastrozole 1 mg was conducted in postmenopausal women with ER-positive and/or PgR-positive locally advanced or metastatic breast cancer who had not previously been treated with any hormonal therapy. A total of 462 patients were randomised 1:1 sequentially to receive either fulvestrant 500 mg or anastrozole 1 mg.

Randomisation was stratified by disease setting (locally advanced or metastatic), prior chemotherapy for advanced disease, and measurable disease.

The primary efficacy endpoint of the study was investigator assessed progression-free survival (PFS) evaluated according to RECIST 1.1 (Response Evaluation Criteria in Solid Tumours). Key secondary efficacy endpoints included overall survival (OS) and objective response rate (ORR).

Patients enrolled in this study had a median age of 63 years (range 36-90). The majority of patients (87.0%) had metastatic disease at baseline. Fifty-five percent (55.0%) of patients had visceral metastasis at baseline. A total of 17.1% of patients received a prior chemotherapy regimen for advanced disease; 84.2% of patients had measurable disease.

Consistent results were observed across the majority of pre-specified patient subgroups. For the subgroup of patients with disease limited to non-visceral metastasis (n=208), the HR was 0.592 (95% CI: 0.419, 0.837) for the Faslodex arm compared to the anastrozole arm. For the subgroup of patients with visceral metastasis (n=254), the HR was 0.993 (95% CI: 0.740, 1.331) for the Faslodex arm compared to the anastrozole arm. The efficacy results of the FALCON study are presented in Table 4 and Figure 1.


**Table 4 Summary of results of the primary efficacy endpoint (PFS) and key secondary efficacy endpoints (Investigator Assessment, Intent-To-Treat Population) — FALCON study**

	<b>Faslodex 500 mg (N=230)</b>	<b>Anastrozole 1 mg (N=232)</b>
<b>Progression-Free Survival</b>		
<b>Number of PFS Events (%)</b>	<b>143 (62.2%)</b>	<b>166 (71.6%)</b>
<b>PFS Hazard Ratio (95% CI) and p-value</b>	<b>HR 0.797 (0.637 - 0.999) p = 0.0486</b>	
<b>PFS Median [months (95% CI)]</b>	<b>16.6 (13.8, 21.0)</b>	<b>13.8 (12.0, 16.6)</b>
<b>Number of OS Events*</b>	<b>67 (29.1%)</b>	<b>75 (32.3%)</b>
<b>OS Hazard Ratio (95% CI) and p-value</b>	<b>HR 0.875 (0.629 – 1.217) p = 0.4277</b>	
<b>ORR**</b>	<b>89 (46.1%)</b>	<b>88 (44.9%)</b>
<b>ORR Odds Ratio (95% CI) and p-value</b>	<b>OR 1.074 (0.716 – 1.614) p = 0.7290</b>	
<b>Median DoR (months)</b>	<b>20.0</b>	<b>13.2</b>
<b>CBR</b>	<b>180 (78.3%)</b>	<b>172 (74.1%)</b>
<b>CBR Odds Ratio (95% CI) and p-value</b>	<b>OR 1.253 (0.815 – 1.932) p = 0.3045</b>	

\*(31% maturity)-not final OS analysis

\*\*for patients with measurable disease

**Figure 1 Kaplan-Meier Plot of Progression-Free Survival (Investigator Assessment, Intent-To-Treat Population) – FALCON Study**



Two Phase 3 clinical studies were completed in a total of 851 postmenopausal women with advanced breast cancer who had disease recurrence on or after adjuvant endocrine therapy or progression following endocrine therapy for advanced disease. Seventy seven percent (77%) of the study population had estrogen receptor positive breast cancer. These studies compared the safety and efficacy of monthly administration of Faslodex 250 mg versus the daily administration of 1 mg anastrozole (aromatase inhibitor). Overall, Faslodex at the 250 mg monthly dose was at least as effective as anastrozole in terms of progression-free survival, objective response, and time to death. There were no statistically significant differences in any of these endpoints between the two treatment groups. Progression-free survival was the primary endpoint. Combined analysis of both studies showed that 83% of patients who received Faslodex progressed, compared with 85% of patients who received anastrozole. Combined analysis of both studies showed the hazard ratio of Faslodex 250 mg to anastrozole for progression-free survival was 0.95 (95% CI 0.82 to 1.10). The objective response rate for Faslodex 250 mg was 19.2% compared with 16.5% for anastrozole. The median time to death was 27.4 months for patients treated with Faslodex and 27.6 months for patients treated with anastrozole. The hazard ratio of Faslodex 250 mg to anastrozole for time to death was 1.01 (95% CI 0.86 to 1.19).

#### *Combination therapy with palbociclib*

A Phase 3, international, randomised, double-blind, parallel-group, multicentre study of Faslodex 500 mg plus palbociclib 125 mg versus Faslodex 500 mg plus placebo was conducted in women with HR-positive, HER2-negative locally advanced breast cancer not amenable to resection or radiation therapy with curative intent or metastatic breast cancer, regardless of their menopausal status, whose disease progressed after prior endocrine therapy in the (neo) adjuvant or metastatic setting.

A total of 521 pre/peri- and postmenopausal women who had progressed on or within 12 months from completion of adjuvant endocrine therapy on or within 1 month from prior endocrine therapy for advanced disease, were randomised 2:1 to Faslodex plus palbociclib or Faslodex plus placebo and stratified by documented sensitivity to prior hormonal therapy, menopausal status at study entry (pre/peri- versus postmenopausal), and presence of visceral metastases. Pre/perimenopausal women received the LHRH agonist goserelin. Patients with advanced/metastatic, symptomatic, visceral spread, that were at risk of life-threatening complications in the short term (including patients with

massive uncontrolled effusions [pleural, pericardial, peritoneal], pulmonary lymphangitis, and over 50% liver involvement), were not eligible for enrolment into the study.

Patients continued to receive assigned treatment until objective disease progression, symptomatic deterioration, unacceptable toxicity, death, or withdrawal of consent, whichever occurred first. Crossover between treatment arms was not allowed.

Patients were well matched for baseline demographics and prognostic characteristics between the Faslodex plus palbociclib arm and the Faslodex plus placebo arm. The median age of patients enrolled in this study was 57 years (range 29, 88). In each treatment arm the majority of patients were White, had documented sensitivity to prior hormonal therapy, and were postmenopausal. Approximately 20% of patients were pre/perimenopausal. All patients had received prior systemic therapy and most patients in each treatment arm had received a previous chemotherapy regimen for their primary diagnosis. More than half (62%) had an ECOG PS of 0, 60% had visceral metastases, and 60% had received more than 1 prior hormonal regimen for their primary diagnosis.

The primary endpoint of the study was investigator-assessed PFS evaluated according to RECIST 1.1. Supportive PFS analyses were based on an Independent Central Radiology Review. Secondary endpoints included OR, CBR, OS, safety, and time-to-deterioration (TTD) in pain endpoint.

The study met its primary endpoint of prolonging investigator-assessed PFS at the interim analysis conducted on 82% of the planned PFS events; the results crossed the pre-specified Haybittle-Peto efficacy boundary ( $\alpha=0.00135$ ), demonstrating a statistically significant prolongation in PFS and a clinically meaningful treatment effect. A more mature update of efficacy data is reported in Table 5.


**Table 5 Efficacy results – PALOMA3 study (Investigator assessment, intent-to-treat population)**

	Updated Analysis (23 October 2015 cut-off)	
	Faslodex plus palbociclib (N=347)	Faslodex plus placebo (N=174)
Progression-Free Survival		
Median [months (95% CI)]	11.2 (9.5, 12.9)	4.6 (3.5, 5.6)
Hazard ratio (95% CI) and p-value	0.497 (0.398, 0.620), p <0.000001	
Secondary end points*		
OR [% (95% CI)]	26.2 (21.7, 31.2)	13.8 (9.0, 19.8)
OR (measurable disease) [% (95% CI)]	33.7 (28.1, 39.7)	17.4 (11.5, 24.8)
DOR (measurable disease) [months (95% CI)]	9.2 (7.2, 10.4)	7.4 (3.9, NE)
CBR [% (95% CI)]	68.0 (62.8, 72.9)	39.7 (32.3, 47.3)

\*Response endpoints based on confirmed and unconfirmed responses.

N=number of patients; CI=confidence interval; NE=not estimable; OR=objective response; CBR=clinical benefit response; DOR=duration of response

**Figure 2. Kaplan-Meier plot of progression-free survival (investigator assessment, intent-to-treat population) – PALOMA3 study**



FUL=fulvestrant; PAL=palbociclib; PCB=placebo.

A reduction in the risk of disease progression or death in the Faslodex plus palbociclib arm was observed in all individual patient subgroups defined by stratification factors and baseline characteristics. This was evident for pre/perimenopausal women (HR of 0.46 [95% CI: 0.28, 0.75]) and postmenopausal women (HR of 0.52 [95% CI: 0.40, 0.66]) and patients with visceral site of metastatic disease (HR of 0.50 [95% CI: 0.38, 0.65]) and non-visceral site of metastatic disease (HR of 0.48 [95% CI: 0.33, 0.71]). Benefit was also observed regardless of lines of prior therapy in the metastatic setting, whether 0 (HR of 0.59 [95% CI: 0.37, 0.93]), 1 (HR of 0.46 [95% CI: 0.32, 0.64]), 2 (HR of 0.48 [95% CI: 0.30, 0.76]), or  $\geq 3$  lines (HR of 0.59 [95% CI: 0.28, 1.22]). Additional efficacy measures (OR and TTR) assessed in the sub-groups of patients with or without visceral disease are displayed in Table 6.

**Table 6 Efficacy results in visceral and non-visceral disease from PALOMA3 study (intent-to-treat population)**

	Visceral Disease		Non-visceral Disease	
	Faslodex plus palbociclib (N=206)	Faslodex plus placebo (N=105)	Faslodex plus palbociclib (N=141)	Faslodex plus placebo (N=69)
<b>OR [% (95% CI)]</b>	35.0 (28.5, 41.9)	13.3 (7.5, 21.4)	13.5 (8.3, 20.2)	14.5 (7.2, 25.0)
<b>TTR*, Median [months (range)]</b>	3.8 (3.5, 16.7)	5.4 (3.5, 16.7)	3.7 (1.9, 13.7)	3.6 (3.4, 3.7)

\*Response results based on confirmed and unconfirmed responses.

N=number of patients; CI=confidence interval; OR= objective response; TTR=time to first tumour response.

Patient-reported symptoms were assessed using the European Organization for Research and Treatment of Cancer (EORTC) quality of life questionnaire (QLQ)-C30 and its Breast Cancer Module (EORTC QLQ-BR23). A total of 335 patients in the Faslodex plus palbociclib arm and 166 patients in the Faslodex plus placebo arm completed the questionnaire at baseline and at least 1 post-baseline visit.

Time-to-Deterioration was pre-specified as time between baseline and first occurrence of  $\geq 10$  points increase from baseline in pain symptom scores. Addition of palbociclib to Faslodex resulted in a symptom benefit by significantly delaying Time-to-Deterioration in pain symptom compared with Faslodex plus placebo (median 8.0 months versus 2.8 months; HR of 0.64 [95% CI: 0.49, 0.85];  $p < 0.001$ ).

#### *Effects on the postmenopausal endometrium*

Preclinical data do not suggest a stimulatory effect of fulvestrant on the postmenopausal endometrium (see section 5.3). A 2-week study in healthy postmenopausal volunteers treated with 20  $\mu\text{g}$  per day ethinylestradiol showed that pretreatment with Faslodex 250 mg resulted in significantly reduced stimulation of the postmenopausal endometrium, compared to pre-treatment with placebo, as judged by ultrasound measurement of endometrium thickness.

Neoadjuvant treatment for up to 16 weeks in breast cancer patients treated with either Faslodex 500 mg or Faslodex 250 mg did not result in clinically significant changes in endometrial thickness, indicating a lack of agonist effect. There is no evidence of adverse endometrial effects in the breast cancer patients studied. No data are available regarding endometrial morphology.

In two short-term studies (1 and 12 weeks) in premenopausal patients with benign gynaecologic disease, no significant differences in endometrial thickness were observed by ultrasound measurement between fulvestrant and placebo groups.

#### *Effects on bone*

There are no long-term data on the effect of fulvestrant on bone. Neoadjuvant treatment for up to 16 weeks in breast cancer patients with either Faslodex 500 mg or Faslodex 250 mg did not result in clinically significant changes in serum bone-turnover markers.

#### Paediatric population

Faslodex is not indicated for use in children. The European Medicines Agency has waived the obligation to submit the results of studies with Faslodex in all subsets of the paediatric population in breast cancer (see section 4.2 for information on paediatric use).

An open-label Phase 2 study investigated the safety, efficacy and pharmacokinetics of fulvestrant in 30 girls aged 1 to 8 years with Progressive Precocious Puberty associated with McCune Albright Syndrome (MAS). The paediatric patients received 4 mg/kg monthly intramuscular dose of fulvestrant. This 12-month study investigated a range of MAS endpoints and showed a reduction in the frequency of vaginal bleeding and a reduction in the rate of bone age advancement. The steady-state trough concentrations of fulvestrant in children in this study were consistent with that in adults (see section 5.2). There were no new safety concerns arising from this small study, but 5-year data are yet not available.

## **5.2 Pharmacokinetic properties**

### Absorption

After administration of Faslodex long-acting intramuscular injection, fulvestrant is slowly absorbed and maximum plasma concentrations ( $C_{\text{max}}$ ) are reached after about 5 days. Administration of Faslodex 500 mg regimen achieves exposure levels at, or close to, steady state within the first month of dosing (mean [CV]: AUC 475 [33.4%] ng.days/ml,  $C_{\text{max}}$  25.1 [35.3%] ng/ml,  $C_{\text{min}}$  16.3 [25.9%] ng/ml, respectively). At steady state, fulvestrant plasma concentrations are maintained within a relatively narrow range with up to an approximately 3-fold difference between maximum and trough concentrations. After intramuscular administration, the exposure is approximately dose-proportional in the dose range 50 to 500 mg.

### Distribution

Fulvestrant is subject to extensive and rapid distribution. The large apparent volume of distribution at steady state ( $V_{\text{dss}}$ ) of approximately 3 to 5 l/kg suggests that distribution is largely extravascular.

Fulvestrant is highly (99%) bound to plasma proteins. Very low density lipoprotein (VLDL), low density lipoprotein (LDL), and high density lipoprotein (HDL) fractions are the major binding components. No interaction studies were conducted on competitive protein binding. The role of sex hormone-binding globulin (SHBG) has not been determined.

#### Biotransformation

The metabolism of fulvestrant has not been fully evaluated, but involves combinations of a number of possible biotransformation pathways analogous to those of endogenous steroids. Identified metabolites (includes 17-ketone, sulphone, 3-sulphate, 3- and 17-glucuronide metabolites) are either less active or exhibit similar activity to fulvestrant in antiestrogen models. Studies using human liver preparations and recombinant human enzymes indicate that CYP3A4 is the only P450 isoenzyme involved in the oxidation of fulvestrant; however, non-P450 routes appear to be more predominant *in vivo*. *In vitro* data suggest that fulvestrant does not inhibit CYP450 isoenzymes.

#### Elimination

Fulvestrant is eliminated mainly in metabolised form. The major route of excretion is via the faeces, with less than 1% being excreted in the urine. Fulvestrant has a high clearance,  $11 \pm 1.7$  ml/min/kg, suggesting a high hepatic extraction ratio. The terminal half-life ( $t_{1/2}$ ) after intramuscular administration is governed by the absorption rate and was estimated to be 50 days.

#### Special populations

In a population pharmacokinetic analysis of data from Phase 3 studies, no difference in fulvestrant's pharmacokinetic profile was detected with regard to age (range 33 to 89 years), weight (40-127 kg) or race.

#### *Renal impairment*

Mild to moderate impairment of renal function did not influence the pharmacokinetics of fulvestrant to any clinically relevant extent.

#### *Hepatic impairment*

The pharmacokinetics of fulvestrant has been evaluated in a single-dose clinical study conducted in women with mild to moderate hepatic impairment (Child-Pugh class A and B). A high dose of a shorter duration intramuscular injection formulation was used. There was up to about 2.5-fold increase in AUC in women with hepatic impairment compared to healthy subjects. In patients administered Faslodex, an increase in exposure of this magnitude is expected to be well tolerated. Women with severe hepatic impairment (Child-Pugh class C) were not evaluated.

#### *Paediatric population*

The pharmacokinetics of fulvestrant has been evaluated in a clinical study conducted in 30 girls with Progressive Precocious Puberty associated with McCune Albright Syndrome (see section 5.1). The paediatric patients were aged 1 to 8 years and received 4 mg/kg monthly intramuscular dose of fulvestrant. The geometric mean (standard deviation) steady state trough concentration ( $C_{\min,ss}$ ) and  $AUC_{ss}$  was 4.2 (0.9) ng/mL and 3680 (1020) ng\*hr/mL, respectively. Although the data collected were limited, the steady-state trough concentrations of fulvestrant in children appear to be consistent with those in adults.

### **5.3 Preclinical safety data**

The acute toxicity of fulvestrant is low.

Faslodex and other formulations of fulvestrant were well tolerated in animal species used in multiple dose studies. Local reactions, including myositis and granulomata at the injection site were attributed to the vehicle but the severity of myositis in rabbits increased with fulvestrant, compared to the saline control. In toxicity studies with multiple intramuscular doses of fulvestrant in rats and dogs, the antiestrogenic activity of fulvestrant was responsible for most of the effects seen, particularly in the

female reproductive system, but also in other organs sensitive to hormones in both sexes. Arteritis involving a range of different tissues was seen in some dogs after chronic (12 months) dosing.

In dog studies following oral and intravenous administration, effects on the cardiovascular system (slight elevations of the S-T segment of the ECG [oral], and sinus arrest in one dog [intravenous]) were seen. These occurred at exposure levels higher than in patients ( $C_{\max}$  >15 times) and are likely to be of limited significance for human safety at the clinical dose.

Fulvestrant showed no genotoxic potential.

Fulvestrant showed effects upon reproduction and embryo/foetal development consistent with its antiestrogenic activity, at doses similar to the clinical dose. In rats, a reversible reduction in female fertility and embryonic survival, dystocia and an increased incidence of foetal abnormalities including tarsal flexure were observed. Rabbits given fulvestrant failed to maintain pregnancy. Increases in placental weight and post-implantation loss of foetuses were seen. There was an increased incidence of foetal variations in rabbits (backwards displacement of the pelvic girdle and 27 pre-sacral vertebrae).

A two-year oncogenicity study in rats (intramuscular administration of Faslodex) showed increased incidence of ovarian benign granulosa cell tumours in female rats at the high dose, 10 mg/rat/15 days and an increased incidence of testicular Leydig cell tumours in males. In a two-year mouse oncogenicity study (daily oral administration) there was an increased incidence of ovarian sex cord stromal tumours (both benign and malignant) at doses of 150 and 500 mg/kg/day. At the no-effect level for these findings, systemic exposure levels (AUC) were, in rats, approximately 1.5-fold the expected human exposure levels in females and 0.8-fold in males, and in mice, approximately 0.8-fold the expected human exposure levels in both males and females. Induction of such tumours is consistent with pharmacology-related endocrine feedback alterations in gonadotropin levels caused by antiestrogens in cycling animals. Therefore these findings are not considered to be relevant to the use of fulvestrant in postmenopausal women with advanced breast cancer.

#### Environmental Risk Assessment (ERA)

Environmental risk assessment studies have shown that fulvestrant may have potential to cause adverse effects to the aquatic environment (see section 6.6).

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Ethanol (96 per cent)  
Benzyl alcohol  
Benzyl benzoate  
Castor oil refined

### **6.2 Incompatibilities**

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

4 years

### **6.4 Special precautions for storage**

Store and transport in a refrigerator (2°C - 8°C).



Temperature excursions outside 2°C - 8°C should be limited. This includes avoiding storage at temperatures exceeding 30°C, and not exceeding a 28 day period where the average storage temperature for the product is below 25°C (but above 2°C - 8°C). After temperature excursions, the product should be returned immediately to the recommended storage conditions (store and transport in a refrigerator 2°C - 8°C). Temperature excursions have a cumulative effect on the product quality and the 28 day time period must not be exceeded over the duration of the 4-year shelf life of Faslodex (see section 6.3). Exposure to temperatures below 2°C will not damage the product providing it is not stored below – 20°C.

Store the pre-filled syringe in the original package in order to protect from light.

## 6.5 Nature and contents of container

The pre-filled syringe presentation consists of:

One clear type 1 glass pre-filled syringe with polystyrene plunger rod, fitted with a tamper-evident closure, containing 5 ml Faslodex solution for injection.

A safety needle (BD SafetyGlide) for connection to the barrel is also provided.

Or

Two clear type 1 glass pre-filled syringes with polystyrene plunger rod, fitted with a tamper-evident closure, each containing 5 ml Faslodex solution for injection. Safety needles (BD SafetyGlide) for connection to each barrel are also provided.

Not all pack sizes may be marketed.

## 6.6 Special precautions for disposal and other handling

### Instructions for administration

Administer the injection according to the local guidelines for performing large volume intramuscular injections.


NOTE: Due to the proximity of the underlying sciatic nerve, caution should be taken if administering Faslodex at the dorsogluteal injection site (see section 4.4).

Warning - Do not autoclave safety needle (BD SafetyGlide Shielding Hypodermic Needle) before use. Hands must remain behind the needle at all times during use and disposal.

For each of the two syringes:


- Remove glass syringe barrel from tray and check that it is not damaged.
- Peel open the safety needle (SafetyGlide) outer packaging.
- Parenteral solutions must be inspected visually for particulate matter and discolouration prior to administration.
- Hold the syringe upright on the ribbed part (C). With the other hand, take hold of the cap (A) and carefully tilt back and forth until the cap disconnects and can be pulled off, do not twist (see Figure 1).

Figure 1




- Remove the cap (A) in a straight upward direction. To maintain sterility do not touch the syringe tip (B) (see Figure 2).

Figure 2




- Attach the safety needle to the Luer-Lok and twist until firmly seated (see Figure 3).
- Check that the needle is locked to the Luer connector before moving out of the vertical plane.
- Pull shield straight off needle to avoid damaging needle point.
- Transport filled syringe to point of administration.
- Remove needle sheath.
- Expel excess gas from the syringe.

Figure 3



- Administer intramuscularly slowly (1-2 minutes/injection) into the buttock (gluteal area). For user convenience, the needle bevel-up position is oriented to the lever arm (see Figure 4).

Figure 4



- After injection, immediately apply a single-finger stroke to the activation assisted lever arm to activate the shielding mechanism (see Figure 5).  
NOTE: Activate away from self and others. Listen for click and visually confirm needle tip is fully covered.

Figure 5



### Disposal

Pre-filled syringes are for single use **only**.

This medicine may pose a risk to the aquatic environment. Any unused medicinal product or waste material should be disposed of in accordance with local requirements (see section 5.3).

## 7. MARKETING AUTHORISATION HOLDER

AstraZeneca UK Limited  
Charter Way,  
Macclesfield,  
Cheshire  
SK10 2NA  
United Kingdom

**8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/03/269/001

EU/1/03/269/002

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 10 March 2004

Date of latest renewal: 10 March 2009

**10. DATE OF REVISION OF THE TEXT**

<{MM/YYYY}>

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## **ANNEX II**

- A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**

### Name and address of the manufacturer responsible for batch release

AstraZeneca UK Limited  
Silk Road Business Park,  
Macclesfield, SK10 2NA  
United Kingdom

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to medical prescription.

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

### **• Periodic Safety Update Reports**

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

### **• Risk Management Plan (RMP)**

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING AND THE IMMEDIATE PACKAGING**

**CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Faslodex 250 mg solution for injection.  
fulvestrant

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

One pre-filled syringe contains 250 mg fulvestrant in 5 ml solution

**3. LIST OF EXCIPIENTS**

Ethanol (96 per cent), benzyl alcohol, benzyl benzoate and castor oil refined. See the package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Solution for injection in a pre-filled syringe.

1 pre-filled syringe (5 ml)  
1 safety needle  
2 pre-filled syringes (5 ml each)  
2 safety needles

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Intramuscular use.  
For single use only.  
For full instructions on the administration of Faslodex and the use of the safety needle see enclosed, Instructions for administration.  
Two syringes must be administered to receive the 500 mg recommended monthly dose.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**



EXP

**9. SPECIAL STORAGE CONDITIONS**

Store and transport in a refrigerator.

Store the pre-filled syringe in the original package in order to protect from light. See package leaflet for information on temperature excursions.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

AstraZeneca UK Limited  
Charter Way,  
Macclesfield,  
Cheshire  
SK10 2NA  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/03/269/001

EU/1/03/269/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC:  
SN:  
NN:

<b>MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS</b> <b>PRE-FILLED SYRINGE LABEL</b>
--

<b>1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION</b>
--

Faslodex 250 mg solution for injection  
fulvestrant  
IM use

<b>2. METHOD OF ADMINISTRATION</b>
------------------------------------

<b>3. EXPIRY DATE</b>
-----------------------

EXP

<b>4. BATCH NUMBER</b>
------------------------

Lot

<b>5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT</b>
--

5 ml

<b>6. OTHER</b>
-----------------

## **B. PACKAGE LEAFLET**

## **Package leaflet: Information for the user**

### **Faslodex 250 mg solution for injection** Fulvestrant

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist, or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist, or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Faslodex is and what it is used for
2. What you need to know before you use Faslodex
3. How to use Faslodex
4. Possible side effects
5. How to store Faslodex
6. Contents of the pack and other information

#### **1. What Faslodex is and what it is used for**

Faslodex contains the active substance fulvestrant, which belongs to the group of estrogen blockers. Estrogens, a type of female sex hormones, can in some cases be involved in the growth of breast cancer.

Faslodex is used either:

- alone, to treat postmenopausal women with a type of breast cancer called estrogen receptor positive breast cancer that is locally advanced or has spread to other parts of the body (metastatic), or
- in combination with palbociclib to treat women with a type of breast cancer called hormone receptor-positive, human epidermal growth factor receptor 2-negative breast cancer, that is locally advanced or has spread to other parts of the body (metastatic). Women who have not reached menopause will also be treated with a medicine called a luteinizing hormone releasing hormone (LHRH) agonist.

When Faslodex is given in combination with palbociclib, it is important that you also read the package leaflet for palbociclib. If you have any questions about palbociclib, please ask your doctor.

#### **2. What you need to know before you use Faslodex**

##### **Do not use Faslodex:**

- if you are allergic to fulvestrant or to any of the other ingredients of this medicine (listed in section 6)
- if you are pregnant or breast-feeding
- if you have severe liver problems

##### **Warnings and precautions**

Talk to your doctor or pharmacist or nurse before using Faslodex if any of these apply to you:

- kidney or liver problems
- low numbers of platelets (which help blood clotting) or bleeding disorders

- previous problems with blood clots
- osteoporosis (loss of bone density)
- alcoholism

### **Children and adolescents**

Faslodex is not indicated in children and adolescents under 18 years.

### **Other medicines and Faslodex**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

In particular, you should tell your doctor if you are using anticoagulants (medicines to prevent blood clots).

### **Pregnancy and breast-feeding**

You must not use Faslodex if you are pregnant. If you can become pregnant, you should use effective contraception while being treated with Faslodex.

You must not breast-feed while on treatment with Faslodex.

### **Driving and using machines**

Faslodex is not expected to affect your ability to drive or use machines. However, if you feel tired after treatment do not drive or use machines.

**Faslodex contains 10% w/v ethanol (alcohol)**, i.e. up to 1000 mg per dose, equivalent to 20 ml beer or 8 ml wine per dose.

Harmful for those suffering from alcoholism.

To be taken into account in pregnant or breast-feeding women, children and high-risk groups such as patients with liver disease, or epilepsy.

## **3. How to use Faslodex**

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is 500 mg fulvestrant (two 250 mg/5 ml injections) given once a month, with an additional 500 mg dose given 2 weeks after the initial dose.

Your doctor or nurse will give you Faslodex as a slow intramuscular injection, one into each of your buttocks.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**You may need immediate medical treatment if you experience any of the following side effects:**

- Allergic (hypersensitivity) reactions, including swelling of the face, lips, tongue and/or throat that may be signs of anaphylactic reactions
- Thromboembolism (increased risk of blood clots)\*
- Inflammation of the liver (hepatitis)
- Liver failure

**Tell your doctor, pharmacist, or nurse if you notice any of the following side effects:**

**Very common side effects** (may affect more than 1 in 10 people)

- Injection site reactions, such as pain and/or inflammation
- Abnormal levels of liver enzymes (in blood tests)\*
- Nausea (feeling sick)
- Weakness, tiredness\*
- Joint and musculoskeletal pain
- Hot flushes
- Skin rash
- Allergic (hypersensitivity) reactions, including swelling of the face, lips, tongue and/or throat

**All other side effects:**

**Common side effects** (may affect up to 1 in 10 people)

- Headache
- Vomiting, diarrhoea, or loss of appetite\*
- Urinary tract infections
- Back pain\*
- Increase of bilirubin (bile pigment produced by the liver)
- Thromboembolism (increased risk of blood clots)\*
- Decreased levels of platelets (thrombocytopenia)
- Vaginal bleeding
- Lower back pain irradiating to leg on one side (sciatica)
- Sudden weakness, numbness, tingling, or loss of movement in your leg, especially on only one side of your body, sudden problems with walking or balance (peripheral neuropathy)

**Uncommon side effects** (may affect up to 1 in 100 people)

- Thick, whitish vaginal discharge and candidiasis (infection)
- Bruising and bleeding at the site of injection
- Increase of gamma-GT, a liver enzyme seen in a blood test
- Inflammation of the liver (hepatitis)
- Liver failure
- Numbness, tingling and pain
- Anaphylactic reactions

\* Includes side effects for which the exact role of Faslodex cannot be assessed due to the underlying disease.

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Faslodex**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton or syringe labels after EXP. The expiry date refers to the last day of that month.

Store and transport in a refrigerator (2°C - 8°C).

Temperature excursions outside 2°C - 8°C should be limited. This includes avoiding storage at temperatures exceeding 30°C, and not exceeding a 28-day period where the average storage temperature for the product is below 25°C (but above 2°C - 8°C). After temperature excursions, the product should be returned immediately to the recommended storage conditions (store and transport in a refrigerator 2°C - 8°C). Temperature excursions have a cumulative effect on the product quality and the 28-day time period must not be exceeded over the duration of the 4-year shelf life of Faslodex. Exposure to temperatures below 2°C will not damage the product providing it is not stored below -20°C.

Keep the pre-filled syringe in the original package, in order to protect from light.

Your healthcare professional will be responsible for the correct storage, use and disposal of Faslodex.

This medicine may pose a risk to the aquatic environment. Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Faslodex contains**

- The active substance is fulvestrant. Each pre-filled syringe (5 ml) contains 250 mg fulvestrant.
- The other ingredients (excipients) are ethanol (96 per cent), benzyl alcohol, benzyl benzoate and castor oil refined.

### **What Faslodex looks like and contents of the pack**

Faslodex is a clear, colourless to yellow, viscous solution in a pre-filled syringe fitted with a tamper-evident closure, containing 5 ml solution for injection. Two syringes must be administered to receive the 500 mg recommended monthly dose.

Faslodex has 2 pack presentations, either a pack containing 1 glass pre-filled syringe or a pack containing 2 glass pre-filled syringes. Safety needles (BD SafetyGlide) for connection to each barrel are also provided.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder**

AstraZeneca UK Limited  
Charter Way,  
Macclesfield,  
Cheshire  
SK10 2NA  
United Kingdom

### **Manufacturer**

AstraZeneca UK Limited  
Silk Road Business Park  
Macclesfield  
Cheshire  
SK10 2NA  
United Kingdom

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

**Lietuva**



AstraZeneca S.A./N.V.  
Tel: +32 2 370 48 11

**България**  
АстраЗенека България ЕООД  
Тел.: +359 24455000

**Česká republika**  
AstraZeneca Czech Republic s.r.o.  
Tel: +420 222 807 111

**Danmark**  
AstraZeneca A/S  
Tlf: +45 43 66 64 62

**Deutschland**  
AstraZeneca GmbH  
Tel: +49 41 03 7080

**Eesti**  
AstraZeneca  
Tel: +372 6549 600

**Ελλάδα**  
AstraZeneca A.E.  
Τηλ: + 30 210 6871500

**España**  
AstraZeneca Farmacéutica Spain, S.A.  
Tel: +34 91 301 91 00

**France**  
AstraZeneca  
Tél: +33 1 41 29 40 00

**Hrvatska**  
AstraZeneca d.o.o.  
Tel: +385 1 4628 000

**Ireland**  
AstraZeneca Pharmaceuticals (Ireland) DAC  
Tel: +353 1609 7100

**Ísland**  
Vistor hf.  
Sími: +354 535 7000

**Italia**  
AstraZeneca S.p.A.  
Tel: +39 02 9801 1

**Κύπρος**  
Αλέκτωρ Φαρμακευτική Λτδ  
Τηλ: +357 22490305

UAB AstraZeneca Lietuva  
Tel: +370 5 2660550

**Luxembourg/Luxemburg**  
AstraZeneca S.A./N.V.  
Tél/Tel: +32 2 370 48 11

**Magyarország**  
AstraZeneca Kft  
Tel: +36 1 883 6500

**Malta**  
Associated Drug Co. Ltd  
Tel: +356 2277 8000

**Nederland**  
AstraZeneca BV  
Tel: +31 79 363 2222

**Norge**  
AstraZeneca AS  
Tlf: +47 21 00 64 00

**Österreich**  
AstraZeneca Österreich GmbH  
Tel: +43 1 711 31 0

**Polska**  
AstraZeneca Pharma Poland Sp. z o.o.  
Tel.: +48 22 245 73 00

**Portugal**  
AstraZeneca Produtos Farmacêuticos, Lda.  
Tel: +351 21 434 61 00

**România**  
AstraZeneca Pharma SRL  
Tel: +40 21 317 60 41

**Slovenija**  
AstraZeneca UK Limited  
Tel: +386 1 51 35 600

**Slovenská republika**  
AstraZeneca AB, o.z.  
Tel: +421 2 5737 7777

**Suomi/Finland**  
AstraZeneca Oy  
Puh/Tel: +358 10 23 010

**Sverige**  
AstraZeneca AB  
Tel: +46 8 553 26 000

**Latvija**  
SIA AstraZeneca Latvija  
Tel: +371 67377100

**United Kingdom**  
AstraZeneca UK Ltd  
Tel: +44 1582 836 836

**This leaflet was last revised in** <{MM/YYYY}>

### Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>

---

### The following information is intended for healthcare professionals only:

Faslodex 500 mg (2 x 250 mg/5 ml solution for injection) should be administered using two pre-filled syringes, see section 3.

#### Instructions for administration

Warning - Do not autoclave safety needle (BD SafetyGlide Shielding Hypodermic Needle) before use. Hands must remain behind the needle at all times during use and disposal.

For each of the two syringes:

- Remove glass syringe barrel from tray and check that it is not damaged.
- Peel open the safety needle (SafetyGlide) outer packaging.
- Parenteral solutions must be inspected visually for particulate matter and discolouration prior to administration.
- Hold the syringe upright on the ribbed part (C). With the other hand, take hold of the cap (A) and carefully tilt back and forth until the cap disconnects and can be pulled off, do not twist (see Figure 1).
- Remove the cap (A) in a straight upward direction. To maintain sterility do not touch the syringe tip (B) (see Figure 2).

Figure 1






Figure 2




- Attach the safety needle to the Luer-Lok and twist until firmly seated (see Figure 3).
- Check that the needle is locked to the Luer connector before moving out of the vertical plane.
- Pull shield straight off needle to avoid damaging needle point.
- Transport filled syringe to point of administration.
- Remove needle sheath.
- Expel excess gas from the syringe.

Figure 3



- Administer intramuscularly slowly (1-2 minutes/injection) into the buttock (gluteal area). For user convenience, the needle bevel-up position is oriented to the lever arm (see Figure 4).

Figure 4



- After injection, immediately apply a single-finger stroke to the activation assisted lever arm to activate the shielding mechanism (see Figure 5).

NOTE: Activate away from self and others. Listen for click and visually confirm needle tip is fully covered.

Figure 5



### Disposal

Pre-filled syringes are for single use **only**.

This medicine may pose a risk to the aquatic environment. Any unused medicinal product or waste material should be disposed of in accordance with local requirements.