

ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Faslodex 250 mg solution for injection

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One pre-filled syringe contains 250 mg fulvestrant in 5 ml solution.

For a full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Solution for injection.

Clear, colourless to yellow, viscous solution.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Faslodex is indicated for the treatment of postmenopausal women with estrogen receptor positive, locally advanced or metastatic breast cancer for disease relapse on or after adjuvant anti-estrogen therapy, or disease progression on therapy with an anti-estrogen.

4.2 Posology and method of administration

Posology

Adult females (including the elderly)

The recommended dose is 500 mg at intervals of one month, with an additional 500 mg dose given two weeks after the initial dose.

Paediatric patients

Faslodex is not recommended for use in children or adolescents, as safety and efficacy have not been established in this age group.

Renal impairment

No dose adjustments are recommended for patients with mild to moderate renal impairment (creatinine clearance ≥ 30 ml/min). Safety and efficacy have not been evaluated in patients with severe renal impairment (creatinine clearance < 30 ml/min), and, therefore, caution is recommended in these patients (see section 4.4).

Hepatic impairment

No dose adjustments are recommended for patients with mild to moderate hepatic impairment. However, as fulvestrant exposure may be increased, Faslodex should be used with caution in these patients. There are no data in patients with severe hepatic impairment (see sections 4.3, 4.4 and 5.2).

Method of administration

Faslodex should be administered as two consecutive 5 ml injections by slow intramuscular injection (1-2 minutes/injection), one in each buttock.

For detailed instructions for administration, see section 6.6.

4.3 Contraindications

Hypersensitivity to the active substance, or to any of the other excipients.

Pregnancy and lactation (see section 4.6)
Severe hepatic impairment (see sections 4.4 and 5.2).

4.4 Special warnings and precautions for use

Faslodex should be used with caution in patients with mild to moderate hepatic impairment (see sections 4.2, 4.3 and 5.2).

Faslodex should be used with caution in patients with severe renal impairment (creatinine clearance less than 30 ml/min).

Due to the intramuscular route of administration, Faslodex should be used with caution if treating patients with bleeding diatheses, thrombocytopenia or those taking anticoagulant treatment.

Thromboembolic events are commonly observed in women with advanced breast cancer and have been observed in clinical trials with Faslodex (see section 4.8). This should be taken into consideration when prescribing Faslodex to patients at risk.

There are no long-term data on the effect of fulvestrant on bone. Due to the mechanism of action of fulvestrant, there is a potential risk of osteoporosis.

4.5 Interaction with other medicinal products and other forms of interaction

A clinical interaction study with midazolam (substrate of CYP3A4) demonstrated that fulvestrant does not inhibit CYP3A4. Clinical interaction studies with rifampicin (inducer of CYP3A4) and ketoconazole (inhibitor of CYP3A4) showed no clinically relevant change in fulvestrant clearance. Dose adjustment is therefore not necessary in patients who are receiving fulvestrant and CYP3A4 inhibitors or inducers concomitantly.

4.6 Pregnancy and lactation

Faslodex is contraindicated in pregnancy (see section 4.3). Fulvestrant has been shown to cross the placenta after single intramuscular doses in rat and rabbit. Studies in animals have shown reproductive toxicity including an increased incidence of foetal abnormalities and deaths (see section 5.3). Patients of child-bearing potential should be advised to use effective contraception while on treatment. If pregnancy occurs while taking Faslodex, the patient must be informed of the potential hazard to the foetus and potential risk for loss of pregnancy.

Breast-feeding must be discontinued during treatment with Faslodex. Fulvestrant is excreted in milk in lactating rats. It is not known whether fulvestrant is excreted in human milk. Considering the potential for serious adverse reactions due to fulvestrant in breast-fed infants, use during lactation is contraindicated (see section 4.3).

4.7 Effects on ability to drive and use machines

Faslodex has no or negligible influence on the ability to drive or use machines. However, since asthenia has been reported very commonly with Faslodex, caution should be observed by those patients who experience this adverse reaction when driving or operating machinery.

4.8 Undesirable effects

This section provides information based on all adverse reactions from clinical trials, post-marketing studies or spontaneous reports. The most frequently reported adverse reactions are injection site reactions, asthenia, nausea, and increased hepatic enzymes.

The following frequency categories for adverse drug reactions (ADRs) were calculated based on the Faslodex 500 mg treatment group in pooled safety analyses of the CONFIRM (Study D6997C00002),

FINDER 1 (Study D6997C00004), FINDER 2 (Study D6997C00006), and NEWEST (Study D6997C00003) studies that compared Faslodex 500 mg with Faslodex 250 mg. The frequencies in the following table were based on all reported adverse drug reactions, regardless of the investigator assessment of causality.

Adverse reactions listed below are classified according to frequency and System Organ Class (SOC). Frequency groupings are defined according to the following convention: Very common ($\geq 1/10$), Common ($\geq 1/100$ to $< 1/10$), Uncommon ($\geq 1/1,000$ to $< 1/100$). Within each frequency grouping adverse reactions are reported in order of decreasing seriousness.

Table 1 Adverse Drug Reactions

SOC	Very common $\geq 10\%$	Common $\geq 1 - < 10\%$	Uncommon $\geq 0.1 - < 1\%$
Nervous system disorders		Headache	
Gastrointestinal disorders	Nausea	Vomiting, diarrhoea	
Infections and infestations		Urinary tract infections	
Skin and subcutaneous tissue disorders		Rash	
Musculoskeletal and connective tissue disorders		Back pain ^a	
Metabolism and nutrition disorders		Anorexia ^a	
Vascular disorders		Venous thromboembolism ^a , hot flushes	
General disorders and administration site conditions	Asthenia ^a , Injection site reactions		
Immune system disorders		Hypersensitivity reactions	
Hepatobiliary disorders	Increased Hepatic Enzymes (ALT, AST, ALP) ^a		
Reproductive system and breast disorders			Vaginal moniliasis Leukorrhea Vaginal haemorrhage

^a Includes adverse drug reactions for which the exact contribution of Faslodex cannot be assessed due to the underlying disease.

4.9 Overdose

There is no human experience of overdose. Animal studies suggest that no effects other than those related directly or indirectly to anti-estrogenic activity were evident with higher doses of fulvestrant (see section 5.3). If overdose occurs, symptomatic supportive treatment is recommended.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Anti-estrogens, ATC code: L02BA03

Mechanism of action

Fulvestrant is a competitive estrogen receptor (ER) antagonist with an affinity comparable to estradiol. Fulvestrant blocks the trophic actions of estrogens without any partial agonist (estrogen-like) activity. The mechanism of action is associated with down-regulation of estrogen receptor protein levels. Clinical trials in postmenopausal women with primary breast cancer have shown that fulvestrant significantly down-regulates ER protein in ER positive tumours compared with placebo. There was also a significant decrease in progesterone receptor expression consistent with a lack of intrinsic estrogen agonist effects.

Clinical safety and efficacy in advanced breast cancer

A phase III clinical trial was completed in 736 postmenopausal women with advanced breast cancer who had disease recurrence on or after adjuvant endocrine therapy or progression following endocrine therapy for advanced disease. This trial compared the efficacy and safety of Faslodex 500 mg (n=362) with Faslodex 250 mg (n=374). Time to progression was the primary endpoint, key secondary efficacy endpoints included objective response rate (ORR), clinical benefit rate (CBR) and overall survival (OS). Efficacy results for the CONFIRM study are summarized in Table 2.

Table 2 Summary of results of the primary efficacy endpoint (TTP) and key secondary efficacy endpoints in the CONFIRM study

Variable	Type of estimate; treatment comparison	Faslodex 500 mg (N=362)	Faslodex 250 mg (N=374)	Comparison between groups (Faslodex 500 mg/Faslodex 250 mg)		
				Hazard ratio / odds ratio	95% CI	p-value
TTP	K-M median in months; hazard ratio	6.5	5.5	0.80	0.68, 0.94	0.006
ORR ^a	% of patients with OR; odds ratio	13.8	14.6	0.94	0.57, 1.55	0.795
CBR	% of patients with CB; odds ratio	45.6	39.6	1.28	0.95, 1.71	0.100
OS	K-M median in months; hazard ratio	25.1	22.8	0.84	0.69, 1.03	0.091

^a ORR was assessed in patients who were evaluable for response at baseline (ie, those with measurable disease at baseline: 240 patients in the Faslodex 500 mg group and 261 patients in the Faslodex 250 mg group). TTP: Time to progression; ORR: Objective response rate; OR: Objective response; CBR: Clinical benefit rate; CB: Clinical benefit; OS: Overall survival; K-M: Kaplan-Meier; CI: Confidence interval.

Two phase III clinical trials were completed in a total of 851 postmenopausal women with advanced breast cancer who had disease recurrence on or after adjuvant endocrine therapy or progression following endocrine therapy for advanced disease. 77% of the study population had estrogen receptor positive breast cancer. These trials compared the safety and efficacy of monthly administration of Faslodex 250 mg versus the daily administration of 1 mg anastrozole (aromatase inhibitor). Overall, Faslodex at the 250 mg monthly dose was at least as effective as anastrozole in terms of time to progression, objective response, and time to death. There were no statistically significant differences in any of these endpoints between the two treatment groups. Time to progression was the primary

endpoint. Combined analysis of both trials showed that 83% of patients who received fulvestrant progressed, compared with 85% of patients who received anastrozole. Combined analysis of both trials showed the hazard ratio of Faslodex 250 mg to anastrozole for time to progression was 0.95 (95% CI 0.82 to 1.10). The objective response rate for Faslodex 250 mg was 19.2% compared with 16.5% for anastrozole. The median time to death was 27.4 months for patients treated with fulvestrant and 27.6 months for patients treated with anastrozole. The hazard ratio of Faslodex 250 mg to anastrozole for time to death was 1.01 (95% CI 0.86 to 1.19).

Effects on the postmenopausal endometrium

Preclinical data do not suggest a stimulatory effect of fulvestrant on the postmenopausal endometrium (see section 5.3). A 2-week study in healthy postmenopausal volunteers treated with 20 µg per day ethinylestradiol showed that pre-treatment with Faslodex 250 mg resulted in significantly reduced stimulation of the postmenopausal endometrium, compared to pre-treatment with placebo, as judged by ultrasound measurement of endometrium thickness

There are no data on the long-term effects of fulvestrant on the postmenopausal endometrium. No data are available regarding endometrial morphology.

In two short-term studies (1 and 12 weeks) with premenopausal patients with benign gynaecologic disease, no significant differences in endometrial thickness were observed by ultrasound measuring between fulvestrant and placebo groups.

Effects on bone

There are no long-term data on the effect of fulvestrant on bone.

5.2 Pharmacokinetic properties

Absorption

After administration of Faslodex long-acting intramuscular injection, fulvestrant is slowly absorbed and maximum plasma concentrations (C_{max}) are reached after about 5 days. Administration of Faslodex 500 mg regimen achieves exposure levels at, or close to, steady state within the first month of dosing (mean [CV]: AUC 475 [33.4%] ng.days/ml, C_{max} 25.1 [35.1%] ng/ml, C_{min} 16.3 [25.9%] ng/ml, respectively). At steady state, fulvestrant plasma concentrations are maintained within a relatively narrow range with up to an approximately 3-fold difference between maximum and trough concentrations. After intramuscular administration, the exposure is approximately dose-proportional in the dose range 50 to 500 mg.

Distribution

Fulvestrant is subject to extensive and rapid distribution. The large apparent volume of distribution at steady state (V_{dss}) of approximately 3 to 5 l/kg suggests that distribution is largely extravascular. Fulvestrant is highly (99%) bound to plasma proteins. Very low density lipoprotein (VLDL), low density lipoprotein (LDL), and high density lipoprotein (HDL) fractions are the major binding components. No interaction studies were conducted on competitive protein binding. The role of sex hormone-binding globulin (SHBG) has not been determined.

Metabolism

The metabolism of fulvestrant has not been fully evaluated, but involves combinations of a number of possible biotransformation pathways analogous to those of endogenous steroids. Identified metabolites (includes 17-ketone, sulphone, 3-sulphate, 3- and 17-glucuronide metabolites) are either less active or exhibit similar activity to fulvestrant in anti-estrogen models. Studies using human liver preparations and recombinant human enzymes indicate that CYP3A4 is the only P450 isoenzyme involved in the oxidation of fulvestrant; however, non-P450 routes appear to be more predominant *in vivo*. *In vitro* data suggest that fulvestrant does not inhibit CYP450 isoenzymes.

Elimination

Fulvestrant is eliminated mainly in metabolised form. The major route of excretion is via the faeces, with less than 1% being excreted in the urine. Fulvestrant has a high clearance, 11 ± 1.7 ml/min/kg, suggesting a high hepatic extraction ratio. The terminal half-life ($t_{1/2}$) after intramuscular administration is governed by the absorption rate and was estimated to be 50 days.

Special populations

In a population pharmacokinetic analysis of data from phase III studies, no difference in fulvestrant's pharmacokinetic profile was detected with regard to age (range 33 to 89 years), weight (40-127 kg) or race.

Renal impairment

Mild to moderate impairment of renal function did not influence the pharmacokinetics of fulvestrant to any clinically relevant extent.

Hepatic impairment

The pharmacokinetics of fulvestrant has been evaluated in a single-dose clinical trial conducted in subjects with mild to moderate hepatic impairment (Child-Pugh class A and B). A high dose of a shorter duration intramuscular injection formulation was used. There was up to about 2.5-fold increase in AUC in subjects with hepatic impairment compared to healthy subjects. In patients administered Faslodex, an increase in exposure of this magnitude is expected to be well tolerated. Subjects with severe hepatic impairment (Child-Pugh class C) were not evaluated.

5.3 Preclinical safety data

The acute toxicity of fulvestrant is low.

Faslodex and other formulations of fulvestrant were well tolerated in animal species used in multiple dose studies. Local reactions, including myositis and granulomatoma at the injection site were attributed to the vehicle but the severity of myositis in rabbits increased with fulvestrant, compared to the saline control. In toxicity studies with multiple intramuscular doses of fulvestrant in rats and dogs, the anti-estrogenic activity of fulvestrant was responsible for most of the effects seen, particularly in the female reproductive system, but also in other organs sensitive to hormones in both sexes.

In dog studies following oral and intravenous administration, effects on the cardiovascular system (slight elevations of the S-T segment of the ECG [oral], and sinus arrest in one dog [intravenous]) were seen. These occurred at exposure levels higher than in patients ($C_{max} > 40$ times) and are likely to be of limited significance for human safety at the clinical dose.

Fulvestrant showed no genotoxic potential.

Fulvestrant showed effects upon reproduction and embryo/foetal development consistent with its anti-estrogenic activity, at doses similar to the clinical dose. In rats, a reversible reduction in female fertility and embryonic survival, dystocia and an increased incidence of foetal abnormalities including tarsal flexure were observed. Rabbits given fulvestrant failed to maintain pregnancy. Increases in placental weight and post-implantation loss of foetuses were seen. There was an increased incidence of foetal variations in rabbits (backwards displacement of the pelvic girdle and 27 pre-sacral vertebrae).

A two-year oncogenicity study in rats (intramuscular administration of Faslodex) showed increased incidence of ovarian benign granulosa cell tumours in female rats at the high dose, 10 mg/rat/15 days and an increased incidence of testicular Leydig cell tumours in males. Induction of such tumours is consistent with pharmacology-related endocrine feedback alterations. These findings are not of clinical relevance for the use of fulvestrant in postmenopausal women with advanced breast cancer.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Ethanol (96 per cent)
Benzyl alcohol
Benzyl benzoate
Castor oil

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

6.3 Shelf life

4 years

6.4 Special precautions for storage

Store in a refrigerator (2°C - 8°C).

Store the pre-filled syringe in the original package in order to protect from light.

6.5 Nature and contents of container

BD SafetyGlide is a trademark of Becton Dickinson and Company and is CE-marked: CE 0050.

The pre-filled syringe presentation consists of:

One clear type 1 glass pre-filled syringe with polystyrene plunger rod, fitted with a tamper-evident closure, containing 5 ml Faslodex solution for injection.

A safety needle (BD SafetyGlide™) for connection to the barrel is also provided.


6.6 Special precautions for disposal and other handling

Instructions for administration

Warning - Do not autoclave safety needle (BD SafetyGlide Shielding Hypodermic Needle) before use. Hands must remain behind the needle at all times during use and disposal.

For each of the two syringes:

- Remove glass syringe barrel from tray and check that it is not damaged. Figure 1
- Break the seal of the white plastic cover on the syringe Luer connector Luer-Lok to remove the cover with the attached rubber tip cap (see Figure 1).



- Peel open the safety needle (BD SafetyGlide) outer packaging. Attach the safety needle to the Luer-Lok (see Figure 2)
- Twist until firmly seated.
- Twist to lock the needle to the Luer connector.
- Pull shield straight off needle to avoid damaging needle point.
- Transport filled syringe to point of administration.
- Remove needle sheath.
- Parenteral solutions must be inspected visually for particulate matter and discolouration prior to administration.
- Expel excess gas from the syringe.
- Administer intramuscularly slowly (1-2 minutes/injection) into the buttock. For user convenience, the needle bevel- up position is oriented to the lever arm (see Figure 3).

Figure 2



Figure 3



- After injection, immediately apply a single-finger stroke to the activation assisted lever arm to activate the shielding mechanism (see Figure 4).

NOTE: Activate away from self and others. Listen for click and visually confirm needle tip is fully covered.

Figure 4



Disposal

Pre-filled syringes are for single use **only**.

Any unused product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

AstraZeneca UK Limited

Alderley Park
Macclesfield
Cheshire
SK10 4TG
United Kingdom

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/03/269/001

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 10 March 2004

Date of last renewal: 10 March 2009

10. DATE OF REVISION OF THE TEXT

ANNEX II

- A. MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OF THE MARKETING AUTHORISATION**

A MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer responsible for batch release

AstraZeneca UK Limited
Silk Road Business Park,
Macclesfield, SK10 2NA
United Kingdom

B CONDITIONS OF THE MARKETING AUTHORISATION

• **CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE IMPOSED ON THE MARKETING AUTHORISATION HOLDER**

Medicinal product subject to medical prescription.

• **OTHER CONDITIONS**

Pharmacovigilance system

The MAH must ensure that the system of pharmacovigilance, as described in version 10.0 presented in Module 1.8.1. of the Marketing Authorisation Application, is in place and functioning before and whilst the product is on the market.

Risk Management Plan

The MAH commits to performing the studies and additional pharmacovigilance activities detailed in the Pharmacovigilance Plan, as agreed in version 1.0 of the Risk Management Plan (RMP) presented in Module 1.8.2. of the Marketing Authorisation Application and any subsequent updates of the RMP agreed by the CHMP.

As per the CHMP Guideline on Risk Management Systems for medicinal products for human use, the updated RMP should be submitted at the same time as the next Periodic Safety Update Report (PSUR).

In addition, an updated RMP should be submitted

- When new information is received that may impact on the current Safety Specification, Pharmacovigilance Plan or risk minimisation activities
- Within 60 days of an important (pharmacovigilance or risk minimisation) milestone being reached
- At the request of the EMEA.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING AND THE IMMEDIATE PACKAGING

CARTON

1. NAME OF THE MEDICINAL PRODUCT

Faslodex 250 mg solution for injection.
fulvestrant

2. STATEMENT OF ACTIVE SUBSTANCE(S)

One pre-filled syringe contains 250 mg fulvestrant in 5 ml solution

3. LIST OF EXCIPIENTS

Ethanol (96 per cent), benzyl alcohol, benzyl benzoate and castor oil. See the package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Solution for injection in a pre-filled syringe.
1 pre-filled syringe (5 ml)
1 safety needle

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Intramuscular use.
For single use only.
For full instructions on the administration of Faslodex and the use of the safety needle see enclosed, Instructions for administration.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN

Keep out of the reach and sight of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store in a refrigerator.

Store the pre-filled syringe in the original package in order to protect from light.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE
--

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

AstraZeneca UK Limited
Alderley Park
Macclesfield
Cheshire
SK10 4TG
United Kingdom

12. MARKETING AUTHORISATION NUMBER(S)
--

EU/1/03/269/001

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY
--

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS PRE-FILLED SYRINGE LABEL
--

1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION
--

Faslodex 250 mg solution for injection
fulvestrant
IM use

2. METHOD OF ADMINISTRATION

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT
--

5 ml

B. PACKAGE LEAFLET

PACKAGE LEAFLET: INFORMATION FOR THE USER

Faslodex 250 mg solution for injection Fulvestrant

Read all of this leaflet carefully before you start using this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, nurse or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor, nurse or pharmacist.

In this leaflet:

1. What Faslodex is and what it is used for
2. Before you use Faslodex
3. How to use Faslodex
4. Possible side effects
5. How to store Faslodex
6. Further information

1. WHAT FASLODEX IS AND WHAT IT IS USED FOR

Faslodex contains the active substance fulvestrant, which belongs to the group of estrogen blockers. Estrogens, a type of female sex hormones, can in some cases be involved in the growth of breast cancer.

Faslodex is used to treat advanced or metastatic breast cancer in postmenopausal women.

2. BEFORE YOU USE FASLODEX

Do not use Faslodex

- if you are allergic (hypersensitive) to fulvestrant or to any of the other ingredients of Faslodex (listed in section 6 'What Faslodex contains')
- if you are pregnant or breast-feeding
- if you have severe liver problems

Take special care with FASLODEX

Tell your doctor if any of these apply to you:

- kidney or liver problems
- low numbers of platelets (which help blood clotting) or bleeding disorders
- previous problems with blood clots
- osteoporosis (loss of bone density)
- Alcoholism

Using other medicines

Please tell your doctor, nurse or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

In particular, you should tell your doctor if you are using anticoagulants (medicines to prevent blood clots).

Pregnancy and breast-feeding

You must not use Faslodex if you are pregnant. If you can become pregnant, you should use effective contraception while being treated with Faslodex.

You must not breast-feed while on treatment with Faslodex.

Driving and using machines

Faslodex is not expected to affect your ability to drive or use machines. However, if you feel tired after treatment do not drive or use machines.

Important information about some of the ingredients of Faslodex

This medicinal product contains 10% w/v ethanol (alcohol), i.e. up to 1000 mg per dose, equivalent to 20 ml beer or 8 ml wine per dose.

Harmful for those suffering from alcoholism.

To be taken into account in pregnant or breast-feeding women, children and high-risk groups such as patients with liver disease, or epilepsy.

3. HOW TO USE FASLODEX

The usual dose is 500 mg fulvestrant (two 250 mg/5 ml injections) given once a month with an additional 500 mg dose given 2 weeks after the initial dose.

Your doctor or nurse will give you Faslodex as a slow intramuscular injection, one into each of your buttocks.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Faslodex can cause side effects, although not everybody gets them.

These side effects may occur with certain frequencies, which are defined as follows:

- very common: affects more than 1 user in 10
- common: affects 1 to 10 users in 100
- uncommon: affects 1 to 10 users in 1,000
- rare: affects 1 to 10 users in 10,000
- very rare: affects less than 1 user in 10,000
- not known: frequency cannot be estimated from the available data.

Very common side effects

- Injection site reactions, such as pain and/or inflammation
- Abnormal levels of liver enzymes (in blood tests)
- Nausea (feeling sick)
- Weakness, tiredness

Common side effects

- Headache
- Hot flushes
- Vomiting, diarrhoea, or loss of appetite
- Rash
- Urinary tract infections
- Back pain
- Thromboembolism (Increased risk of blood clots)
- Allergic (hypersensitivity) reactions, including swelling of the face, lips, tongue and/or throat

Uncommon side effects

- Vaginal bleeding, thick, whitish discharge and candidiasis (infection)

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor, nurse or pharmacist.

5. HOW TO STORE FASLODEX

Store in a refrigerator (2°C – 8°C)

Keep the pre-filled syringe in the original package, in order to protect from light.

Keep out of the reach and sight of children.

Do not use after the expiry date which is stated on the carton or syringe labels after the abbreviation EXP. The expiry date refers to the last day of that month.

Your health care professional will be responsible for the correct storage, use and disposal of Faslodex.

6. FURTHER INFORMATION

What Faslodex contains

- The active substance is fulvestrant. Each pre-filled syringe (5 ml) contains 250 mg fulvestrant.
- The other ingredients are ethanol (96 per cent), benzyl alcohol, benzyl benzoate and castor oil.

What Faslodex looks like and contents of the pack

Faslodex is a clear, colourless to yellow, viscous solution in a pre-filled syringe fitted with a tamper-evident closure, containing 5 ml solution for injection. A safety needle (BD SafetyGlide™) for connection to the barrel is also provided.

Marketing Authorisation Holder

AstraZeneca UK Limited
Alderley Park
Macclesfield
Cheshire
SK10 4TG
United Kingdom

Manufacturer

AstraZeneca UK Limited
Silk Road Business Park
Macclesfield
Cheshire
SK10 2NA
United Kingdom

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder (see contacts list).

België/Belgique/Belgien

NV AstraZeneca SA
Tel: +32 2 370 48 11

България

ТП AstraZeneca UK Limited
Тел.: +359 2 971 25 33

Česká republika

AstraZeneca Czech Republic s.r.o.
Tel: 00420222807232

Danmark

AstraZeneca A/S
Tel: +45 43 66 64 62

Deutschland

AstraZeneca GmbH
Tel: +49 41 03 7080

Ελλάδα

AstraZeneca A.E.
Τηλ: + 30 2 106871500

France

AstraZeneca
Tél: + 33 1 41 29 40 00

Ísland

Vistor hf.
Tel: + 354 535 7000

Κύπρος

Αλέκτωρ Φαρμακευτική Ατδ
Τηλ: +357 22490305

Lietuva

Tel.: +3705 2660550

Magyarország

AstraZeneca Kft
Tel: +36 23 517 300

Nederland

AstraZeneca BV
Tel: +31 79 363 2222

Österreich

AstraZeneca Österreich GmbH
Tel: +43 1 711 31 310

Portugal

AstraZeneca Produtos Farmacêuticos, Lda.
Tel: +351 21 434 61 00

Slovenija

AstraZeneca UK Limited
Tel: +386 1 51 35 600

Suomi/Finland

AstraZeneca Oy
Puh/Tel: + 358 10 23 010

United Kingdom

AstraZeneca UK Ltd
Tel: + 44 (0)1582 836 836

Eesti

AstraZeneca
Tel: +372 6549 600

España

AstraZeneca Farmacéutica Spain, S.A.
Tel: +34 91 301 91 00

Ireland

AstraZeneca Pharmaceuticals (Ireland) Ltd
Tel: + 353 1609 7100

Italia

AstraZeneca S.p.A.
Tel: + 39 02 98011

Latvija

AstraZeneca AB pārstāvniecība Latvijā
Tel: +371 7377100

Luxembourg/Luxemburg

NV AstraZeneca SA
Tel: +32 2 370 48 11

Malta

Associated Drug Co. Ltd
Tel: +356 21232175

Norge

AstraZeneca AS
Tel: +47 21 00 64 00

Polska

AstraZeneca Pharma Poland Sp. z o.o.
Tel.: + 48 22 874 35 00

România

AstraZeneca UK Limited Reprezentanța România
Tel: + 40 21 317 60 41

Slovenská republika

AstraZeneca AB o.z.
Tel.: + 421 2 5737 7777

Sverige

AstraZeneca AB
Tel: +46 8 553 26 000

This leaflet was last approved on

The following information is intended for healthcare professionals only:

Faslodex 500 mg (2 x 250 mg/5 ml solution for injection) should be administered using two pre-filled syringes, see section 3

BD SafetyGlide is a trademark of Becton Dickinson and Company and is CE-marked: CE 0050.


Instructions for administration

Warning - Do not autoclave safety needle (BD SafetyGlide™ Shielding Hypodermic Needle) before use. Hands must remain behind the needle at all times during use and disposal.

For each of the two syringes:

- Remove glass syringe barrel from tray and check that it is not damaged.
- Break the seal of the white plastic cover on the syringe Luer connector Luer-Lok to remove the cover with the attached rubber tip cap (see Figure 1).

Figure 1



- Peel open the safety needle (BD SafetyGlide) outer packaging. Attach the safety needle to the Luer-Lok (see Figure 2)
- Twist until firmly seated.
- Twist to lock the needle to the Luer connector.
- Pull shield straight off needle to avoid damaging needle point.
- Transport filled syringe to point of administration.
- Remove needle sheath.
- Parenteral solutions must be inspected visually for particulate matter and discolouration prior to administration.
- Expel excess gas from the syringe.

Figure 2



- Administer intramuscularly slowly (1-2 minutes/injection) into the buttock. For user convenience, the needle bevel- up position is oriented to the lever arm (see Figure 3).

Figure 3



- After injection, immediately apply a single-finger stroke to the activation assisted lever arm to activate the shielding mechanism (see Figure 4).

Figure 4



NOTE: Activate away from self and others. Listen for click and visually confirm needle tip is fully covered.

Disposal

Pre-filled syringes are for single use **only**.

Any unused product or waste material should be disposed of in accordance with local requirements.