

SEMESTER III

MODULE 2

FOURIER SERIES

Fourier series

Definition

A series of sines and cosines of an angle and its multiples of the form.

$$\begin{aligned} & \frac{a_0}{2} + a_1 \cos x + a_2 \cos 2x + a_3 \cos 3x + \dots + a_n \cos nx + \dots \\ & \quad + b_1 \sin x + b_2 \sin 2x + b_3 \sin 3x + \dots + b_n \sin nx + \dots \\ & = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx. \end{aligned}$$

is called the *Fourier series*, where $a_0, a_1, a_2, \dots, a_n, \dots, b_1, b_2, b_3, \dots, b_n, \dots$ are constants.

Dirichlet's Conditions:

Consider a function $f(x)$ in the interval $(a, a+2l)$ satisfying the following conditions.

1. $f(x)$ is single valued in $(a, a+2l)$
2. $f(x)$ is continuous or has discontinuities finite in number in the interval $(a, a+2l)$
3. $f(x)$ has no maxima or minima or has maxima and minima finite in number in the interval $(a, a+2l)$
4. $f(x)$ is periodic with period $2l$ in the interval $(a, a+2l)$

These conditions are called Dirichlet's Conditions.

Then $f(x)$ in the interval $(a, a+2l)$ can be written in terms of an infinite series as follows.

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left[a_n \cos \frac{n\pi x}{l} + b_n \sin \frac{n\pi x}{l} \right]$$

Where $a_0 = \frac{1}{2l} \int_a^{a+2l} f(x) dx$

$$a_n = \frac{1}{l} \int_a^{a+2l} f(x) \cos \frac{n\pi x}{l} dx$$

$$b_n = \frac{1}{l} \int_a^{a+2l} f(x) \sin \frac{n\pi x}{l} dx$$

Above formulae are called Euler's formulae.

Note that 1) $\sin^{-1} x$ can not be expressed as a Fourier series as it is not a single valued function.

2) $\tan x$ can not be expressed as a Fourier series in $(0, 2\pi)$ as it is not defined at $x = \frac{\pi}{2}$ and $3\frac{\pi}{2}$ which are the points of interval $(0, 2\pi)$.

Advantages of Fourier series

1. Discontinuous function can be represented by Fourier series. Although derivatives of the discontinuous functions do not exist. (This is not true for Taylor's series).
2. The Fourier series is useful in expanding the periodic functions since outside the closed interval, there exists a periodic extension of the function.
3. Expansion of an oscillating function by Fourier series gives all modes of oscillation (fundamental and all overtones) which is extremely useful in physics.
4. Fourier series of a discontinuous function is not uniformly convergent at all points.
5. Term by term integration of a convergent Fourier series is always valid, and it may be valid if the series is not convergent. However, term by term, differentiation of a Fourier series is not valid in most cases.

Useful Integrals

The following integrals are useful in Fourier Series.

$$(i) \int_0^{2\pi} \sin nx \, dx = 0$$

$$(ii) \int_0^{2\pi} \cos nx \, dx = 0$$

$$(iii) \int_0^{2\pi} \sin^2 nx \, dx = \pi$$

$$(iv) \int_0^{2\pi} \cos^2 nx \, dx = \pi$$

$$(v) \int_0^{2\pi} \sin nx \cdot \sin mx \, dx = 0$$

$$(vi) \int_0^{2\pi} \cos nx \cos mx \, dx = 0$$

$$(vii) \int_0^{2\pi} \sin nx \cdot \cos mx \, dx = 0$$

$$(viii) \int_0^{2\pi} \sin nx \cdot \cos nx \, dx = 0$$

$$(ix) [uv] = uv_1 - u'v_2 + u''v_3 - u'''v_4 + \dots$$

where $v_1 = \int v \, dx$, $v_2 = \int v_1 \, dx$ and so on. $u' = \frac{du}{dx}$, $u'' = \frac{d^2u}{dx^2}$ and so on

$$(x) \sin n\pi = 0, \quad \cos n\pi = (-1)^n \text{ where } n \in I$$

- SOME REQUIRED FORMULAE:

$$1) \int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx \text{ if } f(x) \text{ is even}$$
$$= 0 \quad \text{if } f(x) \text{ is odd.}$$

$$2) \int e^{ax} \sin bx \, dx = \frac{e^{ax}}{a^2 + b^2} [a \sin bx - b \cos bx]$$

$$3) \int e^{ax} \cos bx \, dx = \frac{e^{ax}}{a^2 + b^2} [a \cos bx + b \sin bx]$$

$$4) \text{ If } f(x) \text{ is discontinuous at } x = c \text{ then the value of } f(x) \text{ at } x = c \text{ is}$$
$$\text{given by } \frac{1}{2} \left[\lim_{x \rightarrow c^-} f(x) + \lim_{x \rightarrow c^+} f(x) \right]$$

$$5) \text{ Parseval's identity for the function } f(x) \text{ in the interval } (a, a+2l) \text{ is}$$
$$\text{given by } \frac{1}{2l} \int_a^{a+2l} f^2(x) \, dx = a_0^2 + \frac{1}{2} \sum_{n=1}^{\infty} a_n^2 + b_n^2$$

If $f(x)$ is defined in the interval $(a, a+2l)$ Fourier series is $f(x) = a_0 + \sum_{n=1}^{\infty} [a_n \cos \frac{n\pi x}{l} + b_n \sin \frac{n\pi x}{l}]$ where

$$a_0 = \frac{1}{2l} \int_a^{a+2l} f(x) dx ; \quad a_n = \frac{1}{l} \int_a^{a+2l} f(x) \cos \frac{n\pi x}{l} dx$$

$$b_n = \frac{1}{l} \int_a^{a+2l} f(x) \sin \frac{n\pi x}{l} dx$$

If $f(x)$ is defined in the interval $(a, a+2l)$ Fourier series is $f(x) = a_0 + \sum_{n=1}^{\infty} [a_n \cos \frac{n\pi x}{l} + b_n \sin \frac{n\pi x}{l}]$ where

$$a_0 = \frac{1}{2l} \int_a^{a+2l} f(x) dx ; \quad a_n = \frac{1}{l} \int_a^{a+2l} f(x) \cos \frac{n\pi x}{l} dx$$

$$b_n = \frac{1}{l} \int_a^{a+2l} f(x) \sin \frac{n\pi x}{l} dx$$

$(0, 2\pi)$

If $f(x)$ is defined in the interval $(a, a+2l)$ Fourier series is $f(x) = a_0 + \sum_{n=1}^{\infty} [a_n \cos \frac{n\pi x}{l} + b_n \sin \frac{n\pi x}{l}]$ where

$$a_0 = \frac{1}{2l} \int_a^{a+2l} f(x) dx ; \quad a_n = \frac{1}{l} \int_a^{a+2l} f(x) \cos \frac{n\pi x}{l} dx$$

$$b_n = \frac{1}{l} \int_a^{a+2l} f(x) \sin \frac{n\pi x}{l} dx$$

$(0, 2\pi)$

$$f(x) = a_0 + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx]$$

$$a_0 = \frac{1}{2\pi} \int_0^{2\pi} f(x) dx$$

$$a_n = \frac{1}{\pi} \int_a^{2\pi} f(x) \cos nx dx$$

$$b_n = \frac{1}{\pi} \int_a^{2\pi} f(x) \sin nx dx$$

FOURIER SERIES IN THE INTERVAL $(0, 2\pi)$.

EX 1 Obtain the Fourier expansion of $f(x) = \left(\frac{\pi-x}{2}\right)^2$

with $f(x + \pi) = f(x)$. Also deduce that

$$(i) \frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$(ii) \frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$$

$$(iii) \frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots$$

$$(iv) \frac{\pi^4}{90} = \frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots$$

FOURIER SERIES IN THE INTERVAL $(0, 2\pi)$.

EX 1 Obtain the Fourier expansion of $f(x) = \left(\frac{\pi-x}{2}\right)^2$

with $f(x + \pi) = f(x)$. Also deduce that

$$(i) \frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$(ii) \frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$$

$$(iii) \frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots$$

$$(iv) \frac{\pi^4}{90} = \frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots$$

Solution:

$$f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots (1)$$

FOURIER SERIES IN THE INTERVAL $(0, 2\pi)$.

EX 1 Obtain the Fourier expansion of $f(x) = \left(\frac{\pi-x}{2}\right)^2$

with $f(x + \pi) = f(x)$. Also deduce that

$$(i) \frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$(ii) \frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$$

$$(iii) \frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots$$

$$(iv) \frac{\pi^4}{90} = \frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots$$

Solution:

$$f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots (1)$$

$$\therefore a_0 = \frac{1}{2\pi} \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} \frac{(\pi-x)^2}{4} dx$$

FOURIER SERIES IN THE INTERVAL $(0, 2\pi)$.

EX 1 Obtain the Fourier expansion of $f(x) = \left(\frac{\pi-x}{2}\right)^2$

with $f(x + \pi) = f(x)$. Also deduce that

$$(i) \frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$(ii) \frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$$

$$(iii) \frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots$$

$$(iv) \frac{\pi^4}{90} = \frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots$$

Solution:

$$f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots (1)$$

$$\therefore a_0 = \frac{1}{2\pi} \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} \frac{(\pi-x)^2}{4} dx$$

$$\therefore a_0 = \frac{1}{8\pi} \left[\frac{(\pi-x)^3}{-3} \right]_0^{2\pi} = -\frac{1}{24\pi} [-\pi^3 - \pi^3] = \frac{\pi^2}{12} \quad \dots (A)$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \frac{(\pi - x)^2}{4} \cos nx \, dx$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \frac{(\pi - x)^2}{4} \cos nx \, dx \\
 &= \frac{1}{4\pi} \left[(\pi - x)^2 \left(\frac{\sin nx}{n} \right) - 2(\pi - x)(-1) \left(-\frac{\cos nx}{n^2} \right) + 2(-1)(-1) \left(-\frac{\sin nx}{n^3} \right) \right]_0^{2\pi}
 \end{aligned}$$

(By generalised rule of integration by parts.)

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \frac{(\pi - x)^2}{4} \cos nx \, dx \\
 &= \frac{1}{4\pi} \left[(\pi - x)^2 \left(\frac{\sin nx}{n} \right) - 2(\pi - x)(-1) \left(-\frac{\cos nx}{n^2} \right) + 2(-1)(-1) \left(-\frac{\sin nx}{n^3} \right) \right]_0^{2\pi}
 \end{aligned}$$

(By generalised rule of integration by parts.)

$$= \frac{1}{4\pi} \left[\left(0 + 2\pi \frac{\cos 2n\pi}{n^2} - 0 \right) - \left(0 - \frac{2\pi}{n^2} - 0 \right) \right]$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \frac{(\pi - x)^2}{4} \cos nx \, dx$$

$$= \frac{1}{4\pi} \left[(\pi - x)^2 \left(\frac{\sin nx}{n} \right) - 2(\pi - x)(-1) \left(-\frac{\cos nx}{n^2} \right) + 2(-1)(-1) \left(-\frac{\sin nx}{n^3} \right) \right]_0^{2\pi}$$

(By generalised rule of integration by parts.)

$$= \frac{1}{4\pi} \left[\left(0 + 2\pi \frac{\cos 2n\pi}{n^2} - 0 \right) - \left(0 - \frac{2\pi}{n^2} - 0 \right) \right]$$

$$\therefore a_n = \frac{1}{4\pi} \left[\frac{2\pi}{n^2} + \frac{2\pi}{n^2} \right] = \frac{1}{n^2} \quad [\because \cos 2n\pi = 1] \quad \dots\dots\dots (B)$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \frac{(\pi - x)^2}{4} \cdot \sin nx \, dx$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \frac{(\pi - x)^2}{4} \cdot \sin nx \, dx \\
 &= \frac{1}{4\pi} \left[(\pi - x)^2 \left(-\frac{\cos nx}{n} \right) - 2(\pi - x)(-1) \left(-\frac{\sin nx}{n^2} \right) \right. \\
 &\quad \left. + 2(-1)(-1) \left(\frac{\cos nx}{n^3} \right) \right]_0^{2\pi}
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \frac{(\pi - x)^2}{4} \cdot \sin nx \, dx \\
 &= \frac{1}{4\pi} \left[(\pi - x)^2 \left(-\frac{\cos nx}{n} \right) - 2(\pi - x)(-1) \left(-\frac{\sin nx}{n^2} \right) \right. \\
 &\quad \left. + 2(-1)(-1) \left(\frac{\cos nx}{n^3} \right) \right]_0^{2\pi} \\
 &= \frac{1}{4\pi} \left[\left(-\frac{\pi^2 \cos 2m\pi}{n} + 0 + \frac{2 \cos 2m\pi}{n^3} \right) - \left(-\frac{\pi^2}{n} + 0 + \frac{2}{n^3} \right) \right]
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \frac{(\pi - x)^2}{4} \cdot \sin nx \, dx \\
 &= \frac{1}{4\pi} \left[(\pi - x)^2 \left(-\frac{\cos nx}{n} \right) - 2(\pi - x)(-1) \left(-\frac{\sin nx}{n^2} \right) \right. \\
 &\quad \left. + 2(-1)(-1) \left(\frac{\cos nx}{n^3} \right) \right]_0^{2\pi} \\
 &= \frac{1}{4\pi} \left[\left(-\frac{\pi^2 \cos 2m\pi}{n} + 0 + \frac{2 \cos 2m\pi}{n^3} \right) - \left(-\frac{\pi^2}{n} + 0 + \frac{2}{n^3} \right) \right] \\
 \therefore b_n &= \frac{1}{4\pi} \left[-\frac{\pi^2}{n} + \frac{2}{n^3} + \frac{\pi^2}{n} - \frac{2}{n^3} \right] = 0 \quad \text{..... (C)}
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \frac{(\pi - x)^2}{4} \cdot \sin nx \, dx \\
 &= \frac{1}{4\pi} \left[(\pi - x)^2 \left(-\frac{\cos nx}{n} \right) - 2(\pi - x)(-1) \left(-\frac{\sin nx}{n^2} \right) \right. \\
 &\quad \left. + 2(-1)(-1) \left(\frac{\cos nx}{n^3} \right) \right]_0^{2\pi}
 \end{aligned}$$

$$= \frac{1}{4\pi} \left[\left(-\frac{\pi^2 \cos 2m\pi}{n} + 0 + \frac{2 \cos 2m\pi}{n^3} \right) - \left(-\frac{\pi^2}{n} + 0 + \frac{2}{n^3} \right) \right]$$

$$\therefore b_n = \frac{1}{4\pi} \left[-\frac{\pi^2}{n} + \frac{2}{n^3} + \frac{\pi^2}{n} - \frac{2}{n^3} \right] = 0 \quad \text{..... (C)}$$

Putting these values in (1), we get

$$\left(\frac{\pi - x}{2} \right)^2 = \frac{\pi^2}{12} + \sum_{n=1}^{\infty} \frac{1}{n^2} \cos nx$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \frac{(\pi - x)^2}{4} \cdot \sin nx \, dx \\
 &= \frac{1}{4\pi} \left[(\pi - x)^2 \left(-\frac{\cos nx}{n} \right) - 2(\pi - x)(-1) \left(-\frac{\sin nx}{n^2} \right) \right. \\
 &\quad \left. + 2(-1)(-1) \left(\frac{\cos nx}{n^3} \right) \right]_0^{2\pi}
 \end{aligned}$$

$$= \frac{1}{4\pi} \left[\left(-\frac{\pi^2 \cos 2m\pi}{n} + 0 + \frac{2 \cos 2m\pi}{n^3} \right) - \left(-\frac{\pi^2}{n} + 0 + \frac{2}{n^3} \right) \right]$$

$$\therefore b_n = \frac{1}{4\pi} \left[-\frac{\pi^2}{n} + \frac{2}{n^3} + \frac{\pi^2}{n} - \frac{2}{n^3} \right] = 0 \quad \dots\dots\dots (C)$$

Putting these values in (1), we get

$$\begin{aligned}
 \left(\frac{\pi - x}{2} \right)^2 &= \frac{\pi^2}{12} + \sum_{n=1}^{\infty} \frac{1}{n^2} \cos nx \\
 &= \frac{\pi^2}{12} + \frac{1}{1^2} \cos x + \frac{1}{2^2} \cos 2x + \frac{1}{3^2} \cos 3x + \dots \quad \dots\dots\dots (2)
 \end{aligned}$$

(i) Now, put $x = 0$ in (2),

(i) Now, put $x = 0$ in (2),

$$\therefore \frac{\pi^2}{4} = \frac{\pi^2}{12} + \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

(i) Now, put $x = 0$ in (2),

$$\therefore \frac{\pi^2}{4} = \frac{\pi^2}{12} + \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$\therefore \frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots \quad \text{..... (3)}$$

(i) Now, put $x = 0$ in (2),

$$\therefore \frac{\pi^2}{4} = \frac{\pi^2}{12} + \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$\therefore \frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots \quad \text{..... (3)}$$

(ii) Again, put $x = \pi$ in (2).

$$\therefore 0 = \frac{\pi^2}{12} - \frac{1}{1^2} + \frac{1}{2^2} - \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

(i) Now, put $x = 0$ in (2),

$$\therefore \frac{\pi^2}{4} = \frac{\pi^2}{12} + \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$\therefore \frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots \quad \text{..... (3)}$$

(ii) Again, put $x = \pi$ in (2).

$$\therefore 0 = \frac{\pi^2}{12} - \frac{1}{1^2} + \frac{1}{2^2} - \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$\therefore \frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots \quad \text{..... (4)}$$

(i) Now, put $x = 0$ in (2),

$$\therefore \frac{\pi^2}{4} = \frac{\pi^2}{12} + \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$\therefore \frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots \quad \text{..... (3)}$$

(ii) Again, put $x = \pi$ in (2).

$$\therefore 0 = \frac{\pi^2}{12} - \frac{1}{1^2} + \frac{1}{2^2} - \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$\therefore \frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots \quad \text{..... (4)}$$

(iii) To get the last result add (3) and (4).

$$\therefore \frac{\pi^2}{6} + \frac{\pi^2}{12} = \frac{2}{1^2} + \frac{2}{3^2} + \frac{2}{5^2} + \frac{2}{7^2} + \dots$$

(i) Now, put $x = 0$ in (2),

$$\therefore \frac{\pi^2}{4} = \frac{\pi^2}{12} + \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$\therefore \frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots \quad \text{..... (3)}$$

(ii) Again, put $x = \pi$ in (2).

$$\therefore 0 = \frac{\pi^2}{12} - \frac{1}{1^2} + \frac{1}{2^2} - \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$\therefore \frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots \quad \text{..... (4)}$$

(iii) To get the last result add (3) and (4).

$$\therefore \frac{\pi^2}{6} + \frac{\pi^2}{12} = \frac{2}{1^2} + \frac{2}{3^2} + \frac{2}{5^2} + \frac{2}{7^2} + \dots$$

$$\therefore \frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots$$

- (iv) To derive the last result we use Parseval's identity. We know that by Parseval's identity in $(0, 2\pi)$

(iv) To derive the last result we use Parseval's identity. We know that by Parseval's identity in $(0, 2\pi)$

$$\frac{1}{2\pi} \int_0^{2\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \dots\dots\dots (5)$$

(iv) To derive the last result we use Parseval's identity. We know that by Parseval's identity in $(0, 2\pi)$

$$\frac{1}{2\pi} \int_0^{2\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \quad \dots\dots\dots (5)$$

Now,
$$\frac{1}{2\pi} \int_0^{2\pi} [f(x)]^2 dx = \frac{1}{2\pi} \int_0^{2\pi} \left(\frac{\pi - x}{2} \right)^4 dx$$

(iv) To derive the last result we use Parseval's identity. We know that by Parseval's identity in $(0, 2\pi)$

$$\frac{1}{2\pi} \int_0^{2\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \quad \dots\dots\dots (5)$$

$$\begin{aligned} \text{Now, } \frac{1}{2\pi} \int_0^{2\pi} [f(x)]^2 dx &= \frac{1}{2\pi} \int_0^{2\pi} \left(\frac{\pi - x}{2} \right)^4 dx \\ &= \frac{1}{32\pi} \int_0^{2\pi} [\pi^4 - 4\pi^3 x + 6\pi^2 x^2 - 4\pi x^3 + x^4] dx \end{aligned}$$

(iv) To derive the last result we use Parseval's identity. We know that by Parseval's identity in $(0, 2\pi)$

$$\frac{1}{2\pi} \int_0^{2\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \quad \dots\dots\dots (5)$$

$$\begin{aligned} \text{Now, } \frac{1}{2\pi} \int_0^{2\pi} [f(x)]^2 dx &= \frac{1}{2\pi} \int_0^{2\pi} \left(\frac{\pi - x}{2} \right)^4 dx \\ &= \frac{1}{32\pi} \int_0^{2\pi} [\pi^4 - 4\pi^3 x + 6\pi^2 x^2 - 4\pi x^3 + x^4] dx \\ &= \frac{1}{32\pi} \left[\pi^4 x - 2\pi^3 x^2 + 2\pi^2 x^3 - \pi x^4 + \frac{x^5}{5} \right]_0^{2\pi} \end{aligned}$$

(iv) To derive the last result we use Parseval's identity. We know that by Parseval's identity in $(0, 2\pi)$

$$\frac{1}{2\pi} \int_0^{2\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \quad \dots\dots\dots (5)$$

$$\begin{aligned} \text{Now, } \frac{1}{2\pi} \int_0^{2\pi} [f(x)]^2 dx &= \frac{1}{2\pi} \int_0^{2\pi} \left(\frac{\pi - x}{2} \right)^4 dx \\ &= \frac{1}{32\pi} \int_0^{2\pi} [\pi^4 - 4\pi^3 x + 6\pi^2 x^2 - 4\pi x^3 + x^4] dx \\ &= \frac{1}{32\pi} \left[\pi^4 x - 2\pi^3 x^2 + 2\pi^2 x^3 - \pi x^4 + \frac{x^5}{5} \right]_0^{2\pi} \\ &= \frac{1}{32\pi} \left[2\pi^5 - 8\pi^5 + 16\pi^5 - 16\pi^5 + \frac{32\pi^5}{5} \right] \end{aligned}$$

- (iv) To derive the last result we use Parseval's identity. We know that by Parseval's identity in $(0, 2\pi)$

$$\frac{1}{2\pi} \int_0^{2\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \quad \dots\dots\dots (5)$$

$$\begin{aligned} \text{Now, } \frac{1}{2\pi} \int_0^{2\pi} [f(x)]^2 dx &= \frac{1}{2\pi} \int_0^{2\pi} \left(\frac{\pi - x}{2} \right)^4 dx \\ &= \frac{1}{32\pi} \int_0^{2\pi} [\pi^4 - 4\pi^3 x + 6\pi^2 x^2 - 4\pi x^3 + x^4] dx \\ &= \frac{1}{32\pi} \left[\pi^4 x - 2\pi^3 x^2 + 2\pi^2 x^3 - \pi x^4 + \frac{x^5}{5} \right]_0^{2\pi} \\ &= \frac{1}{32\pi} \left[2\pi^5 - 8\pi^5 + 16\pi^5 - 16\pi^5 + \frac{32\pi^5}{5} \right] \\ &= \frac{1}{32\pi} \cdot \frac{2\pi^5}{5} = \frac{\pi^4}{80} \end{aligned}$$

Hence, by (5) using (A), (B) and (C)

Hence, by (5) using (A), (B) and (C)

$$\frac{\pi^4}{80} = \frac{\pi^4}{144} + \frac{1}{2} \left(\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots \right)$$

Hence, by (5) using (A), (B) and (C)

$$\frac{\pi^4}{80} = \frac{\pi^4}{144} + \frac{1}{2} \left(\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots \right)$$

$$\pi^4 \left(\frac{1}{80} - \frac{1}{144} \right) = \frac{1}{2} \left(\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots \right)$$

Hence, by (5) using (A), (B) and (C)

$$\frac{\pi^4}{80} = \frac{\pi^4}{144} + \frac{1}{2} \left(\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots \right)$$

$$\pi^4 \left(\frac{1}{80} - \frac{1}{144} \right) = \frac{1}{2} \left(\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots \right)$$

$$\pi^4 \left(\frac{9-5}{720} \right) = \frac{\pi^4}{180} = \frac{1}{2} \left(\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots \right)$$

Hence, by (5) using (A), (B) and (C)

$$\frac{\pi^4}{80} = \frac{\pi^4}{144} + \frac{1}{2} \left(\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots \right)$$

$$\pi^4 \left(\frac{1}{80} - \frac{1}{144} \right) = \frac{1}{2} \left(\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots \right)$$

$$\pi^4 \left(\frac{9-5}{720} \right) = \frac{\pi^4}{180} = \frac{1}{2} \left(\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots \right)$$

$$\therefore \frac{\pi^4}{90} = \frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots$$

EX.2 Expand $f(x) = x \sin x$ in the interval $(0, 2\pi)$

Deduce that $\sum_{n=2}^{\infty} \frac{1}{n^2-1} = \frac{3}{4}$

EX.2 Expand $f(x) = x \sin x$ in the interval $(0, 2\pi)$

Deduce that $\sum_{n=2}^{\infty} \frac{1}{n^2-1} = \frac{3}{4}$

Solution:

$$\therefore \text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

EX.2 Expand $f(x) = x \sin x$ in the interval $(0, 2\pi)$

Deduce that $\sum_{n=2}^{\infty} \frac{1}{n^2-1} = \frac{3}{4}$

Solution:

$$\therefore \text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\therefore a_0 = \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} (x \sin x) \cdot dx$$

EX.2 Expand $f(x) = x \sin x$ in the interval $(0, 2\pi)$

Deduce that $\sum_{n=2}^{\infty} \frac{1}{n^2-1} = \frac{3}{4}$

Solution:

$$\therefore \text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\begin{aligned} \therefore a_0 &= \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} (x \sin x) \cdot dx \\ &= \frac{1}{2\pi} \left[(x)(-\cos x) - (-\sin x)(1) \right]_0^{2\pi} \end{aligned}$$

EX.2 Expand $f(x) = x \sin x$ in the interval $(0, 2\pi)$

Deduce that $\sum_{n=2}^{\infty} \frac{1}{n^2-1} = \frac{3}{4}$

Solution:

$$\therefore \text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\therefore a_0 = \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} (x \sin x) \cdot dx$$

$$= \frac{1}{2\pi} [(x)(-\cos x) - (-\sin x)(1)]_0^{2\pi}$$

$$= \frac{1}{2\pi} [(-2\pi + 0) - (0 + 0)] = -1$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \cos nx \, dx \quad \dots (2)$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \cos nx \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \cos nx \sin x \, dx$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \cos nx \, dx \quad \dots (2)$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \cos nx \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \cos nx \sin x \, dx$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x [\sin(n+1)x - \sin(n-1)x] \, dx$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \cos nx \, dx \quad \dots (2)$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \cos nx \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \cos nx \sin x \, dx$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x [\sin(n+1)x - \sin(n-1)x] \, dx$$

$$= \frac{1}{2\pi} \left[x \left\{ -\frac{\cos(n+1)x}{n+1} + \frac{\cos(n-1)x}{n-1} \right\} \right. \\ \left. - (1) \left\{ -\frac{\sin(n+1)x}{(n+1)^2} + \frac{\sin(n-1)x}{(n-1)^2} \right\} \right]_0^{2\pi}$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \cos nx \, dx \quad \dots (2)$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \cos nx \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \cos nx \sin x \, dx$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x [\sin(n+1)x - \sin(n-1)x] \, dx$$

$$= \frac{1}{2\pi} \left[x \left\{ -\frac{\cos(n+1)x}{n+1} + \frac{\cos(n-1)x}{n-1} \right\} \right.$$

$$\left. - (1) \left\{ -\frac{\sin(n+1)x}{(n+1)^2} + \frac{\sin(n-1)x}{(n-1)^2} \right\} \right]_0^{2\pi}$$

$$= \frac{1}{2\pi} \left[(2\pi) \left\{ -\frac{\cos 2(n+1)\pi}{(n+1)} + \frac{\cos 2(n-1)\pi}{(n-1)} \right\} - 0 \right]$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \cos nx \, dx \quad \dots (2)$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \cos nx \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \cos nx \sin x \, dx$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x [\sin(n+1)x - \sin(n-1)x] \, dx$$

$$= \frac{1}{2\pi} \left[x \left\{ -\frac{\cos(n+1)x}{n+1} + \frac{\cos(n-1)x}{n-1} \right\} \right.$$

$$\left. - (1) \left\{ -\frac{\sin(n+1)x}{(n+1)^2} + \frac{\sin(n-1)x}{(n-1)^2} \right\} \right]_0^{2\pi}$$

$$= \frac{1}{2\pi} \left[(2\pi) \left\{ -\frac{\cos 2(n+1)\pi}{(n+1)} + \frac{\cos 2(n-1)\pi}{(n-1)} \right\} - 0 \right]$$

$$= -\frac{1}{n+1} + \frac{1}{n-1} = \frac{2}{n^2-1} \quad \text{if } n \neq 1$$

If $n = 1$, the above method fails.

Putting $n = 1$ in (2), we get,

Putting $n = 1$ in (2), we get,

$$a_1 = \frac{1}{\pi} \int_0^{2\pi} x \sin x \cos x \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \cos x \, dx$$

Putting $n = 1$ in (2), we get,

$$\begin{aligned} a_1 &= \frac{1}{\pi} \int_0^{2\pi} x \sin x \cos x \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \cos x \, dx \\ &= \frac{1}{2\pi} \int_0^{2\pi} x \cdot \sin 2x \, dx \end{aligned}$$

Putting $n = 1$ in (2), we get,

$$\begin{aligned}a_1 &= \frac{1}{\pi} \int_0^{2\pi} x \sin x \cos x \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \cos x \, dx \\&= \frac{1}{2\pi} \int_0^{2\pi} x \cdot \sin 2x \, dx \\&= \frac{1}{2\pi} \left[x \left(-\frac{\cos 2x}{2} \right) - (1) \left(-\frac{\sin 2x}{4} \right) \right]_0^{2\pi}\end{aligned}$$

Putting $n = 1$ in (2), we get,

$$\begin{aligned} a_1 &= \frac{1}{\pi} \int_0^{2\pi} x \sin x \cos x \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \cos x \, dx \\ &= \frac{1}{2\pi} \int_0^{2\pi} x \cdot \sin 2x \, dx \\ &= \frac{1}{2\pi} \left[x \left(-\frac{\cos 2x}{2} \right) - (1) \left(-\frac{\sin 2x}{4} \right) \right]_0^{2\pi} \\ &= \frac{1}{2\pi} \left[2\pi \left(-\frac{\cos 4\pi}{2} \right) - 0 \right] = -\frac{1}{2} \end{aligned}$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin nx \, dx \quad \dots\dots\dots (3)$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin nx \, dx \quad \dots\dots\dots (3)$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \sin nx \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin nx \sin x \, dx$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin nx \, dx \quad \dots\dots\dots (3)$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \sin nx \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin nx \sin x \, dx$$

$$= -\frac{1}{2\pi} \int_0^{2\pi} x [\cos(n+1)x - \cos(n-1)x] \, dx$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin nx \, dx \quad \dots\dots\dots (3)$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \sin nx \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin nx \sin x \, dx$$

$$= -\frac{1}{2\pi} \int_0^{2\pi} x [\cos(n+1)x - \cos(n-1)x] \, dx$$

$$\therefore b_n = -\frac{1}{2\pi} \left[x \left\{ \frac{\sin(n+1)x}{n+1} - \frac{\sin(n-1)x}{n-1} \right\} - (1) \left\{ -\frac{\cos(n+1)x}{(n+1)^2} + \frac{\cos(n-1)x}{(n-1)^2} \right\} \right]_0^{2\pi}$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin nx \, dx \quad \dots\dots\dots (3)$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \sin nx \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin nx \sin x \, dx$$

$$= -\frac{1}{2\pi} \int_0^{2\pi} x [\cos(n+1)x - \cos(n-1)x] \, dx$$

$$\therefore b_n = -\frac{1}{2\pi} \left[x \left\{ \frac{\sin(n+1)x}{n+1} - \frac{\sin(n-1)x}{n-1} \right\} - (1) \left\{ -\frac{\cos(n+1)x}{(n+1)^2} + \frac{\cos(n-1)x}{(n-1)^2} \right\} \right]_0^{2\pi}$$

$$= -\frac{1}{2\pi} \left[- (1) \left\{ -\frac{\cos 2(n+1)\pi}{(n+1)^2} + \frac{\cos 2(n-1)\pi}{(n-1)^2} \right\} + (1) \left\{ -\frac{1}{(n+1)^2} + \frac{1}{(n-1)^2} \right\} \right]$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin nx \, dx \quad \dots\dots\dots (3)$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \sin nx \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin nx \sin x \, dx$$

$$= -\frac{1}{2\pi} \int_0^{2\pi} x [\cos(n+1)x - \cos(n-1)x] \, dx$$

$$\therefore b_n = -\frac{1}{2\pi} \left[x \left\{ \frac{\sin(n+1)x}{n+1} - \frac{\sin(n-1)x}{n-1} \right\} - (1) \left\{ -\frac{\cos(n+1)x}{(n+1)^2} + \frac{\cos(n-1)x}{(n-1)^2} \right\} \right]_0^{2\pi}$$

$$= -\frac{1}{2\pi} \left[-(1) \left\{ -\frac{\cos 2(n+1)\pi}{(n+1)^2} + \frac{\cos 2(n-1)\pi}{(n-1)^2} \right\} + (1) \left\{ -\frac{1}{(n+1)^2} + \frac{1}{(n-1)^2} \right\} \right]$$

$$= -\frac{1}{2\pi} \left[\frac{1}{(n+1)^2} - \frac{1}{(n-1)^2} - \frac{1}{(n+1)^2} + \frac{1}{(n-1)^2} \right]$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin nx \, dx \quad \dots\dots\dots (3)$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin x \sin nx \, dx = \frac{1}{2\pi} \int_0^{2\pi} x \cdot 2 \sin nx \sin x \, dx$$

$$= -\frac{1}{2\pi} \int_0^{2\pi} x [\cos(n+1)x - \cos(n-1)x] \, dx$$

$$\therefore b_n = -\frac{1}{2\pi} \left[x \left\{ \frac{\sin(n+1)x}{n+1} - \frac{\sin(n-1)x}{n-1} \right\} - (1) \left\{ -\frac{\cos(n+1)x}{(n+1)^2} + \frac{\cos(n-1)x}{(n-1)^2} \right\} \right]_0^{2\pi}$$

$$= -\frac{1}{2\pi} \left[-(1) \left\{ -\frac{\cos 2(n+1)\pi}{(n+1)^2} + \frac{\cos 2(n-1)\pi}{(n-1)^2} \right\} + (1) \left\{ -\frac{1}{(n+1)^2} + \frac{1}{(n-1)^2} \right\} \right]$$

$$= -\frac{1}{2\pi} \left[\frac{1}{(n+1)^2} - \frac{1}{(n-1)^2} - \frac{1}{(n+1)^2} + \frac{1}{(n-1)^2} \right]$$

$$= 0 \text{ if } n \neq 1$$

If $n = 1$, the above method fails.

Putting $n = 1$ in (3), we get,

$$\frac{1}{2} \int_{-\pi}^{\pi} f(x) dx$$

$$= \frac{1}{2} \int_{-\pi}^{\pi} f(x) dx$$

$$= \frac{1}{2} \int_{-\pi}^{\pi} f(x) dx$$

Putting $n = 1$ in (3), we get,

$$b_1 = \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin x \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin^2 x \, dx = \frac{1}{2\pi} \int_0^{2\pi} x (1 - \cos 2x) \, dx$$

Putting $n = 1$ in (3), we get,

$$\begin{aligned} b_1 &= \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin x \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin^2 x \, dx = \frac{1}{2\pi} \int_0^{2\pi} x (1 - \cos 2x) \, dx \\ &= \frac{1}{2\pi} \left[x \left(x - \frac{\sin 2x}{2} \right) - (1) \left(\frac{x^2}{2} + \frac{\cos 2x}{4} \right) \right]_0^{2\pi} \end{aligned}$$

Putting $n = 1$ in (3), we get,

$$\begin{aligned} b_1 &= \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin x \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin^2 x \, dx = \frac{1}{2\pi} \int_0^{2\pi} x(1 - \cos 2x) \, dx \\ &= \frac{1}{2\pi} \left[x \left(x - \frac{\sin 2x}{2} \right) - (1) \left(\frac{x^2}{2} + \frac{\cos 2x}{4} \right) \right]_0^{2\pi} \\ &= \frac{1}{2\pi} \left[\left\{ 2\pi(2\pi - 0) - \left(\frac{4\pi^2}{2} + \frac{1}{4} \right) \right\} - \left(0 - \frac{1}{4} \right) \right] = \frac{1}{2\pi} [2\pi^2] = \pi \end{aligned}$$

Putting $n = 1$ in (3), we get,

$$\begin{aligned} b_1 &= \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin x \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin^2 x \, dx = \frac{1}{2\pi} \int_0^{2\pi} x(1 - \cos 2x) \, dx \\ &= \frac{1}{2\pi} \left[x \left(x - \frac{\sin 2x}{2} \right) - (1) \left(\frac{x^2}{2} + \frac{\cos 2x}{4} \right) \right]_0^{2\pi} \\ &= \frac{1}{2\pi} \left[\left\{ 2\pi(2\pi - 0) - \left(\frac{4\pi^2}{2} + \frac{1}{4} \right) \right\} - \left(0 - \frac{1}{4} \right) \right] = \frac{1}{2\pi} [2\pi^2] = \pi \end{aligned}$$

Putting these values in (1),

$$x \sin x = -1 - \frac{1}{2} \cos x + \sum_{n=2}^{\infty} \frac{2}{n^2 - 1} \cos nx + \pi \sin x$$

Putting $n = 1$ in (3), we get,

$$\begin{aligned} b_1 &= \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin x \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin^2 x \, dx = \frac{1}{2\pi} \int_0^{2\pi} x (1 - \cos 2x) \, dx \\ &= \frac{1}{2\pi} \left[x \left(x - \frac{\sin 2x}{2} \right) - (1) \left(\frac{x^2}{2} + \frac{\cos 2x}{4} \right) \right]_0^{2\pi} \\ &= \frac{1}{2\pi} \left[\left\{ 2\pi (2\pi - 0) - \left(\frac{4\pi^2}{2} + \frac{1}{4} \right) \right\} - \left(0 - \frac{1}{4} \right) \right] = \frac{1}{2\pi} [2\pi^2] = \pi \end{aligned}$$

Putting these values in (1),

$$x \sin x = -1 - \frac{1}{2} \cos x + \sum_{n=2}^{\infty} \frac{2}{n^2 - 1} \cos nx + \pi \sin x$$

Deduction : Putting $x = 0$, we get $\frac{3}{4} = \sum_{n=2}^{\infty} \frac{1}{n^2 - 1}.$

EX 3 Find the Fourier series for $f(x) = e^{-x}$ in $0 < x < 2\pi$.

Deduce the value of the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{n^2+1}$. Also derive

the series for $\operatorname{csch} \pi$

EX 3 Find the Fourier series for $f(x) = e^{-x}$ in $0 < x < 2\pi$.

Deduce the value of the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{n^2+1}$. Also derive

the series for $\operatorname{csch} \pi$

Solution:

$$\therefore \text{Let } f(x) = e^{-x} = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

EX 3 Find the Fourier series for $f(x) = e^{-x}$ in $0 < x < 2\pi$.

Deduce the value of the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{n^2+1}$. Also derive

the series for $\operatorname{csch} \pi$

Solution:

$$\therefore \text{Let } f(x) = e^{-x} = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\therefore a_0 = \frac{1}{2\pi} \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} e^{-x} dx = \frac{1}{2\pi} \left[-e^{-x} \right]_0^{2\pi}$$

EX 3 Find the Fourier series for $f(x) = e^{-x}$ in $0 < x < 2\pi$.

Deduce the value of the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{n^2+1}$. Also derive

the series for $\operatorname{csch} \pi$

Solution:

$$\therefore \text{Let } f(x) = e^{-x} = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\begin{aligned} \therefore a_0 &= \frac{1}{2\pi} \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} e^{-x} dx = \frac{1}{2\pi} \left[-e^{-x} \right]_0^{2\pi} \\ &= \frac{1}{2\pi} (-e^{-2\pi} + 1) = \frac{1 - e^{-2\pi}}{2\pi} \end{aligned}$$

EX 3 Find the Fourier series for $f(x) = e^{-x}$ in $0 < x < 2\pi$.

Deduce the value of the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{n^2+1}$. Also derive

the series for $\operatorname{csch} \pi$

Solution:

$$\therefore \text{Let } f(x) = e^{-x} = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\begin{aligned} \therefore a_0 &= \frac{1}{2\pi} \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} e^{-x} dx = \frac{1}{2\pi} \left[-e^{-x} \right]_0^{2\pi} \\ &= \frac{1}{2\pi} (-e^{-2\pi} + 1) = \frac{1 - e^{-2\pi}}{2\pi} \end{aligned}$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx = \frac{1}{\pi} \int_0^{2\pi} e^{-x} \cos nx dx$$

EX 3 Find the Fourier series for $f(x) = e^{-x}$ in $0 < x < 2\pi$.

Deduce the value of the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{n^2+1}$. Also derive

the series for $\operatorname{csch} \pi$

Solution:

$$\therefore \text{Let } f(x) = e^{-x} = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\begin{aligned} \therefore a_0 &= \frac{1}{2\pi} \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} e^{-x} dx = \frac{1}{2\pi} \left[-e^{-x} \right]_0^{2\pi} \\ &= \frac{1}{2\pi} (-e^{-2\pi} + 1) = \frac{1 - e^{-2\pi}}{2\pi} \end{aligned}$$

$$\begin{aligned} a_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx = \frac{1}{\pi} \int_0^{2\pi} e^{-x} \cos nx dx \\ &= \frac{1}{\pi} \cdot \frac{1}{1+n^2} \left[e^{-x} (-\cos nx + n \sin nx) \right]_0^{2\pi} \end{aligned}$$

EX 3 Find the Fourier series for $f(x) = e^{-x}$ in $0 < x < 2\pi$.

Deduce the value of the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{n^2+1}$. Also derive

the series for $\operatorname{csch} \pi$

Solution:

$$\therefore \text{Let } f(x) = e^{-x} = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\begin{aligned} \therefore a_0 &= \frac{1}{2\pi} \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} e^{-x} dx = \frac{1}{2\pi} \left[-e^{-x} \right]_0^{2\pi} \\ &= \frac{1}{2\pi} (-e^{-2\pi} + 1) = \frac{1 - e^{-2\pi}}{2\pi} \end{aligned}$$

$$\begin{aligned} a_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx = \frac{1}{\pi} \int_0^{2\pi} e^{-x} \cos nx dx \\ &= \frac{1}{\pi} \cdot \frac{1}{1+n^2} \left[e^{-x} (-\cos nx + n \sin nx) \right]_0^{2\pi} \\ &= \frac{1}{\pi(1+n^2)} \left[e^{-2\pi} (-\cos 2\pi + n \sin 2\pi) - e^0 (-\cos 0 + n \sin 0) \right] \end{aligned}$$

EX 3 Find the Fourier series for $f(x) = e^{-x}$ in $0 < x < 2\pi$.

Deduce the value of the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{n^2+1}$. Also derive

the series for $\operatorname{csch} \pi$

Solution:

$$\therefore \text{Let } f(x) = e^{-x} = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\begin{aligned} \therefore a_0 &= \frac{1}{2\pi} \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} e^{-x} dx = \frac{1}{2\pi} \left[-e^{-x} \right]_0^{2\pi} \\ &= \frac{1}{2\pi} (-e^{-2\pi} + 1) = \frac{1 - e^{-2\pi}}{2\pi} \end{aligned}$$

$$\begin{aligned} a_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx = \frac{1}{\pi} \int_0^{2\pi} e^{-x} \cos nx dx \\ &= \frac{1}{\pi} \cdot \frac{1}{1+n^2} \left[e^{-x} (-\cos nx + n \sin nx) \right]_0^{2\pi} \\ &= \frac{1}{\pi(1+n^2)} \left[e^{-2\pi} (-\cos 2n\pi + n \sin 2n\pi) - e^0 (-\cos 0 + n \sin 0) \right] \\ &= \frac{1}{\pi(1+n^2)} \cdot \left[e^{-2\pi} (-1) - (-1) \right] = \frac{1 - e^{-2\pi}}{\pi(1+n^2)} \end{aligned}$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} e^{-x} \sin nx \, dx$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} e^{-x} \sin nx \, dx \\
 &= \frac{1}{\pi} \cdot \frac{1}{1+n^2} \left[e^{-x} (-\sin nx - n \cos nx) \right]_0^{2\pi}
 \end{aligned}$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} e^{-x} \sin nx \, dx$$

$$= \frac{1}{\pi} \cdot \frac{1}{1+n^2} \left[e^{-x} (-\sin nx - n \cos nx) \right]_0^{2\pi}$$

$$= \frac{1}{\pi(1+n^2)} \left[e^{-2\pi} (-\sin 2n\pi - n \cos 2n\pi) - e^0 (-\sin 0 - n \cos 0) \right]$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} e^{-x} \sin nx \, dx \\
 &= \frac{1}{\pi} \cdot \frac{1}{1+n^2} \left[e^{-x} (-\sin nx - n \cos nx) \right]_0^{2\pi} \\
 &= \frac{1}{\pi(1+n^2)} \left[e^{-2\pi} (-\sin 2n\pi - n \cos 2n\pi) - e^0 (-\sin 0 - n \cos 0) \right] \\
 &= \frac{1}{\pi(1+n^2)} \left[e^{-2\pi} (-n) - 1(-n) \right] = \frac{n}{\pi(1+n^2)} (1 - e^{-2\pi})
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} e^{-x} \sin nx \, dx \\
 &= \frac{1}{\pi} \cdot \frac{1}{1+n^2} \left[e^{-x} (-\sin nx - n \cos nx) \right]_0^{2\pi} \\
 &= \frac{1}{\pi(1+n^2)} \left[e^{-2\pi} (-\sin 2n\pi - n \cos 2n\pi) - e^0 (-\sin 0 - n \cos 0) \right] \\
 &= \frac{1}{\pi(1+n^2)} \left[e^{-2\pi} (-n) - 1(-n) \right] = \frac{n}{\pi(1+n^2)} (1 - e^{-2\pi})
 \end{aligned}$$

Putting the values of a_0 , a_n , b_n in (1), we get

$$e^{-x} = \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[\sum_{n=1}^{\infty} \frac{1}{1+n^2} \cos nx + \sum_{n=1}^{\infty} \frac{n}{1+n^2} \sin nx \right] \dots\dots\dots (2)$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} e^{-x} \sin nx \, dx \\
 &= \frac{1}{\pi} \cdot \frac{1}{1+n^2} \left[e^{-x} (-\sin nx - n \cos nx) \right]_0^{2\pi} \\
 &= \frac{1}{\pi(1+n^2)} \left[e^{-2\pi} (-\sin 2n\pi - n \cos 2n\pi) - e^0 (-\sin 0 - n \cos 0) \right] \\
 &= \frac{1}{\pi(1+n^2)} \left[e^{-2\pi} (-n) - 1(-n) \right] = \frac{n}{\pi(1+n^2)} (1 - e^{-2\pi})
 \end{aligned}$$

Putting the values of a_0 , a_n , b_n in (1), we get

$$\begin{aligned}
 e^{-x} &= \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[\sum_{n=1}^{\infty} \frac{1}{1+n^2} \cos nx + \sum_{n=1}^{\infty} \frac{n}{1+n^2} \sin nx \right] \dots\dots\dots (2) \\
 &= \frac{1 - e^{-2\pi}}{\pi} \left[\frac{1}{2} + \sum_{n=1}^{\infty} \frac{1}{1+n^2} (\cos nx + n \sin nx) \right]
 \end{aligned}$$

Putting $x = \pi$ in (2), we get,

Putting $x = \pi$ in (2), we get,

$$e^{-\pi} = \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[\sum_{n=1}^{\infty} \frac{1}{1 + n^2} \cos n\pi \right] \quad (\because \sin n\pi = 0)$$

Putting $x = \pi$ in (2), we get,

$$\begin{aligned} e^{-\pi} &= \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[\sum_{n=1}^{\infty} \frac{1}{1 + n^2} \cos n\pi \right] \quad (\because \sin n\pi = 0) \\ &= \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[-\frac{1}{2} + \sum_{n=2}^{\infty} \frac{1}{1 + n^2} (-1)^n \right] \end{aligned}$$

Putting $x = \pi$ in (2), we get,

$$\begin{aligned} e^{-\pi} &= \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[\sum_{n=1}^{\infty} \frac{1}{1 + n^2} \cos n\pi \right] \quad (\because \sin n\pi = 0) \\ &= \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[-\frac{1}{2} + \sum_{n=2}^{\infty} \frac{1}{1 + n^2} (-1)^n \right] \\ \therefore e^{-\pi} &= \frac{1 - e^{-2\pi}}{\pi} \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2} \end{aligned}$$

Putting $x = \pi$ in (2), we get,

$$\begin{aligned} e^{-\pi} &= \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[\sum_{n=1}^{\infty} \frac{1}{1 + n^2} \cos n\pi \right] \quad (\because \sin n\pi = 0) \\ &= \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[-\frac{1}{2} + \sum_{n=2}^{\infty} \frac{1}{1 + n^2} (-1)^n \right] \\ \therefore e^{-\pi} &= \frac{1 - e^{-2\pi}}{\pi} \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2} \quad \therefore \frac{\pi}{e^{\pi}(1 - e^{-2\pi})} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2} \end{aligned}$$

Putting $x = \pi$ in (2), we get,

$$e^{-\pi} = \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[\sum_{n=1}^{\infty} \frac{1}{1 + n^2} \cos n\pi \right] \quad (\because \sin n\pi = 0)$$

$$= \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[-\frac{1}{2} + \sum_{n=2}^{\infty} \frac{1}{1 + n^2} (-1)^n \right]$$

$$\therefore e^{-\pi} = \frac{1 - e^{-2\pi}}{\pi} \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2} \quad \therefore \frac{\pi}{e^{\pi}(1 - e^{-2\pi})} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2}$$

$$\therefore \frac{\pi}{e^{\pi} - e^{-\pi}} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2}$$

Putting $x = \pi$ in (2), we get,

$$e^{-\pi} = \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[\sum_{n=1}^{\infty} \frac{1}{1 + n^2} \cos n\pi \right] \quad (\because \sin n\pi = 0)$$

$$= \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[-\frac{1}{2} + \sum_{n=2}^{\infty} \frac{1}{1 + n^2} (-1)^n \right]$$

$$\therefore e^{-\pi} = \frac{1 - e^{-2\pi}}{\pi} \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2} \quad \therefore \frac{\pi}{e^{\pi}(1 - e^{-2\pi})} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2}$$

$$\therefore \frac{\pi}{e^{\pi} - e^{-\pi}} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2} \quad \therefore \frac{\pi}{2} \cdot \frac{2}{e^{\pi} - e^{-\pi}} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2}$$

Putting $x = \pi$ in (2), we get,

$$e^{-\pi} = \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[\sum_{n=1}^{\infty} \frac{1}{1 + n^2} \cos n\pi \right] \quad (\because \sin n\pi = 0)$$

$$= \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[-\frac{1}{2} + \sum_{n=2}^{\infty} \frac{1}{1 + n^2} (-1)^n \right]$$

$$\therefore e^{-\pi} = \frac{1 - e^{-2\pi}}{\pi} \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2} \quad \therefore \frac{\pi}{e^{\pi}(1 - e^{-2\pi})} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2}$$

$$\therefore \frac{\pi}{e^{\pi} - e^{-\pi}} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2} \quad \therefore \frac{\pi}{2} \cdot \frac{2}{e^{\pi} - e^{-\pi}} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2}$$

$$\therefore \frac{\pi}{2} \cdot \frac{1}{\sinh \pi} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1 + n^2}$$

Putting $x = \pi$ in (2), we get,

$$e^{-\pi} = \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[\sum_{n=1}^{\infty} \frac{1}{1+n^2} \cos n\pi \right] \quad (\because \sin n\pi = 0)$$

$$= \frac{1 - e^{-2\pi}}{2\pi} + \frac{(1 - e^{-2\pi})}{\pi} \left[-\frac{1}{2} + \sum_{n=2}^{\infty} \frac{1}{1+n^2} (-1)^n \right]$$

$$\therefore e^{-\pi} = \frac{1 - e^{-2\pi}}{\pi} \sum_{n=2}^{\infty} \frac{(-1)^n}{1+n^2} \quad \therefore \frac{\pi}{e^{\pi}(1 - e^{-2\pi})} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1+n^2}$$

$$\therefore \frac{\pi}{e^{\pi} - e^{-\pi}} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1+n^2} \quad \therefore \frac{\pi}{2} \cdot \frac{2}{e^{\pi} - e^{-\pi}} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1+n^2}$$

$$\therefore \frac{\pi}{2} \cdot \frac{1}{\sinh \pi} = \sum_{n=2}^{\infty} \frac{(-1)^n}{1+n^2}$$

$$\therefore \operatorname{cosec} h\pi = \frac{2}{\pi} \sum_{n=2}^{\infty} \frac{(-1)^n}{1+n^2} = \frac{2}{\pi} \left[\frac{1}{1+2^2} - \frac{1}{1+3^2} + \frac{1}{1+4^2} - \dots \right]$$

EX 4. Find Fourier expansion of $\cos px$ in $(0, 2\pi)$ where p is not an integer .

Hence deduce that $\pi \operatorname{csch} p\pi = \frac{1}{p} + \sum_{n=1}^{\infty} (-1)^n \left[\frac{1}{p+n} + \frac{1}{p-n} \right]$

Also deduce that $\pi \cot 2p\pi = \frac{1}{2p} + p \sum_{n=1}^{\infty} \frac{1}{p^2 - n^2}$

EX 4. Find Fourier expansion of $\cos px$ in $(0, 2\pi)$ where p is not an integer .

Hence deduce that $\pi \operatorname{csch} p\pi = \frac{1}{p} + \sum_{n=1}^{\infty} (-1)^n \left[\frac{1}{p+n} + \frac{1}{p-n} \right]$

Also deduce that $\pi \cot 2p\pi = \frac{1}{2p} + p \sum_{n=1}^{\infty} \frac{1}{p^2 - n^2}$

Solution:

$$: \text{Let } \cos px = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

EX 4. Find Fourier expansion of $\cos px$ in $(0, 2\pi)$ where p is not an integer .

Hence deduce that $\pi \operatorname{csch} p\pi = \frac{1}{p} + \sum_{n=1}^{\infty} (-1)^n \left[\frac{1}{p+n} + \frac{1}{p-n} \right]$

Also deduce that $\pi \cot 2p\pi = \frac{1}{2p} + p \sum_{n=1}^{\infty} \frac{1}{p^2 - n^2}$

Solution:

$$\therefore \text{Let } \cos px = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\therefore a_0 = \frac{1}{2\pi} \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} \cos px dx$$

EX 4. Find Fourier expansion of $\cos px$ in $(0, 2\pi)$ where p is not an integer .

Hence deduce that $\pi \operatorname{csch} p\pi = \frac{1}{p} + \sum_{n=1}^{\infty} (-1)^n \left[\frac{1}{p+n} + \frac{1}{p-n} \right]$

Also deduce that $\pi \cot 2p\pi = \frac{1}{2p} + p \sum_{n=1}^{\infty} \frac{1}{p^2 - n^2}$

Solution:

$$\therefore \text{Let } \cos px = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\therefore a_0 = \frac{1}{2\pi} \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} \cos px dx$$

$$\therefore a_0 = \frac{1}{2\pi} \left[\frac{\sin px}{p} \right]_0^{2\pi}$$

EX 4. Find Fourier expansion of $\cos px$ in $(0, 2\pi)$ where p is not an integer .

Hence deduce that $\pi \operatorname{csch} p\pi = \frac{1}{p} + \sum_{n=1}^{\infty} (-1)^n \left[\frac{1}{p+n} + \frac{1}{p-n} \right]$

Also deduce that $\pi \cot 2p\pi = \frac{1}{2p} + p \sum_{n=1}^{\infty} \frac{1}{p^2 - n^2}$

Solution:

$$\therefore \text{Let } \cos px = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\therefore a_0 = \frac{1}{2\pi} \int_0^{2\pi} f(x) dx = \frac{1}{2\pi} \int_0^{2\pi} \cos px dx$$

$$\therefore a_0 = \frac{1}{2\pi} \left[\frac{\sin px}{p} \right]_0^{2\pi} = \frac{1}{2\pi} \cdot \frac{\sin 2p\pi}{p} = \frac{\sin 2p\pi}{2\pi p}$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \cos nx \, dx$$

$$\begin{aligned}a_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \cos nx \, dx \\&= \frac{1}{2\pi} \int_0^{2\pi} 2 \cos px \cos nx \, dx\end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \cos nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 2 \cos px \cos nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} [\cos(p+n)x + \cos(p-n)x] \, dx
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \cos nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 2 \cos px \cos nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} [\cos(p+n)x + \cos(p-n)x] \, dx \\
 &= \frac{1}{2\pi} \left[\frac{\sin(p+n)x}{p+n} + \frac{\sin(p-n)x}{p-n} \right]_0^{2\pi}
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \cos nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 2 \cos px \cos nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} [\cos (p+n)x + \cos (p-n)x] \, dx \\
 &= \frac{1}{2\pi} \left[\frac{\sin (p+n)x}{p+n} + \frac{\sin (p-n)x}{p-n} \right]_0^{2\pi} \\
 &= \frac{1}{2\pi} \left[\frac{\sin 2\pi (p+n)}{p+n} + \frac{\sin 2\pi (p-n)}{p-n} \right]
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \cos nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 2 \cos px \cos nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} [\cos(p+n)x + \cos(p-n)x] \, dx \\
 &= \frac{1}{2\pi} \left[\frac{\sin(p+n)x}{p+n} + \frac{\sin(p-n)x}{p-n} \right]_0^{2\pi} \\
 &= \frac{1}{2\pi} \left[\frac{\sin 2\pi(p+n)}{p+n} + \frac{\sin 2\pi(p-n)}{p-n} \right] \\
 &= \frac{1}{2\pi} \left[\frac{\sin 2p\pi}{p+n} + \frac{\sin 2p\pi}{p-n} \right]
 \end{aligned}$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \cos nx \, dx$$

$$= \frac{1}{2\pi} \int_0^{2\pi} 2 \cos px \cos nx \, dx$$

$$= \frac{1}{2\pi} \int_0^{2\pi} [\cos(p+n)x + \cos(p-n)x] \, dx$$

$$= \frac{1}{2\pi} \left[\frac{\sin(p+n)x}{p+n} + \frac{\sin(p-n)x}{p-n} \right]_0^{2\pi}$$

$$= \frac{1}{2\pi} \left[\frac{\sin 2\pi(p+n)}{p+n} + \frac{\sin 2\pi(p-n)}{p-n} \right]$$

$$= \frac{1}{2\pi} \left[\frac{\sin 2p\pi}{p+n} + \frac{\sin 2p\pi}{p-n} \right]$$

$$\therefore a_n = \frac{1}{2\pi} \cdot \sin 2p\pi \left[\frac{1}{p+n} + \frac{1}{p-n} \right]$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \cos nx \, dx$$

$$= \frac{1}{2\pi} \int_0^{2\pi} 2 \cos px \cos nx \, dx$$

$$= \frac{1}{2\pi} \int_0^{2\pi} [\cos(p+n)x + \cos(p-n)x] \, dx$$

$$= \frac{1}{2\pi} \left[\frac{\sin(p+n)x}{p+n} + \frac{\sin(p-n)x}{p-n} \right]_0^{2\pi}$$

$$= \frac{1}{2\pi} \left[\frac{\sin 2\pi(p+n)}{p+n} + \frac{\sin 2\pi(p-n)}{p-n} \right]$$

$$= \frac{1}{2\pi} \left[\frac{\sin 2p\pi}{p+n} + \frac{\sin 2p\pi}{p-n} \right]$$

$$\therefore a_n = \frac{1}{2\pi} \cdot \sin 2p\pi \left[\frac{1}{p+n} + \frac{1}{p-n} \right] = \frac{p}{\pi} \cdot \frac{\sin 2p\pi}{p^2 - n^2}$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \sin nx \, dx$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \sin nx \, dx$$

$$= \frac{1}{2\pi} \int_0^{2\pi} 2 \sin nx \cos px \, dx :$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \sin nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 2 \sin nx \cos px \, dx = \frac{1}{2\pi} \int_0^{2\pi} [\sin(n+p)x + \sin(n-p)x] \, dx
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \sin nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 2 \sin nx \cos px \, dx = \frac{1}{2\pi} \int_0^{2\pi} [\sin(n+p)x + \sin(n-p)x] \, dx \\
 &= \frac{1}{2\pi} \left[-\frac{\cos(n+p)x}{n+p} - \frac{\cos(n-p)x}{n-p} \right]_0^{2\pi}
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \sin nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 2 \sin nx \cos px \, dx = \frac{1}{2\pi} \int_0^{2\pi} [\sin(n+p)x + \sin(n-p)x] \, dx \\
 &= \frac{1}{2\pi} \left[-\frac{\cos(n+p)x}{n+p} - \frac{\cos(n-p)x}{n-p} \right]_0^{2\pi} \\
 &= \frac{1}{2\pi} \left[-\frac{\cos 2\pi(n+p)}{n+p} - \frac{\cos 2\pi(n-p)}{n-p} + \frac{1}{n+p} + \frac{1}{n-p} \right]
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \sin nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 2 \sin nx \cos px \, dx = \frac{1}{2\pi} \int_0^{2\pi} [\sin(n+p)x + \sin(n-p)x] \, dx \\
 &= \frac{1}{2\pi} \left[-\frac{\cos(n+p)x}{n+p} - \frac{\cos(n-p)x}{n-p} \right]_0^{2\pi} \\
 &= \frac{1}{2\pi} \left[-\frac{\cos 2\pi(n+p)}{n+p} - \frac{\cos 2\pi(n-p)}{n-p} + \frac{1}{n+p} + \frac{1}{n-p} \right] \\
 &= \frac{1}{2\pi} \left[-\frac{\cos 2p\pi}{n+p} - \frac{\cos 2p\pi}{n-p} + \frac{1}{n+p} + \frac{1}{n-p} \right]
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \sin nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 2 \sin nx \cos px \, dx = \frac{1}{2\pi} \int_0^{2\pi} [\sin(n+p)x + \sin(n-p)x] \, dx \\
 &= \frac{1}{2\pi} \left[-\frac{\cos(n+p)x}{n+p} - \frac{\cos(n-p)x}{n-p} \right]_0^{2\pi} \\
 &= \frac{1}{2\pi} \left[-\frac{\cos 2\pi(n+p)}{n+p} - \frac{\cos 2\pi(n-p)}{n-p} + \frac{1}{n+p} + \frac{1}{n-p} \right] \\
 &= \frac{1}{2\pi} \left[-\frac{\cos 2p\pi}{n+p} - \frac{\cos 2p\pi}{n-p} + \frac{1}{n+p} + \frac{1}{n-p} \right] \\
 &= \frac{1}{2\pi} \left[\frac{1 - \cos 2p\pi}{n+p} - \frac{1 - \cos 2p\pi}{n-p} \right]
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \sin nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 2 \sin nx \cos px \, dx = \frac{1}{2\pi} \int_0^{2\pi} [\sin(n+p)x + \sin(n-p)x] \, dx \\
 &= \frac{1}{2\pi} \left[-\frac{\cos(n+p)x}{n+p} - \frac{\cos(n-p)x}{n-p} \right]_0^{2\pi} \\
 &= \frac{1}{2\pi} \left[-\frac{\cos 2\pi(n+p)}{n+p} - \frac{\cos 2\pi(n-p)}{n-p} + \frac{1}{n+p} + \frac{1}{n-p} \right] \\
 &= \frac{1}{2\pi} \left[-\frac{\cos 2p\pi}{n+p} - \frac{\cos 2p\pi}{n-p} + \frac{1}{n+p} + \frac{1}{n-p} \right] \\
 &= \frac{1}{2\pi} \left[\frac{1 - \cos 2p\pi}{n+p} - \frac{1 - \cos 2p\pi}{n-p} \right] \\
 &= \frac{1 - \cos 2p\pi}{2\pi} \left[\frac{1}{n+p} + \frac{1}{n-p} \right]
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \sin nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 2 \sin nx \cos px \, dx = \frac{1}{2\pi} \int_0^{2\pi} [\sin(n+p)x + \sin(n-p)x] \, dx \\
 &= \frac{1}{2\pi} \left[-\frac{\cos(n+p)x}{n+p} - \frac{\cos(n-p)x}{n-p} \right]_0^{2\pi} \\
 &= \frac{1}{2\pi} \left[-\frac{\cos 2\pi(n+p)}{n+p} - \frac{\cos 2\pi(n-p)}{n-p} + \frac{1}{n+p} + \frac{1}{n-p} \right] \\
 &= \frac{1}{2\pi} \left[-\frac{\cos 2p\pi}{n+p} - \frac{\cos 2p\pi}{n-p} + \frac{1}{n+p} + \frac{1}{n-p} \right] \\
 &= \frac{1}{2\pi} \left[\frac{1 - \cos 2p\pi}{n+p} - \frac{1 - \cos 2p\pi}{n-p} \right] \\
 &= \frac{1 - \cos 2p\pi}{2\pi} \left[\frac{1}{n+p} + \frac{1}{n-p} \right] = \frac{1 - \cos 2p\pi}{2\pi} \cdot \frac{2n}{n^2 - p^2}
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \sin nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 2 \sin nx \cos px \, dx = \frac{1}{2\pi} \int_0^{2\pi} [\sin(n+p)x + \sin(n-p)x] \, dx \\
 &= \frac{1}{2\pi} \left[-\frac{\cos(n+p)x}{n+p} - \frac{\cos(n-p)x}{n-p} \right]_0^{2\pi} \\
 &= \frac{1}{2\pi} \left[-\frac{\cos 2\pi(n+p)}{n+p} - \frac{\cos 2\pi(n-p)}{n-p} + \frac{1}{n+p} + \frac{1}{n-p} \right] \\
 &= \frac{1}{2\pi} \left[-\frac{\cos 2p\pi}{n+p} - \frac{\cos 2p\pi}{n-p} + \frac{1}{n+p} + \frac{1}{n-p} \right] \\
 &= \frac{1}{2\pi} \left[\frac{1 - \cos 2p\pi}{n+p} - \frac{1 - \cos 2p\pi}{n-p} \right] \\
 &= \frac{1 - \cos 2p\pi}{2\pi} \left[\frac{1}{n+p} + \frac{1}{n-p} \right] = \frac{1 - \cos 2p\pi}{2\pi} \cdot \frac{2n}{n^2 - p^2} \\
 \therefore b_n &= -\frac{n}{\pi} \frac{(1 - \cos 2p\pi)}{p^2 - n^2}
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} \cos px \sin nx \, dx \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 2 \sin nx \cos px \, dx = \frac{1}{2\pi} \int_0^{2\pi} [\sin(n+p)x + \sin(n-p)x] \, dx \\
 &= \frac{1}{2\pi} \left[-\frac{\cos(n+p)x}{n+p} - \frac{\cos(n-p)x}{n-p} \right]_0^{2\pi} \\
 &= \frac{1}{2\pi} \left[-\frac{\cos 2\pi(n+p)}{n+p} - \frac{\cos 2\pi(n-p)}{n-p} + \frac{1}{n+p} + \frac{1}{n-p} \right] \\
 &= \frac{1}{2\pi} \left[-\frac{\cos 2p\pi}{n+p} - \frac{\cos 2p\pi}{n-p} + \frac{1}{n+p} + \frac{1}{n-p} \right] \\
 &= \frac{1}{2\pi} \left[\frac{1 - \cos 2p\pi}{n+p} - \frac{1 - \cos 2p\pi}{n-p} \right] \\
 &= \frac{1 - \cos 2p\pi}{2\pi} \left[\frac{1}{n+p} + \frac{1}{n-p} \right] = \frac{1 - \cos 2p\pi}{2\pi} \cdot \frac{2n}{n^2 - p^2}
 \end{aligned}$$

$$\therefore b_n = -\frac{n(1 - \cos 2p\pi)}{\pi(p^2 - n^2)}$$

Putting these values in (1), we get,

$$\cos px = \frac{\sin 2p\pi}{2p\pi} + \frac{p \sin 2p\pi}{\pi} \sum_{n=1}^{\infty} \frac{\cos nx}{p^2 - n^2} - \frac{(1 - \cos 2p\pi)}{\pi} \sum_{n=1}^{\infty} \frac{n \sin nx}{p^2 - n^2} \quad \dots\dots\dots (2)$$

To deduce the first result, put $x = \pi$ in (2).

To deduce the first result, put $x = \pi$ in (2).

$$\begin{aligned}\therefore \cos p\pi &= \frac{\sin 2p\pi}{2p\pi} + \frac{p \sin 2p\pi}{\pi} \sum_{n=1}^{\infty} \frac{\cos n\pi}{p^2 - n^2} \\ &\quad - \frac{(1 - \cos 2p\pi)}{\pi} \sum_{n=1}^{\infty} \frac{n \sin n\pi}{p^2 - n^2}\end{aligned}$$

To deduce the first result, put $x = \pi$ in (2).

$$\begin{aligned}\therefore \cos p\pi &= \frac{\sin 2p\pi}{2p\pi} + \frac{p \sin 2p\pi}{\pi} \sum_{n=1}^{\infty} \frac{\cos n\pi}{p^2 - n^2} \\ &\quad - \frac{(1 - \cos 2p\pi)}{\pi} \sum_{n=1}^{\infty} \frac{n \sin n\pi}{p^2 - n^2}\end{aligned}$$

But since, n is a positive integer, $\sin n\pi = 0$ and $\cos n\pi = (-1)^n$.

To deduce the first result, put $x = \pi$ in (2).

$$\begin{aligned} \therefore \cos p\pi &= \frac{\sin 2p\pi}{2p\pi} + \frac{p \sin 2p\pi}{\pi} \sum_{n=1}^{\infty} \frac{\cos n\pi}{p^2 - n^2} \\ &\quad - \frac{(1 - \cos 2p\pi)}{\pi} \sum_{n=1}^{\infty} \frac{n \sin n\pi}{p^2 - n^2} \end{aligned}$$

But since, n is a positive integer, $\sin n\pi = 0$ and $\cos n\pi = (-1)^n$.

$$\therefore \cos p\pi = \frac{\sin 2p\pi}{2p\pi} + \frac{p \sin 2p\pi}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{p^2 - n^2}$$

To deduce the first result, put $x = \pi$ in (2).

$$\begin{aligned} \therefore \cos p\pi &= \frac{\sin 2p\pi}{2p\pi} + \frac{p \sin 2p\pi}{\pi} \sum_{n=1}^{\infty} \frac{\cos n\pi}{p^2 - n^2} \\ &\quad - \frac{(1 - \cos 2p\pi)}{\pi} \sum_{n=1}^{\infty} \frac{n \sin n\pi}{p^2 - n^2} \end{aligned}$$

But since, n is a positive integer, $\sin n\pi = 0$ and $\cos n\pi = (-1)^n$.

$$\begin{aligned} \therefore \cos p\pi &= \frac{\sin 2p\pi}{2p\pi} + \frac{p \sin 2p\pi}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{p^2 - n^2} \\ &= \frac{2 \sin p\pi \cos p\pi}{2p\pi} + \frac{2p \sin p\pi \cos p\pi}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{p^2 - n^2} \end{aligned}$$

To deduce the first result, put $x = \pi$ in (2).

$$\begin{aligned}\therefore \cos p\pi &= \frac{\sin 2p\pi}{2p\pi} + \frac{p \sin 2p\pi}{\pi} \sum_{n=1}^{\infty} \frac{\cos n\pi}{p^2 - n^2} \\ &\quad - \frac{(1 - \cos 2p\pi)}{\pi} \sum_{n=1}^{\infty} \frac{n \sin n\pi}{p^2 - n^2}\end{aligned}$$

But since, n is a positive integer, $\sin n\pi = 0$ and $\cos n\pi = (-1)^n$.

$$\begin{aligned}\therefore \cos p\pi &= \frac{\sin 2p\pi}{2p\pi} + \frac{p \sin 2p\pi}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{p^2 - n^2} \\ &= \frac{2 \sin p\pi \cos p\pi}{2p\pi} + \frac{2p \sin p\pi \cos p\pi}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{p^2 - n^2} \\ \therefore \pi \operatorname{cosec} p\pi &= \frac{1}{p} + 2p \sum_{n=1}^{\infty} \frac{(-1)^n}{p^2 - n^2} = \frac{1}{p} + \sum_{n=1}^{\infty} (-1)^n \left[\frac{1}{p+n} + \frac{1}{p-n} \right]\end{aligned}$$

Fourier series in the interval $(-\pi, \pi)$

Period of $f(x)$ is 2π

$f(x) = a_0 + \sum a_n \cos nx + \sum b_n \sin nx$, where

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx$$

Fourier series in the interval $(-\pi, \pi)$

Period of $f(x)$ is 2π

$f(x) = a_0 + \sum a_n \cos nx + \sum b_n \sin nx$, where

$$\begin{aligned} a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx \\ a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx \\ b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx \end{aligned}$$

There are three cases in this interval

Fourier series in the interval $(-\pi, \pi)$

Period of $f(x)$ is 2π

$f(x) = a_0 + \sum a_n \cos nx + \sum b_n \sin nx$, where

$$\begin{aligned} a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx \\ a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx \\ b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx \end{aligned}$$

There are three cases in this interval

1. $f(x)$ is neither even nor odd

Fourier series in the interval $(-\pi, \pi)$

Period of $f(x)$ is 2π

$f(x) = a_0 + \sum a_n \cos nx + \sum b_n \sin nx$, where

$$\begin{aligned} a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx \\ a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx \\ b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx \end{aligned}$$

There are three cases in this interval

1. $f(x)$ is neither even nor odd
2. $f(x)$ is even.
3. $f(x)$ is odd.

Fourier series in the interval $(-\pi, \pi)$

Period of $f(x)$ is 2π

$f(x) = a_0 + \sum a_n \cos nx + \sum b_n \sin nx$, where

$$\begin{aligned} a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx \\ a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx \\ b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx \end{aligned}$$

There are three cases in this interval

1. $f(x)$ is neither even nor odd
2. $f(x)$ is even. $[f(-x)=f(x)]$
3. $f(x)$ is odd.

Fourier series in the interval $(-\pi, \pi)$

Period of $f(x)$ is 2π

$f(x) = a_0 + \sum a_n \cos nx + \sum b_n \sin nx$, where

$$\begin{aligned} a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx \\ a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx \\ b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx \end{aligned}$$

There are three cases in this interval

1. $f(x)$ is neither even nor odd
2. $f(x)$ is even. $[f(-x)=f(x)]$ $b_n = 0$
3. $f(x)$ is odd.

Fourier series in the interval $(-\pi, \pi)$

Period of $f(x)$ is 2π

$f(x) = a_0 + \sum a_n \cos nx + \sum b_n \sin nx$, where

$$\begin{aligned} a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx \\ a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx \\ b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx \end{aligned}$$

There are three cases in this interval

1. $f(x)$ is neither even nor odd
2. $f(x)$ is even. $[f(-x)=f(x)]$ $b_n = 0$
3. $f(x)$ is odd. $[f(-x)=-f(x)]$

Fourier series in the interval $(-\pi, \pi)$

Period of $f(x)$ is 2π

$f(x) = a_0 + \sum a_n \cos nx + \sum b_n \sin nx$, where

$$\begin{aligned} a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx \\ a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx \\ b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx \end{aligned}$$

There are three cases in this interval

1. $f(x)$ is neither even nor odd
2. $f(x)$ is even. $[f(-x)=f(x)]$ $b_n = 0$
3. $f(x)$ is odd. $[f(-x)=-f(x)]$ $a_0 = 0, a_n = 0$

Fourier series in the interval $(-\pi, \pi)$

EX1 Find the Fourier series for a periodic function

$$f(x) = \begin{cases} -\pi, & -\pi < x < 0 \\ x, & 0 < x < \pi \end{cases}$$

State the value of $f(x)$ at $x = 0$ and hence deduce that

$$\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}$$

Fourier series in the interval $(-\pi, \pi)$

EX1 Find the Fourier series for a periodic function

$$f(x) = \begin{cases} -\pi, & -\pi < x < 0 \\ x, & 0 < x < \pi \end{cases}$$

State the value of $f(x)$ at $x = 0$ and hence deduce that

$$\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}$$

Solution:

$$\text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

Fourier series in the interval $(-\pi, \pi)$

EX1 Find the Fourier series for a periodic function

$$f(x) = \begin{cases} -\pi, & -\pi < x < 0 \\ x, & 0 < x < \pi \end{cases}$$

State the value of $f(x)$ at $x = 0$ and hence deduce that

$$\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}$$

Solution: Let $f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx$ (1)

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{2\pi} \left[\int_{-\pi}^0 -\pi \cdot dx + \int_0^{\pi} x dx \right]$$

Fourier series in the interval $(-\pi, \pi)$

EX1 Find the Fourier series for a periodic function

$$f(x) = \begin{cases} -\pi, & -\pi < x < 0 \\ x, & 0 < x < \pi \end{cases}$$

State the value of $f(x)$ at $x = 0$ and hence deduce that

$$\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}$$

Solution:

$$\text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\begin{aligned} a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{2\pi} \left[\int_{-\pi}^0 -\pi \cdot dx + \int_0^{\pi} x dx \right] \\ &= \frac{1}{2\pi} \left[-\pi \left\{ x \right\}^0 + \left\{ \frac{x^2}{2} \right\}^{\pi} \right] = \end{aligned}$$

Fourier series in the interval $(-\pi, \pi)$

EX1 Find the Fourier series for a periodic function

$$f(x) = \begin{cases} -\pi, & -\pi < x < 0 \\ x, & 0 < x < \pi \end{cases}$$

State the value of $f(x)$ at $x = 0$ and hence deduce that

$$\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}$$

Solution:

$$\text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \quad \dots\dots\dots (1)$$

$$\begin{aligned} a_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{2\pi} \left[\int_{-\pi}^0 -\pi \cdot dx + \int_0^{\pi} x dx \right] \\ &= \frac{1}{2\pi} \left[-\pi \left\{ x \right\}_{-\pi}^0 + \left\{ \frac{x^2}{2} \right\}_0^{\pi} \right] = \frac{1}{2\pi} \left[-\pi^2 + \frac{\pi^2}{2} \right] = -\frac{\pi}{4} \end{aligned}$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \cos nx \, dx + \int_0^{\pi} x \cos nx \, dx \right]$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \cos nx \, dx + \int_0^{\pi} x \cos nx \, dx \right] \\
 &= \frac{1}{\pi} \left[-\pi \left(\frac{\sin nx}{n} \right)_{-\pi}^0 + \left\{ x \left(\frac{\sin nx}{n} \right) - (1) \left(-\frac{\cos nx}{n^2} \right) \right\}_0^{\pi} \right]
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \cos nx \, dx + \int_0^{\pi} x \cos nx \, dx \right] \\
 &= \frac{1}{\pi} \left[-\pi \left(\frac{\sin nx}{n} \right)_{-\pi}^0 + \left\{ x \left(\frac{\sin nx}{n} \right) - (1) \left(-\frac{\cos nx}{n^2} \right) \right\}_0^{\pi} \right] \\
 &= \frac{1}{\pi} \left[-\pi (0 - 0) + \left\{ \pi (0) + \frac{\cos n\pi}{n^2} - 0 - \frac{1}{n^2} \right\} \right]
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \cos nx \, dx + \int_0^{\pi} x \cos nx \, dx \right] \\
 &= \frac{1}{\pi} \left[-\pi \left(\frac{\sin nx}{n} \right)_{-\pi}^0 + \left\{ x \left(\frac{\sin nx}{n} \right) - (1) \left(-\frac{\cos nx}{n^2} \right) \right\}_0^{\pi} \right] \\
 &= \frac{1}{\pi} \left[-\pi (0 - 0) + \left\{ \pi (0) + \frac{\cos n\pi}{n^2} - 0 - \frac{1}{n^2} \right\} \right] \\
 &= \frac{1}{\pi} \left[\frac{(-1)^n}{n^2} - \frac{1}{n^2} \right] = \frac{1}{\pi n^2} [(-1)^n - 1]
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \cos nx \, dx + \int_0^{\pi} x \cos nx \, dx \right] \\
 &= \frac{1}{\pi} \left[-\pi \left(\frac{\sin nx}{n} \right)_{-\pi}^0 + \left\{ x \left(\frac{\sin nx}{n} \right) - (1) \left(-\frac{\cos nx}{n^2} \right) \right\}_0^{\pi} \right] \\
 &= \frac{1}{\pi} \left[-\pi (0 - 0) + \left\{ \pi (0) + \frac{\cos n\pi}{n^2} - 0 - \frac{1}{n^2} \right\} \right] \\
 &= \frac{1}{\pi} \left[\frac{(-1)^n}{n^2} - \frac{1}{n^2} \right] = \frac{1}{\pi n^2} [(-1)^n - 1]
 \end{aligned}$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \cos nx \, dx + \int_0^{\pi} x \cos nx \, dx \right] \\
 &= \frac{1}{\pi} \left[-\pi \left(\frac{\sin nx}{n} \right)_{-\pi}^0 + \left\{ x \left(\frac{\sin nx}{n} \right) - (1) \left(-\frac{\cos nx}{n^2} \right) \right\}_0^{\pi} \right] \\
 &= \frac{1}{\pi} \left[-\pi (0 - 0) + \left\{ \pi (0) + \frac{\cos n\pi}{n^2} - 0 - \frac{1}{n^2} \right\} \right] \\
 &= \frac{1}{\pi} \left[\frac{(-1)^n}{n^2} - \frac{1}{n^2} \right] = \frac{1}{\pi n^2} [(-1)^n - 1] \\
 b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \sin nx \, dx + \int_0^{\pi} x \sin nx \, dx \right]
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \cos nx \, dx + \int_0^{\pi} x \cos nx \, dx \right] \\
 &= \frac{1}{\pi} \left[-\pi \left(\frac{\sin nx}{n} \right)_{-\pi}^0 + \left\{ x \left(\frac{\sin nx}{n} \right) - (1) \left(-\frac{\cos nx}{n^2} \right) \right\}_0^{\pi} \right] \\
 &= \frac{1}{\pi} \left[-\pi (0 - 0) + \left\{ \pi (0) + \frac{\cos n\pi}{n^2} - 0 - \frac{1}{n^2} \right\} \right] \\
 &= \frac{1}{\pi} \left[\frac{(-1)^n}{n^2} - \frac{1}{n^2} \right] = \frac{1}{\pi n^2} [(-1)^n - 1]
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \sin nx \, dx + \int_0^{\pi} x \sin nx \, dx \right] \\
 &= \frac{1}{\pi} \left[-\pi \left(-\frac{\cos nx}{n} \right)_{-\pi}^0 + \left\{ x \left(-\frac{\cos nx}{n} \right) - (1) \left(-\frac{\sin nx}{n} \right) \right\}_0^{\pi} \right]
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \cos nx \, dx + \int_0^{\pi} x \cos nx \, dx \right] \\
 &= \frac{1}{\pi} \left[-\pi \left(\frac{\sin nx}{n} \right)_{-\pi}^0 + \left\{ x \left(\frac{\sin nx}{n} \right) - (1) \left(-\frac{\cos nx}{n^2} \right) \right\}_0^{\pi} \right] \\
 &= \frac{1}{\pi} \left[-\pi (0 - 0) + \left\{ \pi (0) + \frac{\cos n\pi}{n^2} - 0 - \frac{1}{n^2} \right\} \right] \\
 &= \frac{1}{\pi} \left[\frac{(-1)^n}{n^2} - \frac{1}{n^2} \right] = \frac{1}{\pi n^2} [(-1)^n - 1]
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \sin nx \, dx + \int_0^{\pi} x \sin nx \, dx \right] \\
 &= \frac{1}{\pi} \left[-\pi \left(-\frac{\cos nx}{n} \right)_{-\pi}^0 + \left\{ x \left(-\frac{\cos nx}{n} \right) - (1) \left(-\frac{\sin nx}{n} \right) \right\}_0^{\pi} \right] \\
 &= \frac{1}{\pi} \left[\pi \frac{(1 - \cos n\pi)}{n} + \left\{ \pi \left(-\frac{\cos n\pi}{n} \right) - 0 \right\} \right]
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \cos nx \, dx + \int_0^{\pi} x \cos nx \, dx \right] \\
 &= \frac{1}{\pi} \left[-\pi \left(\frac{\sin nx}{n} \right)_{-\pi}^0 + \left\{ x \left(\frac{\sin nx}{n} \right) - (1) \left(-\frac{\cos nx}{n^2} \right) \right\}_0^{\pi} \right] \\
 &= \frac{1}{\pi} \left[-\pi (0 - 0) + \left\{ \pi (0) + \frac{\cos n\pi}{n^2} - 0 - \frac{1}{n^2} \right\} \right] \\
 &= \frac{1}{\pi} \left[\frac{(-1)^n}{n^2} - \frac{1}{n^2} \right] = \frac{1}{\pi n^2} [(-1)^n - 1]
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \sin nx \, dx + \int_0^{\pi} x \sin nx \, dx \right] \\
 &= \frac{1}{\pi} \left[-\pi \left(-\frac{\cos nx}{n} \right)_{-\pi}^0 + \left\{ x \left(-\frac{\cos nx}{n} \right) - (1) \left(-\frac{\sin nx}{n} \right) \right\}_0^{\pi} \right] \\
 &= \frac{1}{\pi} \left[\pi \frac{(1 - \cos n\pi)}{n} + \left\{ \pi \left(-\frac{\cos n\pi}{n} \right) - 0 \right\} \right] \\
 &= \frac{1}{\pi} \left[\frac{\pi}{n} (1 - \cos n\pi) - \frac{\pi}{n} \cos n\pi \right]
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \cos nx \, dx + \int_0^{\pi} x \cos nx \, dx \right] \\
 &= \frac{1}{\pi} \left[-\pi \left(\frac{\sin nx}{n} \right)_{-\pi}^0 + \left\{ x \left(\frac{\sin nx}{n} \right) - (1) \left(-\frac{\cos nx}{n^2} \right) \right\}_0^{\pi} \right] \\
 &= \frac{1}{\pi} \left[-\pi (0 - 0) + \left\{ \pi (0) + \frac{\cos n\pi}{n^2} - 0 - \frac{1}{n^2} \right\} \right] \\
 &= \frac{1}{\pi} \left[\frac{(-1)^n}{n^2} - \frac{1}{n^2} \right] = \frac{1}{\pi n^2} [(-1)^n - 1]
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 -\pi \sin nx \, dx + \int_0^{\pi} x \sin nx \, dx \right] \\
 &= \frac{1}{\pi} \left[-\pi \left(-\frac{\cos nx}{n} \right)_{-\pi}^0 + \left\{ x \left(-\frac{\cos nx}{n} \right) - (1) \left(-\frac{\sin nx}{n} \right) \right\}_0^{\pi} \right] \\
 &= \frac{1}{\pi} \left[\pi \frac{(1 - \cos n\pi)}{n} + \left\{ \pi \left(-\frac{\cos n\pi}{n} \right) - 0 \right\} \right] \\
 &= \frac{1}{\pi} \left[\frac{\pi}{n} (1 - \cos n\pi) - \frac{\pi}{n} \cos n\pi \right] \\
 &= \frac{1}{n} [1 - 2 \cos n\pi] = \frac{1}{n} [1 - 2(-1)^n]
 \end{aligned}$$

Putting these values in (1),

Putting these values in (1),

$$f(x) = -\frac{\pi}{4} + \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n - 1}{n^2} \cos nx + \sum_{n=1}^{\infty} \frac{1 - 2(-1)^n}{n} \sin nx$$

Putting these values in (1),

$$\begin{aligned} f(x) &= -\frac{\pi}{4} + \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n - 1}{n^2} \cos nx + \sum_{n=1}^{\infty} \frac{1 - 2(-1)^n}{n} \sin nx \\ &= -\frac{\pi}{4} - \frac{2}{\pi} \left[\frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} + \dots \right] \\ &\quad + \left[3 \cdot \frac{\sin x}{1} - \frac{1}{2} \sin 2x + \sin 3x - \dots \right] \end{aligned} \quad \text{..... (2)}$$

deduction

State the value of $f(x)$ at $x = 0$ and hence deduce that

$$\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}$$

Putting these values in (1),

$$\begin{aligned}
 f(x) &= -\frac{\pi}{4} + \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n - 1}{n^2} \cos nx + \sum_{n=1}^{\infty} \frac{1 - 2(-1)^n}{n} \sin nx \\
 &= -\frac{\pi}{4} - \frac{2}{\pi} \left[\frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} + \dots \right] \\
 &\quad + \left[3 \cdot \frac{\sin x}{1} - \frac{1}{2} \sin 2x + \sin 3x - \dots \right] \dots\dots\dots (2)
 \end{aligned}$$

Now, $f(x)$ is discontinuous at $x = 0$. At a point of discontinuity $x = c$,

Putting these values in (1),

$$\begin{aligned}
 f(x) &= -\frac{\pi}{4} + \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n - 1}{n^2} \cos nx + \sum_{n=1}^{\infty} \frac{1 - 2(-1)^n}{n} \sin nx \\
 &= -\frac{\pi}{4} - \frac{2}{\pi} \left[\frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} + \dots \right] \\
 &\quad + \left[3 \cdot \frac{\sin x}{1} - \frac{1}{2} \sin 2x + \sin 3x - \dots \right] \dots\dots\dots (2)
 \end{aligned}$$

Now, $f(x)$ is discontinuous at $x = 0$. At a point of discontinuity $x = c$,

$$f(x) = \frac{1}{2} \left[\lim_{x \rightarrow c^-} f(x) + \lim_{x \rightarrow c^+} f(x) \right]$$

Putting these values in (1),

$$\begin{aligned}
 f(x) &= -\frac{\pi}{4} + \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n - 1}{n^2} \cos nx + \sum_{n=1}^{\infty} \frac{1 - 2(-1)^n}{n} \sin nx \\
 &= -\frac{\pi}{4} - \frac{2}{\pi} \left[\frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} + \dots \right] \\
 &\quad + \left[3 \cdot \frac{\sin x}{1} - \frac{1}{2} \sin 2x + \sin 3x - \dots \right] \dots\dots\dots (2)
 \end{aligned}$$

Now, $f(x)$ is discontinuous at $x = 0$. At a point of discontinuity $x = c$,

$$f(x) = \frac{1}{2} \left[\lim_{x \rightarrow c^-} f(x) + \lim_{x \rightarrow c^+} f(x) \right]$$

$$\therefore f(0) = \frac{1}{2} [-\pi + 0] = -\frac{\pi}{2}$$

Putting these values in (1),

$$\begin{aligned}
 f(x) &= -\frac{\pi}{4} + \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n - 1}{n^2} \cos nx + \sum_{n=1}^{\infty} \frac{1 - 2(-1)^n}{n} \sin nx \\
 &= -\frac{\pi}{4} - \frac{2}{\pi} \left[\frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} + \dots \right] \\
 &\quad + \left[3 \cdot \frac{\sin x}{1} - \frac{1}{2} \sin 2x + \sin 3x - \dots \right] \dots\dots\dots (2)
 \end{aligned}$$

Now, $f(x)$ is discontinuous at $x = 0$. At a point of discontinuity $x = c$,

$$f(x) = \frac{1}{2} \left[\lim_{x \rightarrow c^-} f(x) + \lim_{x \rightarrow c^+} f(x) \right]$$

$$\therefore f(0) = \frac{1}{2} [-\pi + 0] = -\frac{\pi}{2}$$

Hence, putting $x = 0$ in (2),

$$-\frac{\pi}{2} = -\frac{\pi}{4} - \frac{2}{\pi} \left[\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right]$$

Putting these values in (1),

$$\begin{aligned}
 f(x) &= -\frac{\pi}{4} + \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n - 1}{n^2} \cos nx + \sum_{n=1}^{\infty} \frac{1 - 2(-1)^n}{n} \sin nx \\
 &= -\frac{\pi}{4} - \frac{2}{\pi} \left[\frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} + \dots \right] \\
 &\quad + \left[3 \cdot \frac{\sin x}{1} - \frac{1}{2} \sin 2x + \sin 3x - \dots \right] \dots\dots\dots (2)
 \end{aligned}$$

Now, $f(x)$ is discontinuous at $x = 0$. At a point of discontinuity $x = c$,

$$f(x) = \frac{1}{2} \left[\lim_{x \rightarrow c^-} f(x) + \lim_{x \rightarrow c^+} f(x) \right]$$

$$\therefore f(0) = \frac{1}{2} [-\pi + 0] = -\frac{\pi}{2}$$

Hence, putting $x = 0$ in (2),

$$-\frac{\pi}{2} = -\frac{\pi}{4} - \frac{2}{\pi} \left[\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right]$$

$$\therefore \frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \sum_{n=1}^{\infty} \frac{1}{(2n-1)^2}$$

EX 2 Find the Fourier series for $f(x) = e^{-|x|}$ in $(0, 2\pi)$

EX 2 Find the Fourier series for $f(x) = e^{-|x|}$ in $(0, 2\pi)$

Solution:

: We have $f(x) = e^{-|x|}$.

EX 2 Find the Fourier series for $f(x) = e^{-|x|}$ in $(0, 2\pi)$

Solution:

: We have $f(x) = e^{-|x|}$.

$$\therefore f(-x) = e^{-|-x|} = e^{-|x|} = f(x)$$

EX 2 Find the Fourier series for $f(x) = e^{-|x|}$ in $(0, 2\pi)$

Solution:

: We have $f(x) = e^{-|x|}$.

$$\therefore f(-x) = e^{-|-x|} = e^{-|x|} = f(x)$$

Hence, $f(x)$ is an even function and hence $b_n = 0$.

EX 2 Find the Fourier series for $f(x) = e^{-|x|}$ in $(0, 2\pi)$

Solution:

: We have $f(x) = e^{-|x|}$.

$$\therefore f(-x) = e^{-|-x|} = e^{-|x|} = f(x)$$

Hence, $f(x)$ is an even function and hence $b_n = 0$.

$$\text{Further, } f(x) = \begin{cases} e^x, & -\pi < x < 0 \\ e^{-x}, & 0 < x < \pi \end{cases}$$

EX 2 Find the Fourier series for $f(x) = e^{-|x|}$ in $(0, 2\pi)$

Solution:

: We have $f(x) = e^{-|x|}$.

$$\therefore f(-x) = e^{-|-x|} = e^{-|x|} = f(x)$$

Hence, $f(x)$ is an even function and hence $b_n = 0$.

$$\text{Further, } f(x) = \begin{cases} e^x, & -\pi < x < 0 \\ e^{-x}, & 0 < x < \pi \end{cases}$$

Now, using (A) above,

$$a_0 = \frac{1}{\pi} \int_0^\pi f(x) dx, \quad a_n = \frac{2}{\pi} \int_0^\pi f(x) \cos nx dx \quad \text{and} \quad b_n = 0.$$

EX 2 Find the Fourier series for $f(x) = e^{-|x|}$ in $(0, 2\pi)$

Solution:

: We have $f(x) = e^{-|x|}$.

$$\therefore f(-x) = e^{-|-x|} = e^{-|x|} = f(x)$$

Hence, $f(x)$ is an even function and hence $b_n = 0$.

$$\text{Further, } f(x) = \begin{cases} e^x, & -\pi < x < 0 \\ e^{-x}, & 0 < x < \pi \end{cases}$$

Now, using (A) above,

$$a_0 = \frac{1}{\pi} \int_0^\pi f(x) dx, \quad a_n = \frac{2}{\pi} \int_0^\pi f(x) \cos nx dx \quad \text{and} \quad b_n = 0.$$

$$\therefore a_0 = \frac{1}{\pi} \int_0^\pi e^{-x} dx = \frac{1}{\pi} \left[-e^{-x} \right]_0^\pi = \frac{1}{\pi} (1 - e^{-\pi})$$

EX 2 Find the Fourier series for $f(x) = e^{-|x|}$ in $(0, 2\pi)$

Solution:

: We have $f(x) = e^{-|x|}$.

$$\therefore f(-x) = e^{-|-x|} = e^{-|x|} = f(x)$$

Hence, $f(x)$ is an even function and hence $b_n = 0$.

$$\text{Further, } f(x) = \begin{cases} e^x, & -\pi < x < 0 \\ e^{-x}, & 0 < x < \pi \end{cases}$$

Now, using (A) above,

$$a_0 = \frac{1}{\pi} \int_0^\pi f(x) dx, \quad a_n = \frac{2}{\pi} \int_0^\pi f(x) \cos nx dx \text{ and } b_n = 0.$$

$$\therefore a_0 = \frac{1}{\pi} \int_0^\pi e^{-x} dx = \frac{1}{\pi} \left[-e^{-x} \right]_0^\pi = \frac{1}{\pi} (1 - e^{-\pi})$$

$$a_n = \frac{2}{\pi} \int_0^\pi f(x) \cos nx dx = \frac{2}{\pi} \int_0^\pi e^{-x} \cos nx dx$$

EX 2 Find the Fourier series for $f(x) = e^{-|x|}$ in $(0, 2\pi)$

Solution:

: We have $f(x) = e^{-|x|}$.

$$\therefore f(-x) = e^{-|-x|} = e^{-|x|} = f(x)$$

Hence, $f(x)$ is an even function and hence $b_n = 0$.

$$\text{Further, } f(x) = \begin{cases} e^x, & -\pi < x < 0 \\ e^{-x}, & 0 < x < \pi \end{cases}$$

Now, using (A) above,

$$a_0 = \frac{1}{\pi} \int_0^\pi f(x) dx, \quad a_n = \frac{2}{\pi} \int_0^\pi f(x) \cos nx dx \quad \text{and} \quad b_n = 0.$$

$$\therefore a_0 = \frac{1}{\pi} \int_0^\pi e^{-x} dx = \frac{1}{\pi} \left[-e^{-x} \right]_0^\pi = \frac{1}{\pi} (1 - e^{-\pi})$$

$$\begin{aligned} a_n &= \frac{2}{\pi} \int_0^\pi f(x) \cos nx dx = \frac{2}{\pi} \int_0^\pi e^{-x} \cos nx dx \\ &= \frac{2}{\pi} \left[\frac{e^{-x}}{1+n^2} (-\cos nx + n \sin nx) \right]_0^\pi \end{aligned}$$

EX 2 Find the Fourier series for $f(x) = e^{-|x|}$ in $(0, 2\pi)$

Solution:

: We have $f(x) = e^{-|x|}$.

$$\therefore f(-x) = e^{-|-x|} = e^{-|x|} = f(x)$$

Hence, $f(x)$ is an even function and hence $b_n = 0$.

$$\text{Further, } f(x) = \begin{cases} e^x, & -\pi < x < 0 \\ e^{-x}, & 0 < x < \pi \end{cases}$$

Now, using (A) above,

$$a_0 = \frac{1}{\pi} \int_0^\pi f(x) dx, \quad a_n = \frac{2}{\pi} \int_0^\pi f(x) \cos nx dx \quad \text{and} \quad b_n = 0.$$

$$\therefore a_0 = \frac{1}{\pi} \int_0^\pi e^{-x} dx = \frac{1}{\pi} \left[-e^{-x} \right]_0^\pi = \frac{1}{\pi} (1 - e^{-\pi})$$

$$a_n = \frac{2}{\pi} \int_0^\pi f(x) \cos nx dx = \frac{2}{\pi} \int_0^\pi e^{-x} \cos nx dx$$

$$= \frac{2}{\pi} \left[\frac{e^{-x}}{1+n^2} (-\cos nx + n \sin nx) \right]_0^\pi$$

$$= \frac{2}{\pi(1+n^2)} \left[-e^{-\pi} \cos n\pi - (-1) \right]$$

EX 2 Find the Fourier series for $f(x) = e^{-|x|}$ in $(0, 2\pi)$

Solution:

: We have $f(x) = e^{-|x|}$.

$$\therefore f(-x) = e^{-|-x|} = e^{-|x|} = f(x)$$

Hence, $f(x)$ is an even function and hence $b_n = 0$.

$$\text{Further, } f(x) = \begin{cases} e^x, & -\pi < x < 0 \\ e^{-x}, & 0 < x < \pi \end{cases}$$

Now, using (A) above,

$$a_0 = \frac{1}{\pi} \int_0^{\pi} f(x) dx, \quad a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx dx \text{ and } b_n = 0.$$

$$\therefore a_0 = \frac{1}{\pi} \int_0^{\pi} e^{-x} dx = \frac{1}{\pi} \left[-e^{-x} \right]_0^{\pi} = \frac{1}{\pi} (1 - e^{-\pi})$$

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx dx = \frac{2}{\pi} \int_0^{\pi} e^{-x} \cos nx dx$$

$$= \frac{2}{\pi} \left[\frac{e^{-x}}{1+n^2} (-\cos nx + n \sin nx) \right]_0^{\pi}$$

$$= \frac{2}{\pi(1+n^2)} \left[-e^{-\pi} \cos n\pi - (-1) \right]$$

$$\therefore a_n = \frac{2}{\pi(1+n^2)} \left[1 - (-1)^n e^{-\pi} \right]$$

EX 2 Find the Fourier series for $f(x) = e^{-|x|}$ in $(-\pi, \pi)$

Solution:

: We have $f(x) = e^{-|x|}$.

$$\therefore f(-x) = e^{-|-x|} = e^{-|x|} = f(x)$$

Hence, $f(x)$ is an even function and hence $b_n = 0$.

$$\text{Further, } f(x) = \begin{cases} e^x, & -\pi < x < 0 \\ e^{-x}, & 0 < x < \pi \end{cases}$$

Now, using (A) above,

$$a_0 = \frac{1}{\pi} \int_0^{\pi} f(x) dx, \quad a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx dx \text{ and } b_n = 0.$$

$$\therefore a_0 = \frac{1}{\pi} \int_0^{\pi} e^{-x} dx = \frac{1}{\pi} \left[-e^{-x} \right]_0^{\pi} = \frac{1}{\pi} (1 - e^{-\pi})$$

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx dx = \frac{2}{\pi} \int_0^{\pi} e^{-x} \cos nx dx$$

$$= \frac{2}{\pi} \left[\frac{e^{-x}}{1+n^2} (-\cos nx + n \sin nx) \right]_0^{\pi}$$

$$= \frac{2}{\pi(1+n^2)} \left[-e^{-\pi} \cos n\pi - (-1) \right]$$

$$\therefore a_n = \frac{2}{\pi(1+n^2)} \left[1 - (-1)^n e^{-\pi} \right]$$

$$\therefore f(x) = \frac{1}{\pi} (1 - e^{-\pi}) + \frac{2}{\pi} \sum_{n=1}^{\infty} \left[\frac{1 - (-1)^n e^{-\pi}}{1+n^2} \right] \cdot \cos nx$$

EX 3 Obtain Fourier expansion of $f(x) = |\cos x|$ in $(0, 2\pi)$

EX 3 Obtain Fourier expansion of $f(x) = |\cos x|$ in $(0, 2\pi)$

Solution:

Here $f(x)$ is an even function because $f(-x) = |\cos(-x)| = f(x)$. $[\therefore b_n = 0]$

EX 3 Obtain Fourier expansion of $f(x) = |\cos x|$ in $(0, 2\pi)$

Solution:

Here $f(x)$ is an even function because $f(-x) = |\cos(-x)| = f(x)$. $[\therefore b_n = 0]$

$$\text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx \quad \dots\dots\dots (1)$$

EX 3 Obtain Fourier expansion of $f(x) = |\cos x|$ in $(0, 2\pi)$

Solution:

Here $f(x)$ is an even function because $f(-x) = |\cos(-x)| = f(x)$. [$\therefore b_n = 0$]

$$\text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx \quad \dots\dots\dots (1)$$

$$\therefore a_0 = \frac{1}{\pi} \int_0^{\pi} f(x) dx = \frac{1}{\pi} \int_0^{\pi} |\cos x| dx$$

EX 3 Obtain Fourier expansion of $f(x) = |\cos x|$ in $(0, 2\pi)$

Solution:

Here $f(x)$ is an even function because $f(-x) = |\cos(-x)| = f(x)$. [$\therefore b_n = 0$]

$$\text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx \quad \dots\dots\dots (1)$$

$$\begin{aligned} \therefore a_0 &= \frac{1}{\pi} \int_0^{\pi} f(x) dx = \frac{1}{\pi} \int_0^{\pi} |\cos x| dx \\ &= \frac{1}{\pi} \left[\int_0^{\pi/2} \cos x dx - \int_{\pi/2}^{\pi} \cos x \cdot dx \right] \end{aligned}$$

EX 3 Obtain Fourier expansion of $f(x) = |\cos x|$ in $(0, 2\pi)$

Solution:

Here $f(x)$ is an even function because $f(-x) = |\cos(-x)| = f(x)$. [$\therefore b_n = 0$]

$$\text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx \quad \dots\dots\dots (1)$$

$$\begin{aligned} \therefore a_0 &= \frac{1}{\pi} \int_0^{\pi} f(x) dx = \frac{1}{\pi} \int_0^{\pi} |\cos x| dx \\ &= \frac{1}{\pi} \left[\int_0^{\pi/2} \cos x dx - \int_{\pi/2}^{\pi} \cos x \cdot dx \right] \end{aligned}$$

$$\begin{aligned} &[\because |\cos x| = \cos x, \quad \text{for } 0 < x < \pi/2 \\ &\text{and } |\cos x| = -\cos x, \quad \text{for } \pi/2 < x < \pi.] \end{aligned}$$

EX 3 Obtain Fourier expansion of $f(x) = |\cos x|$ in $(0, 2\pi)$

Solution:

Here $f(x)$ is an even function because $f(-x) = |\cos(-x)| = f(x)$. [$\therefore b_n = 0$]

$$\text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx \quad \dots\dots\dots (1)$$

$$\begin{aligned} \therefore a_0 &= \frac{1}{\pi} \int_0^{\pi} f(x) dx = \frac{1}{\pi} \int_0^{\pi} |\cos x| dx \\ &= \frac{1}{\pi} \left[\int_0^{\pi/2} \cos x dx - \int_{\pi/2}^{\pi} \cos x \cdot dx \right] \end{aligned}$$

[$\therefore |\cos x| = \cos x$, for $0 < x < \pi/2$
and $|\cos x| = -\cos x$, for $\pi/2 < x < \pi$.]

$$= \frac{1}{\pi} \left[\{\sin x\}_0^{\pi/2} - \{\sin x\}_{\pi/2}^{\pi} \right]$$

EX 3 Obtain Fourier expansion of $f(x) = |\cos x|$ in $(0, 2\pi)$

Solution:

Here $f(x)$ is an even function because $f(-x) = |\cos(-x)| = f(x)$. [$\therefore b_n = 0$]

$$\text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos nx \quad \dots\dots\dots (1)$$

$$\begin{aligned} \therefore a_0 &= \frac{1}{\pi} \int_0^{\pi} f(x) dx = \frac{1}{\pi} \int_0^{\pi} |\cos x| dx \\ &= \frac{1}{\pi} \left[\int_0^{\pi/2} \cos x dx - \int_{\pi/2}^{\pi} \cos x \cdot dx \right] \end{aligned}$$

[$\therefore |\cos x| = \cos x$, for $0 < x < \pi/2$
and $|\cos x| = -\cos x$, for $\pi/2 < x < \pi$.]

$$= \frac{1}{\pi} \left[\{\sin x\}_0^{\pi/2} - \{\sin x\}_{\pi/2}^{\pi} \right]$$

$$\therefore a_0 = \frac{1}{\pi} [\{1 - 0\} - \{0 - 1\}] = \frac{2}{\pi} \quad \dots\dots\dots (2)$$

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx$$

$$\begin{aligned}
 a_n &= \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx \\
 &= \frac{2}{\pi} \left[\int_0^{\pi/2} \cos x \cos nx \, dx - \int_{\pi/2}^{\pi} \cos x \cos nx \, dx \right] \dots\dots\dots (3)
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx \\
 &= \frac{2}{\pi} \left[\int_0^{\pi/2} \cos x \cos nx \, dx - \int_{\pi/2}^{\pi} \cos x \cos nx \, dx \right] \\
 &= \frac{1}{\pi} \left[\int_0^{\pi/2} 2 \cos nx \cos x \, dx - \int_{\pi/2}^{\pi} 2 \cos nx \cos x \, dx \right]
 \end{aligned}
 \dots\dots\dots (3)$$

$$\begin{aligned}
 a_n &= \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx \\
 &= \frac{2}{\pi} \left[\int_0^{\pi/2} \cos x \cos nx \, dx - \int_{\pi/2}^{\pi} \cos x \cos nx \, dx \right] \dots\dots\dots (3) \\
 &= \frac{1}{\pi} \left[\int_0^{\pi/2} 2 \cos nx \cos x \, dx - \int_{\pi/2}^{\pi} 2 \cos nx \cos x \, dx \right] \\
 &= \frac{1}{\pi} \left[\int_0^{\pi/2} \{ \cos (n+1)x + \cos (n-1)x \} \, dx - \int_{\pi/2}^{\pi} \{ \cos (n+1)x + \cos (n-1)x \} \, dx \right]
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx \\
 &= \frac{2}{\pi} \left[\int_0^{\pi/2} \cos x \cos nx \, dx - \int_{\pi/2}^{\pi} \cos x \cos nx \, dx \right] \dots\dots\dots (3) \\
 &= \frac{1}{\pi} \left[\int_0^{\pi/2} 2 \cos nx \cos x \, dx - \int_{\pi/2}^{\pi} 2 \cos nx \cos x \, dx \right] \\
 &= \frac{1}{\pi} \left[\int_0^{\pi/2} \{ \cos (n+1)x + \cos (n-1)x \} \, dx - \int_{\pi/2}^{\pi} \{ \cos (n+1)x + \cos (n-1)x \} \, dx \right] \\
 &= \frac{1}{\pi} \left[\left\{ \frac{\sin (n+1)x}{n+1} + \frac{\sin (n-1)x}{n-1} \right\}_0^{\pi/2} - \left\{ \frac{\sin (n+1)x}{n+1} + \frac{\sin (n-1)x}{n-1} \right\}_{\pi/2}^{\pi} \right]
 \end{aligned}$$

$$\begin{aligned}
a_n &= \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx \\
&= \frac{2}{\pi} \left[\int_0^{\pi/2} \cos x \cos nx \, dx - \int_{\pi/2}^{\pi} \cos x \cos nx \, dx \right] \dots\dots\dots (3) \\
&= \frac{1}{\pi} \left[\int_0^{\pi/2} 2 \cos nx \cos x \, dx - \int_{\pi/2}^{\pi} 2 \cos nx \cos x \, dx \right] \\
&= \frac{1}{\pi} \left[\int_0^{\pi/2} \{ \cos (n+1)x + \cos (n-1)x \} \, dx - \int_{\pi/2}^{\pi} \{ \cos (n+1)x + \cos (n-1)x \} \, dx \right] \\
&= \frac{1}{\pi} \left[\left\{ \frac{\sin (n+1)x}{n+1} + \frac{\sin (n-1)x}{n-1} \right\}_0^{\pi/2} - \left\{ \frac{\sin (n+1)x}{n+1} + \frac{\sin (n-1)x}{n-1} \right\}_{\pi/2}^{\pi} \right] \\
&= \frac{1}{\pi} \left[\left\{ \frac{\sin (n+1)\pi/2}{n+1} + \frac{\sin (n-1)\pi/2}{n-1} - 0 \right\} - \left\{ 0 - \frac{\sin (n+1)\pi/2}{n+1} - \frac{\sin (n-1)\pi/2}{n-1} \right\} \right]
\end{aligned}$$

$$\begin{aligned}
a_n &= \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx \\
&= \frac{2}{\pi} \left[\int_0^{\pi/2} \cos x \cos nx \, dx - \int_{\pi/2}^{\pi} \cos x \cos nx \, dx \right] \dots\dots\dots (3) \\
&= \frac{1}{\pi} \left[\int_0^{\pi/2} 2 \cos nx \cos x \, dx - \int_{\pi/2}^{\pi} 2 \cos nx \cos x \, dx \right] \\
&= \frac{1}{\pi} \left[\int_0^{\pi/2} \{ \cos (n+1)x + \cos (n-1)x \} \, dx - \int_{\pi/2}^{\pi} \{ \cos (n+1)x + \cos (n-1)x \} \, dx \right] \\
&= \frac{1}{\pi} \left[\left\{ \frac{\sin (n+1)x}{n+1} + \frac{\sin (n-1)x}{n-1} \right\}_0^{\pi/2} - \left\{ \frac{\sin (n+1)x}{n+1} + \frac{\sin (n-1)x}{n-1} \right\}_{\pi/2}^{\pi} \right] \\
&= \frac{1}{\pi} \left[\left\{ \frac{\sin (n+1)\pi/2}{n+1} + \frac{\sin (n-1)\pi/2}{n-1} - 0 \right\} - \left\{ 0 - \frac{\sin (n+1)\pi/2}{n+1} - \frac{\sin (n-1)\pi/2}{n-1} \right\} \right] \\
&= \frac{2}{\pi} \left[\frac{\sin (n+1)\pi/2}{n+1} + \frac{\sin (n-1)\pi/2}{n-1} \right]
\end{aligned}$$

$$\begin{aligned}
a_n &= \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx \\
&= \frac{2}{\pi} \left[\int_0^{\pi/2} \cos x \cos nx \, dx - \int_{\pi/2}^{\pi} \cos x \cos nx \, dx \right] \dots\dots\dots (3) \\
&= \frac{1}{\pi} \left[\int_0^{\pi/2} 2 \cos nx \cos x \, dx - \int_{\pi/2}^{\pi} 2 \cos nx \cos x \, dx \right] \\
&= \frac{1}{\pi} \left[\int_0^{\pi/2} \{ \cos (n+1)x + \cos (n-1)x \} \, dx - \int_{\pi/2}^{\pi} \{ \cos (n+1)x + \cos (n-1)x \} \, dx \right] \\
&= \frac{1}{\pi} \left[\left\{ \frac{\sin (n+1)x}{n+1} + \frac{\sin (n-1)x}{n-1} \right\}_0^{\pi/2} - \left\{ \frac{\sin (n+1)x}{n+1} + \frac{\sin (n-1)x}{n-1} \right\}_{\pi/2}^{\pi} \right] \\
&= \frac{1}{\pi} \left[\left\{ \frac{\sin (n+1)\pi/2}{n+1} + \frac{\sin (n-1)\pi/2}{n-1} - 0 \right\} - \left\{ 0 - \frac{\sin (n+1)\pi/2}{n+1} - \frac{\sin (n-1)\pi/2}{n-1} \right\} \right] \\
&= \frac{2}{\pi} \left[\frac{\sin (n+1)\pi/2}{n+1} + \frac{\sin (n-1)\pi/2}{n-1} \right] \\
&= \frac{2}{\pi} \left[\frac{\cos n(\pi/2)}{n+1} - \frac{\cos n(\pi/2)}{n-1} \right]
\end{aligned}$$

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx$$

$$= \frac{2}{\pi} \left[\int_0^{\pi/2} \cos x \cos nx \, dx - \int_{\pi/2}^{\pi} \cos x \cos nx \, dx \right] \dots\dots\dots (3)$$

$$= \frac{1}{\pi} \left[\int_0^{\pi/2} 2 \cos nx \cos x \, dx - \int_{\pi/2}^{\pi} 2 \cos nx \cos x \, dx \right]$$

$$= \frac{1}{\pi} \left[\int_0^{\pi/2} \{ \cos (n+1)x + \cos (n-1)x \} \, dx - \int_{\pi/2}^{\pi} \{ \cos (n+1)x + \cos (n-1)x \} \, dx \right]$$

$$= \frac{1}{\pi} \left[\left\{ \frac{\sin (n+1)x}{n+1} + \frac{\sin (n-1)x}{n-1} \right\}_0^{\pi/2} - \left\{ \frac{\sin (n+1)x}{n+1} + \frac{\sin (n-1)x}{n-1} \right\}_{\pi/2}^{\pi} \right]$$

$$= \frac{1}{\pi} \left[\left\{ \frac{\sin (n+1)\pi/2}{n+1} + \frac{\sin (n-1)\pi/2}{n-1} - 0 \right\} - \left\{ 0 - \frac{\sin (n+1)\pi/2}{n+1} - \frac{\sin (n-1)\pi/2}{n-1} \right\} \right]$$

$$= \frac{2}{\pi} \left[\frac{\sin (n+1)\pi/2}{n+1} + \frac{\sin (n-1)\pi/2}{n-1} \right]$$

$$= \frac{2}{\pi} \left[\frac{\cos n(\pi/2)}{n+1} - \frac{\cos n(\pi/2)}{n-1} \right]$$

$$\therefore a_n = \begin{cases} -\frac{4}{\pi(n^2-1)} \cos\left(\frac{n\pi}{2}\right) & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd and } n \neq 1 \end{cases} \dots\dots\dots (4)$$

To find a_1 , we put $n = 1$ in (3)

To find a_1 , we put $n = 1$ in (3)

$$\therefore a_1 = \frac{2}{\pi} \left[\int_0^{\pi/2} \cos^2 x \, dx - \int_{\pi/2}^{\pi} \cos^2 x \, dx \right]$$

To find a_1 , we put $n = 1$ in (3)

$$\begin{aligned}\therefore a_1 &= \frac{2}{\pi} \left[\int_0^{\pi/2} \cos^2 x \, dx - \int_{\pi/2}^{\pi} \cos^2 x \, dx \right] \\ &= \frac{2}{\pi} \left[\int_0^{\pi/2} \left(\frac{1 + \cos 2x}{2} \right) dx - \int_{\pi/2}^{\pi} \left(\frac{1 + \cos 2x}{2} \right) dx \right]\end{aligned}$$

To find a_1 , we put $n = 1$ in (3)

$$\begin{aligned}\therefore a_1 &= \frac{2}{\pi} \left[\int_0^{\pi/2} \cos^2 x \, dx - \int_{\pi/2}^{\pi} \cos^2 x \, dx \right] \\ &= \frac{2}{\pi} \left[\int_0^{\pi/2} \left(\frac{1 + \cos 2x}{2} \right) dx - \int_{\pi/2}^{\pi} \left(\frac{1 + \cos 2x}{2} \right) dx \right] \\ &= \frac{1}{\pi} \left[\left\{ x + \frac{\sin 2x}{2} \right\}_0^{\pi/2} - \left\{ x + \frac{\sin 2x}{2} \right\}_{\pi/2}^{\pi} \right]\end{aligned}$$

To find a_1 , we put $n = 1$ in (3)

$$\begin{aligned}\therefore a_1 &= \frac{2}{\pi} \left[\int_0^{\pi/2} \cos^2 x \, dx - \int_{\pi/2}^{\pi} \cos^2 x \, dx \right] \\&= \frac{2}{\pi} \left[\int_0^{\pi/2} \left(\frac{1 + \cos 2x}{2} \right) dx - \int_{\pi/2}^{\pi} \left(\frac{1 + \cos 2x}{2} \right) dx \right] \\&= \frac{1}{\pi} \left[\left\{ x + \frac{\sin 2x}{2} \right\}_0^{\pi/2} - \left\{ x + \frac{\sin 2x}{2} \right\}_{\pi/2}^{\pi} \right] \\&= \frac{1}{\pi} \left[\left\{ \frac{\pi}{2} \right\} - \left\{ \pi - \frac{\pi}{2} \right\} \right] = 0\end{aligned}$$

To find a_1 , we put $n = 1$ in (3)

$$\begin{aligned}\therefore a_1 &= \frac{2}{\pi} \left[\int_0^{\pi/2} \cos^2 x \, dx - \int_{\pi/2}^{\pi} \cos^2 x \, dx \right] \\&= \frac{2}{\pi} \left[\int_0^{\pi/2} \left(\frac{1 + \cos 2x}{2} \right) dx - \int_{\pi/2}^{\pi} \left(\frac{1 + \cos 2x}{2} \right) dx \right] \\&= \frac{1}{\pi} \left[\left\{ x + \frac{\sin 2x}{2} \right\}_0^{\pi/2} - \left\{ x + \frac{\sin 2x}{2} \right\}_{\pi/2}^{\pi} \right] \\&= \frac{1}{\pi} \left[\left\{ \frac{\pi}{2} \right\} - \left\{ \pi - \frac{\pi}{2} \right\} \right] = 0\end{aligned}$$

Putting these values from (2) and (4) in (1), we get,

$$|\cos x| = \frac{2}{\pi} + \frac{4}{\pi} \left[\frac{\cos 2x}{3} - \frac{\cos 4x}{15} + \dots \right]$$

EX 4. Obtain Fourier series for $f(x) = \begin{cases} x + \frac{\pi}{2}, & -\pi < x < 0 \\ \frac{\pi}{2} - x, & 0 < x < \pi \end{cases}$

Hence deduce that (i) $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$

(ii) $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$

EX 4. Obtain Fourier series for $f(x) = \begin{cases} x + \frac{\pi}{2}, & -\pi < x < 0 \\ \frac{\pi}{2} - x, & 0 < x < \pi \end{cases}$

Hence deduce that (i) $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$

(ii) $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$

Solution:

Clearly $f(x)$ is an even function

$$f(-x) = \begin{cases} -x + \frac{\pi}{2}, & -\pi < -x < 0 \\ \frac{\pi}{2} + x, & 0 < -x < \pi \end{cases}$$

EX 4. Obtain Fourier series for $f(x) = \begin{cases} x + \frac{\pi}{2}, & -\pi < x < 0 \\ \frac{\pi}{2} - x, & 0 < x < \pi \end{cases}$

Hence deduce that (i) $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$

(ii) $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$

Solution:

Clearly $f(x)$ is an even function

$$f(-x) = \begin{cases} -x + \frac{\pi}{2}, & -\pi < -x < 0 \\ \frac{\pi}{2} + x, & 0 < -x < \pi \end{cases} = \begin{cases} \frac{\pi}{2} - x, & \pi > x > 0 \\ \frac{\pi}{2} + x, & 0 < x < -\pi \end{cases}$$

EX 4. Obtain Fourier series for $f(x) = \begin{cases} x + \frac{\pi}{2}, & -\pi < x < 0 \\ \frac{\pi}{2} - x, & 0 < x < \pi \end{cases}$

Hence deduce that (i) $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$

(ii) $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$

Solution:

Clearly $f(x)$ is an even function

$$f(-x) = \begin{cases} -x + \frac{\pi}{2}, & -\pi < -x < 0 \\ \frac{\pi}{2} + x, & 0 < -x < \pi \end{cases} = \begin{cases} \frac{\pi}{2} - x, & \pi > x > 0 \\ \frac{\pi}{2} + x, & 0 < x < -\pi \end{cases}$$

$$= f(x) \quad [\because b_n = 0]$$

EX 4. Obtain Fourier series for $f(x) = \begin{cases} x + \frac{\pi}{2}, & -\pi < x < 0 \\ \frac{\pi}{2} - x, & 0 < x < \pi \end{cases}$

Hence deduce that (i) $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$

(ii) $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$

Solution:

Clearly $f(x)$ is an even function

$$f(-x) = \begin{cases} -x + \frac{\pi}{2}, & -\pi < -x < 0 \\ \frac{\pi}{2} + x, & 0 < -x < \pi \end{cases} = \begin{cases} \frac{\pi}{2} - x, & \pi > x > 0 \\ \frac{\pi}{2} + x, & 0 < x < -\pi \end{cases}$$

$$= f(x) \quad [\because b_n = 0]$$

$$a_0 = \frac{1}{\pi} \int_0^{\pi} f(x) dx = \frac{1}{\pi} \int_0^{\pi} \left(\frac{\pi}{2} - x \right) dx$$

EX 4. Obtain Fourier series for $f(x) = \begin{cases} x + \frac{\pi}{2}, & -\pi < x < 0 \\ \frac{\pi}{2} - x, & 0 < x < \pi \end{cases}$

Hence deduce that (i) $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$

(ii) $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$

Solution:

Clearly $f(x)$ is an even function

$$f(-x) = \begin{cases} -x + \frac{\pi}{2}, & -\pi < -x < 0 \\ \frac{\pi}{2} + x, & 0 < -x < \pi \end{cases} = \begin{cases} \frac{\pi}{2} - x, & \pi > x > 0 \\ \frac{\pi}{2} + x, & 0 < x < -\pi \end{cases}$$

$$= f(x) \quad [\because b_n = 0]$$

$$a_0 = \frac{1}{\pi} \int_0^{\pi} f(x) dx = \frac{1}{\pi} \int_0^{\pi} \left(\frac{\pi}{2} - x \right) dx = \frac{1}{\pi} \left[\frac{\pi}{2} x - \frac{x^2}{2} \right]_0^{\pi} = 0 \quad \dots (1)$$

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx = \frac{2}{\pi} \int_0^{\pi} \left(\frac{\pi}{2} - x \right) \cos nx \, dx$$

$$\begin{aligned}
 a_n &= \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx = \frac{2}{\pi} \int_0^{\pi} \left(\frac{\pi}{2} - x \right) \cos nx \, dx \\
 &= \frac{2}{\pi} \left[\left(\frac{\pi}{2} - x \right) \left(\frac{\sin nx}{n} \right) - (-1) \cdot \left(-\frac{\cos nx}{n^2} \right) \right]_0^{\pi}
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx = \frac{2}{\pi} \int_0^{\pi} \left(\frac{\pi}{2} - x \right) \cos nx \, dx \\
 &= \frac{2}{\pi} \left[\left(\frac{\pi}{2} - x \right) \left(\frac{\sin nx}{n} \right) - (-1) \cdot \left(-\frac{\cos nx}{n^2} \right) \right]_0^{\pi} \\
 &= \frac{2}{\pi} \left[-(-1)^n \cdot \frac{1}{n^2} + \frac{1}{n^2} \right] :
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx = \frac{2}{\pi} \int_0^{\pi} \left(\frac{\pi}{2} - x \right) \cos nx \, dx \\
 &= \frac{2}{\pi} \left[\left(\frac{\pi}{2} - x \right) \left(\frac{\sin nx}{n} \right) - (-1) \cdot \left(-\frac{\cos nx}{n^2} \right) \right]_0^{\pi} \\
 &= \frac{2}{\pi} \left[-(-1)^n \cdot \frac{1}{n^2} + \frac{1}{n^2} \right] = \frac{2}{\pi} \left[\frac{1}{n^2} - \frac{(-1)^n}{n^2} \right]
 \end{aligned}$$

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx = \frac{2}{\pi} \int_0^{\pi} \left(\frac{\pi}{2} - x \right) \cos nx \, dx$$

$$= \frac{2}{\pi} \left[\left(\frac{\pi}{2} - x \right) \left(\frac{\sin nx}{n} \right) - (-1) \cdot \left(-\frac{\cos nx}{n^2} \right) \right]_0^{\pi}$$

$$= \frac{2}{\pi} \left[-(-1)^n \cdot \frac{1}{n^2} + \frac{1}{n^2} \right] = \frac{2}{\pi} \left[\frac{1}{n^2} - \frac{(-1)^n}{n^2} \right]$$

$$\therefore f(x) = \frac{4}{\pi} \left[\frac{1}{1^2} \cos x + \frac{1}{3^2} \cos 3x + \frac{1}{5^2} \cos 5x + \dots \right] \quad \dots\dots\dots (2)$$

Hence deduce that (i) $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} \dots\dots\dots$
(ii) $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} \dots\dots\dots$

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx = \frac{2}{\pi} \int_0^{\pi} \left(\frac{\pi}{2} - x \right) \cos nx \, dx$$

$$= \frac{2}{\pi} \left[\left(\frac{\pi}{2} - x \right) \left(\frac{\sin nx}{n} \right) - (-1) \cdot \left(-\frac{\cos nx}{n^2} \right) \right]_0^{\pi}$$

$$= \frac{2}{\pi} \left[-(-1)^n \cdot \frac{1}{n^2} + \frac{1}{n^2} \right] = \frac{2}{\pi} \left[\frac{1}{n^2} - \frac{(-1)^n}{n^2} \right]$$

$$\therefore f(x) = \frac{4}{\pi} \left[\frac{1}{1^2} \cos x + \frac{1}{3^2} \cos 3x + \frac{1}{5^2} \cos 5x + \dots \right] \quad \dots\dots\dots (2)$$

(i) Now $f(x)$ is discontinuous at $x = 0$

But at a point of discontinuity $x = c$.

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx = \frac{2}{\pi} \int_0^{\pi} \left(\frac{\pi}{2} - x \right) \cos nx \, dx$$

$$= \frac{2}{\pi} \left[\left(\frac{\pi}{2} - x \right) \left(\frac{\sin nx}{n} \right) - (-1) \cdot \left(-\frac{\cos nx}{n^2} \right) \right]_0^{\pi}$$

$$= \frac{2}{\pi} \left[-(-1)^n \cdot \frac{1}{n^2} + \frac{1}{n^2} \right] = \frac{2}{\pi} \left[\frac{1}{n^2} - \frac{(-1)^n}{n^2} \right]$$

$$\therefore f(x) = \frac{4}{\pi} \left[\frac{1}{1^2} \cos x + \frac{1}{3^2} \cos 3x + \frac{1}{5^2} \cos 5x + \dots \right] \quad \dots\dots\dots (2)$$

(i) Now $f(x)$ is discontinuous at $x = 0$

But at a point of discontinuity $x = c$.

$$f(x) = \frac{1}{2} \left[\lim_{x \rightarrow c^-} f(x) + \lim_{x \rightarrow c^+} f(x) \right]$$

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx = \frac{2}{\pi} \int_0^{\pi} \left(\frac{\pi}{2} - x \right) \cos nx \, dx$$

$$= \frac{2}{\pi} \left[\left(\frac{\pi}{2} - x \right) \left(\frac{\sin nx}{n} \right) - (-1) \cdot \left(-\frac{\cos nx}{n^2} \right) \right]_0^{\pi}$$

$$= \frac{2}{\pi} \left[-(-1)^n \cdot \frac{1}{n^2} + \frac{1}{n^2} \right] = \frac{2}{\pi} \left[\frac{1}{n^2} - \frac{(-1)^n}{n^2} \right]$$

$$\therefore f(x) = \frac{4}{\pi} \left[\frac{1}{1^2} \cos x + \frac{1}{3^2} \cos 3x + \frac{1}{5^2} \cos 5x + \dots \right] \quad \dots\dots\dots (2)$$

(i) Now $f(x)$ is discontinuous at $x = 0$

But at a point of discontinuity $x = c$.

$$f(x) = \frac{1}{2} \left[\lim_{x \rightarrow c^-} f(x) + \lim_{x \rightarrow c^+} f(x) \right] \quad \therefore f(0) = \frac{1}{2} \left[\frac{\pi}{2} + \frac{\pi}{2} \right] = \frac{\pi}{2}$$

\therefore Putting $x = 0$,

$$\frac{\pi}{2} = \frac{4}{\pi} \left[\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right]$$

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx \, dx = \frac{2}{\pi} \int_0^{\pi} \left(\frac{\pi}{2} - x \right) \cos nx \, dx$$

$$= \frac{2}{\pi} \left[\left(\frac{\pi}{2} - x \right) \left(\frac{\sin nx}{n} \right) - (-1) \cdot \left(-\frac{\cos nx}{n^2} \right) \right]_0^{\pi}$$

$$= \frac{2}{\pi} \left[-(-1)^n \cdot \frac{1}{n^2} + \frac{1}{n^2} \right] = \frac{2}{\pi} \left[\frac{1}{n^2} - \frac{(-1)^n}{n^2} \right]$$

$$\therefore f(x) = \frac{4}{\pi} \left[\frac{1}{1^2} \cos x + \frac{1}{3^2} \cos 3x + \frac{1}{5^2} \cos 5x + \dots \right] \quad \dots\dots\dots (2)$$

(i) Now $f(x)$ is discontinuous at $x = 0$

But at a point of discontinuity $x = c$.

$$f(x) = \frac{1}{2} \left[\lim_{x \rightarrow c^-} f(x) + \lim_{x \rightarrow c^+} f(x) \right] \quad \therefore f(0) = \frac{1}{2} \left[\frac{\pi}{2} + \frac{\pi}{2} \right] = \frac{\pi}{2}$$

\therefore Putting $x = 0$,

$$\frac{\pi}{2} = \frac{4}{\pi} \left[\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right]$$

$$\therefore \frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$$

(ii) We now use Parseval's identity in $(-\pi, \pi)$

(ii) We now use Parseval's identity in $(-\pi, \pi)$

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \quad \dots\dots\dots (3)$$

(ii) We now use Parseval's identity in $(-\pi, \pi)$

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \quad \dots\dots\dots (3)$$

$$\therefore \frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx = \frac{1}{2\pi} \left[\int_{-\pi}^0 \left(x + \frac{\pi}{2} \right)^2 dx + \int_0^{\pi} \left(\frac{\pi}{2} - x \right)^2 dx \right]$$

(ii) We now use Parseval's identity in $(-\pi, \pi)$

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \quad \dots\dots\dots (3)$$

$$\begin{aligned} \therefore \frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx &= \frac{1}{2\pi} \left[\int_{-\pi}^0 \left(x + \frac{\pi}{2} \right)^2 dx + \int_0^{\pi} \left(\frac{\pi}{2} - x \right)^2 dx \right] \\ &= \frac{1}{2\pi} \left[\int_{-\pi}^0 \left(x^2 + \pi x + \frac{\pi^2}{4} \right) dx + \int_0^{\pi} \left(\frac{\pi^2}{4} - \pi x + x^2 \right) dx \right] \end{aligned}$$

(ii) We now use Parseval's identity in $(-\pi, \pi)$

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \quad \dots\dots\dots (3)$$

$$\begin{aligned} \therefore \frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx &= \frac{1}{2\pi} \left[\int_{-\pi}^0 \left(x + \frac{\pi}{2} \right)^2 dx + \int_0^{\pi} \left(\frac{\pi}{2} - x \right)^2 dx \right] \\ &= \frac{1}{2\pi} \left[\int_{-\pi}^0 \left(x^2 + \pi x + \frac{\pi^2}{4} \right) dx + \int_0^{\pi} \left(\frac{\pi^2}{4} - \pi x + x^2 \right) dx \right] \\ &= \frac{1}{2\pi} \left[\left\{ \frac{x^3}{3} + \frac{\pi x^2}{2} + \frac{\pi^2}{4} x \right\}_{-\pi}^0 + \left\{ \frac{\pi^2}{4} x - \frac{\pi x^2}{2} + \frac{x^3}{3} \right\}_0^{\pi} \right] \end{aligned}$$

(ii) We now use Parseval's identity in $(-\pi, \pi)$

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \quad \dots\dots\dots (3)$$

$$\begin{aligned} \therefore \frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx &= \frac{1}{2\pi} \left[\int_{-\pi}^0 \left(x + \frac{\pi}{2} \right)^2 dx + \int_0^{\pi} \left(\frac{\pi}{2} - x \right)^2 dx \right] \\ &= \frac{1}{2\pi} \left[\int_{-\pi}^0 \left(x^2 + \pi x + \frac{\pi^2}{4} \right) dx + \int_0^{\pi} \left(\frac{\pi^2}{4} - \pi x + x^2 \right) dx \right] \\ &= \frac{1}{2\pi} \left[\left\{ \frac{x^3}{3} + \frac{\pi x^2}{2} + \frac{\pi^2}{4} x \right\}_{-\pi}^0 + \left\{ \frac{\pi^2}{4} x - \frac{\pi x^2}{2} + \frac{x^3}{3} \right\}_0^{\pi} \right] \\ &= \frac{1}{2\pi} \left[0 - \left\{ -\frac{\pi^3}{3} + \frac{\pi^3}{2} - \frac{\pi^3}{4} \right\} + \left\{ \frac{\pi^3}{4} - \frac{\pi^3}{2} + \frac{\pi^3}{3} \right\} - 0 \right] \end{aligned}$$

(ii) We now use Parseval's identity in $(-\pi, \pi)$

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \quad \dots\dots\dots (3)$$

$$\begin{aligned} \therefore \frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx &= \frac{1}{2\pi} \left[\int_{-\pi}^0 \left(x + \frac{\pi}{2} \right)^2 dx + \int_0^{\pi} \left(\frac{\pi}{2} - x \right)^2 dx \right] \\ &= \frac{1}{2\pi} \left[\int_{-\pi}^0 \left(x^2 + \pi x + \frac{\pi^2}{4} \right) dx + \int_0^{\pi} \left(\frac{\pi^2}{4} - \pi x + x^2 \right) dx \right] \\ &= \frac{1}{2\pi} \left[\left\{ \frac{x^3}{3} + \frac{\pi x^2}{2} + \frac{\pi^2}{4} x \right\}_{-\pi}^0 + \left\{ \frac{\pi^2}{4} x - \frac{\pi x^2}{2} + \frac{x^3}{3} \right\}_0^{\pi} \right] \\ &= \frac{1}{2\pi} \left[0 - \left\{ -\frac{\pi^3}{3} + \frac{\pi^3}{2} - \frac{\pi^3}{4} \right\} + \left\{ \frac{\pi^3}{4} - \frac{\pi^3}{2} + \frac{\pi^3}{3} \right\} - 0 \right] \\ &= \frac{2\pi^3}{2\pi} \left[\frac{1}{4} - \frac{1}{2} + \frac{1}{3} \right] = \frac{\pi^2}{12} \end{aligned}$$

(ii) We now use Parseval's identity in $(-\pi, \pi)$

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \quad \dots\dots\dots (3)$$

$$\begin{aligned} \therefore \frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx &= \frac{1}{2\pi} \left[\int_{-\pi}^0 \left(x + \frac{\pi}{2} \right)^2 dx + \int_0^{\pi} \left(\frac{\pi}{2} - x \right)^2 dx \right] \\ &= \frac{1}{2\pi} \left[\int_{-\pi}^0 \left(x^2 + \pi x + \frac{\pi^2}{4} \right) dx + \int_0^{\pi} \left(\frac{\pi^2}{4} - \pi x + x^2 \right) dx \right] \\ &= \frac{1}{2\pi} \left[\left\{ \frac{x^3}{3} + \frac{\pi x^2}{2} + \frac{\pi^2}{4} x \right\}_{-\pi}^0 + \left\{ \frac{\pi^2}{4} x - \frac{\pi x^2}{2} + \frac{x^3}{3} \right\}_0^{\pi} \right] \\ &= \frac{1}{2\pi} \left[0 - \left\{ -\frac{\pi^3}{3} + \frac{\pi^3}{2} - \frac{\pi^3}{4} \right\} + \left\{ \frac{\pi^3}{4} - \frac{\pi^3}{2} + \frac{\pi^3}{3} \right\} - 0 \right] \\ &= \frac{2\pi^3}{2\pi} \left[\frac{1}{4} - \frac{1}{2} + \frac{1}{3} \right] = \frac{\pi^2}{12} \end{aligned}$$

Hence, from (1), (2) and (3),

$$\frac{\pi^2}{12} = \frac{1}{2} \cdot \frac{16}{\pi^2} \left[\frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots \right]$$

(ii) We now use Parseval's identity in $(-\pi, \pi)$

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx = a_0^2 + \frac{1}{2} \sum (a_n^2 + b_n^2) \quad \dots\dots\dots (3)$$

$$\begin{aligned} \therefore \frac{1}{2\pi} \int_{-\pi}^{\pi} [f(x)]^2 dx &= \frac{1}{2\pi} \left[\int_{-\pi}^0 \left(x + \frac{\pi}{2} \right)^2 dx + \int_0^{\pi} \left(\frac{\pi}{2} - x \right)^2 dx \right] \\ &= \frac{1}{2\pi} \left[\int_{-\pi}^0 \left(x^2 + \pi x + \frac{\pi^2}{4} \right) dx + \int_0^{\pi} \left(\frac{\pi^2}{4} - \pi x + x^2 \right) dx \right] \\ &= \frac{1}{2\pi} \left[\left\{ \frac{x^3}{3} + \frac{\pi x^2}{2} + \frac{\pi^2}{4} x \right\}_{-\pi}^0 + \left\{ \frac{\pi^2}{4} x - \frac{\pi x^2}{2} + \frac{x^3}{3} \right\}_0^{\pi} \right] \\ &= \frac{1}{2\pi} \left[0 - \left\{ -\frac{\pi^3}{3} + \frac{\pi^3}{2} - \frac{\pi^3}{4} \right\} + \left\{ \frac{\pi^3}{4} - \frac{\pi^3}{2} + \frac{\pi^3}{3} \right\} - 0 \right] \\ &= \frac{2\pi^3}{2\pi} \left[\frac{1}{4} - \frac{1}{2} + \frac{1}{3} \right] = \frac{\pi^2}{12} \end{aligned}$$

Hence, from (1), (2) and (3),

$$\begin{aligned} \frac{\pi^2}{12} &= \frac{1}{2} \cdot \frac{16}{\pi^2} \left[\frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots \right] \\ \frac{\pi^4}{96} &= \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots \end{aligned}$$

Fourier series in the interval $(0, 2l)$

Fourier series in the interval $(0, 2l)$

EX 1. Find the Fourier expansion of $f(x) = 2x - x^2$ in the interval $(0, 3)$ and whose period is 3.

Fourier series in the interval $(0, 2l)$

EX 1. Find the Fourier expansion of $f(x) = 2x - x^2$ in the interval $(0, 3)$ and whose period is 3.

Solution:

Here period $2l = 3 \quad \therefore l = 3/2.$

Fourier series in the interval $(0, 2l)$

EX 1. Find the Fourier expansion of $f(x) = 2x - x^2$ in the interval $(0, 3)$ and whose period is 3.

Solution:

Here period $2l = 3 \quad \therefore l = 3/2$.

$$\text{Let } f(x) = a_0 + \sum a_n \cos \frac{n\pi x}{l} + \sum b_n \sin \frac{n\pi x}{l}$$

Fourier series in the interval $(0, 2l)$

EX 1. Find the Fourier expansion of $f(x) = 2x - x^2$ in the interval $(0, 3)$ and whose period is 3.

Solution:

Here period $2l = 3 \quad \therefore l = 3/2$.

$$\begin{aligned} \text{Let } f(x) &= a_0 + \sum a_n \cos \frac{n\pi x}{l} + \sum b_n \sin \frac{n\pi x}{l} \\ &= a_0 + \sum a_n \cos \frac{2n\pi x}{3} + \sum b_n \sin \frac{2n\pi x}{3} \quad \dots\dots\dots (1) \end{aligned}$$

Fourier series in the interval $(0, 2l)$

EX 1. Find the Fourier expansion of $f(x) = 2x - x^2$ in the interval $(0, 3)$ and whose period is 3.

Solution:

Here period $2l = 3 \quad \therefore l = 3/2$.

$$\begin{aligned} \text{Let } f(x) &= a_0 + \sum a_n \cos \frac{n\pi x}{l} + \sum b_n \sin \frac{n\pi x}{l} \\ &= a_0 + \sum a_n \cos \frac{2n\pi x}{3} + \sum b_n \sin \frac{2n\pi x}{3} \quad \dots\dots\dots (1) \end{aligned}$$

$$\text{Now } a_0 = \frac{1}{2l} \int_0^{2l} f(x) dx = \frac{1}{3} \int_0^3 (2x - x^2) dx$$

Fourier series in the interval $(0, 2l)$

EX 1. Find the Fourier expansion of $f(x) = 2x - x^2$ in the interval $(0, 3)$ and whose period is 3.

Solution:

Here period $2l = 3 \quad \therefore l = 3/2$.

$$\begin{aligned} \text{Let } f(x) &= a_0 + \sum a_n \cos \frac{n\pi x}{l} + \sum b_n \sin \frac{n\pi x}{l} \\ &= a_0 + \sum a_n \cos \frac{2n\pi x}{3} + \sum b_n \sin \frac{2n\pi x}{3} \quad \dots\dots\dots (1) \end{aligned}$$

$$\begin{aligned} \text{Now } a_0 &= \frac{1}{2l} \int_0^{2l} f(x) dx = \frac{1}{3} \int_0^3 (2x - x^2) dx \\ &= \frac{1}{3} \left[x^2 - \frac{x^3}{3} \right]_0^3 = \frac{1}{3} \left[9 - \frac{27}{3} \right] = 0 \end{aligned}$$

$$a_n = \frac{1}{l} \int_0^{2l} f(x) \cos \frac{n\pi x}{l} dx = \frac{1}{(3/2)} \int_0^3 (2x - x^2) \cos \frac{2n\pi x}{3} dx$$

$$\begin{aligned}
 a_n &= \frac{1}{l} \int_0^{2l} f(x) \cos \frac{n\pi x}{l} dx = \frac{1}{(3/2)} \int_0^3 (2x - x^2) \cos \frac{2n\pi x}{3} dx \\
 &= \frac{2}{3} \left[(2x - x^2) \cdot \left(\frac{3}{2n\pi} \cdot \sin \frac{2n\pi x}{3} \right) \right. \\
 &\quad \left. - (2 - 2x) \left(-\frac{9}{4n^2\pi^2} \cdot \cos \frac{2n\pi x}{3} \right) + (-2) \left(-\frac{27}{8n^3\pi^3} \cdot \sin \frac{2n\pi x}{3} \right) \right]_0^3
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{1}{l} \int_0^{2l} f(x) \cos \frac{n\pi x}{l} dx = \frac{1}{(3/2)} \int_0^3 (2x - x^2) \cos \frac{2n\pi x}{3} dx \\
 &= \frac{2}{3} \left[(2x - x^2) \cdot \left(\frac{3}{2n\pi} \cdot \sin \frac{2n\pi x}{3} \right) \right. \\
 &\quad \left. - (2 - 2x) \left(-\frac{9}{4n^2\pi^2} \cdot \cos \frac{2n\pi x}{3} \right) + (-2) \left(-\frac{27}{8n^3\pi^3} \cdot \sin \frac{2n\pi x}{3} \right) \right]_0^3 \\
 &= \frac{2}{3} \left[\left\{ 0 - 4 \cdot \frac{9}{4n^2\pi^2} \cos 2n\pi + 0 \right\} - \left\{ 0 + 2 \cdot \frac{9}{4n^2\pi^2} + 0 \right\} \right]
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{1}{l} \int_0^{2l} f(x) \cos \frac{n\pi x}{l} dx = \frac{1}{(3/2)} \int_0^3 (2x - x^2) \cos \frac{2n\pi x}{3} dx \\
 &= \frac{2}{3} \left[(2x - x^2) \cdot \left(\frac{3}{2n\pi} \cdot \sin \frac{2n\pi x}{3} \right) \right. \\
 &\quad \left. - (2 - 2x) \left(-\frac{9}{4n^2\pi^2} \cdot \cos \frac{2n\pi x}{3} \right) + (-2) \left(-\frac{27}{8n^3\pi^3} \cdot \sin \frac{2n\pi x}{3} \right) \right]_0^3 \\
 &= \frac{2}{3} \left[\left\{ 0 - 4 \cdot \frac{9}{4n^2\pi^2} \cos 2n\pi + 0 \right\} - \left\{ 0 + 2 \cdot \frac{9}{4n^2\pi^2} + 0 \right\} \right] \\
 &= \frac{2}{3} \cdot \frac{9}{4n^2\pi^2} [-4 - 2] = -\frac{9}{n^2\pi^2}
 \end{aligned}$$

$$b_n = \frac{1}{l} \int_0^{2l} f(x) \sin \frac{n\pi x}{l} dx = \frac{1}{(3/2)} \int_0^3 (2x - x^2) \cdot \sin \frac{2n\pi x}{3} dx$$

$$\begin{aligned}
 b_n &= \frac{1}{l} \int_0^{2l} f(x) \sin \frac{n\pi x}{l} dx = \frac{1}{(3/2)} \int_0^3 (2x - x^2) \cdot \sin \frac{2n\pi x}{3} dx \\
 &= \frac{2}{3} \left[(2x - x^2) \cdot \left(-\frac{3}{2n\pi} \cdot \cos \frac{2n\pi x}{3} \right) \right. \\
 &\quad \left. - (2 - 2x) \left(-\frac{9}{4n^2\pi^2} \cdot \sin \frac{2n\pi x}{3} \right) + (-2) \left(\frac{27}{8n^3\pi^3} \cdot \cos \frac{2n\pi x}{3} \right) \right]_0^3
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{l} \int_0^{2l} f(x) \sin \frac{n\pi x}{l} dx = \frac{1}{(3/2)} \int_0^3 (2x - x^2) \cdot \sin \frac{2n\pi x}{3} dx \\
 &= \frac{2}{3} \left[(2x - x^2) \cdot \left(-\frac{3}{2n\pi} \cdot \cos \frac{2n\pi x}{3} \right) \right. \\
 &\quad \left. - (2 - 2x) \left(-\frac{9}{4n^2\pi^2} \cdot \sin \frac{2n\pi x}{3} \right) + (-2) \left(\frac{27}{8n^3\pi^3} \cdot \cos \frac{2n\pi x}{3} \right) \right]_0^3 \\
 &= \frac{2}{3} \left[\left\{ \frac{9}{2n\pi} - 0 - \frac{27}{4n^3\pi^3} \right\} - \left\{ 0 - 0 - \frac{27}{4n^3\pi^3} \right\} \right]
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{l} \int_0^{2l} f(x) \sin \frac{n\pi x}{l} dx = \frac{1}{(3/2)} \int_0^3 (2x - x^2) \cdot \sin \frac{2n\pi x}{3} dx \\
 &= \frac{2}{3} \left[(2x - x^2) \cdot \left(-\frac{3}{2n\pi} \cdot \cos \frac{2n\pi x}{3} \right) \right. \\
 &\quad \left. - (2 - 2x) \left(-\frac{9}{4n^2\pi^2} \cdot \sin \frac{2n\pi x}{3} \right) + (-2) \left(\frac{27}{8n^3\pi^3} \cdot \cos \frac{2n\pi x}{3} \right) \right]_0^3 \\
 &= \frac{2}{3} \left[\left\{ \frac{9}{2n\pi} - 0 - \frac{27}{4n^3\pi^3} \right\} - \left\{ 0 - 0 - \frac{27}{4n^3\pi^3} \right\} \right] \\
 &= \frac{2}{3} \left[\frac{9}{2n\pi} \right] = \frac{3}{n\pi} .
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{l} \int_0^{2l} f(x) \sin \frac{n\pi x}{l} dx = \frac{1}{(3/2)} \int_0^3 (2x - x^2) \cdot \sin \frac{2n\pi x}{3} dx \\
 &= \frac{2}{3} \left[(2x - x^2) \cdot \left(-\frac{3}{2n\pi} \cdot \cos \frac{2n\pi x}{3} \right) \right. \\
 &\quad \left. - (2 - 2x) \left(-\frac{9}{4n^2\pi^2} \cdot \sin \frac{2n\pi x}{3} \right) + (-2) \left(\frac{27}{8n^3\pi^3} \cdot \cos \frac{2n\pi x}{3} \right) \right]_0^3 \\
 &= \frac{2}{3} \left[\left\{ \frac{9}{2n\pi} - 0 - \frac{27}{4n^3\pi^3} \right\} - \left\{ 0 - 0 - \frac{27}{4n^3\pi^3} \right\} \right] \\
 &= \frac{2}{3} \left[\frac{9}{2n\pi} \right] = \frac{3}{n\pi} .
 \end{aligned}$$

Putting these values in (1),

$$f(x) = 0 + \sum \left(-\frac{9}{n^2\pi^2} \right) \cos \frac{2n\pi x}{3} + \sum \frac{3}{n\pi} \sin \frac{2n\pi x}{3}$$

$$\begin{aligned}
 b_n &= \frac{1}{l} \int_0^{2l} f(x) \sin \frac{n\pi x}{l} dx = \frac{1}{(3/2)} \int_0^3 (2x - x^2) \cdot \sin \frac{2n\pi x}{3} dx \\
 &= \frac{2}{3} \left[(2x - x^2) \cdot \left(-\frac{3}{2n\pi} \cdot \cos \frac{2n\pi x}{3} \right) \right. \\
 &\quad \left. - (2 - 2x) \left(-\frac{9}{4n^2\pi^2} \cdot \sin \frac{2n\pi x}{3} \right) + (-2) \left(\frac{27}{8n^3\pi^3} \cdot \cos \frac{2n\pi x}{3} \right) \right]_0^3 \\
 &= \frac{2}{3} \left[\left\{ \frac{9}{2n\pi} - 0 - \frac{27}{4n^3\pi^3} \right\} - \left\{ 0 - 0 - \frac{27}{4n^3\pi^3} \right\} \right] \\
 &= \frac{2}{3} \left[\frac{9}{2n\pi} \right] = \frac{3}{n\pi} .
 \end{aligned}$$

Putting these values in (1),

$$\begin{aligned}
 f(x) &= 0 + \sum \left(-\frac{9}{n^2\pi^2} \right) \cos \frac{2n\pi x}{3} + \sum \frac{3}{n\pi} \sin \frac{2n\pi x}{3} \\
 &= -\frac{9}{\pi^2} \sum \frac{1}{n^2} \cos \frac{2n\pi x}{3} + \frac{3}{\pi} \sum \frac{1}{n} \sin \frac{2n\pi x}{3}
 \end{aligned}$$

EX 2 If $f(x) = \begin{cases} \pi x, & 0 < x < 1 \\ \pi(2 - x), & 1 < x < 2 \end{cases}$ With period 2

show that $f(x) = \frac{\pi}{2} - \frac{4}{\pi} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} \cos(2n+1)\pi x$

EX 2 If $f(x) = \begin{cases} \pi x, & 0 < x < 1 \\ \pi(2 - x), & 1 < x < 2 \end{cases}$ With period 2

show that $f(x) = \frac{\pi}{2} - \frac{4}{\pi} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} \cos(2n+1)\pi x$

Solution: Here $2l = 2 \quad \therefore l = 1$

$$\text{Let } f(x) = a_0 + \sum a_n \cos \frac{n\pi x}{l} + \sum b_n \sin \frac{n\pi x}{l}$$

EX 2 If $f(x) = \begin{cases} \pi x, & 0 < x < 1 \\ \pi(2 - x), & 1 < x < 2 \end{cases}$ With period 2

show that $f(x) = \frac{\pi}{2} - \frac{4}{\pi} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} \cos(2n+1)\pi x$

Solution: Here $2l = 2 \quad \therefore l = 1$

$$\text{Let } f(x) = a_0 + \sum a_n \cos \frac{n\pi x}{l} + \sum b_n \sin \frac{n\pi x}{l}$$

$$\text{i.e. } f(x) = a_0 + \sum a_n \cos n\pi x + \sum b_n \sin n\pi x \quad \dots\dots\dots (1)$$

EX 2 If $f(x) = \begin{cases} \pi x, & 0 < x < 1 \\ \pi(2 - x), & 1 < x < 2 \end{cases}$ With period 2

show that $f(x) = \frac{\pi}{2} - \frac{4}{\pi} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} \cos(2n+1)\pi x$

Solution: Here $2l = 2 \quad \therefore l = 1$

$$\text{Let } f(x) = a_0 + \sum a_n \cos \frac{n\pi x}{l} + \sum b_n \sin \frac{n\pi x}{l}$$

$$\text{i.e. } f(x) = a_0 + \sum a_n \cos n\pi x + \sum b_n \sin n\pi x \quad \dots\dots\dots (1)$$

$$\therefore a_0 = \frac{1}{2l} \int_0^{2l} f(x) dx = \frac{1}{2} \left[\int_0^1 \pi x dx + \int_1^2 \pi(2 - x) dx \right]$$

EX 2 If $f(x) = \begin{cases} \pi x, & 0 < x < 1 \\ \pi(2 - x), & 1 < x < 2 \end{cases}$ With period 2

show that $f(x) = \frac{\pi}{2} - \frac{4}{\pi} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} \cos(2n+1)\pi x$

Solution: Here $2l = 2 \quad \therefore l = 1$

$$\text{Let } f(x) = a_0 + \sum a_n \cos \frac{n\pi x}{l} + \sum b_n \sin \frac{n\pi x}{l}$$

$$\text{i.e. } f(x) = a_0 + \sum a_n \cos n\pi x + \sum b_n \sin n\pi x \quad \dots\dots\dots (1)$$

$$\therefore a_0 = \frac{1}{2l} \int_0^{2l} f(x) dx = \frac{1}{2} \left[\int_0^1 \pi x dx + \int_1^2 \pi(2 - x) dx \right]$$

$$= \frac{1}{2} \left[\pi \left\{ \frac{x^2}{2} \right\}_0^1 + \pi \left\{ 2x - \frac{x^2}{2} \right\}_1^2 \right]$$

EX 2 If $f(x) = \begin{cases} \pi x, & 0 < x < 1 \\ \pi(2 - x), & 1 < x < 2 \end{cases}$ With period 2

show that $f(x) = \frac{\pi}{2} - \frac{4}{\pi} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} \cos(2n+1)\pi x$

Solution: Here $2l = 2 \quad \therefore l = 1$

$$\text{Let } f(x) = a_0 + \sum a_n \cos \frac{n\pi x}{l} + \sum b_n \sin \frac{n\pi x}{l}$$

$$\text{i.e. } f(x) = a_0 + \sum a_n \cos n\pi x + \sum b_n \sin n\pi x \quad \dots\dots\dots (1)$$

$$\therefore a_0 = \frac{1}{2l} \int_0^{2l} f(x) dx = \frac{1}{2} \left[\int_0^1 \pi x dx + \int_1^2 \pi(2 - x) dx \right]$$

$$= \frac{1}{2} \left[\pi \left\{ \frac{x^2}{2} \right\}_0^1 + \pi \left\{ 2x - \frac{x^2}{2} \right\}_1^2 \right]$$

$$= \frac{\pi}{2} \left[\left\{ \frac{1}{2} \right\} + \left\{ 4 - 2 - 2 + \frac{1}{2} \right\} \right] = \frac{\pi}{2}$$

$$a_n = \frac{1}{l} \int_0^{2l} f(x) \cos \frac{n\pi x}{l} dx$$

$$a_n = \frac{1}{l} \int_0^{2l} f(x) \cos \frac{n\pi x}{l} dx$$

$$= \int_0^1 \pi x \cdot \cos n\pi x dx + \int_1^2 \pi (2 - x) \cos n\pi x dx$$

$$a_n = \frac{1}{l} \int_0^{2l} f(x) \cos \frac{n\pi x}{l} dx$$

$$= \int_0^1 \pi x \cdot \cos n\pi x dx + \int_1^2 \pi (2 - x) \cos n\pi x dx$$

$$= \pi \left[\left\{ x \left(\frac{\sin n\pi x}{n\pi} \right) - (1) \left(-\frac{\cos n\pi x}{n^2 \pi^2} \right) \right\}_0^1 + \left\{ (2 - x) \left(\frac{\sin n\pi x}{n\pi} \right) - (-1) \left(-\frac{\cos n\pi x}{n^2 \pi^2} \right) \right\}_1^2 \right]$$

$$a_n = \frac{1}{l} \int_0^{2l} f(x) \cos \frac{n\pi x}{l} dx$$

$$= \int_0^1 \pi x \cdot \cos n\pi x dx + \int_1^2 \pi (2 - x) \cos n\pi x dx$$

$$= \pi \left[\left\{ x \left(\frac{\sin n\pi x}{n\pi} \right) - (1) \left(-\frac{\cos n\pi x}{n^2 \pi^2} \right) \right\}_0^1 + \left\{ (2 - x) \left(\frac{\sin n\pi x}{n\pi} \right) - (-1) \left(-\frac{\cos n\pi x}{n^2 \pi^2} \right) \right\}_1^2 \right]$$

$$= \pi \left[\left\{ \frac{\cos n\pi}{n^2 \pi^2} - \frac{1}{n^2 \pi^2} \right\} + \left\{ -\frac{1}{n^2 \pi^2} + \frac{\cos n\pi}{n^2 \pi^2} \right\} \right]$$

$$a_n = \frac{1}{l} \int_0^{2l} f(x) \cos \frac{n\pi x}{l} dx$$

$$= \int_0^1 \pi x \cdot \cos n\pi x dx + \int_1^2 \pi (2 - x) \cos n\pi x dx$$

$$= \pi \left[\left\{ x \left(\frac{\sin n\pi x}{n\pi} \right) - (1) \left(-\frac{\cos n\pi x}{n^2 \pi^2} \right) \right\}_0^1 + \left\{ (2 - x) \left(\frac{\sin n\pi x}{n\pi} \right) - (-1) \left(-\frac{\cos n\pi x}{n^2 \pi^2} \right) \right\}_1^2 \right]$$

$$= \pi \left[\left\{ \frac{\cos n\pi}{n^2 \pi^2} - \frac{1}{n^2 \pi^2} \right\} + \left\{ -\frac{1}{n^2 \pi^2} + \frac{\cos n\pi}{n^2 \pi^2} \right\} \right]$$

$$= \frac{2\pi}{n^2 \pi^2} [\cos n\pi - 1]$$

$$a_n = \frac{1}{l} \int_0^{2l} f(x) \cos \frac{n\pi x}{l} dx$$

$$= \int_0^1 \pi x \cdot \cos n\pi x dx + \int_1^2 \pi (2 - x) \cos n\pi x dx$$

$$= \pi \left[\left\{ x \left(\frac{\sin n\pi x}{n\pi} \right) - (1) \left(-\frac{\cos n\pi x}{n^2 \pi^2} \right) \right\}_0^1 + \left\{ (2 - x) \left(\frac{\sin n\pi x}{n\pi} \right) - (-1) \left(-\frac{\cos n\pi x}{n^2 \pi^2} \right) \right\}_1^2 \right]$$

$$= \pi \left[\left\{ \frac{\cos n\pi}{n^2 \pi^2} - \frac{1}{n^2 \pi^2} \right\} + \left\{ -\frac{1}{n^2 \pi^2} + \frac{\cos n\pi}{n^2 \pi^2} \right\} \right]$$

$$= \frac{2\pi}{n^2 \pi^2} [\cos n\pi - 1] = \frac{2}{n^2 \pi} [(-1)^n - 1]$$

$$\begin{aligned}
a_n &= \frac{1}{l} \int_0^{2l} f(x) \cos \frac{n\pi x}{l} dx \\
&= \int_0^1 \pi x \cdot \cos n\pi x dx + \int_1^2 \pi (2-x) \cos n\pi x dx \\
&= \pi \left[\left\{ x \left(\frac{\sin n\pi x}{n\pi} \right) - (1) \left(-\frac{\cos n\pi x}{n^2 \pi^2} \right) \right\}_0^1 \right. \\
&\quad \left. + \left\{ (2-x) \left(\frac{\sin n\pi x}{n\pi} \right) - (-1) \left(-\frac{\cos n\pi x}{n^2 \pi^2} \right) \right\}_1^2 \right] \\
&= \pi \left[\left\{ \frac{\cos n\pi}{n^2 \pi^2} - \frac{1}{n^2 \pi^2} \right\} + \left\{ -\frac{1}{n^2 \pi^2} + \frac{\cos n\pi}{n^2 \pi^2} \right\} \right] \\
&= \frac{2\pi}{n^2 \pi^2} [\cos n\pi - 1] = \frac{2}{n^2 \pi} [(-1)^n - 1] \\
&= \begin{cases} 0 & \text{if } n \text{ is even} \\ -\frac{4}{n^2 \pi} & \text{if } n \text{ is odd} \end{cases}
\end{aligned}$$

$$b_n = \frac{1}{l} \int_0^{2l} f(x) \sin \frac{n\pi x}{l} dx$$

$$\begin{aligned} b_n &= \frac{1}{l} \int_0^{2l} f(x) \sin \frac{n\pi x}{l} dx \\ &= \int_0^1 \pi x \cdot \sin n\pi x dx + \int_1^2 \pi (2-x) \sin n\pi x dx \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{l} \int_0^{2l} f(x) \sin \frac{n\pi x}{l} dx \\
 &= \int_0^1 \pi x \cdot \sin n\pi x dx + \int_1^2 \pi (2-x) \sin n\pi x dx \\
 &= \pi \left[\left\{ x \left(-\frac{\cos n\pi x}{n\pi} \right) - (1) \left(-\frac{\sin n\pi x}{n^2 \pi^2} \right) \right\}_0^1 \right. \\
 &\quad \left. + \left\{ (2-x) \left(-\frac{\cos n\pi x}{n\pi} \right) - (-1) \left(-\frac{\sin n\pi x}{n^2 \pi^2} \right) \right\}_1^2 \right]
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{l} \int_0^{2l} f(x) \sin \frac{n\pi x}{l} dx \\
 &= \int_0^1 \pi x \cdot \sin n\pi x dx + \int_1^2 \pi (2-x) \sin n\pi x dx \\
 &= \pi \left[\left\{ x \left(-\frac{\cos n\pi x}{n\pi} \right) - (1) \left(-\frac{\sin n\pi x}{n^2 \pi^2} \right) \right\}_0^1 \right. \\
 &\quad \left. + \left\{ (2-x) \left(-\frac{\cos n\pi x}{n\pi} \right) - (-1) \left(-\frac{\sin n\pi x}{n^2 \pi^2} \right) \right\}_1^2 \right] \\
 &= \pi \left[\left\{ -\frac{\cos n\pi}{n\pi} \right\} + \left\{ +\frac{\cos n\pi}{n\pi} \right\} \right] = 0
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{l} \int_0^{2l} f(x) \sin \frac{n\pi x}{l} dx \\
 &= \int_0^1 \pi x \cdot \sin n\pi x dx + \int_1^2 \pi (2-x) \sin n\pi x dx \\
 &= \pi \left[\left\{ x \left(-\frac{\cos n\pi x}{n\pi} \right) - (1) \left(-\frac{\sin n\pi x}{n^2 \pi^2} \right) \right\}_0^1 \right. \\
 &\quad \left. + \left\{ (2-x) \left(-\frac{\cos n\pi x}{n\pi} \right) - (-1) \left(-\frac{\sin n\pi x}{n^2 \pi^2} \right) \right\}_1^2 \right] \\
 &= \pi \left[\left\{ -\frac{\cos n\pi}{n\pi} \right\} + \left\{ +\frac{\cos n\pi}{n\pi} \right\} \right] = 0
 \end{aligned}$$

Putting these values in (1)

$$f(x) = \frac{\pi}{2} - \frac{4}{\pi} \left[\frac{\cos \pi x}{1^2} + \frac{\cos 3\pi x}{3^2} + \frac{\cos 5\pi x}{5^2} + \dots \right]$$

$$\begin{aligned}
 b_n &= \frac{1}{l} \int_0^{2l} f(x) \sin \frac{n\pi x}{l} dx \\
 &= \int_0^1 \pi x \cdot \sin n\pi x dx + \int_1^2 \pi (2-x) \sin n\pi x dx \\
 &= \pi \left[\left\{ x \left(-\frac{\cos n\pi x}{n\pi} \right) - (1) \left(-\frac{\sin n\pi x}{n^2 \pi^2} \right) \right\}_0^1 \right. \\
 &\quad \left. + \left\{ (2-x) \left(-\frac{\cos n\pi x}{n\pi} \right) - (-1) \left(-\frac{\sin n\pi x}{n^2 \pi^2} \right) \right\}_1^2 \right] \\
 &= \pi \left[\left\{ -\frac{\cos n\pi}{n\pi} \right\} + \left\{ +\frac{\cos n\pi}{n\pi} \right\} \right] = 0
 \end{aligned}$$

Putting these values in (1)

$$\begin{aligned}
 f(x) &= \frac{\pi}{2} - \frac{4}{\pi} \left[\frac{\cos \pi x}{1^2} + \frac{\cos 3\pi x}{3^2} + \frac{\cos 5\pi x}{5^2} + \dots \right] \\
 &= \frac{\pi}{2} - \frac{4}{\pi} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} \cos (2n+1) \pi x
 \end{aligned}$$

Fourier series in the interval $(-l, l)$


Fourier series in the interval $(-l, l)$

EX 1. Find the Fourier expansion of $f(x) = \begin{cases} 0, & -2 < x < -1 \\ 1 + x, & -1 < x < 0 \\ 1 - x, & 0 < x < 1 \\ 0, & 1 < x < 2 \end{cases}$

Fourier series in the interval $(-l, l)$

EX 1. Find the Fourier expansion of $f(x) = \begin{cases} 0, & -2 < x < -1 \\ 1 + x, & -1 < x < 0 \\ 1 - x, & 0 < x < 1 \\ 0, & 1 < x < 2 \end{cases}$


Solution: The graph of the function is shown below,



Fourier series in the interval $(-l, l)$

EX 1. Find the Fourier expansion of $f(x) = \begin{cases} 0, & -2 < x < -1 \\ 1+x, & -1 < x < 0 \\ 1-x, & 0 < x < 1 \\ 0, & 1 < x < 2 \end{cases}$

Solution: The graph of the function is shown below,




$$\text{Further } f(-x) = \begin{cases} 0, & -2 < -x < -1 \\ 1-x, & -1 < -x < 0 \\ 1+x, & 0 < -x < 1 \\ 0, & 1 < -x < 2 \end{cases}$$

Fourier series in the interval $(-l, l)$

EX 1. Find the Fourier expansion of $f(x) = \begin{cases} 0, & -2 < x < -1 \\ 1+x, & -1 < x < 0 \\ 1-x, & 0 < x < 1 \\ 0, & 1 < x < 2 \end{cases}$

Solution: The graph of the function is shown below,




$$\begin{aligned} \text{Further } f(-x) &= \begin{cases} 0, & -2 < -x < -1 \\ 1-x, & -1 < -x < 0 \\ 1+x, & 0 < -x < 1 \\ 0, & 1 < -x < 2 \end{cases} \\ &= \begin{cases} 0, & 2 > x > 1 \\ 1-x, & 1 > x > 0 \\ 1+x, & 0 > x > -1 \\ 0, & -1 > x > -2 \end{cases} \end{aligned}$$

Fourier series in the interval $(-l, l)$

EX 1. Find the Fourier expansion of $f(x) = \begin{cases} 0, & -2 < x < -1 \\ 1+x, & -1 < x < 0 \\ 1-x, & 0 < x < 1 \\ 0, & 1 < x < 2 \end{cases}$

Solution: The graph of the function is shown below,




$$\begin{aligned} \text{Further } f(-x) &= \begin{cases} 0, & -2 < -x < -1 \\ 1-x, & -1 < -x < 0 \\ 1+x, & 0 < -x < 1 \\ 0, & 1 < -x < 2 \end{cases} \\ &= \begin{cases} 0, & 2 > x > 1 \\ 1-x, & 1 > x > 0 \\ 1+x, & 0 > x > -1 \\ 0, & -1 > x > -2 \end{cases} = \begin{cases} 0, & -2 < x < -1 \\ 1+x, & -1 < x < 0 \\ 1-x, & 0 < x < 1 \\ 0, & 1 < x < 2 \end{cases} \\ &= f(x) \end{aligned}$$

Fourier series in the interval $(-l, l)$

EX 1. Find the Fourier expansion of $f(x) = \begin{cases} 0, & -2 < x < -1 \\ 1+x, & -1 < x < 0 \\ 1-x, & 0 < x < 1 \\ 0, & 1 < x < 2 \end{cases}$

Solution: The graph of the function is shown below,



$$\begin{aligned} \text{Further } f(-x) &= \begin{cases} 0, & -2 < -x < -1 \\ 1-x, & -1 < -x < 0 \\ 1+x, & 0 < -x < 1 \\ 0, & 1 < -x < 2 \end{cases} \\ &= \begin{cases} 0, & 2 > x > 1 \\ 1-x, & 1 > x > 0 \\ 1+x, & 0 > x > -1 \\ 0, & -1 > x > -2 \end{cases} = \begin{cases} 0, & -2 < x < -1 \\ 1+x, & -1 < x < 0 \\ 1-x, & 0 < x < 1 \\ 0, & 1 < x < 2 \end{cases} \\ &= f(x) \end{aligned}$$

$\therefore f(x)$ is an even function.

$\therefore b_n = 0$ and $l = 2$.

$$\therefore f(x) = a_0 + \sum a_n \cos \frac{n\pi x}{l}$$

$$\begin{aligned}\therefore f(x) &= a_0 + \sum a_n \cos \frac{n\pi x}{l} \\ &= a_0 + \sum a_n \cos \frac{n\pi x}{2}\end{aligned}$$

$$\therefore f(x) = a_0 + \sum a_n \cos \frac{n\pi x}{l}$$

$$= a_0 + \sum a_n \cos \frac{n\pi x}{2}$$

$$\therefore a_0 = \frac{1}{l} \int_0^l f(x) dx = \frac{1}{2} \int_0^2 f(x) dx$$

$$\therefore f(x) = a_0 + \sum a_n \cos \frac{n\pi x}{l}$$

$$= a_0 + \sum a_n \cos \frac{n\pi x}{2}$$

$$\therefore a_0 = \frac{1}{l} \int_0^l f(x) dx = \frac{1}{2} \int_0^2 f(x) dx$$

$$= \frac{1}{2} \left[\int_0^1 (1-x) dx + \int_1^2 0 \cdot dx \right]$$

$$\therefore f(x) = a_0 + \sum a_n \cos \frac{n\pi x}{l}$$

$$= a_0 + \sum a_n \cos \frac{n\pi x}{2}$$

$$\therefore a_0 = \frac{1}{l} \int_0^l f(x) dx = \frac{1}{2} \int_0^2 f(x) dx$$

$$= \frac{1}{2} \left[\int_0^1 (1-x) dx + \int_1^2 0 \cdot dx \right]$$

$$= \frac{1}{2} \left[x - \frac{x^2}{2} \right]_0^1 = \frac{1}{4}$$

$$a_n = \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx = \frac{2}{2} \int_0^2 f(x) \cos \frac{n\pi x}{2} dx$$

$$\begin{aligned} a_n &= \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx = \frac{2}{2} \int_0^2 f(x) \cos \frac{n\pi x}{2} dx \\ &= \int_0^1 (1-x) \cos \frac{n\pi x}{2} dx + \int_1^2 0 \cdot dx \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx = \frac{2}{2} \int_0^2 f(x) \cos \frac{n\pi x}{2} dx \\
 &= \int_0^1 (1-x) \cos \frac{n\pi x}{2} dx + \int_1^2 0 \cdot dx \\
 &= \left[(1-x) \left(\frac{2}{n\pi} \sin \frac{n\pi x}{2} \right) - (-1) \left(-\frac{4}{n^2 \pi^2} \cos \frac{n\pi x}{2} \right) \right]_0^1
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx = \frac{2}{2} \int_0^2 f(x) \cos \frac{n\pi x}{2} dx \\
 &= \int_0^1 (1-x) \cos \frac{n\pi x}{2} dx + \int_1^2 0 \cdot dx \\
 &= \left[(1-x) \left(\frac{2}{n\pi} \sin \frac{n\pi x}{2} \right) - (-1) \left(-\frac{4}{n^2 \pi^2} \cos \frac{n\pi x}{2} \right) \right]_0^1 \\
 &= \left[\left(0 - \frac{4}{n^2 \pi^2} \cos \frac{n\pi}{2} \right) - \left(0 - \frac{4}{n^2 \pi^2} \right) \right]
 \end{aligned}$$

$$\begin{aligned}
 a_n &= \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx = \frac{2}{2} \int_0^2 f(x) \cos \frac{n\pi x}{2} dx \\
 &= \int_0^1 (1-x) \cos \frac{n\pi x}{2} dx + \int_1^2 0 \cdot dx \\
 &= \left[(1-x) \left(\frac{2}{n\pi} \sin \frac{n\pi x}{2} \right) - (-1) \left(-\frac{4}{n^2 \pi^2} \cos \frac{n\pi x}{2} \right) \right]_0^1 \\
 &= \left[\left(0 - \frac{4}{n^2 \pi^2} \cos \frac{n\pi}{2} \right) - \left(0 - \frac{4}{n^2 \pi^2} \right) \right] \\
 &= \frac{4}{n^2 \pi^2} \left(1 - \cos \frac{n\pi}{2} \right)
 \end{aligned}$$

$$a_n = \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx = \frac{2}{2} \int_0^2 f(x) \cos \frac{n\pi x}{2} dx$$

$$= \int_0^1 (1-x) \cos \frac{n\pi x}{2} dx + \int_1^2 0 \cdot dx$$

$$= \left[(1-x) \left(\frac{2}{n\pi} \sin \frac{n\pi x}{2} \right) - (-1) \left(-\frac{4}{n^2 \pi^2} \cos \frac{n\pi x}{2} \right) \right]_0^1$$

$$= \left[\left(0 - \frac{4}{n^2 \pi^2} \cos \frac{n\pi}{2} \right) - \left(0 - \frac{4}{n^2 \pi^2} \right) \right]$$

$$= \frac{4}{n^2 \pi^2} \left(1 - \cos \frac{n\pi}{2} \right)$$

$$\therefore f(x) = \frac{1}{4} + \frac{4}{\pi^2} \sum \frac{1}{n^2} \left(1 - \cos \frac{n\pi}{2} \right) \cos \frac{n\pi x}{2}$$