derive posterior distribution

Renjie Wei

2022-05-02

The suggested Bayesian model

$$Y_i(t+6) = \beta_{0,i} + \beta_{1,i}Y_i(t) + \beta_{2,i}\Delta_{i,1}(t) + \beta_{3,i}\Delta_{i,2}(t) + \beta_{4,i}\Delta_{i,3}(t) + \epsilon_i(t)$$

where $Y_i(t)$ the wind speed at time t (i.e. 6 hours earlier), $\Delta_{i,1}(t)$, $\Delta_{i,2}(t)$ and $\Delta_{i,3}(t)$ are the changes of latitude, longitude and wind speed between t and t+6, and $\epsilon_{i,t}$ follows a normal distributions with mean zero and variance σ^2 , independent across t.

In the model, $\beta_i = (\beta_{0,i}, \beta_{1,i}, ..., \beta_{5,i})$ are the random coefficients associated the *i*th hurricane, we assume that

$$\boldsymbol{\beta}_i \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

follows a multivariate normal distributions with mean μ and covariance matrix Σ .

We assume the following non-informative or weak prior distributions for σ^2 , β and Σ .

$$P(\sigma^2) \propto \frac{1}{\sigma^2}; \quad P(\mu) \propto 1; \quad P(\Sigma^{-1}) \propto |\Sigma|^{-(d+1)} \exp(-\frac{1}{2}\Sigma^{-1})$$

d is dimension of β .

1

Let $\mathbf{B} = (\boldsymbol{\beta}_1^\top, ..., \boldsymbol{\beta}_n^\top)^\top$, derive the posterior distribution of the parameters $\Theta = (\mathbf{B}^\top, \boldsymbol{\mu}^\top, \sigma^2, \Sigma)$. Note from given Bayesian model, let

$$\epsilon_{i}(t) = Y_{i}(t+6) - \left(\beta_{0,i} + \beta_{1,i}Y_{i}(t) + \beta_{2,i}\Delta_{i,1}(t) + \beta_{3,i}\Delta_{i,2}(t) + \beta_{4,i}\Delta_{i,3}(t)\right) \stackrel{i.i.d}{\sim} N(0,\sigma^{2})$$
or
$$Y_{i}(t+6) \sim N(\boldsymbol{X}_{i}(t)\boldsymbol{\beta}_{i}^{\top},\sigma^{2})$$

where $\boldsymbol{X}_{i}(t)=(1,Y_{i}(t),\Delta_{i,1}(t),\Delta_{i,2}(t),\Delta_{i,3}(t))$, and $\boldsymbol{\beta}_{i}=(\beta_{0,i},\beta_{1,i},\beta_{2,i},\beta_{3,i},\beta_{4,i})$. Therefore, the wind speed of i^{th} hurricane at time t follows the normal distribution with the pdf below

$$f_{Y_i(t+6)}(y_i(t+6) \mid \boldsymbol{X}_i(t), \boldsymbol{\beta}_i, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left\{-\frac{1}{2\sigma^2} \left(y_i(t+6) - \boldsymbol{X}_i(t)\boldsymbol{\beta}_i^{\top}\right)^2\right\}$$

Therefore, the conditional distribution of Y_i , the wind speed of i^{th} hurricane follows the multivariate normal distribution below, (since $Y_i(t)$'s are independent across t)

$$(\boldsymbol{Y}_i \mid \boldsymbol{X}_i, \boldsymbol{\beta}_i, \sigma^2) \sim \mathcal{N}(\boldsymbol{X}_i \boldsymbol{\beta}_i^\top, \sigma^2 I)$$

where Y_i is an m_i -dimensional vector and \boldsymbol{X}_i is a $m_i \times d$ matrix.

Hence, the joint likelihood function of all i's hurricanes can be expresses as

$$L_Y(\mathbf{B}^\top, \sigma^2) = \prod_{i=1}^n \left\{ \det(2\pi\sigma^2 I)^{-\frac{1}{2}} \exp\left(-\frac{1}{2}(\boldsymbol{Y}_i - \boldsymbol{X}_i \boldsymbol{\beta}_i^\top)^\top (\sigma^2 I)^{-1} (\boldsymbol{Y}_i - \boldsymbol{X}_i \boldsymbol{\beta}_i^\top)\right) \right\}$$

where I is an identical matrix with dimension consistent with Y_i .

From Bayesian theorem, the posterior distribution for Θ is

$$\pi(\boldsymbol{\Theta}|\boldsymbol{Y}) = \pi(\boldsymbol{B}^{\top}, \boldsymbol{\mu}^{\top}, \sigma^{2}, \Sigma \mid \boldsymbol{Y}) \propto L_{Y}(\boldsymbol{B}^{\top}, \sigma^{2}) \times \pi(\boldsymbol{B} \mid \boldsymbol{\mu}, \boldsymbol{\Sigma}) \times \pi(\boldsymbol{\mu}) \times \pi(\sigma^{2}) \times \pi(\boldsymbol{\Sigma}),$$

where $\pi(\mathbf{B} \mid \boldsymbol{\beta}, \Sigma)$ is the joint multivariate normal density of β , since

$$\boldsymbol{\beta}_i \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

Therefore

$$\pi(\mathbf{B} \mid \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \prod_{i=1}^n \Big\{ \det(2\pi\boldsymbol{\Sigma})^{-\frac{1}{2}} \exp(-\frac{1}{2}(\boldsymbol{\beta}_i - \boldsymbol{\mu})\boldsymbol{\Sigma}^{-1}(\boldsymbol{\beta}_i - \boldsymbol{\mu})^\top) \Big\}.$$

So we have the following posterior distribution:

$$\pi(\mathbf{B}^{\top}, \boldsymbol{\mu}^{\top}, \sigma^{2}, \boldsymbol{\Sigma} \mid \boldsymbol{Y}) \propto \prod_{i=1}^{n} \left\{ (2\pi\sigma^{2})^{-m_{i}/2} \exp\left\{-\frac{1}{2}(\boldsymbol{Y}_{i} - \boldsymbol{X}_{i}\boldsymbol{\beta}_{i}^{\top})^{\top}(\sigma^{2}I)^{-1}(\boldsymbol{Y}_{i} - \boldsymbol{X}_{i}\boldsymbol{\beta}_{i}^{\top})\right\} \right\}$$

$$\times \prod_{i=1}^{n} \left\{ \det(2\pi\boldsymbol{\Sigma})^{-\frac{1}{2}} \exp\left\{-\frac{1}{2}(\boldsymbol{\beta}_{i} - \boldsymbol{\mu})\boldsymbol{\Sigma}^{-1}(\boldsymbol{\beta}_{i} - \boldsymbol{\mu})^{\top}\right\} \right\} \times \frac{1}{\sigma^{2}} \times \det(\boldsymbol{\Sigma})^{-(d+1)} \exp\left\{-\frac{1}{2}\boldsymbol{\Sigma}^{-1}\right\}$$

To apply MCMC, we need to find conditional posterior distribution of each parameter.

1. For $\pi(\mathbf{B} \mid .)$

$$\begin{split} \pi(\mathbf{B}\mid.) &\propto L_{Y}(\mathbf{B}^{\top}, \sigma^{2}) \times \pi(\mathbf{B}\mid\boldsymbol{\mu}, \boldsymbol{\Sigma}) \\ &\propto \prod_{i=1}^{n} \left\{ \det(2\pi\sigma^{2}I)^{-\frac{1}{2}} \exp\left(-\frac{1}{2}(\boldsymbol{Y}_{i} - \boldsymbol{X}_{i}\boldsymbol{\beta}_{i}^{\top})^{\top}(\sigma^{2}I)^{-1}(\boldsymbol{Y}_{i} - \boldsymbol{X}_{i}\boldsymbol{\beta}_{i}^{\top})\right) \right\} \\ &\times \prod_{i=1}^{n} \left\{ \det(2\pi\boldsymbol{\Sigma})^{-\frac{1}{2}} \exp(-\frac{1}{2}(\boldsymbol{\beta}_{i} - \boldsymbol{\mu})\boldsymbol{\Sigma}^{-1}(\boldsymbol{\beta}_{i} - \boldsymbol{\mu})^{\top}) \right\} \\ &\propto \prod_{i=1}^{n} \exp\{-\frac{1}{2}\left((\boldsymbol{Y}_{i} - \boldsymbol{X}_{i}\boldsymbol{\beta}_{i}^{\top})^{\top}(\sigma^{2}I)^{-1}(\boldsymbol{Y}_{i} - \boldsymbol{X}_{i}\boldsymbol{\beta}_{i}^{\top}) + (\boldsymbol{\beta}_{i} - \boldsymbol{\mu})\boldsymbol{\Sigma}^{-1}(\boldsymbol{\beta}_{i} - \boldsymbol{\mu})^{\top}) \right) \} \end{split}$$

Considering the exponential term in each component in the product,

$$\begin{split} &(\boldsymbol{Y}_{i}-\boldsymbol{X}_{i}\boldsymbol{\beta}_{i}^{\top})^{\top}(\sigma^{2}I)^{-1}(\boldsymbol{Y}_{i}-\boldsymbol{X}_{i}\boldsymbol{\beta}_{i}^{\top})+(\boldsymbol{\beta}_{i}-\boldsymbol{\mu})\boldsymbol{\Sigma}^{-1}(\boldsymbol{\beta}_{i}-\boldsymbol{\mu})^{\top})\\ &=\boldsymbol{Y}_{i}^{\top}(\sigma^{2}I)^{-1}\boldsymbol{Y}_{i}+\boldsymbol{\beta}_{i}\boldsymbol{X}_{i}^{\top}(\sigma^{2}I)^{-1}\boldsymbol{X}_{i}\boldsymbol{\beta}_{i}^{\top}-2\boldsymbol{Y}_{i}(\sigma^{2}I)^{-1}\boldsymbol{X}_{i}\boldsymbol{\beta}_{i}^{\top}\\ &+\boldsymbol{\beta}_{i}\boldsymbol{\Sigma}^{-1}\boldsymbol{\beta}_{i}^{\top}+\boldsymbol{\mu}\boldsymbol{\Sigma}^{-1}\boldsymbol{\mu}^{\top}-2\boldsymbol{\mu}\boldsymbol{\Sigma}^{-1}\boldsymbol{\beta}_{i}^{\top}\\ &=\boldsymbol{\beta}_{i}\boldsymbol{V}\boldsymbol{\beta}_{i}^{\top}-2\boldsymbol{M}\boldsymbol{\beta}_{i}^{\top}+\boldsymbol{R} \end{split}$$

where,

$$\begin{split} \boldsymbol{V} &= \boldsymbol{\Sigma}^{-1} + \boldsymbol{X}_i^\top (\sigma^2 I)^{-1} \boldsymbol{X}_i \\ \boldsymbol{M} &= \boldsymbol{Y}_i (\sigma^2 I)^{-1} \boldsymbol{X}_i + \boldsymbol{\mu} \boldsymbol{\Sigma}^{-1} \\ \boldsymbol{R} &= \boldsymbol{Y}_i^\top (\sigma^2 I)^{-1} \boldsymbol{Y}_i + \boldsymbol{\mu} \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}^\top \end{split}$$

re-writing the conditional posterior distribution,

$$\pi(\mathbf{B} \mid .) \propto$$

```
# betavec: 1*5
# sigma2: 1*1
# mu: 1*5
# Sigma: 5*5
\# d = 5
# n = 702
loglik_sigma2 <- function(sigma2){</pre>
    if (sigma2 <= 0) {</pre>
        return(-Inf)
    } else {
        return(log(1/sigma2^2))
    }
}
loglik_Sigma <- function(Sigma, d){</pre>
    if (min(eigen(Sigma)$values) <= 0) {</pre>
         return(-Inf)
    } else {
```

```
loglik = log(det(Sigma)^(-(d+1))*exp(-(1/2)*solve(Sigma)))
                                                          return(loglik)
                             }
}
 \# log likelihood for y given a specific hurricane's data given sigma2 and betavec
loglik_Y <- function(y, X, betavec, sigma2){</pre>
                             m = length(y)
                             loglik = (2*pi*sigma2^2)^(-(m/2))*exp(-(1/2)*t(X \%*\% t(betavec)) \%*\% solve(diag(sigma2, m)) \%*\% (X \%*\% t(betavec)) \%*\% solve(diag(sigma2, m)) \%*\% (X \%*\% t(betavec)) \%*\% solve(diag(sigma2, m)) \%*\% (X \%*\% t(betavec)) %*% solve(diag(sigma2, m)) %*% (X \%*\% t(betavec)) %*% (X \%*\% t(betav
                             loglik = log(loglik)
                             return(loglik)
}
 # log likelihood for betavec given a specific hurricane's mu and Sigma
loglik_betavec <- function(betavec, mu, Sigma){</pre>
                             if (min(eigen(Sigma)$values) <= 0) {</pre>
                                                          return(-Inf)
                             } else {
                                                           loglik = log(det(2*pi*Sigma)^(-(1/2))*exp(-(1/2)*(betavec - mu) \%*\% solve(Sigma) \%*\% t(betavec - mu) \%*\% solve(Sigma) %*% t(betavec - mu) %*% solve(Sigma) %*% solve(Sigma
                                                           return(loglik)
                             }
```