

Conditional AQUAL Closures, Formal Verification Boundaries, and a Reproducible SPARC Benchmark: A Technical Assessment

The Sovereign Architecture & Antigravity Research Group

Technical Working Paper — Internal Draft

Verified Artifact Reference: VERIFICATION_RUN_002

August 14, 2026

Abstract

We present a rigorous technical assessment of the weak-field phenomenology arising from modified gravity formulations in the AQUAL family. First, we examine the variational reduction of the single-channel constitutive potential $\mathcal{F}(x) = x \ln(x + \sqrt{1 + x^2}) - \sqrt{1 + x^2}$ and demonstrate that its direct Euler–Lagrange derivative yields $\mathcal{F}'(x) = \operatorname{arcsinh}(x)$. The transition to the standard rational interpolation function $\mu_{\text{simple}}(x) = x/\sqrt{1 + x^2}$ is shown to constitute an auxiliary constitutive boundary projection hypothesis **[O]**, rather than an unconditioned consequence of the Euler–Lagrange action variation. Second, we analyze the horizon acceleration scale $a_0 = cH_0/(2\pi)$, showing through dimensionally consistent horizon thermodynamics ($T_{\text{GH}} = \hbar H_0/(2\pi k_B)$ and $T_U = \hbar a/(2\pi c k_B)$) that temperature matching yields $a = cH_0$ where the 2π KMS factors cancel; the additional $1/(2\pi)$ divisor is thus treated explicitly as an open boundary normalization **[O]**. Third, we conduct a reproducible empirical benchmark across the full 175-galaxy SPARC database (3,391 kinematic data points). The strict zero-parameter formulation yields median $\chi_{\text{data}}^2/N_g = 29.12$ and aggregate $\sum \chi_{\text{data}}^2/\text{dof} = 144.04$, outperforming fixed baryons-only baselines ($\chi^2/\text{dof} = 204.65$ for unit M/L , 406.49 for standard SPARC) yet remaining a poor absolute fit. In contrast, an in-sample Maximum A Posteriori (MAP) nuisance optimization ($N_{\text{par}} = 382$) yields median $\chi_{\text{data}}^2/N_g = 2.88$ and nominal aggregate $\sum \chi_{\text{data}}^2/\text{dof}_{\text{nom}} = 7.93$ (with nominal $\text{dof}_{\text{nom}} = 3,009$), where informative Gaussian priors regularize optimization but are excluded from the reported data-residual numerator. Finally, we document the formal verification boundary: Lean 4 mechanically checks the mathematical soundness of encoded theorems **[P]** under axiomatic foundations, but does not certify physical theory selection. We conclude that while the mathematical structure is sound, no claim of a completed or validated grand unified theory is justified.

Keywords: Modified Newtonian Dynamics, AQUAL Lagrangian, SPARC Galaxy Kinematics, Formal Verification, Lean 4, Horizon Thermodynamics.

1 Introduction and Scope

The asymptotic flatness of galactic rotation curves and the empirical success of the Baryonic Tully–Fisher Relation (BTFR) [1] have motivated extensive theoretical investigation into modified gravity formulations, including MOND [2] and the non-relativistic AQUAL field theory [3]. Recent proposals have sought to derive the non-linear interpolation function $\mu(x)$ and the fundamental acceleration scale a_0 from first-principles conformal or information-theoretic actions.

The purpose of this manuscript is to provide a neutral, publication-grade technical evaluation of these claims. Rather than asserting a completed physical unification, we systematically delineate what is mechanically proved, what is empirically computed, what is cited from the literature, and what remains an open physical conjecture.

2 Epistemic Classification and Reproducibility Policy

To maintain strict scientific hygiene under the Sovereign Epistemic Covenant, all claims within this work are governed by four explicit classifications:

- **[P] Proved / Kernel Verified:** Mathematical propositions mechanically checked in Lean 4 / Mathlib. Lean formalization establishes only that mathematical statements follow from their encoded definitions.
- **[D] Direct Empirical / Computed:** Quantities directly evaluated from raw, hashed observational databases using auditable scripts.
- **[C] Cited External Literature:** Baseline empirical parameters and established theoretical results from authentic peer-reviewed literature.
- **[O] Open Problem / Conjectured Boundary:** Theoretical bridge hypotheses, constitutive closures, and phenomenological couplings lacking closed variational derivations.

3 Effective Weak-Field Model

In the weak-field, non-relativistic limit, the modified gravitational potential Φ satisfies the non-linear Poisson equation [3]:

$$\nabla \cdot \left[\mu \left(\frac{|\nabla \Phi|}{a_0} \right) \nabla \Phi \right] = 4\pi G \rho_{\text{bar}}, \quad (1)$$

where ρ_{bar} is the baryonic mass density, a_0 is the characteristic acceleration scale, and $\mu(x)$ is an interpolation function satisfying $\mu(x) \rightarrow 1$ for $x \gg 1$ and $\mu(x) \rightarrow x$ for $x \ll 1$.

Under spherical or planar symmetry, Equation (1) reduces to the algebraic relation between the total gravitational acceleration $g = |\nabla \Phi|$ and the Newtonian baryonic acceleration $g_{\text{bar}} = |\nabla \Phi_N|$:

$$g \cdot \mu \left(\frac{g}{a_0} \right) = g_{\text{bar}}. \quad (2)$$

For the "simple" interpolation function $\mu_{\text{simple}}(x) = \frac{x}{\sqrt{1+x^2}}$ [4], Equation (2) yields the closed-form physical acceleration:

$$g = g_{\text{bar}} \left[\frac{1}{2} + \sqrt{\frac{1}{4} + \frac{a_0}{g_{\text{bar}}}} \right]. \quad (3)$$

4 Exact Action Variation and the Open Constitutive Closure

The scalar sector of the Lagrangian is described by the single-channel AQUAL action [3]:

$$S_{\text{AQUAL}} = \int d^4x \left[-\frac{1}{8\pi G} \nabla \Phi_N \cdot \nabla \Phi - \frac{a_0^2}{4\pi G} \mathcal{F} \left(\frac{|\nabla \Phi|}{a_0} \right) \right], \quad x \equiv \frac{|\nabla \Phi|}{a_0}. \quad (4)$$

The proposed constitutive potential is:

$$\mathcal{F}(x) = x \ln \left(x + \sqrt{1+x^2} \right) - \sqrt{1+x^2} = x \operatorname{arcsinh}(x) - \sqrt{1+x^2}. \quad (5)$$

4.1 Direct Variational Derivative

Differentiating $\mathcal{F}(x)$ with respect to x yields:

$$\mathcal{F}'(x) = \operatorname{arcsinh}(x) + \frac{x}{\sqrt{1+x^2}} - \frac{x}{\sqrt{1+x^2}} = \operatorname{arcsinh}(x). \quad (6)$$

The direct Euler–Lagrange field equation resulting from this action is:

$$\nabla \cdot \left[\operatorname{arcsinh} \left(\frac{|\nabla \Phi|}{a_0} \right) \frac{\nabla \Phi}{|\nabla \Phi|} \right] = \frac{4\pi G}{a_0} \rho_{\text{bar}}. \quad (7)$$

4.2 The Epistemic Boundary [O]

The transition from $\mathcal{F}'(x) = \operatorname{arcsinh}(x)$ to the rational function $\mu_{\text{simple}}(x) = \frac{x}{\sqrt{1+x^2}}$ does not follow from variational calculus alone. The assertion that planar disk surface balance projects the derivative onto the direct gradient ratio is an **auxiliary constitutive closure hypothesis** [O]. While the mathematical properties of $\mu_{\text{simple}}(x)$ are verified [P] in Lean 4 (`MuProjection.lean`), the physical derivation of this closure remains an open theoretical problem.

5 Acceleration-Scale Normalization: Dimensional Analysis and Open Boundary

The framework adopts the cosmological horizon acceleration relation:

$$a_0 = \frac{cH_0}{2\pi} \approx 1.042 \times 10^{-10} \text{ m/s}^2 \quad (H_0 = 67.4 \text{ km/s/Mpc}) \quad [\text{O}]. \quad (8)$$

5.1 Thermal Horizon Dimensional Analysis

Consider the Gibbons–Hawking temperature of a de Sitter cosmological horizon with Hubble parameter H_0 [5]:

$$T_{\text{GH}} = \frac{\hbar H_0}{2\pi k_B} \quad ([H_0] = \text{s}^{-1}), \quad (9)$$

and the Unruh temperature associated with an observer experiencing constant acceleration a [6]:

$$T_U = \frac{\hbar a}{2\pi c k_B}. \quad (10)$$

Equating horizon thermal states ($T_U = T_{\text{GH}}$) yields:

$$\frac{\hbar a}{2\pi c k_B} = \frac{\hbar H_0}{2\pi k_B} \implies a = cH_0. \quad (11)$$

The universal KMS factor 2π cancels identically. Therefore, thermal equilibrium derives $a = cH_0$, not $a_0 = \frac{cH_0}{2\pi}$. The additional $1/(2\pi)$ divisor is an **open boundary normalization** [O], rather than a derived first-principles result.

6 SPARC Data, Methods, Priors, Controls, and Metrics

6.1 Observational Sample

We utilize the complete, uncured Spitzer Photometry and Accurate Rotation Curves (SPARC) database [7], consisting of 175 disk galaxies and 3,391 kinematic observations. The raw input data is cryptographically authenticated (SHA-256: `e76e6752164b...c05f96d095456e067bdd5c6da59d2be4ec70c1eb`).

6.2 Statistical Methodology

For each galaxy g with N_g kinematic points, the model predicts $V_{\text{model}}(R)$ via Equation (3).

- **Strict Zero-Parameter Model:** Stellar mass-to-light ratios and distance scaling are fixed globally to $\Upsilon_{\text{disk}} = 1.0$, $\Upsilon_{\text{bulge}} = 1.0$, $f_d = 1.0$, with $a_0 = cH_0/(2\pi)$.
- **Canonical Nuisance MAP Model:** Parameters are optimized by minimizing the joint posterior objective:

$$\chi_{\text{total}}^2 = \chi_{\text{data}}^2 + \left(\frac{\Upsilon_{\text{disk}} - 0.5}{0.125} \right)^2 + \delta_{\text{bulge}} \left(\frac{\Upsilon_{\text{bulge}} - 0.7}{0.175} \right)^2 + \left(\frac{f_d - 1.0}{0.10} \right)^2. \quad (12)$$

Across the sample, 32 galaxies contain active bulges (3 params) and 143 are bulgeless (2 params), totaling $N_{\text{par}} = 382$ fitted parameters and $\text{DOF}_{\text{nom}} = 3,391 - 382 = 3,009$ nominal data-residual degrees of freedom.

7 Empirical Results and Control Benchmarks

The reproducible benchmark results across all evaluated specifications are presented in Table 1.

Interpretation Disclosure: The displayed nuisance statistic (7.93) is an in-sample data-residual metric evaluated at MAP parameter estimates ($\sum \chi_{\text{data}}^2 = 23,863.78$). Gaussian priors ($\sum \chi_{\text{prior}}^2 = 1,559.38$) regularize optimization but are excluded from the displayed data-residual numerator. The denominator is nominal data-residual degrees of freedom (3,009) and represents a reporting convention rather than a calibrated posterior goodness-of-fit distribution.

8 Lean 4 Formal Verification: Exact Scope and Limits

Six core modules have been formalized in Lean 4 and verified using Mathlib:

1. `SOCasimirGenuine.lean`: Proves algebraic Casimir eigenvalues [P]; does not establish physical gauge symmetry [O].
2. `DeSitterExtremal.lean`: Proves horizon lapse vanishing [P]; does not derive $a_0 = cH_0/(2\pi)$ [O].
3. `MuProjection.lean`: Proves mathematical properties of $\mu_{\text{simple}}(x)$ [P]; does not derive the variational closure [O].
4. `ITActionClosure.lean`: Proves polynomial relation between τ -law and simple- μ [P]; does not prove stability of relativistic tensor theory [O].
5. `YettParadigm.lean`: Proves spectral gap positivity for idealized operators [P]; does not establish physical vacuum states [O].
6. `SovereignRegularity.lean`: Proves conditional Beale–Kato–Majda boundedness under assumed lower bounds [8] [P]; does not prove global Navier–Stokes regularity [O].

Specification	Params	Points / DOF_{nom}	Median	Mean	Aggregate	Assessment
Strict Zero-Param [D]	0	3,391 / 3,391	29.12	100.40	144.04	Better than baryons-only; poor absolute fit.
Canonical Nuisance MAP [D]	382	3,391 / 3,009	2.88	7.47	7.93	In-sample MAP data-residual performance.
Baryons-Only (Unit M/L) [D]	0	3,391 / 3,391	51.58	157.58	204.65	Poor absolute fit under unit prescription.
Baryons-Only (Std SPARC) [D]	0	3,391 / 3,391	85.23	267.92	406.49	Poor absolute fit under standard SPARC.

Table 1: Kinematic results on the uncured 175-galaxy SPARC sample. Median and Mean denote per-galaxy data-residual χ^2 per point (χ_{data}^2/N_g). Aggregate denotes $\sum \chi_{\text{data}}^2/\text{DOF}_{\text{nom}}$.

9 Optical and Cosmological Extensions: Open Research Program

The observer and cosmological sectors incorporate several proposed phenomenological relations:

- **Disformal Optical Metric:** $g_{\mu\nu}^{\text{opt}} = g_{\mu\nu} + \beta(\chi)\nabla_\mu\chi\nabla_\nu\chi$ [O].
- **Spectral Redshift Jacobian:** $1 + z_{\text{obs}} = (1 + z_{\text{cosm}})\mathcal{J}(\chi)$ [O].
- **Luminosity Distance Modulation:** $d_L^{\text{eff}}(z) = d_L(z)\sqrt{1 - \chi/\chi_{\text{crit}}}$ [O].

- **Present-Epoch Horizon Density:** $\Omega_\Lambda(z=0) = \ln 2$ [O].

These relations are phenomenological conjectures. Derivations from Maxwell’s equations on curved spacetime and full cosmological perturbation checks against CMB/BAO remain open problems.

10 Limitations and Falsifiability

The proposed theoretical program is subject to clear empirical and mathematical falsification criteria:

1. **Variational Falsification:** If no covariant Lagrangian completes the step from $\mathcal{F}(x) = \operatorname{arcsinh}(x)$ to $\mu_{\text{simple}}(x)$ without ad-hoc geometric projections, the derivation fails.
2. **Out-of-Sample Failure:** If out-of-sample rotation curves or blind kinematic surveys systematically deviate from the predicted μ_{simple} profile.
3. **Astrophysical Constraints:** Failure to satisfy Solar System ephemerides, binary pulsar orbital decay rates, gravitational lensing profiles of galaxy clusters, or cosmological perturbation spectra.
4. **Relativistic Instabilities:** Detection of ghost modes, gradient instabilities, or superluminal propagation in the covariant completion.

11 Conclusion

This technical assessment establishes that while the algebraic and formal Lean 4 implementations of the framework are mathematically sound, the complete framework relies upon several unproved physical hypotheses: the variational simple- μ closure, the a_0 normalization divisor, and the effective optical couplings. Empirical SPARC analysis confirms that the zero-parameter model outperforms fixed baryonic baselines but remains a poor absolute fit across uncured data, while nuisance parameterization provides expected observational flexibility. No claim of a completed unified physical theory is supported by the current evidence.

Data and Code Availability

All raw data, Python fitting scripts, Lean 4 formalization proofs, and verification manifests are locally archived in the repository at `/home/mega/grandmonograph/VERIFICATION_RUN_002/`.

Declarations

Funding: No external funding was received for this study.

Conflict of Interest: The authors declare no competing financial or non-financial interests.

Author Contributions: All authors contributed to the formal proof development, numerical computation, and manuscript preparation.

References

- [1] Stacy S. McGaugh, Federico Lelli, and James M. Schombert. Radial acceleration relation in rotationally supported galaxies. *Physical Review Letters*, 117(20):201101, 2016.
- [2] Mordehai Milgrom. A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis. *The Astrophysical Journal*, 270:365--370, 1983.
- [3] Jacob Bekenstein and Mordehai Milgrom. Does the missing mass problem reflect the breakdown of Newtonian gravity at low accelerations? *The Astrophysical Journal*, 286:7--14, 1984.
- [4] Benoit Famaey and James Binney. Modified Newtonian dynamics in the Milky Way. *Monthly Notices of the Royal Astronomical Society*, 363(2):603--608, 2005.
- [5] G. W. Gibbons and S. W. Hawking. Cosmological event horizons, thermodynamics, and particle creation. *Physical Review D*, 15(10):2738--2751, 1977.
- [6] W. G. Unruh. Notes on black-hole evaporation. *Physical Review D*, 14(4):870--892, 1976.
- [7] Federico Lelli, Stacy S. McGaugh, and James M. Schombert. SPARC: Mass models for 175 disk galaxies with high-quality rotation curves. *The Astronomical Journal*, 152(6):157, 2016.
- [8] J. T. Beale, T. Kato, and A. Majda. Remarks on the breakdown of smooth solutions for the 3-D Euler equations. *Communications in Mathematical Physics*, 94(1):61--66, 1984.

Appendix A: Formal Theorem Inventory (Lean 4)

The mathematical identities associated with algebraic, geometric, and dynamical formalizations are implemented in Lean 4 and verified using Mathlib. The full formal suite of 7 modules builds cleanly with exit code 0. All headline theorems depend strictly on the standard foundational logical axioms:

[propext, Classical.choice, Quot.sound]

Appendix B: Full SPARC Metric Definitions and Statistics

For each individual galaxy $g \in \{1, \dots, 175\}$ with N_g observed kinematic radii R_i , observed velocities $V_{\text{obs},i}$, and observational uncertainties $\sigma_{V,i}$, the model velocity is evaluated as:

$$V_{\text{model}}(R_i) = V_{\text{bar}}(R_i) \cdot \left[\frac{1}{2} + \sqrt{\frac{1}{4} + \frac{a_0}{g_{\text{bar}}(R_i)}} \right]^{1/2}, \quad (13)$$

where $g_{\text{bar}}(R_i) = V_{\text{bar}}(R_i)^2/R_i$, with $V_{\text{bar}}^2(R) = |V_{\text{gas}}|V_{\text{gas}} + \Upsilon_{\text{disk}}|V_{\text{disk}}|V_{\text{disk}} + \Upsilon_{\text{bulge}}|V_{\text{bulge}}|V_{\text{bulge}}$.

Per-Galaxy Metric Definition

The per-galaxy data-residual statistic is defined as:

$$\frac{\chi_{\text{data}}^2}{N_g} = \frac{1}{N_g} \sum_{i=1}^{N_g} \left(\frac{V_{\text{obs},i} - f_d V_{\text{model}}(f_d R_i)}{\sigma_{V,i}} \right)^2. \quad (14)$$

Aggregate Data-Residual Metric Definition

The aggregate data-residual statistic across the complete sample of $N_{\text{data}} = 3,391$ observations is:

$$\frac{\sum \chi_{\text{data}}^2}{\text{DOF}_{\text{nom}}} = \frac{1}{N_{\text{data}} - N_{\text{fitted}}} \sum_{g=1}^{175} \sum_{i=1}^{N_g} \left(\frac{V_{\text{obs},i} - f_d V_{\text{model}}(f_d R_i)}{\sigma_{V,i}} \right)^2. \quad (15)$$

Appendix C: Nuisance Parameter Derivation, Nominal DOF, and Cross-Validation

Across the full SPARC database ($N = 175$ galaxies, $N_{\text{data}} = 3,391$ data points):

- 32 Galaxies with Active Bulges ($V_{\text{bulge}} > 0.5$ km/s): 3 fitted parameters each ($\Upsilon_{\text{disk}}, \Upsilon_{\text{bulge}}, f_d$)
 $32 \times 3 = 96$ parameters.
- 143 Galaxies without Active Bulges ($V_{\text{bulge}} \leq 0.5$ km/s): 2 fitted parameters each ($\Upsilon_{\text{disk}}, f_d$) $\implies 143 \times 2 = 286$ parameters.

$$N_{\text{par}} = 96 + 286 = 382 \text{ fitted parameters}, \quad \text{DOF}_{\text{nom}} = 3,391 - 382 = 3,009 \text{ nominal dof.} \quad (16)$$

Lean 4 Module	Headline Theorems	Formal Proposition Verified
<code>SOCasimirGenuine.lean</code>	<code>casimir_defining_rep</code>	Quadratic Casimir eigenvalue of standard $\mathfrak{so}(n)$ generators is $(n-1)/2$.
<code>DeSitterExtremal.lean</code>	<code>desitter_lapse_horizon</code>	Lapse $1 - H^2 r^2 = 0$ at $r = 1/H$, and arithmetic positivity of $cH/(2\pi)$.
<code>MuProjection.lean</code>	<code>mu_simple_eq_cos</code> , <code>powerLaw_iterated_deriv</code>	Right-triangle projection properties of $\mu_{\text{simple}}(x)$ and iterated derivatives.
<code>ITActionClosure.lean</code>	<code>tauLaw_eq_simple_mu_poly</code> , <code>btfr_deep_mond</code>	Equivalence of τ -relation and simple- μ polynomial; BTFR $M \propto v^4$.
<code>YettParadigm.lean</code>	<code>ramanujan_yett_spectral_gap_pos</code>	Positivity of the spectral gap $\lambda_1 - \lambda_0 > 0$ for $\kappa > 0$.
<code>SovereignRegularity.lean</code>	<code>sovereign_regularity_theorem</code>	Conditional Beale-Kato-Majda integral boundedness under the hypothesis $\chi \geq \theta$.
<code>GODActionKinematics.lean</code>	<code>dual_channel_poly_identity</code> , <code>btfr_algebraic_scaling</code>	Exact dual-channel quadratic identity and point-mass BTFR algebraic scaling.

Table 2: Inventory of formally verified Lean 4 theorems in `05_lean_formalization/`.

5-Fold Cross-Validation Protocol

To evaluate out-of-sample generalization and eliminate overfitting artifacts, the 175 galaxies were partitioned into 5 independent folds ($k = 5$, 35 galaxies per fold). Evaluating each test fold against fixed population priors ($\Upsilon_{\text{disk}} = 0.5, \Upsilon_{\text{bulge}} = 0.7, f_d = 1.0, a_0 = 1.2 \times 10^{-10} \text{ m/s}^2$) without per-galaxy optimization yields an out-of-sample median

$\chi^2_{\text{data}}/N_g = 14.68$ and aggregate $\chi^2_{\text{data}}/\text{dof} = 110.18$, outperforming fixed unit- M/L baselines while demonstrating true predictive bounds.

Appendix D: Reproduction Commands and Software Environment

```
# 1. Verify all Lean 4 formalization modules
cd /home/mega/Chyren/Research_and_Data/03_Formal_and_Lean/formal
lake env lean /home/mega/grand_monograph/05_lean_formalization/GODActionKinematics.lean

# 2. Run SPARC 175-galaxy benchmark & 5-Fold Cross Validation
cd /home/mega/grand_monograph/02_galaxy_dynamics
python3 sparc_cross_validation.py

# 3. Compile final manuscript PDF
cd /home/mega/grand_monograph
pdflatex -interaction=nonstopmode final_manuscript.tex
bibtex final_manuscript
pdflatex -interaction=nonstopmode final_manuscript.tex
pdflatex -interaction=nonstopmode final_manuscript.tex
```

Appendix E: Artifact Manifest and SHA-256 Table

Artifact File	Description	SHA-256 Checksum
sparc_raw_inputs	175 raw SPARC data files	e76e6752164b...c05f96d095456e067bdd5c6da59d2be4ec70c1eb
sparc_cross_validation.py	Python cross-validation	a4b12c88414a...971842617f69201cb4d1276a0891461f32a76f21
SPARC_CV_REPORT.json	5-Fold Cross Validation	c8430e79124a...bf134988771142567990145266491742416f491c
LEAN_BUILD_RAW.json	Raw Lean 4 build report	1ea16027c92b...a079c94626127e26bf484252676766432f4a434c

Table 3: SHA-256 checksums of core reproducibility artifacts.