

Stability of various systems — Solution

X18 (2018) — English translation

A1 Express the angular acceleration of rod $\ddot{\varphi}$ in terms of φ , l and g . Show that the time dependence of the inclination angle φ is given by $\varphi = Ae^{t/\tau} + Be^{-t/\tau}$, where A and B are constants depending on the initial position and initial angular velocity of the rod, and τ is the characteristic time of fall. Express τ in terms of l and g .

The
mo-
ment
equa-
tion
for
the
rod
about
the
hinge
is:

$$I = M\ddot{\varphi},$$

where

$$I = \frac{1}{3}ml^2$$

is
the
mo-
ment
of
in-
er-
tia
of
the
rod
about
its
end,
and
 $M =$

$$mg \cdot \left(\frac{1}{2}l \sin \varphi\right) \simeq \frac{1}{2}mgl\varphi$$

is
the
mo-
ment
of
grav-
ity.
Let's
check
that

$$\varphi = Ae^{t/\tau} + Be^{-t/\tau}$$

is a
so-
lu-

A2 Let the boy now try to support a long thin stick in an upright position on his hand. As soon as the stick begins to fall, for example, to the left, he moves his hand to the left an even greater distance so that the center of mass of the stick is to the right of its fulcrum. As a result, the moment of gravity begins to rotate the stick to the right, reducing its speed of rotation directed to the left. Estimate how long a stick the boy can hold vertically if his reaction time is approximately $\tau_r = 0.2$ s. (Reaction time is the time delay between the command sent by the brain to the hands and the corresponding movement of the hands.)

The
boy's
re-
ac-
tion
time

τ_r
must
be
of
the
or-
der
of
the
char-
ac-
ter-
istic
fall
time

τ .
Us-
ing
this,
we
find
the
length
of
the
stick:

$$l_r = \tau_r^2 \frac{3g}{2}.$$

Answer: 0.50

$$l_r = \tau_r^2 \frac{3g}{2} = 0.59 \text{ m}$$

A3 Humans and birds maintain a standing position in a similar way. They move the fulcrum (the point at the bottom of the foot where normal ground reaction force is applied), for example, changing the angle between the foot and leg to counteract the fall of the upper body. A small bird of height $l = 6$ cm can stand on its feet. Estimate her reaction time from above.

The
bird's
re-
ac-
tion
time

τ_b
must
be
of
the
or-
der
of
the
char-
ac-
ter-
istic
fall
time

τ
Us-
ing
this,
we
find
the
bird's
re-
ac-
tion
time:

$$\tau_b = \sqrt{\frac{2l_b}{3g}}.$$

Answer: 50

$$\tau_b = \sqrt{\frac{2l_b}{3g}} = 0.065 \text{ s}$$

A4 Balance on a bicycle is also maintained by moving the fulcrum, which lies on the line between the points of contact of the wheels and the ground. This line can be moved by turning the steering wheel while driving. Estimate the minimum speed of the cyclist v_m at which he can maintain balance in this way. Consider that its characteristic time of fall is the same as that of a rod of length $L = 2$ m; distance between wheel axles $d = 1$ m.

The
time

τ_m
dur-
ing
which
the
cy-
clist's
ful-
crum
can
be
moved
de-
pends
on
the
base
 d of
the
bi-
cy-
cle
and
its
speed

v_m
as
fol-
lows

$$\tau_m = \frac{d}{v_m}.$$

This
time
must
be
on
the
or-
der
of
the
char-
ac-
ter-
istic
fall
time

τ .

B1 Suppose that initially the tightrope walker stood in almost perfect balance ($\alpha_1 = \alpha_2 = 0$). Due to the instability of the equilibrium, it slowly begins to fall clockwise. He notices this when $t = t_0$, $\alpha_1 = \alpha_2 = \alpha_0 > 0$. He quickly bends to stop falling. Consider that angle β instantly takes on the value β_0 . Express the new values of angles α_1 and α_2 in terms of β_0 and α_0 .

In
the
pro-
cess
of
chang-
ing
the
con-
figu-
ra-
tion,
the
tightrope
walker
is af-
fected
by
the
mo-
ment
of
grav-
ity,
how-
ever,
this
does
not
lead
to a
change
in
the
an-
gu-
lar
mo-
men-
tum
(changes
in
an-
gles
 α_1
and
 α_2
oc-
cur
at a
very
high

B2 Thus, the tightrope walker is now bent and maintains the same body shape ($\beta = \beta_0$) for a time T_b , after which he almost instantly straightens up ($\beta = 0$). He wants to stand upright again $\alpha_1 = \alpha_2 = 0$. Should it have folded clockwise ($\beta_0 > 0$) or counterclockwise ($\beta_0 < 0$)? Explain your answer.

Moment

of

grav-

ity

$M =$

$mg(1.4H)\alpha_1 +$

$mgH \cdot$

$\alpha_2 \cdot$

Let's

sub-

sti-

tute

here

α_1

and

α_2

from

the

pre-

vi-

ous

para-

graph

and

get

$$M = mgH(2.4\alpha_0 - \frac{7}{37}\beta_0).$$

In

or-

der

for

the

tightrope

walker

to

re-

turn

to

the

pre-

vi-

ous

po-

si-

tion,

he

needs

to

re-

verse

the

B3 Now we will assume that $\alpha_0 \ll \beta_0$. Immediately after it straightens up, neither its angular velocity ($\dot{\alpha}_1 = \dot{\alpha}_2$) nor its angle α_1 are zero (they will become zero much later). Find the meaning of the expression $\dot{\alpha}_1/\alpha_1$. Express your answer using H and g .

The
mo-
tion
of a
straight-
ened
tightrope
walker

is
de-
scribed
by
the
equa-
tion
 $2.96mH^2\ddot{\alpha}_1 =$
 $2.4mgH\alpha_1$.

We
look
for
its
so-
lu-
tion
in
the
form

$$\alpha_1 = Ae^{t/\tau} + Be^{-t/\tau}.$$

In
or-
der
for
the
tightrope
walker

to
re-
turn
to
the
start-
ing
po-
si-
tion,

$$A = 0.$$

There-
fore

$$\frac{\dot{\alpha}_1}{\alpha_1} =$$

B4 Find the time T_b during which the tightrope walker is bent. Express your answer in terms of α_0 , β_0 , H and g . We can assume that $\alpha_0 \ll \beta_0$.

As
shown
above,
the
speed
of
the
tightrope
walker
im-
me-
di-
ately
af-
ter
straight-
en-
ing
(af-
ter
the
sec-
ond
change
of
form)
is
 $\dot{\alpha}_f =$
 $-\frac{\alpha}{\tau}$.
The
speed
right
be-
fore
bend-
ing
is
equal
to
 $\dot{\alpha}_i =$
 $\frac{\alpha_0}{\tau}$,
since
this
pro-
cess
is
equiv-
a-
lent,
up
to a
change
in
the


C1 Consider that at moment $t = T/2$ the pendulum was motionless and inclined at a small angle φ_0 . Draw a graph of the angle of inclination φ versus time, and determine the angular displacement of the pendulum $\Delta\varphi$ at time $t = T$, that is, $\Delta\varphi = \varphi(T) - \varphi(T/2)$. We can assume that $\Delta\varphi \ll \varphi_0$.

In 1.50

the
case
where
the
ac-
cel-
era-
tion
is
di-
rected
up-
ward
(for
ex-
am-
ple,
 $0 <$
 $t <$
 $T)$,
the
in-
er-
tial
force
is
di-
rected
down-
ward.
The
equa-
tion
of
mo-
tion
for
this
case
is:

$$\ddot{\varphi} = \frac{2v_0}{lT} \varphi.$$

Since
the
dis-
place-
ments
from
po-
si-
ti-



C2 Since we are still neglecting gravity, only the inertial force has a non-zero moment about the hinge. Find the time average value of this moment of inertia over time $2T$.

Let's
cal-
cu-
late
the
av-
er-
age
mo-
ment
"head-
on":

$$\langle M \rangle = \langle mla\varphi \rangle = ml\langle a \cdot [(\varphi - \langle \varphi \rangle) + \langle \varphi \rangle] \rangle = ml\langle a(\varphi - \langle \varphi \rangle) \rangle + ml\langle \varphi \rangle \langle a \rangle = -mla_0\langle |\varphi - \langle \varphi \rangle| \rangle + 0$$

Let's
cal-
cu-
late
the
av-
er-
age
de-
flec-
tion
mod-
u-
lus:

$$\langle |\varphi(t) - \varphi| \rangle = \frac{2}{T} \int_0^{\frac{T}{2}} \Delta\varphi \left(1 - \frac{4t^2}{T^2} \right) dt = \frac{2}{3} \Delta\varphi = \frac{v_0 T}{6l} \varphi_0$$

Then
the
ex-
pres-
sion
for
the
av-
er-
age
mo-
ment
of
force

$$\langle M \rangle = -\frac{mv_0^2}{3} \varphi_0$$

Answer: 50

$$\langle M \rangle = -\frac{mv_0^2}{3} \varphi_0$$

C3 Now let's take gravity into account. Write the inequality that must hold for g , T , l , and v_0 for vertical position to exist.

Gravity

does

not

af-

fect

the

value

of

the

av-

er-

age

mo-

ment

of

in-

er-

tia

ob-

tained

ear-

lier.

A

mo-

ment

of

grav-

ity

equal

to

$mgl\varphi_0$

is

added.

Then

the

equa-

tion

of

mo-

tion

has

the

fol-

low-

ing

form

$$l^2\ddot{\varphi} = \left(gl - \frac{v_0^2}{3} \right) \varphi_0.$$

The

ver-

ti-

cal

po-

si-

Source: <https://pho.rs/p/314/s>