

Tidal locking phenomenon — Solution

X23 (2023) — English translation

A1

0.70 pts

Obtain an exact expression for the angular velocity of orbital rotation ω . Express your answer in terms of G, m, M and L .

Let $\vec{e}_1$ be a unit vector drawn towards the smaller body from the center of the larger one. Acceleration of bodies due to gravity:

$$\vec{a}_m = -\frac{GM}{L^2}\vec{e}_1 \quad \vec{a}_M = \frac{Gm}{L^2}\vec{e}_1; \quad \vec{a}_M - \vec{a}_m = \frac{G(M+m)}{L^3}\vec{e}_1.$$

On the other hand:

$$\vec{a}_M - \vec{a}_m = \omega^2 L \vec{e}_1,$$

from where finally:

Answer:

$$\omega = \sqrt{\frac{G(M+m)}{L^3}}$$

A2

0.20 pts

Obtain an exact expression for the potential difference of the gravitational field of a small body $\Delta\varphi_{gr} = \varphi_{grP} - \varphi_{grO}$ at points P and O . Express your answer using G, m, L, θ and $r(\theta)$.

The field potential of a small body at a point whose distance from it is L_m is equal to:

$$\varphi_{gr} = -\frac{Gm}{L_m}$$

Find distance from small body to point A by theorem cosine:

$$L_m^2 = L^2 + r^2(\theta) - 2Lr(\theta)\cos\theta$$

Finally we have:

Answer:

$$\Delta\varphi_{gr} = Gm \left(\frac{1}{L} - \frac{1}{\sqrt{L^2 + r^2 - 2rL\cos\theta}} \right)$$

A3

0.50 pts

Obtain an exact expression for the potential difference of inertial forces $\Delta\varphi_i = \varphi_{iP} - \varphi_{iO}$ at points P and O . Express your answer using G, m, M, L, θ and $r(\theta)$.

Since the fluid is stationary, and the larger body rotates with a constant angular velocity, the fields of inertial forces $-\Delta m \vec{a}_0$ and $\Delta m \omega^2 \vec{r}_\perp$, acting on a particle of mass Δm , contribute to the potential difference. For the potential difference between points A and O we have:

$$\varphi_{iA} - \varphi_{iO} = - \int_O^A (\omega^2 \vec{r}_\perp - \vec{a}_0) \cdot d\vec{r}$$

Taking into account that $\vec{a}_0 = Gm\vec{e}_1/L^2$, and $\vec{r}_\perp = \vec{r}$:

$$\varphi_{iA} - \varphi_{iO} = - \int_O^A (\omega^2 \vec{r} - \frac{Gm}{L^2} \vec{e}_1) \cdot d\vec{r} = \frac{Gmr \cos \theta}{L^2} - \frac{\omega^2 r^2}{2}$$

Finally:

Answer:

$$\Delta\varphi_i = \frac{Gmr \cos \theta}{L^2} - \frac{G(M+m)r^2}{2L^3}$$

A4

1.30 pts

Get addicted $h(\theta)$. Express your answer in terms of m , M , R , L and θ . Keep your answer as simple as possible. Qualitatively depict the shape of the surface in the section under consideration. In the same figure, depict the unperturbed shape of the surface. Note: Use the following approximation:

$$\frac{1}{\sqrt{1+a^2-2a\cos\theta}} \approx 1 + a\cos\theta + \frac{a^2(3\cos^2\theta-1)}{2}$$

Note: Use the following approximation:

$$\frac{1}{\sqrt{1+a^2-2a\cos\theta}} \approx 1 + a\cos\theta + \frac{a^2(3\cos^2\theta-1)}{2}$$

The surface of the larger body at equilibrium is equipotential. Since the surface disturbances are small, we will assume that the potential of the gravitational field of the larger body on the surface is equal to:

$$\varphi_M = -\frac{GM}{r(\theta)}$$

Since surface is equipotential:

$$\varphi_A = Gm \left(\frac{1}{L} - \frac{1}{\sqrt{L^2 + r^2 - 2rL\cos\theta}} \right) + \frac{Gmr \cos \theta}{L^2} - \frac{G(M+m)r^2}{2L^3} - \frac{GM}{r(\theta)} + C_1 = C_2$$

where C_1 and C_2 - some constant quantity. Expand expression under root:

$$\frac{1}{\sqrt{L^2 + r^2 - 2rL\cos\theta}} \approx \frac{1}{L} \left(1 + a\cos\theta + \frac{a^2(3\cos^2\theta-1)}{2} \right); \quad a = \frac{r}{L}.$$

Thus, have:

$$-\frac{GM}{r} - \frac{G(M+3m\cos^2\theta)r^2}{2L^3} = \text{const}$$

Note, that term, proportional $\cos\theta$ completely cancelled. Frodue to this and it was required expand potential small body to second order. Further substitute $r = R + h$ and expand to first order potential larger body. in potential small body one can immediately consider/count $r = R$, since it itself be small quantity frodue to condition $R \ll L$. Obtain

$$-\frac{GM}{R} + \frac{GMh}{R^2} - \frac{G(M+3m\cos^2\theta)R^2}{2L^3} = \text{const.}$$


Keep only variable term:

$$h \frac{GM}{R^2} - \frac{3GmR^2 \cos^2 \theta}{2L^3} = \text{const}$$

Since $h(\pi/2) = 0$, the last constant is zero. Then we have:

Answer:

$$h(\theta) = \frac{3mR^4 \cos^2 \theta}{2Md^3}$$



B1

0.50 pts

Show that the tangential component of the force $F_\tau(t, \theta)$ acting on a particle of mass Δm located on the surface of a larger body at an angle θ can be represented as:

$$F_\tau(t, \theta) = \Delta m \alpha \sin(2\omega_1 t - 2\theta)$$

Find α . Answer express through G , m , L and $r(\theta)$. Note: use result, obtain for solution part A4.

Note: Use the results obtained when solving item A4.

In point A4 we obtained an approximate expression for the potential of inertial forces and the

gravitational field of a smaller body:

$$\varphi_{\text{ext}} = -\frac{G(M + 3m \cos^2 \theta)r^2}{2L^3}$$

Then for component force F_τ obtain:

$$F_\tau = -\frac{\Delta m}{r} \frac{\partial \varphi_{\text{ext}}}{\partial \theta} = -\frac{3Gm\Delta m r \sin 2\theta}{2L^3}$$

Note, that this component force arise from inhomogeneity gravitational field small body. Since $\omega \neq \Omega$, in arbitrary moment time t parameter θ equal:

$$\theta = \theta_0 - \omega_1 t, \quad \omega_1 = \omega - \Omega.$$

Also account for, that with required accuracy $r \approx R$. Then finally have:

$$F_\tau(t, \theta_0) = \frac{3Gm\Delta m R}{2L^3} \sin(2\omega_1 t - 2\theta_0)$$

Answer:

$$\alpha = \frac{3GmR}{2L^3}.$$

B2

0.80 pts

Show that the dependence $\theta(t, \theta_0)$ has the following form:

$$\theta(t) = \theta_0 + A \sin(2\omega_1 t - 2\theta_0 - \varphi_0)$$

Find A and φ_0 . Answer express through $G, m, L, \gamma, \omega_0, \omega, \Omega$ and θ_0 .

Let's substitute $F_\tau(t, \theta)$ into the equation of motion:

$$\Delta m r^2 (\ddot{\theta} + 2\gamma \dot{\theta} + \omega_0^2 (\theta - \theta_0)) = r F_\tau(t, \theta) = \Delta m r^2 \cdot \frac{3Gm}{2L^3} \sin(2\omega_1 t - 2\theta)$$

Since $\omega_0^2 \gg Gm/L^3$ - one can consider/count, that $F_\tau(t, \theta) \approx F_\tau(t, \theta_0)$. Then:

$$\ddot{\theta} + 2\gamma \dot{\theta} + \omega_0^2 (\theta - \theta_0) = \frac{3Gm}{2L^3} \sin(2\omega_1 t - 2\theta_0)$$

Solution equation be seek in form $\theta(t) = \theta_0 + \text{Re}(\hat{A} e^{2i(\omega - \Omega)t})$:

$$(\omega_0^2 - 4\omega_1^2 + 4i\omega_1\gamma) \hat{A} = \frac{3Gm}{2L^3} e^{-i(2\theta_0 + \pi/2)}$$

We can represent $\hat{A}$ in trigonometric form $\hat{A} = A e^{-i\varphi_0}$, where:

Answer:

$$A = \frac{3Gm}{2L^3 \sqrt{(\omega_0^2 - 4\omega_1^2)^2 + 16\gamma^2 \omega_1^2}}; \quad \varphi_0 = \arctan \frac{4\omega_1\gamma}{\omega_0^2 - 4\omega_1^2}.$$

B3

0.40 pts

Obtain an expression for the rate of growth of height $\dot{h}'(t, \theta_0)$ at time t at angle θ_0 . Express your answer using h_0 and $\frac{d\theta}{d\theta_0}$

The volume of liquid flowing through one of the faces hdz during time dt is equal to the product of

the flow velocity $\dot{\theta}R$ by the area hdz and the considered time dt :

$$dV_1 = \dot{\theta}Rh dz dt \approx \dot{\theta}Rh_0 dz dt.$$

Here we account for, that height surface water change small, and this practically not affects on flux/flow water. Velocity flow, correspond angle θ_0 and $\theta_0 + d\theta_0$ somewhat differ, therefore difference inflowing and follow from volume water

$$dV = \dot{\theta}(\theta_0)Rh_0 dz dt - \dot{\theta}(\theta_0 + d\theta_0)Rh_0 dz dt = -Rh_0 \frac{\partial \dot{\theta}}{\partial \theta_0} d\theta_0 dz dt.$$

Produe to this change height surface liquid:

$$dV = Rd\theta_0 dz dh = Rd\theta_0 dz \dot{h} dt = -Rh_0 \frac{\partial \dot{\theta}}{\partial \theta_0} d\theta_0 dz.$$

Hence $\dot{h} = -h_0 \frac{\partial \dot{\theta}}{\partial \theta_0}$, and since $h_0 = \text{const}$

Answer:

$$\dot{h}'(t, \theta) = -h_0 \frac{\partial \dot{\theta}(t, \theta)}{\partial \theta_0}.$$

B4

0.30 pts

Assuming that the amplitude of oscillations h' is the same for all values of θ_0 , you obtain the dependence $h(t)$. Express your answer using h_0 , A , ω_1 , φ_1 and t .

From the results of point B2 we get:

$$\theta = \theta_0 + A \sin(2\omega_1 t - \varphi_1), \quad \dot{\theta} = 2\omega_1 A \cos(2\omega_1 t - \varphi_1).$$

Then find

$$\frac{\partial \dot{\theta}(t, \theta_0)}{\partial \theta_0} = 4\omega_1 A \sin(2\omega_1 t - \varphi_1),$$

from which:

$$\dot{h}' = -4h_0\omega_1 A \sin(2\omega_1 t - \varphi_1)$$

Integrating, we find:

Answer:

$$h(t) = h_0(1 + 2A \cos(2\omega_1 t - \varphi_1))$$

B5

0.40 pts

For the moment of time t , determine the values of the angles θ_0 corresponding to the maximum value $h(t, \theta_0)$. Express your answers using ω_1 , t and φ_0 .

At the maximum points, the cosine is equal to one, therefore:

$$2\omega_1 t - \varphi_1 = 0; 2\pi.$$

Substitute φ_1 , we find:

Answer:

$$\theta_{1,2} = \omega_1 t - \frac{\varphi_0}{2}; \pi + \omega_1 t - \frac{\varphi_0}{2}.$$

Find the moment of force M_z acting from the small body on the surface of the large body relative to the axis z . Express your answer using G , m , ρ , R , h_0 , L , A and φ_0 .

The force arm corresponding to the angle β is equal to $R \sin \beta$. Then for the moment element dM_z we have:

$$dM_z = R \sin \beta dF_\tau(t, \theta, \beta)$$

For dF_τ also have:

$$dF_\tau = \frac{3GmR \sin \beta}{2L^3} \Delta m \sin(2\omega_1 t - 2\theta).$$

Mass consider volume

$$\Delta m = \rho dV = \rho h(t, \theta, \beta) dS = \rho h(t, \theta, \beta) R^2 \sin \beta d\beta d\theta.$$

Then obtain:

$$dF_\tau = \frac{3GmR \sin \beta}{2L^3} \sin(2\omega_1 t - 2\theta) \rho h(t, \theta, \beta) R^2 \sin \beta d\beta d\theta.$$

For $h(t, \theta, \beta)$ have:

$$h(t, \theta, \beta) = h_0(1 + 2A \cos(2\omega_1 t - \varphi_1))$$

since A not depend from distance to axis z . We can represent expression for M_z :

$$M_z = \frac{3Gm\rho R^4 h_0}{2L^3} \int_0^\pi \sin^3 \beta d\beta \int_0^{2\pi} (1 + 2A \cos(2\omega_1 t - \varphi_1)) \sin(2\omega_1 t - 2\theta) d\theta$$

For the integral by β we have:

$$\int_0^\pi \sin^3 \beta d\beta = \int_{-1}^1 (1 - \cos^2 \beta) d \cos \beta = \cos \beta - \frac{\cos^3 \beta}{3} \Big|_{-1}^1 = \frac{4}{3}$$

Therefore:

$$M_z = \frac{2Gm\rho R^4 h_0}{L^3} \int_0^{2\pi} (1 + 2A \cos(2\omega_1 t - \varphi_1)) \sin(2\omega_1 t - 2\theta) d\theta$$

The second integral is taken over period, therefore for the function $\sin(2\omega_1 t - 2\theta)$ it equals zero. Let us pass to the last integral:

$$\int_0^{2\pi} \cos(2\omega_1 t - \varphi_1) \sin(2\omega_1 t - 2\theta) d\theta = \int_0^{2\pi} \cos(2\omega_1 t - 2\theta - \varphi_0) \sin(2\omega_1 t - 2\theta) d\theta$$

We will use trigonometric formula:

$$\sin \alpha \cos \beta = \frac{\sin(\alpha + \beta) + \sin(\alpha - \beta)}{2}$$

We obtain:

$$M_z = \frac{2Gm\rho R^4 h_0 A}{L^3} \int_0^{2\pi} [\sin(4\omega_1 t - 4\theta - \varphi_0) + \sin \varphi_0] d\theta$$

Final:

Answer:

$$M_z = \frac{4\pi Gm\rho R^4 h_0 A \sin \varphi_0}{L^3}$$

C1

0.80 pts

Find the orbital angular velocity ω_{sync} and the distance L_{sync} between the Earth and the Moon during synchronous rotation. Express the answers in terms of $m, M, R, G, L_0, \Omega_0$ and find their numerical values. Note: with synchronous rotation, the angular momentum of the Earth associated with rotation around its axis can be neglected.

Note: with synchronous rotation, the angular momentum of the Earth associated with rotation around its axis can be neglected.

Let us use the law of conservation of angular momentum, which we denote as K :

$$K = I\Omega_0 + \frac{mML_0^2\omega_0}{M+m} = \text{const}$$

where $I = 2MR^2/5$ - moment of inertia Earth's sphere about the diameter. In the established regime $\Omega = \omega = \omega_{\text{sync}}$, therefore:

$$K = \omega_{\text{sync}} \left(I + \frac{mML_{\text{sync}}^2}{m+M} \right)$$

Since for synchronous rotation Earth can be considered as a material point:

$$K \approx \frac{mML_{\text{sync}}^2\omega_{\text{sync}}}{m+M}$$

Use relation

$$\omega^2 = \frac{G(M+m)}{L^3}$$

obtain:

$$K \approx \frac{G^{2/3}Mm}{(m+M)^{1/3}\omega_{\text{sync}}^{1/3}}$$

from where finally:

Answer:

$$\omega_{\text{sync}} \approx \frac{G^2 m^3 M^3}{(M+m) \left(\frac{2MR^2\Omega_0}{5} + \frac{mM}{m+M} \sqrt{G(M+m)L_0} \right)^3} \approx 1,376 \cdot 10^{-6} \text{ s}^{-1} L_{\text{sync}} = \frac{(m+M) \left(\frac{2MR^2\Omega_0}{5} + \frac{mM}{m+M} \sqrt{G} \right)}{Gm^2M^2}$$

C2

0.70 pts

The distance between the Earth and the Moon is increasing at a rate of $\dot{L}_0 = 1 \text{ cm/year}$. Find the value of the average angular acceleration of the Earth $\dot{\Omega}_0$. Express the answer in terms of $m, M, R, G, L_0, \Omega_0, \dot{L}_0$ and find its numerical value.

The angular momentum of the system K using the connection ω and L is reduced to the form:

$$K = I\Omega + \frac{mML^2}{m+M} \sqrt{\frac{G(m+M)}{L^3}}$$

Differentiate, find:

$$I\dot{\Omega} + \frac{mM}{2(m+M)} \sqrt{\frac{G(m+M)}{L}} \dot{L} = 0$$

from where:

Answer:

$$\dot{\Omega}_0 = -\frac{5m\dot{L}_0}{4(m+M)R^2} \sqrt{\frac{G(m+M)}{L_0}} \approx -1,22 \cdot 10^{-22} \text{ s}^{-1}$$

C3

0.80 pts

Find the numerical values of ω_0 and γ .

At point B5 it was found that the position of the tide lags behind the Moon by an angle $\varphi_0/2$ in the direction of rotation of the Moon relative to the Earth. Then we have:

$$2\beta = \arctan \frac{-4\omega_1\gamma}{\omega_0^2 - 4\omega_1^2} = -\varphi_0$$

Also for difference height of the tide and ebb we have:

$$\Delta h = h_{\text{high}} - h_{\text{low}} = 4Ah_0$$

or, after substitution expression for A :

$$\Delta h = \frac{6Gmh_0}{L_0^3 \sqrt{(\omega_0^2 - 4\omega_1^2)^2 + 16\omega_1^2\gamma^2}} = \frac{6Gmh_0 \cos 2\beta}{L_0^3 (\omega_0^2 - 4\omega_1^2)}$$

We find:

Answer:

$$\omega_0 = \sqrt{4\omega_1^2 + \frac{6Gmh_0 \cos 2\beta}{L_0^3 \Delta h}} \approx 1,664 \cdot 10^{-4} \text{ s}^{-1} \quad \gamma = \frac{3Gmh_0 \sin 2\beta}{2(-\omega_1)L_0^3 \Delta h} \approx 3,025 \cdot 10^{-6} \text{ s}^{-1}$$

C4

0.50 pts

Within the framework of the described model, find the angular acceleration of the Earth $\dot{\Omega}_{0(\text{mod})}$, which is obtained from the results of point B6. Compare it with the value $\dot{\Omega}_0$ obtained in point C2 and draw a conclusion about the applicability of the model under consideration (consider the model applicable if Ω_0 and $\Omega_{0(\text{mod})}$ differ by no more than 10 times).

From the equation of the dynamics of rotational motion relative to the center of mass we obtain:

$$I\dot{\Omega} = M_z = \frac{4\pi Gm\rho R^4 h_0 A \sin \varphi_0}{L^3}$$

Further, note that $h_0 A = \Delta h/4$, and $\varphi_0 = -2\theta$. Then we have:

Answer:

$$\dot{\Omega}_{0(\text{mod})} = -\frac{5\pi Gm\rho R^2 \Delta h \sin 2\theta}{2ML_0^3} = -1,16 \cdot 10^{-22} \text{ s}^{-1}$$

The model under consideration gives a value of the required order and is applicable.

C5

0.70 pts

Estimate the numerical value of τ_2 . Express your answer in years.

After switching to the synchronous rotation mode, we can assume that:

$$\omega = \text{const} \quad L = \text{const} \quad \omega_0^2 \gg \omega_1^2$$

Therefore:

$$I\dot{\Omega} = M_z = \frac{4\pi Gm\rho R^4 h_0 A \sin \varphi_0}{L^3} \approx \frac{4\pi Gm\rho R^4 h_0}{L_{\text{sync}}^3} \cdot \frac{3Gm}{2L_{\text{sync}}^3 \omega_0^2} \cdot \frac{4(\omega_{\text{sync}} - \Omega)\gamma}{\omega_0^2}$$

from which:

$$d\tau_2 = -\frac{d\Omega}{\Omega - \omega_{\text{sync}}} \cdot \frac{ML_{\text{sync}}^6 \omega_0^4}{120\pi G^2 m^2 \rho R^2 h_0 \gamma}$$

from where:

Answer:

$$\tau_2 \approx \frac{ML_{\text{sync}}^6 \omega_0^4}{120\pi G^2 m^2 \rho R^2 h_0 \gamma} \approx 3,14 \cdot 10^{12} \text{ year}$$