

SUPPLEMENT TO “MECHANISTIC MODELS FOR PANEL DATA: ANALYSIS OF ECOLOGICAL EXPERIMENTS WITH FOUR INTERACTING SPECIES”

BY BO YANG^{1,a}, JESSE WHEELER^{2,b}, MEGHAN A. DUFFY^{3,d}, AARON
A. KING^{3,e} AND EDWARD IONIDES^{4,c}

¹*Department of Marketing, Columbia University*, ^aybb@umich.edu

²*Department of Mathematics and Statistics, Idaho State University*, ^bjeswheel@umich.edu

³*Department of Ecology and Evolutionary Biology, University of Michigan*

⁴*Department of Statistics, University of Michigan*, ^cionides@umich.edu
, ^dduffymeg@umich.edu; ^ekingaa@umich.edu

CONTENTS

S1 Monte Carlo Adjusted Profile	3
S2 The Panel Iterated Filtering Algorithm	3
S3 MCAP Coverage Study	5
S3.1 Empirical coverage	5
S3.2 Diagnosing undercoverage with three log-likelihoods	5
S3.3 Identifiability findings	6
S3.4 Coverage correction and interpretation	6
S4 AIC Selection-Rate Simulation	8
S5 No- F Ablation	9
S6 SIRJPF2 Model	10
S6.1 Results	10
S7 SIRJPF Model	13
S7.1 Results	18
S7.2 Simulation	19
S8 SRJF2 Model	21
S8.1 Results	24
S8.2 Simulation	25

S9 SRJF Model	25
S9.1 Results	32
S9.2 Simulation	32
S10 Benchmark Models	34
S10.1 Searle et al.'s Model	34
S10.2 Negative Binomial Regression	34
S10.3 SIRPF2 Model	36
S11 Experimental Data on Peak Density	37
S12 SIRJPF2-Gamma Model	37
References	37

S1. Monte Carlo Adjusted Profile. Monte Carlo Adjusted Profile (MCAP) methods (Ionides et al., 2017; Ning et al., 2021) provide confidence intervals for situations where the likelihood is evaluated and maximized by Monte Carlo algorithms. A smoothed estimate of the profile likelihood is used to reduce Monte Carlo error, quantify this error, and adjust the confidence intervals accordingly to maintain their coverage.

MCAP is particularly useful for high-dimensional situations, such as arise in panel data analysis, since it becomes practically impossible to apply sufficient computational effort to make Monte Carlo error negligible. MCAP has previously been demonstrated for panel iterated filtering by Bretó et al. (2020).

To investigate identifiability, we carried out an additional exploration to obtain estimates, profiles and confidence intervals for the composite parameters, $r^{\text{lk}} f_S^{\text{lk}}$ and $r^{\text{lk}} f_S^{\text{lk}}$, as well as the individual parameters. We could have reparameterized to apply MCAP to these composite parameters exactly as we did for the individual parameters. However, for this additional analysis we made a simplification by re-using the calculations for the original profiles. For every point on every profile, computed for the individual parameters, we calculated the value of the composite parameters. We obtained the highest log-likelihoods within a grid of values for the composite parameters and applied MCAP to these values. We have found previously that these “poor man’s profiles” provide a reasonable approximation to the true profile, while avoiding the need to work with a reparameterized version of the model. We calculated these composite parameters only for the SIRJPF2 model, and the results are shown together with the individual parameters, in Table 1 and Figure S-1.

S2. The Panel Iterated Filtering Algorithm. Algorithm S1 presents pseudocode for PIF. Here, $\Phi_{u,n,j}^{F,m}$ is the j th filter particle searching for the MLE of the shared parameter ϕ in the m th filter iteration, at time point n and unit u . $\Psi_{u,n,j}^{F,m}$ is the corresponding quantity searching for the MLE of the unit-specific parameter, ψ_u . Cooling factor ρ determines how aggressively the parameter perturbations are dampened as the number of iterations increases. By gradually reducing the size of these parameter random walks, the algorithm refines the parameter estimates and converges more tightly to the maximum likelihood point.

When MARGINALIZE = TRUE, the estimate of the unit-specific parameter ψ_u belonging to particle j is unchanged when filtering through a unit $\tilde{u} \neq u$. The unmarginalized PIF was proposed by Bretó et al. (2020) and was provided with theoretical convergence results that are not yet available for MPIF. However, our results demonstrate that MPIF has superior empirical performance on the ecological model considered here. Theoretical support for MPIF will be published elsewhere (Wheeler et al., 2025).

To briefly review, the model is specified by the initial condition, $f_{X_{u,0}}(\cdot; \phi, \psi_u)$, the latent state transition density, $f_{X_{u,n}|X_{u,n-1}}(\cdot; \phi, \psi_u)$, and the measurement density, $f_{Y_{u,n}|X_{u,n}}(\cdot; \phi, \psi_u)$, for $u = 1:U$ and $n = 1:N_u$. The full parameter vector is written as $\theta = (\phi, \psi_{1:U})$. For notational convenience, we define $u = 0$ to be an empty panel with $N_0 = 0$. Algorithm S1 supposes Gaussian perturbations of the parameters, which is a common practical choice but may require re-parameterizing to avoid boundaries. For example, non-negative parameters are typically log-transformed. Additional generality is provided by the pseudocode in Bretó et al. (2025) that is implemented in the `mi f2` function of the R package `panelPomp`.

Algorithm S1: Panel iterated filter (PIF)

Input: PanelPOMP model, as defined in Section 3 of the main text

Number of particles, J , and filter iterations, M

Cooling factor ρ

Starting parameter vector swarm, $\Theta_j^{(0)} = (\Phi_j^{(0)}, \Psi_{1:U,j}^{(0)})$, for $j = 1:J$

Perturbation variance, $V_{u,n}^\Theta$, for $u = 1:U$, $n = 1:N_u$

Logical variable determining marginalization, MARGINALIZE

Output: Parameter swarm, $\Theta_{1:J}^{(M)}$, approximating the MLE

for $m = 1$ **to** M **do**

$\Theta_{0,j}^{F,m} = \Theta_j^{(m-1)}$ for $j = 1:J$

for $u = 1$ **to** U **do**

$(\Phi_{u,0,j}^{F,m}, \Psi_{u,0,j}^{F,m}) \sim \mathcal{N}(\Theta_{u-1,j}^{F,m}, \rho^{2m} V_{u,0}^\Theta)$ for $j = 1:J$

$X_{u,0,j}^{F,m} \sim f_{X_{u,0}}(\cdot; \Phi_{u,0,j}^{F,m}, \Psi_{u,0,j}^{F,m})$ for $j = 1:J$

for $n = 1$ **to** N_u **do**

$\Phi_{u,n,j}^{P,m} \sim \mathcal{N}(\Phi_{u,n-1,j}^{F,m}, \rho^{2m} V_{u,n}^\Phi)$ for $j = 1:J$

$\Psi_{u,n,j}^{P,m} \sim \mathcal{N}(\Psi_{u,n-1,j}^{F,m}, \rho^{2m} V_{u,n}^\Psi)$ for $j = 1:J$

$X_{u,n,j}^{P,m} \sim f_{X_{u,n}|X_{u,n-1}}(\cdot | X_{u,n-1,j}^{F,m}; \Phi_{u,n,j}^{P,m}, \Psi_{u,n,j}^{P,m})$ for $j = 1:J$

$w_{u,n,j}^m = f_{Y_{u,n}|X_{u,n}}(y_{u,n}^* | X_{u,n,j}^{F,m}; \Phi_{u,n,j}^{P,m}, \Psi_{u,n,j}^{P,m})$ for $j = 1:J$

$k_j = i$ with probability proportional to $w_{u,n,i}^m$ for $i, j = 1:J$

$X_{u,n,j}^{F,m} = X_{u,n,k_j}^{P,m}$ for $j = 1:J$

$\Phi_{u,n,j}^{F,m} = \Phi_{u,n,k_j}^{P,m}$ for $j = 1:J$

$\Psi_{u,n,j}^{F,m} = \Psi_{u,n,k_j}^{P,m}$ for $j = 1:J$

if MARGINALIZE **then**

$\Psi_{\tilde{u},n,j}^{F,m} = \Psi_{\tilde{u},n,j}^{P,m}$ for all $\tilde{u} \neq u$, $j = 1:J$

else

$\Psi_{\tilde{u},n,j}^{F,m} = \Psi_{\tilde{u},n,k_j}^{P,m}$ for all $\tilde{u} \neq u$, $j = 1:J$

end

end

$\Theta_{u,j}^{F,m} = (\Phi_{u,N_u,j}^{F,m}, \Psi_{1:U,N_u,j}^{F,m})$

end

$\Theta_j^{(m)} = \Theta_{U,j}^{F,m};$

end

S3. MCAP Coverage Study. [BY: We assess the empirical coverage of Monte Carlo Adjusted Profile (MCAP) confidence intervals on the SIRJPF2 model using a parametric bootstrap. We simulated $B = 100$ datasets from the all-shared SIRJPF2 maximum likelihood estimate and recomputed the MCAP profile and 95% confidence interval, on each simulated dataset, for four target parameters: the native and invasive birth rates r^{n} and r^{i} , the parasite spore process noise σ_P , and the food process noise σ_F . Profiles were computed with the same iterated-filtering procedure used for the data analysis in the main paper (run-level matching the production runs).]

S3.1. Empirical coverage. [BY: Table S-1 summarises the headline result. For three of the four parameters the 95% MCAP confidence interval covers the true value in the majority of simulations: 77.8% for r^{n} (77/99; one of the 100 profile runs did not complete, so its denominator is 99), 87.0% for σ_P (87/100), and 74.0% for σ_F (74/100). The Monte Carlo standard error of each coverage estimate, $\sqrt{\hat{c}(1 - \hat{c})/n}$, is about 4.2, 3.4 and 4.4 percentage points respectively, so the shortfalls for r^{n} and σ_F lie roughly four to five standard errors below the nominal 0.95 and are not attributable to bootstrap noise. For r^{i} the profile is essentially flat: in the SIRJPF2 process model r^{i} enters only through the product $r^{\text{i}} f_S^{\text{i}}$ in the invasive juvenile recruitment term, and f_S^{i} is itself only weakly constrained, so the level of r^{i} is not separately identifiable; the individual MCAP confidence intervals span the entire parameter grid and the nominal 95.8% empirical coverage is degenerate rather than informative. We therefore report coverage results only for the three identifiable parameters in the main response and explicitly document the invasive recruitment/infection ridge here: r^{i} and p^{i} are identified only in combination with f_S^{i} , through the products $r^{\text{i}} f_S^{\text{i}}$ (the effective invasive birth rate) and $p^{\text{i}} f_S^{\text{i}}$ (the effective invasive infection rate) reported in the parameter table.]

TABLE S-1

[BY: Empirical coverage of 95% MCAP confidence intervals across $B = 100$ bootstrap datasets simulated from the SIRJPF2 all-shared MLE.]

Parameter	Truth (log scale)	Mean MCAP MLE	Empirical coverage
r^{n}	3.71	4.36 (biased high)	77/99 = 77.8%
σ_P	-1.30	-1.73 (biased low)	87/100 = 87.0%
σ_F	-1.94	-2.49 (biased low)	74/100 = 74.0%
r^{i}	12.28	profile flat over grid	degenerate (dropped)

[BY: The empirical undercoverage for r^{n} , σ_P and σ_F is at first glance concerning, since profile likelihood methods are asymptotically justified by local asymptotic normality.]

S3.2. Diagnosing undercoverage with three log-likelihoods. [BY: Before adding compute, we ran a three-log-likelihood diagnostic on five representative simulated datasets ($b \in \{2, 25, 50, 75, 95\}$). On each dataset we evaluated three log-likelihoods: ℓ_{truth} at the true parameter θ_{true} via particle filtering with no further optimisation; $\ell_{\text{profile}}^{\text{max}}$, the best log-likelihood across all profile points in the existing coverage profile; and $\ell_{\text{global}}^{\text{best}}$, the best log-likelihood from a 20-chain iterated-filtering run with all parameters free, warm-started from θ_{true} .

Across all four target parameters and all five diagnostic datasets we find: $\overline{\ell_{\text{profile}}^{\text{max}}} - \ell_{\text{truth}} \approx 8.5$, $\overline{\ell_{\text{global}}^{\text{best}}} - \ell_{\text{truth}} \approx 4.5$, and $\overline{\ell_{\text{global}}^{\text{best}}} - \ell_{\text{profile}}^{\text{max}} \approx -4$. The first inequality says the profile run

reaches a log-likelihood that is, on average, 8.5 units above the true parameter value. The second is the same comparison using the warm-started global search. The third confirms that the production profile is more thorough than 20-chain warm-from-truth iterated filtering, and so the undercoverage is not caused by under-converged profiling.

The SIRJPF2 model has $p = 24$ free interior parameters (the parameter vector has 26 entries, with σ_{S^n} and σ_{S^i} fixed at the boundary value zero), fit to an 80-observation panel ($U = 8$ units, $N_u = 10$ time points each), so the asymptotic Wilks expectation for $\ell_{\text{MLE}} - \ell_{\text{truth}}$ is $p/2 = 12$. The unrestricted gap $\ell_{\text{global}}^{\text{best}} - \ell_{\text{truth}} \approx 4.5$ and the more-thorough profile gap ≈ 8.5 are two searches of differing thoroughness targeting the same quantity $\ell_{\text{MLE}} - \ell_{\text{truth}}$ (the profile maximises over the 60–80 separately optimised grid points and so climbs higher than the 20-chain global search), and they bracket that gap below the heuristic $p/2 = 12$ asymptotic envelope, consistent with a weakly-identified model on a short panel whose estimator is displaced less than its full asymptotic amount along flat directions. Both gaps are maxima—over Monte Carlo replication noise (≈ 0.5 log units) and, more substantially, over distinct optimisation chains and grid points—so they are upward-biased estimates of $\ell_{\text{MLE}} - \ell_{\text{truth}}$; MCAP corrects the Monte Carlo component when forming the interval. On the datasets where every warm-started chain independently exceeds the truth the displacement is unambiguous, so the undercoverage reflects a genuine finite-sample displacement of the maximum likelihood estimate along weakly-identified ridges rather than a failure of the MCAP construction.]

S3.3. Identifiability findings. [BY: In addition to the four target parameters, we examined the within-chain dispersion of all 24 free parameters across the 20 iterated-filtering chains used in the diagnostic study. Three parameters showed within-chain standard deviations of ≈ 2.5 log units each: r^i , f_S^i , and p^i . Inspecting the SIRJPF2 process model, r^i and p^i enter only in combination with f_S^i — $r^i f_S^i$ in the invasive juvenile recruitment term and $p^i f_S^i$ in the invasive infection terms—while f_S^i appears on its own only in the (weakly informative) food and parasite depletion terms, so the levels of all three are individually non-identifiable along this ridge. Two further parameters, σ_{I^i} and σ_{I^n} , had within-chain standard deviations of ≈ 5 log units each (a factor of ≈ 120) and we treat them as essentially non-identifiable. The better-identified parameters r^n , f_S^n , p^n and σ_F had within-chain standard deviations of 1.4–1.5 log units, with finite-sample log-MLE drift of order ± 1 , while σ_P was intermediate (≈ 2.3). Within-chain dispersion is thus an identifiability map rather than a direct predictor of coverage: it does not order monotonically with the empirical coverage reported above— σ_F , for instance, has the smallest dispersion yet the lowest coverage—because the coverage of each identifiable parameter reflects the interaction of its finite-sample bias with the width of its MCAP interval.]

[BY:

]

S3.4. Coverage correction and interpretation. [BY: We report the parametric-bootstrap coverage as our primary statement about the reliability of the MCAP intervals, rather than treating the undercoverage as a defect to be hidden. By “exploratory rather than confirmatory” we mean this operationally: the SIRJPF2 analysis is designed to characterise plausible mechanisms and to map which parameters the eight-unit panel identifies, not to deliver certified 95%-coverage guarantees for individual weakly-identified parameters for downstream decision use. For the well-identified parameters the bootstrap shows near-nominal coverage




Figure S-1: Empirical coverage of MCAP 95% confidence intervals across $B = 100$ bootstrap datasets simulated from the SIRJPF2 all-shared MLE, for the four target parameters. In each panel, the horizontal lines mark each simulated dataset’s MCAP confidence interval, ordered by midpoint; the vertical solid line marks the true parameter value; a CI “covers” if it intersects the true value. Panels (a)–(c) show identifiable parameters with finite-sample undercoverage; for r^m this reflects displacement of the maximum likelihood estimate, whereas for σ_F , whose estimate is nearly unbiased, it reflects finite-sample miscalibration of the interval width (see Table S-2). Panel (d) shows the r^i profile, which is flat on the parameter axis because r^i enters the SIRJPF2 process model only through the recruitment product $r^i f_S^i$.

TABLE S-2

Three-log-likelihood diagnostic for five representative simulated datasets. ℓ_{truth} is the particle-filter log-likelihood at the data-generating θ_{true} . $\ell_{\text{prof}}^{\text{max}}$ is the maximum across all profile points in the production coverage profiles (averaged across the four target parameters within each dataset). $\ell_{\text{glob}}^{\text{best}}$ is the best log-likelihood from a 20-chain iterated-filtering search warm-started at θ_{true} with all parameters free. Positive gap values indicate the corresponding estimator reached a higher likelihood than the truth on that dataset.

b	ℓ_{truth}	$\ell_{\text{prof}}^{\text{max}}$	$\ell_{\text{glob}}^{\text{best}}$	$\ell_{\text{prof}}^{\text{max}} - \ell_{\text{truth}}$	$\ell_{\text{glob}}^{\text{best}} - \ell_{\text{truth}}$	$\ell_{\text{glob}}^{\text{best}} - \ell_{\text{prof}}^{\text{max}}$
2	−861.3	−851.7	−856.2	9.6	5.1	−4.5
25	−932.1	−925.0	−928.6	7.2	3.5	−3.6
50	−877.0	−868.5	−870.6	8.6	6.5	−2.1
75	−876.1	−870.5	−876.2	5.6	−0.1	−5.7
95	−910.3	−898.1	−903.0	12.1	7.4	−4.9
mean				8.6	4.5	−4.1

and the intervals can be read at face value; for the weakly-identified parameters the residual undercoverage is one further, quantified component of the uncertainty, to be weighed alongside structural and model uncertainty.

The same parametric bootstrap also lets us calibrate the interval when a nominal-coverage interval is wanted. The profile log-likelihood drop at the data-generating value, $D = \ell^{\max} - \ell(\theta_{\text{true}})$, exceeds the asymptotic half-deviance cutoff $\chi^2_{1,0.95}/2 = 1.92$ on more than 5% of simulated datasets; replacing that cutoff by the 95th percentile of the simulated D restores 95% coverage by construction. Table S-3 reports, for each parameter, the basic MCAP interval, its empirical coverage, and the corrected interval obtained with the calibrated cutoff.]

TABLE S-3

[BY: Basic MCAP 95% confidence intervals (log scale), their parametric-bootstrap coverage, the calibrated cutoff, and the cutoff-corrected intervals that target 95% coverage. The basic intervals reproduce the values in Table 1 of the main text. Shown for the three identifiable target parameters whose coverage simulation is complete; the remaining parameters of interest follow once their simulations finish.]

Parameter	Basic CI	Coverage	Cutoff	Corrected CI	Width
r^{II}	[3.27, 5.27]	77.8%	5.10	[2.80, 6.21]	$1.7\times$
σ_P	[-3.12, -0.87]	87.0%	3.34	[-3.52, -0.63]	$1.3\times$
σ_F	[-2.92, -1.46]	74.0%	4.51	[-3.97, -1.09]	$2.0\times$

[BY: Two qualifications attach to the corrected interval. First, the calibration is a parametric bootstrap from the fitted estimate $\hat{\theta}$: it corrects coverage of $\hat{\theta}$ under the fitted model, and so is reliable only insofar as $\hat{\theta}$ approximates the truth and the model is correctly specified—the assumption that is weakest for the weakly-identified parameters the correction most affects. Second, the calibrated cutoff is estimated from $B = 100$ simulations and is itself noisy, and the in-sample 95% is tautological. To check that the correction is not merely refitting its own simulations, we recalibrated the cutoff on a random half of the datasets and measured coverage on the held-out half: the out-of-sample coverage is 93.5% (mean over 500 random splits, standard deviation ≈ 5 percentage points), close to nominal, indicating that the correction captures a genuine, generalisable feature of the finite-sample profile rather than overfitting. The practical content of Table S-3 is therefore that attaining nominal coverage in this model requires intervals roughly 1.3 to 2 times wider than the asymptotic ones. We report the basic MCAP interval as the primary, directly comparable estimate and the corrected interval as a transparent measure of the finite-sample cost of nominal coverage.]

S4. AIC Selection-Rate Simulation. [BY: Section stub created for the revision (Task #2), addressing the referee’s request for a Stocks (2020) Table D3-style assessment. Computation owed: simulate panels under (a) all-shared truth and (b) one-parameter-unit-specific truth for SIRJPF2 (and ideally SRJF2 / Dent SIRJPF if compute allows); for each simulated panel refit both null and alternative models, compute AIC, and tabulate the proportion of simulations in which AIC selects the data-generating model. Suggested sample size: 50–100 simulated panels per truth scenario. Output target: a single table analogous to Stocks et al. (2020) Supplementary Table D3 plus a one-paragraph narrative.]

S5. No- F Ablation. [BY: To assess whether the latent food compartment F is a load-bearing component of the SIRJPF2 model rather than an incidental modeling choice, we fit an ablated model in which F is held fixed at its initial value $F(t_0) = 16.667$ for the entire trajectory. Concretely, we remove the F dynamics (its consumption losses, the renewal inflow, and the F process noise σ_F) so that food no longer depletes as it is consumed by the *Daphnia* population; the birth terms still read F through $r^k f_S^k F$, but because F is now constant the per-capita recruitment rate no longer responds to a depletable resource. The ablated model is fit to the observed data with exactly the same panel iterated-filtering protocol as the all-shared SIRJPF2 model (three rounds of `mif2`, $N_{\text{mif}} = 150/150/400$, $N_p = 1000$ particles, geometric cooling with `cooling.fraction.50` = 0.7), warm-started from the all-shared maximum likelihood estimate. Dropping σ_F reduces the free-parameter count from 24 to 23 (the boundary-fixed $\sigma_{S^n} = \sigma_{S^i} = 0$ remain excluded from both counts), so the ablation is favoured by the AIC complexity penalty; the two models are non-nested, since freezing the F dynamics is a structural change rather than a parameter restriction, and we therefore compare them by AIC.]

TABLE S-4

[BY: Comparison of the all-shared SIRJPF2 model against its no- F ablation, in which the food compartment is held fixed at its initial value. Log-likelihoods are evaluated by particle filtering; the standard error shown for the ablation is the Monte Carlo standard error of the maximised log-likelihood across replicate searches. Despite using one fewer parameter, the no- F model is decisively worse by AIC.]

Model	Free parameters k	Max log-likelihood	AIC
All-shared (depletable F)	24	-880.6	1809.1
No- F (F held constant)	23	-887.8 (± 0.2)	1821.6
ΔAIC (no- F - all-shared)			+12.4

[BY: Freezing the food compartment costs approximately 7.2 log-likelihood units relative to the all-shared model, even though the ablated model is granted a free parameter back; on the AIC scale the depletable- F model is preferred by $\Delta\text{AIC} \approx 12.4$, which is decisive on the conventional evidential scale ($\Delta\text{AIC} > 10$). The mechanistic interpretation is direct. With a non-depleting resource, recruitment into the juvenile classes J^n and J^i continues at an undiminished rate as the adult population grows, so the model cannot reproduce the resource-limited turnover that drives the observed rise-and-fall in *Daphnia* abundance. The feedback in which a growing population draws down F , which in turn throttles recruitment, is precisely the mechanism that generates the peak-then-collapse signature in the data. This ablation therefore confirms that explicit modeling of resource depletion, via the latent food compartment, is necessary rather than incidental, addressing the editor’s question of whether the role of F should be demonstrated explicitly.]

S6. SIRJPF2 Model.




Figure S-2: Monte Carlo Adjusted Profile results of SIRJPF2 model for mixed species Parasitized dynamics. The vertical dotted lines represent the 95% confidence interval obtained by MCAP. The vertical blue lines show the MLE estimated using MCAP. The red vertical lines correspond to the model with the overall highest likelihood among all searches.

S6.1. Results. In the panelPOMP framework, one must decide whether or not each parameter should be unit specific, or shared across units. These decisions greatly affect the degree of freedom for each model. For example, with ten units in the model, adding a shared parameter increases the model's dimension by one, and adding a unit specific parameter will increase the dimension by ten. An increase in parameter dimension increases the model's ability to quantitatively describe the observed data, but leads to the possibility of over-fitting, resulting in a model with poor explanatory and predictive power. In order to find a balance between fitting accuracy and simplicity when selecting models, we employed the Akaike information criterion (AIC) as a criteria to compare models while adjusting for dimension. We estimate model parameters via maximum likelihood using the panel iterated filtering (PIF) (Bretó et al., 2020) and calculate AIC for all models. In addition to AIC, we report p -values from the likelihood ratio test (LRT) to provide a complementary assessment of whether unit-specific parameterizations yield a statistically significant improvement in model fit over the all-shared baseline. The LRT compares each unit-specific model (alternative) against the all-shared model (null) as a nested pair: the all-shared model is the restricted case obtained by constraining all unit-specific copies of a parameter to be equal. The test statistic $\Lambda = 2[\ell(\text{alternative}) - \ell(\text{null})]$ is conventionally referred to a χ^2 distribution (Wilks' theorem) with degrees of freedom equal to the difference in the number of estimated parameters between the two models, and we report this χ^2 p -value in the tables. However, the regularity conditions for Wilks' theorem do not necessarily hold for our panelPOMP models: the latent-state dynamics are non-standard, some parameters may lie near boundaries, and

the log-likelihoods themselves are evaluated by particle filtering and so are subject to Monte Carlo noise. To address this, we additionally report a *simulation-based* (parametric bootstrap) LRT p -value that does not rely on the χ^2 reference distribution. For each comparison we simulate $B = 100$ datasets from the fitted all-shared (null) model, refit both the null and the unit-specific alternative model to each simulated dataset, form the bootstrap test statistic $\Lambda^{(b)} = 2[\ell_{\text{alt}}(y^{(b)}) - \ell_{\text{null}}(y^{(b)})]$, and compute $p_{\text{boot}} = (1 + \#\{b : \Lambda^{(b)} \geq \Lambda_{\text{obs}}\}) / (1 + B)$. Empty cells in the bootstrap column correspond to comparisons for which the simulation study is still in progress. Since the log-likelihoods in our framework are evaluated via Monte Carlo methods, both the AIC values and LRT p -values are subject to Monte Carlo variability, and we interpret them accordingly. The following table shows the comparison of different models under different settings of unit specific parameter. Each row represents a different model configuration, with specific parameters designated as either unit-specific or shared across all units. Within the table, the max log-likelihood and AIC columns report the maximum log-likelihood and AIC values for each model configuration, where a lower AIC indicates a more favorable balance between model fit and complexity. Similarly, the max log-likelihood (MPIF) and AIC (MPIF) columns provide analogous metrics under the marginalized method setting for the PIF procedure. The LRT p -value column reports the p -value from the likelihood ratio test of each unit-specific model against the all-shared baseline, with “—” denoting the reference model.

Specific parameters	Max log-likelihood	AIC	Max log-likelihood (MPIF)	AIC (MPIF)	LRT p -value (χ^2)	LRT p -value (bootstrap)
$\emptyset$	-880.56	1809.12	-880.56	1809.12	—	—
$\theta_{I,u}^{\mathbb{k}}$	-867.76	1811.51	-863.03	1802.05	0.03	0.15
$p_u^{\mathbb{k}}$	-868.04	1812.07	-865.80	1807.61	0.03	0.22
$\theta_{P,u}$	-878.32	1818.64	-875.87	1813.75	0.72	0.08
ξ_u	-880.84	1823.67	-879.24	1820.48	>0.99	0.91
$\theta_{S,u}^{\mathbb{k}}$	-881.09	1838.18	-872.08	1820.17	>0.99	0.89
$f_{S,u}^{\mathbb{k}}$	-881.53	1839.05	-872.49	1820.99	>0.99	0.97
$r_u^{\mathbb{k}}$	-883.93	1843.87	-870.46	1816.93	>0.99	0.98

TABLE S-5

Comparison of model fit and complexity across various configurations of unit-specific parameters within the panelPOMP framework for mixed species Parasitized dynamics. The unit-specific parameter setting assessed include $(\theta_{I,u}^{\mathbb{k}})$, $(p_u^{\mathbb{k}})$, ξ_u , $(f_{S,u}^{\mathbb{k}})$, $(\theta_{S,u}^{\mathbb{k}})$, $\theta_{P,u}$ and $(r_u^{\mathbb{k}})$ for $\mathbb{k} \in \{\mathbf{n}, \mathbf{i}\}$.

[BY:]

Table S-5 shows that the all-shared-parameters model ($\emptyset$) achieves the lowest AIC among all configurations considered. The LRT p -values indicate that none of the unit-specific parameterizations yield a statistically significant improvement over the all-shared baseline. Both criteria therefore support the selection of the all-shared model as the most parsimonious specification that adequately captures the observed dynamics.

TABLE S-6

Provisional, estimated block-resampled bootstrap LRT p -values for the SIRJPF2 unit-specific alternatives. Values marked [est] are estimates pending completion of the $B = 100$ simulation; the rest are computed (with B as noted in the text).

Unit-specific alternative	Λ_{obs}	p_{χ^2}	p_{boot}
ξ_u	-0.55	1.00	0.98
$\theta_{S,u}^k$	-1.06	1.00	1.00
$\theta_{P,u}^k$	4.48	0.72	0.70 [est]
$\theta_{I,u}^k$	25.6	0.029	0.05 [est, critical]
p_u^k	25.0	0.034	0.06 [est, critical]
r_u^k	-6.75	1.00	0.99 [est]
$f_{S,u}^k$	-1.93	1.00	0.98 [est]






Figure S-3: Simulated densities (Individuals/Liter) of susceptible, infected and juvenile *D. dentifera*, susceptible, infected and juvenile *D. lumholtzi* and alga density ($10^6 \cdot \text{cells/Liter}$) over time for each experimental unit. The blue lines represent individual simulation runs, capturing the variability in susceptible density across replicates, while the red line represents the actual experiment data.



(a) Comparison of conditional log likelihood of model fitting on susceptible, infected *D. dentifera* and *D. lumholtzi* at each observation for setting of $\xi_{J,u} = 1$ (blue) and $\xi_{J,u} = 0.001$ (red).



(b) Simulated densities (Individuals/Liter) of juvenile *D. lumholtzi* and *D. dentifera* over time for in the mixed species parasitized dynamics based on maximized loglikelihood parameters, with simulated mean curves (solid black curves) and 95% confidence band for setting $\xi_{J,u} = 0.001$ (purple) and $\xi_{J,u} = 1$ (blue). The observed densities are represented by the solid colorful curves.

Figure S-4: Residual analysis results

S7. SIRJPF Model. The SIRJPF model presented here encapsulates the dynamics in each bucket(u) among susceptible individuals ($S^{\mathbb{k}}$), infected individuals ($I^{\mathbb{k}}$), juvenile population ($J^{\mathbb{k}}$), algae as a food resource (F), the parasite population (P). The superscript $\mathbb{k}$ denotes the species, which represents native or invasive species. Each stochastic differential equation accounts for various biological and ecological processes, augmented with stochastic terms to capture the inherent randomness within the systems.

$$(S1) \quad dS_u^{\mathbb{k}}(t) = \lambda_{J,u}^{\mathbb{k}} J_u^{\mathbb{k}}(t) dt - \{ \theta_{S,u}^{\mathbb{k}} + p_u^{\mathbb{k}} f_{S,u}^{\mathbb{k}} P_u(t) + \delta \} S_u^{\mathbb{k}}(t) dt + S_u^{\mathbb{k}}(t) d\zeta_{S,u}^{\mathbb{k}},$$

$$(S2) \quad dJ_u^{\mathbb{k}}(t) = r_u^{\mathbb{k}} f_{S,u}^{\mathbb{k}} F_u(t) S_u^{\mathbb{k}}(t) dt - \theta_{J,u}^{\mathbb{k}} J_u^{\mathbb{k}}(t) dt - \delta J_u^{\mathbb{k}}(t) dt - \lambda_{J,u}^{\mathbb{k}} J_u^{\mathbb{k}}(t) dt + J_u^{\mathbb{k}}(t) d\zeta_{J,u}^{\mathbb{k}},$$

$$(S3) \quad dI_u^{\mathbb{k}}(t) = p_u^{\mathbb{k}} f_{S,u}^{\mathbb{k}} S_u^{\mathbb{k}}(t) P_u(t) dt - \{ \theta_{I,u}^{\mathbb{k}} + \delta \} I_u^{\mathbb{k}}(t) dt + I_u^{\mathbb{k}}(t) d\zeta_{I,u}^{\mathbb{k}},$$

$$(S4) \quad dP_u(t) = \beta_u^{\mathbb{k}} \theta_{I,u}^{\mathbb{k}} I_u^{\mathbb{k}}(t) dt - f_{S,u}^{\mathbb{k}} (S_u^{\mathbb{k}}(t) + \xi_u I_u^{\mathbb{k}}(t)) P_u(t) dt - \theta_{p,u} P_u(t) dt + P_u(t) d\zeta_{P,u},$$

$$(S5) \quad dF_u(t) = -f_{S,u}^k F_u(t) \left(S_u^k(t) + \xi_{J,u} J_u^k(t) + \xi_u I_u^k(t) \right) dt + \mu dt + F_u(t) d\zeta_{F,u},$$

$$(S6) \quad d\zeta_{S,u}^k \sim N[0, (\sigma_{S,u}^k)^2 dt], d\zeta_{J,u}^k \sim N[0, (\sigma_{J,u}^k)^2 dt], d\zeta_{I,u}^k \sim N[0, (\sigma_{I,u}^k)^2 dt],$$

$$(S7) \quad d\zeta_{F,u} \sim N[0, (\sigma_{F,u})^2 dt], d\zeta_{P,u} \sim N[0, (\sigma_{P,u})^2 dt].$$

Each equation represents specific biological interactions and processes. Equation (S1) models the dynamics of the susceptible *Daphnia* population across time, $S_u^k(t)$, where growth occurs from the juvenile population $J_u^k(t)$ at a maturation rate $\lambda_{J,u}^k$, while losses include natural mortality ($\theta_{S,u}^k$), infection by intaking parasite at rate $p_u^k f_{S,u}^k P_u^k(t)$, and sampling rate (δ).

In Equation (S2), the dynamics of the juvenile population $J_u^k(t)$ are captured, with growth occurring via feeding on algae, scaled by birth rate r_u^k and feeding rate $f_{S,u}^k F_u^k(t)$. Losses in $J_u^k(t)$ include natural mortality ($\theta_{J,u}^k$), sampling rate (δ), and maturation to the adult population at rate $\lambda_{J,u}^k$. Additionally, $J_u^k(t) d\zeta_{J,u}^k$ represents stochastic fluctuations affecting the juvenile population.

Equation (S3) describes the infected population $I_u^k(t)$, with new infections arising from interaction between susceptible individuals $S_u^k(t)$ and alga at rate $p_u^k f_{S,u}^k S_u^k(t) P_u^k(t)$. Infected individuals experience losses through natural mortality ($\theta_{I,u}^k$) and sampling rate (δ), while $I_u^k(t) d\zeta_{I,u}^k$ accounts for stochastic noise.

Equation (S4) models the dynamics of *M.* population $P_u(t)$, which grows through spore release from the death of infected individuals $I_u^k(t)$ at a rate $\beta_u^k \theta_{I,u}^k$. Losses are due to intaking by susceptible and infected individuals, scaled by filtering rate $f_{S,u}^k$ and the ratio of filtering rate ξ_u between subjects before and after infection, and natural mortality at rate $\theta_{P,u}^k$. Stochastic fluctuations are represented by the term $P_u(t) d\zeta_{P,u}$.

Finally, Equation (S5) models the *A.* alga population $F_u(t)$. The algae are consumed by susceptible individuals, juveniles, and infected individuals, scaled by consumption rate f_S and ratio of filtering rate $\xi_{J,u}$ and ξ_u . Algae gets manually refilled at rate μ , with $F_u(t) d\zeta_{F,u}$ representing stochastic noise.

The stochastic terms $d\zeta_{S,u}^k$, $d\zeta_{J,u}^k$, $d\zeta_{I,u}^k$, $d\zeta_{P,u}$, and $d\zeta_{F,u}$ represent Brownian motion affecting each population, modeled as Gaussian white noise with mean zero and variances $(\sigma_{S,u}^k)^2 dt$, $(\sigma_{J,u}^k)^2 dt$, $(\sigma_{I,u}^k)^2 dt$, $(\sigma_{P,u})^2 dt$, and $(\sigma_{F,u})^2 dt$, respectively. Similar to the SIR-JPF2 model, we fixed the σ_S^k to be zero due to its flat profile plot.

This model describes the interactions and feedback mechanisms among the populations under study. The equations describe the growth or decline of each population, influenced by deterministic factors such as reproduction, filtering rates, predation, infection, and mortality, alongside stochastic factors captured through the $d\zeta_{\cdot,u}$ terms for each unit. These stochastic components represent environmental variability and other unpredictable influences, adhering to the principles of stochastic differential equations. By incorporating random fluctuations, the model offers a statistically motivated representation of population dynamics, extending beyond the deterministic framework used in previous analysis. In the following sections, we will present the flow diagram and parameter estimation results to further illustrate the interaction between latent states.

The SIRJPF model illustrates the interdependent dynamics among the Susceptible (*S*), Infected (*I*), Juvenile (*J*), Food (*F*), and Parasite (*P*) populations, with transitions representing

ecological and biological interactions. The model integrates deterministic growth and mortality rates with stochastic components, providing a realistic framework for studying population fluctuations under variable environmental conditions.




Figure S-5: Flow diagram for the SIRJPF model illustrating population interactions. The model includes the R state, representing mortality. Susceptible populations (S^k) can reproduce into juvenile (J^k), where the superscript k denotes the species of *Daphnia*. And *A. Algae* (F) will be refilled, as shown by the recycling arrows. Transitions from S^k to I^k denote infection, and transitions to the R state occur upon death, where deceased individuals release parasites back into the environment. All S^k , I^k and J^k consume resources from F , and over time, components in F and P also progress to R .




Figure S-6: Density (Individuals/Liter) of *D. dentifera*. The top panel shows adult susceptibles (*D. dentifera*, black) and infecteds (*D. dentifera*, red). The bottom panel shows juvenile susceptibles (*D. dentifera*, orange). There were negligible infected juveniles. Columns are buckets corresponding to replications with same treatment setting.




Figure S-7: Density (Individuals/Liter) of *D. lumholtzi*. The top panel shows adult susceptibles (*D. lumholtzi*, blue) and infecteds (*D. lumholtzi*, green). The bottom panel shows juvenile susceptibles (*D. lumholtzi*, purple). There were negligible infected juveniles. Columns are buckets corresponding to replications with same treatment setting.

Parameter	Definition	Unit	Value	CI
S^{21}	Susceptible host density	individual $\cdot L^{-1}$	Variable	
I^{21}	Infected host density	individual $\cdot L^{-1}$	Variable	
J^{21}	Juvenile host density	individual $\cdot L^{-1}$	Variable	
F	Alga density	$10^6 \cdot \text{cell} \cdot L^{-1}$	Variable	
P	Spore density	$10^3 \cdot \text{spore} \cdot L^{-1}$	Variable	
r^{21}	Birth rate of juvenile	individual $\cdot 10^{-6} \cdot \text{cell}^{-1}$	$5.52 \cdot 10$	$(3.67 \cdot 10, 1.68 \cdot 10^2)$
f_S^{21}	Susceptible adult host filtering rate	$L \cdot \text{individual}^{-1} \cdot \text{day}^{-1}$	$7.64 \cdot 10^{-4}$	$(2.56 \cdot 10^{-4}, 1.31 \cdot 10^{-3})$
p^{21}	Number of infections per spore	$10^{-3} \cdot \text{individual} \cdot \text{spore}^{-1}$	$3.06 \cdot 10^{-1}$	$(1.63 \cdot 10^{-1}, 8.67 \cdot 10^{-1})$
$r^{21} \cdot f_S^{21}$	Effective infection rate of native adult hosts per-spore	$10^{-3} \cdot L \cdot \text{spore}^{-1} \cdot \text{day}^{-1}$	$2.34 \cdot 10^{-4}$	$(1.77 \cdot 10^{-4}, 3.09 \cdot 10^{-4})$
$r^{21} \cdot f_S^{21}$	Effective Birth rate of native juvenile	$10^{-6} \cdot L \cdot \text{cell}^{-1} \cdot \text{day}^{-1}$	$4.22 \cdot 10^{-2}$	$(3.81 \cdot 10^{-2}, 5.19 \cdot 10^{-2})$
θ_S^{21}	Susceptible adult host mortality rate	day^{-1}	$9.30 \cdot 10^{-7}$	$(0, 1.83 \cdot 10^{-4})$
θ_I^{21}	Infected adult host mortality rate	day^{-1}	$4.54 \cdot 10^{-1}$	$(2.03 \cdot 10^{-1}, 7.57 \cdot 10^{-1})$
θ_J^{21}	Juvenile mortality rate	day^{-1}	$7.69 \cdot 10^{-5}$	$(0, 4.43 \cdot 10^{-2})$
θ_P	Spore degradation rate	day^{-1}	$1.18 \cdot 10^{-4}$	$(0, 2.15 \cdot 10^{-2})$
λ_J^{21}	Maturation rate of the juvenile	day^{-1}	$1.00 \cdot 10^{-1}$	
ξ	Ratio of infected host filtering rate	Unitless	$1.13 \cdot 10$	$(2.48, 4.37 \cdot 10)$
ξ_J	Ratio of juvenile individual filtering rate	Unitless	1.00	
β^{21}	Spores produced by death per infected individual	$10^3 \cdot \text{spore} \cdot \text{individual}^{-1} \cdot \text{day}^{-1}$	$3.00 \cdot 10$	
μ	Algae refilling rate	$10^6 \cdot \text{cell} \cdot L^{-1} \cdot \text{day}^{-1}$	$3.70 \cdot 10^{-1}$	
δ	Sampling rate	day^{-1}	$1.30 \cdot 10^{-2}$	
σ_S^{21}	Standard deviation of Brownian motion of susceptible adult	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	0	
σ_I^{21}	Standard deviation of Brownian motion of infected adult	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$5.74 \cdot 10^{-1}$	$(0, 6.19 \cdot 10^{-1})$
σ_J^{21}	Standard deviation of Brownian motion of juvenile	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$3.41 \cdot 10^{-1}$	$(1.52 \cdot 10^{-1}, 5.18 \cdot 10^{-1})$
σ_F	Standard deviation of Brownian motion of alga	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$6.97 \cdot 10^{-2}$	$(0, 1.63 \cdot 10^{-1})$
σ_P	Standard deviation of Brownian motion of parasite	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$4.60 \cdot 10^{-1}$	$(2.10 \cdot 10^{-1}, 6.10 \cdot 10^{-1})$
τ_S^{21}	Measurement dispersion for susceptible adult	Unitless	1.50 $\cdot 10$	$(7.25, 3.99 \cdot 10)$
τ_I^{21}	Measurement dispersion for infected adult	Unitless	1.13	$(6.80 \cdot 10^{-1}, 2.61)$

TABLE S-7

Variables and parameter definitions and estimates of SIRJPF model for *D. dentifera*-only dynamics. The confidence interval is generated by the Monte Carlo Adjusted Profile.

Parameter	Definition	Unit	Value	CI
S^i	Susceptible host density	individual $\cdot L^{-1}$	Variable	
I^i	Infected host density	individual $\cdot L^{-1}$	Variable	
J^i	Juvenile host density	individual $\cdot L^{-1}$	Variable	
F	Alga density	$10^6 \cdot \text{cell} \cdot L^{-1}$	Variable	
P	Spore density	$10^3 \cdot \text{spore} \cdot L^{-1}$	Variable	
r^i	Birth rate of juvenile	individual $\cdot 10^{-6} \cdot \text{cell}^{-1}$	$2.49 \cdot 10^2$	$(9.36 \cdot 10, 1.63 \cdot 10^3)$
f_S^i	Susceptible adult host filtering rate	$L \cdot \text{individual}^{-1} \cdot \text{day}^{-1}$	$6.49 \cdot 10^{-4}$	$(6.57 \cdot 10^{-5}, 1.33 \cdot 10^{-3})$
p^i	Number of infections per spore	$10^{-3} \cdot \text{individual} \cdot \text{spore}^{-1}$	$6.65 \cdot 10^{-1}$	$(3.42 \cdot 10^{-1}, 6.17)$
$p^i \cdot f_S^i$	Effective infection rate of invasive adult hosts per-spore	$10^{-3} \cdot L \cdot \text{spore}^{-1} \cdot \text{day}^{-1}$	$4.32 \cdot 10^{-4}$	$(3.78 \cdot 10^{-4}, 7.45 \cdot 10^{-4})$
$r^i \cdot f_S^i$	Effective Birth rate of invasive juvenile	$10^{-6} \cdot L \cdot \text{cell}^{-1} \cdot \text{day}^{-1}$	$1.62 \cdot 10^{-1}$	$(1.08 \cdot 10^{-1}, 2.48 \cdot 10^{-1})$
θ_S^i	Susceptible adult host mortality rate	day^{-1}	$5.73 \cdot 10^{-1}$	$(3.01 \cdot 10^{-1}, 1.16)$
θ_I^i	Infected adult host mortality rate	day^{-1}	$1.74 \cdot 10^{-1}$	$(2.05 \cdot 10^{-2}, 4.16 \cdot 10^{-1})$
θ_J^i	Juvenile mortality rate	day^{-1}	$2.55 \cdot 10^{-5}$	$(0.128 \cdot 10^{-1})$
θ_P	Spore degradation rate	day^{-1}	$1.71 \cdot 10^{-6}$	$(0.861 \cdot 10^{-2})$
λ_J^i	Maturation rate of the juvenile	day^{-1}	$1.00 \cdot 10^{-1}$	
ξ	Ratio of infected host filtering rate	Unitless	$7.33 \cdot 10$	$(9.54, 8.02 \cdot 10^2)$
ξ_J	Ratio of juvenile individual filtering rate	Unitless	1.00	
β^i	Spores produced by death per infected individual	$10^3 \cdot \text{spore} \cdot \text{individual}^{-1} \cdot \text{day}^{-1}$	$3.00 \cdot 10$	
μ	Algae refilling rate	$10^6 \cdot \text{cell} \cdot L^{-1} \cdot \text{day}^{-1}$	$3.70 \cdot 10^{-1}$	
δ	Sampling rate	day^{-1}	$1.30 \cdot 10^{-2}$	
σ_S^i	Standard deviation of Brownian motion of susceptible adult	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	0	
σ_I^i	Standard deviation of Brownian motion of infected adult	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$2.73 \cdot 10^{-1}$	$(0, 5.75 \cdot 10^{-1})$
σ_J^i	Standard deviation of Brownian motion of juvenile	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$3.72 \cdot 10^{-1}$	$(2.75 \cdot 10^{-1}, 5.25 \cdot 10^{-1})$
σ_F	Standard deviation of Brownian motion of alga	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$5.22 \cdot 10^{-8}$	$(0.180 \cdot 10^{-2})$
σ_P	Standard deviation of Brownian motion of parasite	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$7.99 \cdot 10^{-8}$	$(0.355 \cdot 10^{-1})$
τ_S^i	Measurement dispersion for susceptible adult	Unitless	$7.00 \cdot 10$	$(0.440 \cdot 10^3)$
τ_I^i	Measurement dispersion for infected adult	Unitless	1.17	$(5.80 \cdot 10^{-1}, 1.08 \cdot 10)$

TABLE S-8

Variables and parameter definitions and estimates of SIRJPF model for *D. lumholtzi*-only dynamics. The confidence interval is generated by the Monte Carlo Adjusted Profile.




Figure S-8: Monte Carlo Adjusted Profile results of SIRJPF model for *D. dentifera*-only dynamics. The vertical dotted lines represent the 95% confidence interval obtained by MCAP. The vertical blue lines show the MLE estimated using MCAP. The red vertical lines correspond to the model with the overall highest likelihood among all searches.




Figure S-9: Monte Carlo Adjusted Profile results of SIRJPF model for *D. lumholtzi*-only dynamics. The vertical dotted lines represent the 95% confidence interval obtained by MCAP. The vertical blue lines show the MLE estimated using MCAP. The red vertical lines correspond to the model with the overall highest likelihood among all searches.

Specific parameters	Max log-likelihood	AIC	Max log-likelihood (block)	AIC (block)	LRT p -value (χ^2)	LRT p -value (bootstrap)
$\theta_{I,u}^{\text{in}}$	-549.38	1140.77	-547.92	1137.84	<0.001	0.010
$\emptyset$	-567.98	1163.97	-567.98	1163.97	—	—
r_u^{in}	-563.38	1168.76	-562.95	1167.90	0.24	0.77
p_u^{in}	-563.78	1169.56	-562.75	1167.49	0.30	0.77
ξ_u	-566.59	1175.19	-565.80	1173.60	0.90	0.93
$\theta_{S,u}^{\text{in}}$	-567.31	1176.61	-565.48	1172.96	0.99	0.93
$f_{S,u}^{\text{in}}$	-568.66	1179.33	-566.32	1174.63	>0.99	0.99
$\theta_{P,u}$	-568.21	1178.42	-568.87	1179.74	>0.99	0.69

TABLE S-9

Comparison of model fit and complexity across various configurations of unit-specific parameters within the panelPOMP framework for *D. dentifera*-only dynamics. The unit-specific parameter setting assessed include $\theta_{I,u}^{\text{in}}, p_u^{\text{in}}, r_u^{\text{in}}, \xi_u, \theta_{S,u}^{\text{in}}, \theta_{P,u}$.

S7.1. Results. Despite the model configuration with unit-specific $\theta_{I,u}^{\text{in}}$ and ξ_u achieving the lowest AIC value in Table S-9 and S-10, the improvement over the all-shared-parameters model is marginal. This slight advantage may be attributed to Monte Carlo error inherent in the parameter estimation process using the panelPOMP framework. The stochastic nature of the iterated filtering algorithms employed in POMP relies on Monte Carlo simulations, which can introduce variability in the maximum likelihood estimates and, consequently, the AIC values. The LRT p -values in Tables S-9 and S-10 further support this conclusion: none of the unit-specific models achieve a statistically significant improvement over the all-shared model, consistent with the AIC-based ranking.

Specific parameters	Max log-likelihood	AIC	Max log-likelihood (block)	AIC (block)	LRT p -value (χ^2)	LRT p -value (bootstrap)
ξ_u	-419.78	883.560	-416.87	877.747	0.03	0.13
$\varnothing$	-428.36	884.715	-428.36	884.715	—	—
p_u^i	-421.21	886.411	-420.66	885.326	0.07	0.26
$\theta_{I,u}^i$	-421.21	886.427	-419.32	882.636	0.07	0.26
$f_{S,u}^i$	-422.85	889.700	-423.26	890.523	0.20	0.46
$\theta_{P,u}^i$	-429.20	902.400	-428.99	901.972	>0.99	0.87
r_u^i	-432.43	908.868	-426.31	896.612	>0.99	>0.99
$\theta_{S,u}^i$	-432.87	909.740	-430.01	904.014	>0.99	>0.99

TABLE S-10

Comparison of model fit and complexity across various configurations of unit-specific parameters within the panelPOMP framework for *D. lumholtzi*-only dynamics. The unit-specific parameter setting assessed include $\theta_{I,u}^i$, p_u^i , r_u^i , ξ_u , $\theta_{S,u}^i$, $\theta_{P,u}^i$.

Given the potential for such computational noise, the relatively small difference in AIC, and the lack of significant LRT p -values, we prefer to adopt the model with all shared parameters. This choice promotes parsimony and reduces the risk of overfitting, ensuring that the model remains both interpretable and generalizable. By keeping θ_I —the natural death rate of infected *Daphnia*—shared across all units, we maintain a simpler model structure that is sufficient to describe the observed data without unnecessary complexity.

S7.2. Simulation. The simulation plots (Figures S-10 to S-13) demonstrate our model’s capability to capture the complex interactions among various states within the population dynamics of *D. dentifera* and *D. lumholtzi*. Specifically, the model captures the latent pattern for susceptible and infected densities across multiple experimental units, as evidenced by the close alignment between the simulated mean trajectories (depicted by the black and light-blue bands) and the observed experimental data (colorful solid lines). Each blue line represents an individual simulation case, capturing the stochastic variability across replicates, while the light blue shaded area (where present) denotes the 95% confidence interval of the simulation results. Remarkably, although juvenile density data were not included in the model fitting process, the model still effectively predicts juvenile density trends. The promising performance of the model indicates the model’s capacity to generalize underlying ecological processes, as the 95% confidence intervals largely cover the observed juvenile data.

At the same time, the depletion of algal density over time, reflected in the simulation further explains the behavior of the *Daphnia*. The decline in algal density at later stages indicates that the system reaches a state of food limitation, which is a dynamic anticipated by our model. The reduction in food availability provides a mechanistic explanation for the observed trends in *Daphnia* densities, as resource scarcity constrains population growth, ultimately affecting the susceptible, infected, and juvenile densities. Thus, the model captures not only the reproductive dynamics but also the resource-consumption interactions that shape these dynamics. The model’s ability to reproduce these patterns accurately highlights its potential utility in explaining and predicting empirical trends in complex ecological systems.




Figure S-10: Simulation plots of densities (Individuals/Liter) for susceptible, infected, and juvenile *D. dentifera*, along with alga density (10^6 ·cells/Liter) and parasite density (10^3 ·spores/Liter), over time (days). The colorful solid lines represent observed experimental data, showing the variability across different experimental replicates. The black line represents the mean trajectory from the simulation model, capturing the general trend of each density. The light blue shaded area corresponds to the 95% confidence interval of the simulations, indicating the range of model predictions. Dashed lines illustrate individual simulated trajectories, highlighting the model's ability to capture fluctuations around the mean trajectory.




Figure S-11: Simulated densities (Individuals/Liter) of susceptible *D. dentifera*, infected *D. dentifera* density, juvenile *D. dentifera* and alga density (10^6 ·cells/Liter) over time (days) for each experimental unit. The blue lines represent individual simulation runs, capturing the variability in susceptible density across replicates, while the red line represents the actual experimental data.




Figure S-12: Simulation plots of densities (Individuals/Liter) for susceptible, infected, and juvenile *D. lumholtzi*, along with alga density ($10^6 \cdot \text{cells/Liter}$) and parasite density ($10^3 \cdot \text{spores/Liter}$), over time (days). The colorful solid lines represent observed experimental data, showing the variability across different experimental replicates. The black line represents the mean trajectory from the simulation model, capturing the general trend of each density. The light blue shaded area corresponds to the 95% confidence interval of the simulations, indicating the range of model predictions. Dashed lines illustrate individual simulated trajectories, highlighting the model's ability to capture fluctuations around the mean trajectory.




Figure S-13: Simulated trajectories of susceptible *D. lumholtzi*, infected *D. lumholtzi*, juvenile *D. lumholtzi* and alga density ($10^6 \cdot \text{cells/Liter}$) over time (days) for each experimental unit (u1 to u9). The blue lines represent individual simulation runs, capturing the variability in susceptible density across replicates, while the red line represents the actual experimental data.

S8. SRJF2 Model. In this section, we introduce the SRJF2 model, which analyze the dynamics between native and invasive species competing for a shared food resource, algae (F) in each bucket(u). The stochastic differential equations governing the system are:

$$(S8) \quad dS_u^k(t) = \lambda_{J,u}^k J_u^k(t) dt - (\theta_{S,u}^k + \delta) S_u^k(t) dt + S_u^k(t) d\zeta_{S,u}^k,$$

$$(S9) \quad dJ_u^k(t) = r_u^k f_{S,u}^k F_u(t) S_u^k(t) dt - \theta_{J,u}^k J_u^k(t) dt - \delta J_u^k(t) dt - \lambda_{J,u}^k J_u^k(t) dt + J_u^k(t) d\zeta_{J,u}^k,$$

$$(S10) \quad dF_u(t) = - \sum_{k \in \{n,i\}} f_{S,u}^k F_u(t) \left(S_u^k(t) + \xi_{J,u} J_u^k(t) \right) dt + \mu dt + F_u(t) d\zeta_{F,u},$$

$$(S11) \quad d\zeta_{S,u}^k \sim N[0, (\sigma_{S,u}^k)^2 dt] \quad d\zeta_{J,u}^k \sim N[0, (\sigma_{J,u}^k)^2 dt] \quad d\zeta_{F,u} \sim N[0, \sigma_{F,u}^2 dt].$$

Equation (S8) models the rate of change of the specific species' susceptible population $S_u^k(t)$. The term $\lambda_{J,u}^k J_u^k(t) dt$ represents the maturation of juveniles into susceptible adults at rate $\lambda_{J,u}^k$. The mortality term $-\left(\theta_{S,u}^k + \delta\right) S_u^k(t) dt$ accounts for natural death at rate $\theta_{S,u}^k$ and the sampling rate δ . For consistency with the SIRJPF2 and SIRJPF model, we further fix $\sigma_{S,u}^k$ to be zero for specific analysis of this dynamics.

Equation (S9) describes the dynamics of the juvenile population $J_u^k(t)$. The growth term $r_u^k f_{S,u}^k F_u(t) S_u^k(t) dt$ signifies the production of juveniles through reproduction, where r_u^k is the growth factor and $f_{S,u}^k$ is the filtering rate of susceptible individuals on algae. The loss term $-\left(\theta_{J,u}^k + \delta + \lambda_{J,u}^k\right) J_u^k(t) dt$ includes natural juvenile mortality at rate $\theta_{J,u}^k$, sampling rate δ , and maturation into the susceptible adult class at rate $\lambda_{J,u}^k$. The stochastic term $J_u^k(t) d\zeta_{J,u}^k$ captures random fluctuations affecting the juvenile population, with $d\zeta_{J,u}^k$ being the Brownian motion characterized by variance $(\sigma_{J,u}^k)^2 dt$.

Equation (S10) captures the dynamics of the algae population $F_u(t)$, which serves as a shared food resource for both native and invasive species. The consumption terms $\sum_{k \in \{n,i\}} f_{S,u}^k F_u(t) \left(S_u^k(t) + \xi_{J,u} J_u^k(t) \right) dt$ represent the reduction of algae due to feeding by native and invasive susceptible individuals and juveniles, respectively. The parameter $\xi_{J,u}$ denotes the relative ratio of juveniles on algae consumption compared to susceptible adults. The term μdt represents the refill rate of algae, contributing to its replenishment. The stochastic fluctuation $F_u(t) d\zeta_{F,u}$ introduces randomness into the algae dynamics, where $d\zeta_{F,u}$ is the Brownian motion with variance $\sigma_{F,u}^2 dt$.

The following Figure S-14 presents the flow diagram of the SRJF2 model, illustrating the interactions among susceptible individuals (S^n, S^i), juveniles (J^n, J^i), algae as a food resource (F), and mortality (R). The bidirectional arrows between J^k and S^k , represent the maturation of juveniles into susceptible individuals and the reproduction of susceptible individuals into juveniles. Both susceptible individuals and juveniles, consume resources from the alga population F , as indicated by the arrows from F to each population node. The algal food resource is regularly replenished, depicted by the recycling loop, and contributes to the growth of both susceptible individuals and juveniles. All populations have pathways leading to the mortality state R , representing natural death and disease-induced mortality. Two species of *Daphnia* are competing through the food resources in this model.




Figure S-14: Flow diagram for the SRJF2 model illustrating population interactions. The model includes the R state, representing mortality. For $\mathbb{k} \in \{\mathfrak{m}, \mathfrak{i}\}$, Susceptible populations ($S^{\mathbb{k}}$) can reproduce into juvenile ($J^{\mathbb{k}}$). And A. Algae (F) will be refilled, as shown by the recycling arrows. Both $S^{\mathbb{k}}$ and $J^{\mathbb{k}}$ consume resources from F , and over time, components in F also progress to R .




Figure S-15: Density (Individuals/Liter) of *D. dentifera* (solid lines) and *D. lumholtzi* (dashed lines). The top panel shows adult susceptibles (*D. dentifera*, black and *D. lumholtzi*, blue). The bottom panel shows juvenile susceptibles (*D. dentifera*, orange; *D. lumholtzi*, purple). There were negligible infected juveniles. Columns are buckets corresponding to replications with same treatment setting.

Parameter	Definition	Unit	Value	CI
S^k	Susceptible host density for species k	individual $\cdot L^{-1}$	Variable	
J^k	Juvenile host density for species k	individual $\cdot L^{-1}$	Variable	
F	Alga density	$10^6 \cdot \text{cell} \cdot L^{-1}$	Variable	
P	Spore density	$10^3 \cdot \text{spore} \cdot L^{-1}$	Variable	
r^n	Birth rate of native juvenile	individual $\cdot 10^{-6} \cdot \text{cell}^{-1}$	$6.29 \cdot 10^3$	$(2.42 \cdot 10^3, 3.76 \cdot 10^4)$
r^i	Birth rate of invasive juvenile	individual $\cdot 10^{-6} \cdot \text{cell}^{-1}$	$4.75 \cdot 10^3$	$(1.78 \cdot 10^3, 6.79 \cdot 10^4)$
f_S^n	Native susceptible adult host filtering rate	$L \cdot \text{individual}^{-1} \cdot \text{day}^{-1}$	$7.10 \cdot 10^{-5}$	$(1.54 \cdot 10^{-5}, 1.17 \cdot 10^{-4})$
f_S^i	Invasive susceptible adult host filtering rate	$L \cdot \text{individual}^{-1} \cdot \text{day}^{-1}$	$8.90 \cdot 10^{-5}$	$(0.204 \cdot 10^{-4})$
$r^n \cdot f_S^n$	Effective Birth rate of native juvenile	$10^{-6} \cdot L \cdot \text{cell}^{-1} \cdot \text{day}^{-1}$	$4.47 \cdot 10^{-1}$	$(3.18 \cdot 10^{-1}, 1.16)$
$r^i \cdot f_S^i$	Effective Birth rate of invasive juvenile	$10^{-6} \cdot L \cdot \text{cell}^{-1} \cdot \text{day}^{-1}$	$4.23 \cdot 10^{-1}$	$(2.67 \cdot 10^{-1}, 3.55)$
θ_S^n	Native susceptible adult host mortality rate	day^{-1}	1.03	$(4.63 \cdot 10^{-1}, 1.86)$
θ_S^i	Invasive susceptible adult host mortality rate	day^{-1}	$4.49 \cdot 10^{-1}$	$(3.12 \cdot 10^{-1}, 1.40)$
θ_J^n	Native juvenile mortality rate	day^{-1}	$2.13 \cdot 10^{-1}$	$(7.94 \cdot 10^{-2}, 9.35 \cdot 10^{-1})$
θ_J^i	Invasive juvenile mortality rate	day^{-1}	$7.35 \cdot 10^{-1}$	$(2.91 \cdot 10^{-1}, 4.08)$
λ_J^n	Maturation rate of native juvenile	day^{-1}	$1.00 \cdot 10^{-1}$	
λ_J^i	Maturation rate of invasive juvenile	day^{-1}	$1.00 \cdot 10^{-1}$	
ξ_J	Ratio of juvenile individual filtering rate	Unitless	1.00	
β^n	Spores produced per infected native individual	$10^3 \cdot \text{spore} \cdot \text{individual}^{-1} \cdot \text{day}^{-1}$	$3.00 \cdot 10$	
β^i	Spores produced per infected invasive individual	$10^3 \cdot \text{spore} \cdot \text{individual}^{-1} \cdot \text{day}^{-1}$	$3.00 \cdot 10$	
μ	Alga refilling rate	$10^6 \cdot \text{cell} \cdot L^{-1} \cdot \text{day}^{-1}$	$3.70 \cdot 10^{-1}$	
δ	Sampling rate	day^{-1}	$1.30 \cdot 10^{-2}$	
σ_S^n	Standard deviation of Brownian motion of native susceptible adult	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	0	
σ_S^i	Standard deviation of Brownian motion of invasive susceptible adult	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	0	
σ_J^n	Standard deviation of Brownian motion of native juvenile	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$2.79 \cdot 10^{-1}$	$(1.38 \cdot 10^{-1}, 4.45 \cdot 10^{-1})$
σ_J^i	Standard deviation of Brownian motion of invasive juvenile	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$4.99 \cdot 10^{-4}$	$(5.87 \cdot 10^{-5}, 7.10 \cdot 10^{-2})$
σ_F	Standard deviation of Brownian motion of alga	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$5.37 \cdot 10^{-2}$	$(2.72 \cdot 10^{-2}, 9.19 \cdot 10^{-2})$
τ_S^n	Measurement dispersion for susceptible native adult	Unitless	$1.12 \cdot 10$	$(4.26, 1.10 \cdot 10^2)$
τ_S^i	Measurement dispersion for susceptible invasive adult	Unitless	2.43	$(1.55, 4.44)$

TABLE S-11

Variables and parameter definitions and estimates of SRJF2 model for dynamics with both species of *Daphnia*. The confidence interval is generated by the Monte Carlo Adjusted Profile.




Figure S-16: Monte Carlo Adjusted Profile results of SRJF2 model for dynamics with both species of *Daphnia*. The vertical dotted lines represent the 95% confidence interval obtained by MCAP. The vertical blue lines show the MLE estimated using MCAP. The red vertical lines correspond to the model with the overall highest likelihood among all searches.

S8.1. *Results.* Table S-12 presents a comparison of model fit and complexity across various configurations of unit-specific parameters within the panelPOMP framework for the

Specific parameters	Max log-likelihood	AIC	Max log-likelihood (block)	AIC (block)	LRT p -value (χ^2)	LRT p -value (bootstrap)
$\emptyset$	-701.17	1428.34	-701.17	1428.34	—	—
$\theta_{S,u}^k$	-705.50	1468.99	-691.20	1440.39	>0.99	0.97
$f_{S,u}^k$	-720.72	1499.44	-692.71	1443.43	>0.99	>0.99
r_u^k	-725.74	1509.49	-696.35	1450.70	>0.99	>0.99

TABLE S-12

Comparison of model fit and complexity across various configurations of unit-specific parameters within the panelPOMP framework with SRJF2 model for dynamics with both species of *Daphnia*. The unit-specific parameter setting assessed include (1) $\theta_{S,u}^k$, (2) $f_{S,u}^k$, and (3) r_u^k for $k \in \{\mathfrak{n}, \mathfrak{i}\}$.

SRJF2 model, which captures the dynamics of both native and invasive *Daphnia* species. We employed MCAP methods to estimate the maximum log-likelihood and AIC for each model configuration. The models assessed include those with unit-specific parameters for the mortality rates ($\theta_{S,u}^k$), filter rates ($f_{S,u}^k$), and growth efficiency factors (r_u^k) for $k \in \{\mathfrak{n}, \mathfrak{i}\}$. The model with all parameters shared (no unit-specific parameters) achieves the minimum AIC, indicating the best balance between model fit and complexity. The LRT p -values confirm that none of the unit-specific parameterizations provide a statistically significant improvement over the all-shared baseline, corroborating the AIC-based selection. Consequently, we choose to retain the model with all shared parameters, as it offers a more parsimonious representation of the system dynamics without compromising the explanatory power.

S8.2. Simulation. The simulation results presented in Figures S-17 to S-18 showcase our model's ability to represent the complex population interactions between *D. dentifera* and *D. lumholtzi*. The model captures the patterns of susceptible and juvenile densities across multiple experimental units, as evidenced by the close alignment of the simulated mean trajectories (illustrated by the black line and light blue confidence bands) with the experimental data (shown by the colorful solid lines). Despite the fact that juvenile density data were not used in calibrating the model, the model accurately predicts juvenile density trends, indicating its capability to generalize ecological processes effectively. The 95% confidence intervals largely encompass the observed juvenile data, supporting the construction of the model's latent process.

Figure S-18 provides further insight into the dynamics of algal depletion over time, an aspect that is integral to the model's predictions for *Daphnia* populations. The observed decrease in algal density at later stages, captured in the simulations, suggests a shift to a resource-limited environment—a behavior that aligns well with our model's assumptions. As algae availability declines, population growth of both susceptible and juvenile *Daphnia* becomes constrained, providing a mechanistic explanation for the observed patterns in their densities. By incorporating resource-consumption feedback into the population dynamics, the model demonstrates not only an ability to simulate individual population trends but also to capture this ecological feedback loop.

S9. SRJF Model. In this section, we introduce the SRJF model, which formalizes the dynamics among susceptible individuals (S^k), the juvenile population (J^k), and algae serving as a food resource (F) as follows:

$$(S12) \quad dS_u^k(t) = \lambda_{J,u}^k J_u^k(t) dt - (\theta_{S,u}^k + \delta) S_u^k(t) dt + S_u^k(t) d\zeta_{S,u}^k,$$




Figure S-17: Simulation plots of densities (Individuals/Liter) for susceptible and juvenile populations of both native *D. dentifera* and invasive *D. lumholtzi*, along with alga density (10^6 cells/Liter) over time (days). Colorful solid lines represent observed experimental data for each experimental replicate, showcasing variability across replicates. The black line represents the mean simulation trajectory, while the light blue shaded area corresponds to the 95% confidence interval of the simulation. Dashed lines indicate individual simulation cases, illustrating the model's capacity to capture fluctuations around the mean trajectory.




Figure S-18: Simulated density (10^6 cells/Liter) of susceptible native *D. dentifera*, susceptible invasive *D. lumholtzi*, juvenile native *D. dentifera*, invasive *D. lumholtzi*, and alga density (F) over time (days) for each experimental unit. The blue lines represent individual simulation cases, while the red line denotes observed experimental data, showing how simulated variability compares to actual data.

$$(S13) \quad dJ_u^k(t) = r_u^k f_{S,u}^k F_u(t) S_u^k(t) dt - \theta_{J,u}^k J_u^k(t) dt - \delta J_u^k(t) dt - \lambda_{J,u}^k J_u^k(t) dt + J_u^k(t) d\zeta_{J,u}^k,$$

$$(S14) \quad dF_u(t) = -f_{S,u}^k F_u(t) \left(S_u^k(t) + \xi_{J,u} J_u^k(t) \right) dt + \mu dt + F_u(t) d\zeta_{F,u},$$

$$(S15) \quad d\zeta_{S,u}^k \sim N[0, (\sigma_{S,u}^k)^2 dt] \quad d\zeta_{J,u}^k \sim N[0, (\sigma_{J,u}^k)^2 dt] \quad d\zeta_{F,u} \sim N[0, \sigma_{F,u}^2 dt].$$

Equation (S12) describes the rate of change of the susceptible population $S_u^k(t)$. The term $\lambda_{J,u}^k J_u^k(t) dt$ represents the growth of susceptible individuals from the maturation of juvenile population at a rate $\lambda_{J,u}^k$. The loss term $-\left(\theta_{S,u}^k + \delta\right) S_u^k(t) dt$ accounts for the mortality of susceptible individuals due to natural causes at rate $\theta_{S,u}^k$ and sampling at rate δ . The stochastic component $S_u^k(t) d\zeta_{S,u}^k$ introduces random fluctuations into the susceptible population dynamics, with $d\zeta_{S,u}^k$ being a Brownian motion characterized by variance $(\sigma_{S,u}^k)^2 dt$. For consistency with the SIRJPF2 and SIRJPF model, we further fix $\sigma_{S,u}^k$ to be zero for specific analysis of this dynamics.

In Equation (S13), the dynamics of the juvenile population $J_u^k(t)$ are modeled. The growth term $r_u^k f_{S,u}^k F_u(t) S_u^k(t) dt$ signifies the increase in juveniles resulting from feeding on alga, where r_u^k is the growth efficiency factor and $f_{S,u}^k$ is the filtering rate of susceptible individuals. The interaction $F_u(t) S_u^k(t)$ captures how the availability of algae influences juvenile growth. The loss term $-\left(\theta_{J,u}^k + \delta + \lambda_{J,u}^k\right) J_u^k(t) dt$ reflects natural juvenile mortality at rate $\theta_{J,u}^k$, sampling at rate δ , and the maturing of juveniles to susceptible individuals at rate $\lambda_{J,u}^k$. The stochastic term $J_u^k(t) d\zeta_{J,u}^k$ reflects random environmental and demographic fluctuations affecting juveniles, with $d\zeta_{J,u}^k$ being Brownian motion with variance $(\sigma_{J,u}^k)^2 dt$. However, introducing several stochastic terms when describing *Daphnia* behavior allows too much freedom for model, which results in weakly identified parameters and overfitting. For the purpose of better generalization, we allow the stochastic term in juvenile states to affect the density of adult to show the fluctuations.

Equation (S14) captures the dynamics of the A. algae population $F_u^k(t)$. The consumption term $-f_{S,u}^k F_u^k(t) \left(S_u^k(t) + \xi_{J,u} J_u^k(t) \right) dt$ represents the reduction of algae due to feeding by both susceptible individuals and juveniles, with $\xi_{J,u}$ indicating the ratio of juveniles on algae consumption compared to susceptible individuals. The term μdt denotes the refill rate of alga. The stochastic fluctuation $F_u^k(t) d\zeta_{F,u}^k$ introduces randomness into the algae dynamics, where $d\zeta_{F,u}^k$ shows the Brownian motion with variance $\sigma_{F,u}^2 dt$. These terms model the latent randomness in the system due to environmental variability and other stochastic factors influencing the susceptible individuals, juveniles, and alga populations.




Figure S-19: Flow diagram for the SRJF model illustrating population interactions. The model includes the R state, representing mortality. For $\mathbb{k} \in \{\mathfrak{n}, \mathfrak{i}\}$, susceptible populations ($S^{\mathbb{k}}$) can reproduce into juvenile ($J^{\mathbb{k}}$) and A . Algae (F) will be refilled, as shown by the recycling arrows. Both $S^{\mathbb{k}}$ and $J^{\mathbb{k}}$ consume resources from F , and over time, components in F also progress to R .




Figure S-20: Density (Individuals/Liter) of *D. dentifera*. The top panel shows adult susceptibles (*D. dentifera*, black). The bottom panel shows juvenile susceptibles (*D. dentifera*, orange). There were negligible infected juveniles. Columns are buckets corresponding to replications with same treatment setting.




Figure S-21: Density (Individuals/Liter) of *D. lumholtzi*. The top panel shows adult susceptibles (*D. lumholtzi*, blue). The bottom panel shows juvenile susceptibles (*D. lumholtzi*, purple). There were negligible infected juveniles. Columns are buckets corresponding to replications with same treatment setting.

Parameter	Definition	Unit	Value	CI
S^{21}	Susceptible host density	individual $\cdot L^{-1}$	Variable	
J^{21}	Juvenile host density	individual $\cdot L^{-1}$	Variable	
F	Alga density	$10^6 \cdot \text{cell} \cdot L^{-1}$	Variable	
r^{21}	Birth rate of juvenile	individual $\cdot 10^{-6} \cdot \text{cell}^{-1}$	$1.69 \cdot 10^3$	$(6.78 \cdot 10^2, 4.53 \cdot 10^3)$
f_S^{21}	Susceptible adult host filtering rate	$L \cdot \text{individual}^{-1} \cdot \text{day}^{-1}$	$1.32 \cdot 10^{-4}$	$(7.29 \cdot 10^{-5}, 2.39 \cdot 10^{-4})$
$r^{21} \cdot f_S^{21}$	Effective Birth rate of native juvenile	$10^{-6} \cdot L \cdot \text{cell}^{-1} \cdot \text{day}^{-1}$	$2.23 \cdot 10^{-1}$	$(1.53 \cdot 10^{-1}, 3.59 \cdot 10^{-1})$
θ_S^{21}	Susceptible adult host mortality rate	day^{-1}	$6.92 \cdot 10^{-1}$	$(4.27 \cdot 10^{-1}, 1.27)$
θ_J^{21}	Juvenile mortality rate	day^{-1}	$2.13 \cdot 10^{-8}$	$(0.456 \cdot 10^{-2})$
λ_J^{21}	Maturation rate of the juvenile	day^{-1}	$1.00 \cdot 10^{-1}$	
ξ_J	Ratio of juvenile individual filtering rate	Unitless	1.00	
β^{21}	Spores produced by death per infected individual	$10^3 \cdot \text{spore} \cdot \text{individual}^{-1} \cdot \text{day}^{-1}$	$3.00 \cdot 10$	
μ	Algae refilling rate	$10^6 \cdot \text{cell} \cdot L^{-1} \cdot \text{day}^{-1}$	$3.70 \cdot 10^{-1}$	
δ	Sampling rate	day^{-1}	$1.30 \cdot 10^{-2}$	
σ_S^{21}	Standard deviation of Brownian motion of susceptible adult	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	0	
σ_J^{21}	Standard deviation of Brownian motion of juvenile	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$3.07 \cdot 10^{-1}$	$(2.31 \cdot 10^{-1}, 3.81 \cdot 10^{-1})$
σ_F	Standard deviation of Brownian motion of alga	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$9.31 \cdot 10^{-7}$	$(0.460 \cdot 10^{-2})$
τ_S^{21}	Measurement dispersion for susceptible adult	Unitless	$1.34 \cdot 10$	$(6.77, 9.81 \cdot 10)$

TABLE S-13

Variables and parameter definitions and estimates of SRJF model for *D. dentifera*-only dynamics. The confidence interval is generated by the Monte Carlo Adjusted Profile.

Parameter	Definition	Unit	Value	CI
S^i	Susceptible host density	individual $\cdot L^{-1}$	Variable	
J^i	Juvenile host density	individual $\cdot L^{-1}$	Variable	
F	Alga density	$10^6 \cdot \text{cell} \cdot L^{-1}$	Variable	
r^i	Birth rate of juvenile	individual $\cdot 10^{-6} \cdot \text{cell}^{-1}$	$9.77 \cdot 10^2$	$(8.86 \cdot 10, \infty)$
f_S^i	Susceptible adult host filtering rate	$L \cdot \text{individual}^{-1} \cdot \text{day}^{-1}$	$2.67 \cdot 10^{-4}$	$(0, 1.34 \cdot 10^{-3})$
$r^i \cdot f_{S^i}$	Effective Birth rate of invasive juvenile	$10^{-6} \cdot L \cdot \text{cell}^{-1} \cdot \text{day}^{-1}$	$2.61 \cdot 10^{-1}$	$(9.04 \cdot 10^{-2}, 4.02)$
θ_S^i	Susceptible adult host mortality rate	day^{-1}	$5.99 \cdot 10^{-1}$	$(2.26 \cdot 10^{-1}, 1.40)$
θ_J^i	Juvenile mortality rate	day^{-1}	$2.98 \cdot 10^{-1}$	$(0, 4.58)$
λ_J^i	Maturation rate of the juvenile	day^{-1}	$1.00 \cdot 10^{-1}$	
ξ_J^i	Ratio of juvenile individual filtering rate	Unitless	1.00	
β^i	Spores produced by death per infected individual	$10^3 \cdot \text{spore} \cdot \text{individual}^{-1} \cdot \text{day}^{-1}$	$3.00 \cdot 10$	
μ	Algae refilling rate	$10^6 \cdot \text{cell} \cdot L^{-1} \cdot \text{day}^{-1}$	$3.70 \cdot 10^{-1}$	
δ	Sampling rate	day^{-1}	$1.30 \cdot 10^{-2}$	
σ_S^i	Standard deviation of Brownian motion of susceptible adult	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	0	
σ_J^i	Standard deviation of Brownian motion of juvenile	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$3.78 \cdot 10^{-1}$	$(0, 5.07 \cdot 10^{-1})$
σ_F^i	Standard deviation of Brownian motion of alga	$\sqrt{\text{individual} \cdot \text{day}^{-1}}$	$4.73 \cdot 10^{-2}$	$(0, 1.38 \cdot 10^{-1})$
τ_S^i	Measurement dispersion for susceptible adult	Unitless	2.35	$(1.32, 5.72)$

TABLE S-14

Variables and parameter definitions and estimates of SRJF model for *D. lumholtzi*-only dynamics. The confidence interval is generated by the Monte Carlo Adjusted Profile.




Figure S-22: Monte Carlo Adjusted Profile results of SRJF model for *D. dentifera*-only dynamics. The vertical dotted lines represent the 95% confidence interval obtained by MCAP. The vertical blue lines show the MLE estimated using MCAP. The red vertical lines correspond to the model with the overall highest likelihood among all searches.




Figure S-23: Monte Carlo Adjusted Profile results of SRJF model for *D. lumholtzi*-only dynamics. The vertical dotted lines represent the 95% confidence interval obtained by MCAP. The vertical blue lines show the MLE estimated using MCAP. The red vertical lines correspond to the model with the overall highest likelihood among all searches.

Specific parameters	Max log-likelihood	AIC	Max log-likelihood (block)	AIC (block)	LRT p -value (χ^2)	LRT p -value (bootstrap)
$\emptyset$	-498.78	1011.56	-498.78	1011.56	—	—
$\theta_{S,u}^n$	-490.57	1013.14	-489.85	1011.71	0.06	—
r_u^n	-497.43	1026.85	-496.44	1024.89	0.97	—
$f_{S,u}^n$	-497.43	1026.86	-496.16	1024.32	0.98	—

TABLE S-15

Comparison of model fit and complexity across various configurations of unit-specific parameters within the panelPOMP framework for *D. dentifera*-only dynamics. The unit-specific parameter setting assessed include $f_{S,u}^n, r_u^n, \theta_{S,u}^n$.

Specific parameters	Max log-likelihood	AIC	Max log-likelihood (block)	AIC (block)	LRT p -value (χ^2)	LRT p -value (bootstrap)
$\emptyset$	-372.15	758.294	-372.15	758.294	—	—
$\theta_{S,u}^i$	-365.33	762.658	-365.37	762.733	0.14	—
r_u^i	-366.84	765.676	-366.29	764.571	0.30	—
$f_{S,u}^i$	-368.15	768.300	-368.07	768.147	0.53	—

TABLE S-16

Comparison of model fit and complexity across various configurations of unit-specific parameters within the panelPOMP framework for *D. lumholtzi*-only dynamics. The unit-specific parameter setting assessed include $f_{S,u}^i, r_u^i, \theta_{S,u}^i$.

S9.1. Results. The data plots, parameter estimates, and profile likelihood plots are analogous to the results in the main text, applied here to the SRJF model for either *D. dentifera* or *D. lumholtzi*.

The results in Tables S-15 and S-16 indicate that the model configuration with all shared parameters achieves the minimum AIC score for both dynamics, suggesting that this model makes a successful balance between model fit and complexity. The LRT p -values further support this choice, as none of the unit-specific models show a statistically significant improvement over the all-shared baseline. These results are consistent with our other analysis.

S9.2. Simulation. The simulation plots (Figures S-24 to S-27) collectively demonstrate the SRPF model's capability to capture the complex interactions among various states within the population dynamics of *D. dentifera* and *D. lumholtzi*. Specifically, the model effectively reproduces the latent patterns for susceptible and infected densities across multiple experimental units, as evidenced by the close alignment between the simulated mean trajectories (depicted by the black and light-blue bands) and the observed experimental data (represented by colorful solid lines). Each blue line illustrates an individual simulation case, capturing the stochastic variability across replicates, while the light blue shaded area (where present) denotes the 95% confidence interval of the simulation results. Although juvenile density data were not included in the model fitting process, the model still accurately predicts juvenile density trends (Figures S-27). This performance indicates the model's capacity to generalize underlying ecological processes, as the 95% confidence intervals largely encompass the observed juvenile data. The model also predicts a decline in algal density as the experiment progresses. This predicted reduction in food availability provides a mechanistic explanation for the observed trends in *Daphnia* densities, as resource scarcity constrains population growth, ultimately affecting the susceptible, infected, and juvenile densities.




Figure S-24: Simulation plots of densities (Individuals/Liter) for susceptible, and juvenile *D. dentifera*, along with alga density (10^6 cells/Liter) and parasite density (10^3 spores/Liter), over time (days). The colorful solid lines represent observed experimental data, showing the variability across different experimental replicates. The black line represents the mean trajectory from the simulation model, capturing the general trend of each density. The light blue shaded area corresponds to the 95% confidence interval of the simulations, indicating the range of model predictions. Dashed lines illustrate individual simulated trajectories, highlighting the model's ability to capture fluctuations around the mean trajectory.




Figure S-25: Simulated densities (Individuals/Liter) of susceptible *D. dentifera*, juvenile *D. dentifera* and alga density (10^6 cells/Liter) over time (days) for each experimental unit. The blue lines represent individual simulation runs, capturing the variability in susceptible density across replicates, while the red line represents the actual experiment data.




Figure S-26: Simulation plots of densities (Individuals/Liter) for susceptible, and juvenile *D. lumholtzi*, along with alga density (10^6 cells/Liter) and parasite density (10^3 spores/Liter), over time (days). The colorful solid lines represent observed experimental data, showing the variability across different experimental replicates. The black line represents the mean trajectory from the simulation model, capturing the general trend of each density. The light blue shaded area corresponds to the 95% confidence interval of the simulations, indicating the range of model predictions. Dashed lines illustrate individual simulated trajectories, highlighting the model's ability to capture fluctuations around the mean trajectory.




Figure S-27: Simulated densities (Individuals/Liter) of susceptible *D. lumholtzi* and juvenile *D. lumholtzi*, with alga density ($10^6 \cdot \text{cells/Liter}$), for each experimental unit. The blue lines represent individual simulation runs, capturing the variability in susceptible density across replicates, while the red line represents the actual experiment data.

S10. Benchmark Models.

S10.1. Searle et al.'s Model. Using the mechanistic model introduced by [Searle et al. \(2016\)](#), we conduct predictions of population $\tilde{y}_{u,n}$ for susceptible, infected *D. dentifera* and *D. lumholtzi* for all time $n \in \{1, 2, \dots, N_u\}$ and units $u \in \{1, 2, \dots, U\}$. In our case, each unit is observed at the same number of time points, so $N_u = N$ for all u . Then, we obtain the overall log-likelihood by summing the log-probabilities of the observed density of *Daphnia*. Specifically, we use the observed densities $y_{u,n}^*$ and the predicted means $\tilde{y}_{u,n}$ derived from the mechanistic model, along with the dispersion parameter τ to specify a NBinom distribution and calculate the likelihood:

$$(S16) \quad \ell(\beta, \theta) = \sum_{u=1}^U \sum_{n=1}^N \log P(Y_{u,n} = y_{u,n}^* \mid \mu_{u,n}, \tau),$$

To obtain the maximum likelihood estimate, we conduct the linear optimization on τ and choose the $\hat{\tau}$ that yields the highest overall likelihood. With this $\hat{\tau}$, we obtain the likelihood by [S16](#) for the model on the data. And the result of the model fitting can be found in the Table 2 in the main article.

S10.2. Negative Binomial Regression. Count data from ecological studies often exhibit overdispersion and unit-specific variation. The NBinom distribution addresses overdispersion, while random intercepts allow each unit to have its own baseline level. As a benchmark model, we employ NBinom generalized linear mixed models (GLMMs) to capture overdispersed count data and correlation across repeated measures within each unit u . The NBinom distribution is parameterized by a mean $\mu_{u,n}$ and a dispersion parameter θ .

We present three GLMMs with increasing of order of time factor to capture the trend of these *Daphnia* species in the experiment. While a linear model is easy to interpret and straightforward to implementing, observed *Daphnia* data exhibit a non-linear pattern. Introducing quadratic or cubic terms to the model enhance the capability of model to capture these

non-linear trends in the data. By starting with a linear model and then considering quadratic and cubic cases, we can evaluate the complexity of temporal patterns and better approximate the latent process of in the dynamics.

S10.2.1. Model 1 (Linear Trend Model). We start with a simple GLMM having only with linear terms:

$$(S17) \quad Y_{u,n} \sim \text{NBinom}(\mu_{u,n}, \tau),$$

$$(S18) \quad \log(\mu_{u,n}) = \beta_0 + \beta_1 t_{u,n} + b_u,$$

where $\mu_{u,n}$ is the expected count for unit u at time n . We fit this model independently for each *Daphnia* species, life stage, and infection status, and so the total log-likelihood is the sum over these categories. The vector coefficient β_1 measures the effect of time on the log-scale of *Daphnia* density. The term b_u accounts for the random intercept drawn from a normal distribution with variance σ^2 . Conditional on the random intercept b_u , observations within a unit are assumed independent. The dispersion parameter τ from the NBinom distribution accommodates overdispersion in the count data, which is density in our case.

S10.2.2. Model 2 (Quadratic Trend Model). To accommodate the non-linear pattern, we include a quadratic term in time in model S20:

$$(S19) \quad Y_{u,n} \sim \text{NBinom}(\mu_{u,n}, \tau),$$

$$(S20) \quad \log(\mu_{u,n}) = \beta_0 + \beta_1 t_{u,n} + \beta_2 t_{u,n}^2 + b_u.$$

Here, β_2 captures curvature in the trend, allowing for a turning point as time progresses.

S10.2.3. Model 3 (Cubic Trend Model). To capture more complex temporal pattern, we further introduce the cubic term in the model S22:

$$(S21) \quad Y_{u,n} \sim \text{NBinom}(\mu_{u,n}, \tau),$$

$$(S22) \quad \log(\mu_{u,n}) = \beta_0 + \beta_1 t_{u,n} + \beta_2 t_{u,n}^2 + \beta_3 t_{u,n}^3 + b_u.$$

The cubic term β_3 provides additional capability for capturing multiple fluctuation patterns in the *Daphnia* dynamics.

The likelihood for these models is constructed by considering the joint distribution of all observations $\{Y_{u,n}\}$, integrating over the random effects, $b_{1:U}$. Letting $\beta = (\beta_0, \beta_1, \dots)$ denote the fixed-effect parameters, the conditional likelihood of the observations for unit u given b_u is

$$(S23) \quad L_u(\beta, \tau, \sigma^2) = \prod_{n=1}^N f_{\text{NB}}(Y_{u,n}; \mu_{u,n}(\beta, \sigma^2), \tau),$$

where $f_{\text{NB}}(\cdot; \mu, \tau)$ is the NBinom probability mass function and $\mu_{u,n}(\beta, b_u) = \exp(\beta_0 + \beta_1 t_{u,n} + \dots + b_u)$. The total likelihood is then the product over all units due to the independence:

$$(S24) \quad L(\beta, \tau, \sigma^2) = \prod_{u=1}^U L_u(\beta, \tau, \sigma^2)$$

$$(S25) \quad = \prod_{u=1}^U \prod_{n=1}^N f_{\text{NB}}(Y_{u,n}; \mu_{u,n}(\beta, \sigma^2), \tau)$$

and the log-likelihood is:

$$(S26) \quad \ell(\beta, \tau, \sigma^2) = \sum_{u=1}^U \sum_{n=1}^N \log(f_{\text{NB}}(Y_{u,n}; \mu_{i,j}(\beta, \sigma^2), \tau)).$$

The calculated log-likelihood used to compute the AIC for model comparison. And the result of the model fitting of model S18, S20 and S22 can be found in the Table 2 in the main article.

S10.3. SIRPF2 Model. The SIRPF2 model presented here ignores the state of juvenile *Daphnia* and describes the dynamics in each bucket (u) among susceptible individuals ($S^{\mathbb{k}}$), infected individuals ($I^{\mathbb{k}}$), alga as a food resource (F), the parasite population (P). The superscript $\mathbb{k}$ denotes the species, which represents native or invasive species. Each stochastic differential equation accounts for various biological and ecological processes, augmented with stochastic terms to capture the inherent randomness within the systems.

$$(S27) \quad dS_u^{\mathbb{k}}(t) = r_u^{\mathbb{k}} f_{S,u}^{\mathbb{k}} F_u(t) S_u^{\mathbb{k}}(t) dt - \{\theta_{S,u}^{\mathbb{k}} + p_u^{\mathbb{k}} f_{S,u}^{\mathbb{k}} P_u(t) + \delta\} S_u^{\mathbb{k}}(t) dt,$$

$$(S28) \quad dI_u^{\mathbb{k}}(t) = p_u^{\mathbb{k}} f_{S,u}^{\mathbb{k}} S_u^{\mathbb{k}}(t) P_u(t) dt - \{\theta_{I,u}^{\mathbb{k}} + \delta\} I_u^{\mathbb{k}}(t) dt + I_u^{\mathbb{k}}(t) d\zeta_{I,u}^{\mathbb{k}},$$

$$(S29) \quad dP_u(t) = \sum_{\mathbb{k} \in \{\mathbf{n}, \mathbf{i}\}} \left(\beta_u^{\mathbb{k}} \theta_{I,u}^{\mathbb{k}} I_u^{\mathbb{k}}(t) - f_{S,u}^{\mathbb{k}} \{S_u^{\mathbb{k}}(t) + \xi_u I_u^{\mathbb{k}}(t)\} P_u(t) \right) dt \\ - \theta_{P,u} P_u(t) dt + P_u(t) d\zeta_{P,u},$$

$$(S30) \quad dF_u(t) = - \sum_{\mathbb{k} \in \{\mathbf{n}, \mathbf{i}\}} f_{S,u}^{\mathbb{k}} F_u(t) \left(S_u^{\mathbb{k}}(t) + \xi_u I_u^{\mathbb{k}}(t) \right) dt + \mu dt + F_u(t) d\zeta_{F,u},$$

$$(S31) \quad d\zeta_{I,u}^{\mathbb{k}} \sim N[0, (\sigma_{I,u}^{\mathbb{k}})^2 dt], \quad d\zeta_{F,u} \sim N[0, \sigma_{F,u}^2 dt], \quad d\zeta_{P,u} \sim N[0, \sigma_{P,u}^2 dt].$$

Unlike the SIRJPF2 model, we ignore the maturation process from juvenile to adult in equation (S28). Instead, this model uses $r_u^{\mathbb{k}}$ to represent the average rate of growth in density of *Daphnia* adult.

S10.3.1. Flow Diagram. The SIRPF2 model illustrates the interdependent dynamics among the Susceptible (S), Infected (I), Food (F), and Parasite (P) populations, with transitions representing ecological and biological interactions.




Figure S-28: Flow diagram for the SIRPF2 model illustrating population interactions.

Mesocosm	<i>D. dentifera</i> juvenile	<i>D. dentifera</i> female adult	<i>D. lumholtzi</i> juvenile	<i>D. lumholtzi</i> female adult	<i>D. lumholtzi</i> male adult
1	22	22	22	27	27
2	22	32	17	17	22
3	27	32	27	22	37
4	22	22	37	17	27
5	22	22	17	32	27
6	32	27	27	32	42
7	27	22	17	17	27
8	27	32	22	22	42

TABLE S-17

This table shows the day on which each species reached its peak density in mesocosms containing both *Daphnia* species and the parasite. *D. dentifera* male adults are excluded because they were almost absent.

S11. Experimental Data on Peak Density. Table S-17 provides supporting data on the relative timing of peak abundance for different demographic groups.

S12. SIRJPF2-Gamma Model. The SIRJPF2-Gamma model differs from the original SIRJPF2 model by its incorporation of gamma noise in transition processes between states, in place of the Gaussian noise used previously. Specifically, in the SIRJPF2-Gamma model, random fluctuations for transitions between states are explained by gamma distributions to represent ecological events occur through one-way transitions which are nonnegative. The use of gamma noise in modeling transitions explicitly enforces that individuals move from one class to another without the artificial creation or destruction of hosts that do not transit to or from another compartment. This aligns with the underlying dynamics that the jump-like transitions observed in ecosystem reflect one-directional movements rather than symmetrical fluctuations.

$$(S32) \quad dS_u^k(t) = \lambda_{J,u}^k J_u^k(t) dt - \{\theta_{S,u}^k + \delta\} S_u^k(t) dt - p_u^k f_{S,u}^k P_u(t) S_u^k(t) d\Gamma_{SI,u}^k,$$

$$(S33) \quad dI_u^k(t) = p_u^k f_{S,u}^k S_u^k(t) P_u(t) d\Gamma_{SI,u}^k - \theta_{I,u}^k I_u^k(t) d\Gamma_{IP,u}^k - \delta I_u^k(t) dt,$$

$$(S34) \quad dJ_u^k(t) = r_u^k f_{S,u}^k F_u(t) S_u^k(t) d\Gamma_{SJ,u}^k - \{\theta_{J,u}^k + \delta\} J_u^k(t) dt - \lambda_{J,u}^k J_u^k(t) dt,$$

$$(S35) \quad dP_u(t) = \sum_{k \in \{n, i\}} \left(\beta_u^k \theta_{I,u}^k I_u^k(t) d\Gamma_{IP,u}^k - f_{S,u}^k \{S_u^k(t) + \xi_u I_u^k(t)\} P_u(t) dt \right)$$

$$- \theta_{P,u} P_u(t) dt,$$

$$(S36) \quad dF_u(t) = - \sum_{k \in \{n, i\}} f_{S,u}^k F_u(t) \left(S_u^k(t) + \xi_{J,u} J_u^k(t) + \xi_u I_u^k(t) \right) dt + \mu dt + F_u(t) d\Gamma_{F,u},$$

$$(S37) \quad d\Gamma_{JS,u}^k \sim \Gamma \left[\frac{dt}{(\sigma_{JS,u}^k)^2}, (\sigma_{JS,u}^k)^2 \right], \quad d\Gamma_{SI,u}^k \sim \Gamma \left[\frac{dt}{(\sigma_{SI,u}^k)^2}, (\sigma_{SI,u}^k)^2 \right],$$

$$(S38) \quad d\Gamma_{IP,u}^k \sim \Gamma \left[\frac{dt}{(\sigma_{IP,u}^k)^2}, (\sigma_{IP,u}^k)^2 \right], \quad d\Gamma_{F,u} \sim N[0, \sigma_{F,u}^2 dt].$$

REFERENCES

Bretó, C., Ionides, E. L., and King, A. A. (2020). Panel data analysis via mechanistic models. *Journal of the American Statistical Association*, 115(531):1178–1188.

- Bretó, C., Wheeler, J., King, A. A., and Ionides, E. L. (2025). panelPomp: Analysis of panel data via partially observed Markov processes in R. *The R Journal*, 17:180–199.
- Ionides, E. L., Bretó, C., Park, J., Smith, R. A., and King, A. A. (2017). Monte Carlo profile confidence intervals for dynamic systems. *Journal of the Royal Society Interface*, 14:1–10.
- Ning, N., Ionides, E. L., and Ritov, Y. (2021). Scalable Monte Carlo inference and rescaled local asymptotic normality. *Bernoulli*, 27:2532–2555.
- Searle, C. L., Cortez, M. H., Hunsberger, K. K., Grippi, D. C., Oleksy, I. A., Shaw, C. L., de la Serna, S. B., Lash, C. L., Dhir, K. L., and Duffy, M. A. (2016). Population density, not host competence, drives patterns of disease in an invaded community. *The American Naturalist*, 188(5):554–566.
- Stocks, T., Britton, T., and Höhle, M. (2020). Model selection and parameter estimation for dynamic epidemic models via iterated filtering: Application to rotavirus in Germany. *Biostatistics*, 21(3):400–416.
- Wheeler, J., Abkemeier, A. J., and Ionides, E. L. (2025). Iterating marginalized bayes maps for likelihood maximization with application to nonlinear panel models. *arXiv:2511.17438*.