

15-puzzle

Alexander Shen

LIRMM / CNRS, University of Montpellier

Outline

The Game


Permutations

Proof: The Difficult Part

Mission Impossible

Classify a Permutation

15-Puzzle



By Micha L. Rieser, <https://commons.wikimedia.org/w/index.php?curid=3104433>

Game Rules

Game Rules

- move pieces (fill the empty square moving one of the neighbors)

Game Rules

- move pieces (fill the empty square moving one of the neighbors)
- goal: to reach some configuration, e.g., the initial configuration with 14 and 15 interchanged

Game Rules

- move pieces (fill the empty square moving one of the neighbors)
- goal: to reach some configuration, e.g., the initial configuration with 14 and 15 interchanged
- reverse: return to the standard position from a given one

Game Rules

- move pieces (fill the empty square moving one of the neighbors)
- goal: to reach some configuration, e.g., the initial configuration with 14 and 15 interchanged
- reverse: return to the standard position from a given one
- try yourself: if you succeed, we pay you \$1000

History

History

- invented in nineteenth century

History

- invented in nineteenth century
- complicated history with conflicting claims (Wikipedia)

History

- invented in nineteenth century
- complicated history with conflicting claims (Wikipedia)
- was popular at some time, and cash prizes were announced

History

- invented in nineteenth century
- complicated history with conflicting claims (Wikipedia)
- was popular at some time, and cash prizes were announced
- why did the organizers take the risk?

History

- invented in nineteenth century
- complicated history with conflicting claims (Wikipedia)
- was popular at some time, and cash prizes were announced
- why did the organizers take the risk?
- a *proof* of impossibility existed

History

- invented in nineteenth century
- complicated history with conflicting claims (Wikipedia)
- was popular at some time, and cash prizes were announced
- why did the organizers take the risk?
- a *proof* of impossibility existed
- challenge: reinvent this proof

Outline

The Game

Permutations

Proof: The Difficult Part

Mission Impossible

Classify a Permutation

Another View of the Game

Another View of the Game

- empty cell is active: it moves around, exchanging places with the neighbors

Another View of the Game

- empty cell is active: it moves around, exchanging places with the neighbors
- more general question: permutations of n objects obtained by a sequence of pair exchanges (=transpositions)

Another View of the Game

- empty cell is active: it moves around, exchanging places with the neighbors
- more general question: permutations of n objects obtained by a sequence of pair exchanges (=transpositions)
- STOP \rightarrow SPOT: one transposition is enough

Another View of the Game

- empty cell is active: it moves around, exchanging places with the neighbors
- more general question: permutations of n objects obtained by a sequence of pair exchanges (=transpositions)
- STOP \rightarrow SPOT: one transposition is enough
- STOP \rightarrow POST: how many transpositions needed?

Another View of the Game

- empty cell is active: it moves around, exchanging places with the neighbors
- more general question: permutations of n objects obtained by a sequence of pair exchanges (=transpositions)
- STOP \rightarrow SPOT: one transposition is enough
- STOP \rightarrow POST: how many transpositions needed?
- cat and permutations: <https://www.youtube.com/watch?v=MDhsT0nd3J8>

Even and Odd Permutations

Even and Odd Permutations

- STOP \rightarrow SPOT: 1, 3, 5, 7 ...

Even and Odd Permutations

- STOP \rightarrow SPOT: 1, 3, 5, 7 ...
- STOP \rightarrow POST: 3, 5, 7, ...

Even and Odd Permutations

- STOP \rightarrow SPOT: 1, 3, 5, 7 ...
- STOP \rightarrow POST: 3, 5, 7, ...
- STOP \rightarrow POTS: 2, 4, 6, ...

Even and Odd Permutations


- $\text{STOP} \rightarrow \text{SPOT}: 1, 3, 5, 7 \dots$
- $\text{STOP} \rightarrow \text{POST}: 3, 5, 7, \dots$
- $\text{STOP} \rightarrow \text{POTS}: 2, 4, 6, \dots$
- $n \rightarrow n + 2$: transposition: twice = nothing

Even and Odd Permutations

- STOP \rightarrow SPOT: 1, 3, 5, 7 ...
- STOP \rightarrow POST: 3, 5, 7, ...
- STOP \rightarrow POTS: 2, 4, 6, ...
- $n \rightarrow n + 2$: transposition: twice = nothing
- conjecture: permutations are of two types: even or odd


A Counterexample?

TOTEM



MOTET

TOTEM




TOMET



MOTET


A Counterexample?

TOTEM



MOTET

TOTEM



TOMET




MOTET

even and odd at the same time?


A Counterexample?

TOTEM



MOTET

TOTEM



TOMET




MOTET

even and odd at the same time?

spoiler:


A Counterexample?

TOTEM



MOTET

TOTEM



TOMET



MOTET

even and odd at the same time?

spoiler: two T's are mixed (we assumed that all letters are different)

Theorem

Theorem

- each permutation can be obtained by transpositions

Theorem

- each permutation can be obtained by transpositions
- some permutations can be obtained only by even number of transpositions, while others can be obtained only by odd number of transpositions

Proof: The Easy Part

Proof: The Easy Part

- claim: each permutation can be obtained by transpositions

Proof: The Easy Part

- claim: each permutation can be obtained by transpositions
- proof: put letters to the right places one by one using one transposition per letter

Proof: The Easy Part

- claim: each permutation can be obtained by transpositions
- proof: put letters to the right places one by one using one transposition per letter

STOP \rightarrow POST

Proof: The Easy Part

- claim: each permutation can be obtained by transpositions
- proof: put letters to the right places one by one using one transposition per letter

STOP \rightarrow POST

STOP

Proof: The Easy Part

- claim: each permutation can be obtained by transpositions
- proof: put letters to the right places one by one using one transposition per letter

STOP \rightarrow POST

S^TOP

Proof: The Easy Part

- claim: each permutation can be obtained by transpositions
- proof: put letters to the right places one by one using one transposition per letter

STOP \rightarrow POST

S~~T~~OP

PTOS

Proof: The Easy Part

- claim: each permutation can be obtained by transpositions
- proof: put letters to the right places one by one using one transposition per letter

STOP \rightarrow POST

S**T**OP

P**T**OS

Proof: The Easy Part

- claim: each permutation can be obtained by transpositions
- proof: put letters to the right places one by one using one transposition per letter

STOP \rightarrow POST

STOP

POTS

POTS

Proof: The Easy Part

- claim: each permutation can be obtained by transpositions
- proof: put letters to the right places one by one using one transposition per letter

STOP \rightarrow POST

S**T**OP

P**T**OS

P**O**TS

Proof: The Easy Part

- claim: each permutation can be obtained by transpositions
- proof: put letters to the right places one by one using one transposition per letter

STOP \rightarrow POST

STOP

PTOS

POTS

POST

Outline

The Game

Permutations

Proof: The Difficult Part

Mission Impossible

Classify a Permutation

Even Is Not Odd

Even Is Not Odd

- second claim: the same permutation cannot be even and odd at the same time

Even Is Not Odd

- second claim: the same permutation cannot be even and odd at the same time
- special case: *an identity permutation* (no changes) *cannot be obtained by odd number of transpositions*

Even Is Not Odd

- second claim: the same permutation cannot be even and odd at the same time
- special case: *an identity permutation* (no changes) *cannot be obtained by odd number of transpositions*
- reduction to this special case:

Even Is Not Odd

- second claim: the same permutation cannot be even and odd at the same time
- special case: *an identity permutation* (no changes) *cannot be obtained by odd number of transpositions*
- reduction to this special case:
 $A \rightarrow \dots \rightarrow B$ (even), $A \rightarrow \dots \rightarrow B$ (odd).

Even Is Not Odd

- second claim: the same permutation cannot be even and odd at the same time
- special case: *an identity permutation* (no changes) *cannot be obtained by odd number of transpositions*
- reduction to this special case:
 $A \rightarrow \dots \rightarrow B$ (even), $A \rightarrow \dots \rightarrow B$ (odd).
Back and forth: even + odd = odd number of transpositions brings us back;

Even Is Not Odd

- second claim: the same permutation cannot be even and odd at the same time
- special case: *an identity permutation (no changes) cannot be obtained by odd number of transpositions*
- reduction to this special case:
 $A \rightarrow \dots \rightarrow B$ (even), $A \rightarrow \dots \rightarrow B$ (odd).
Back and forth: even + odd = odd number of transpositions brings us back; not possible if we believe in the special case

Transpositions of Neighbors

Transpositions of Neighbors

- claim: after odd number of *neighbor* transpositions we cannot return to original position

Transpositions of Neighbors

- claim: after odd number of *neighbor* transpositions we cannot return to original position
- look at every pair of letters. Why the number of transpositions for this pair is even?

Transpositions of Neighbors


- claim: after odd number of *neighbor* transpositions we cannot return to original position
- look at every pair of letters. Why the number of transpositions for this pair is even?
- easy: each transposition of this pair changes the order (who is on the left).

Transpositions of Neighbors


- claim: after odd number of *neighbor* transpositions we cannot return to original position
- look at every pair of letters. Why the number of transpositions for this pair is even?
- easy: each transposition of this pair changes the order (who is on the left).
- note that *neighbor* transposition does not change order in other pairs

Reduction to the Neighbor Case

Reduction to the Neighbor Case



Reduction to the Neighbor Case



$$\begin{aligned} k + 1 + k &= 2k + 1 \text{ neighbor transpositions} \\ &= 1 \pmod{2} \end{aligned}$$

Where We Are

Where We Are

- in 15-puzzle one cannot exchange 14 and 15. Why?

Where We Are

- in 15-puzzle one cannot exchange 14 and 15. Why?
- generalization: permutations, transpositions

Where We Are

- in 15-puzzle one cannot exchange 14 and 15. Why?
- generalization: permutations, transpositions
- classification theorem: some permutations need even number of transpositions, other require an odd number

Where We Are

- in 15-puzzle one cannot exchange 14 and 15. Why?
- generalization: permutations, transpositions
- classification theorem: some permutations need even number of transpositions, other require an odd number
- OK — so what?

Outline

The Game

Permutations

Proof: The Difficult Part

Mission Impossible

Classify a Permutation

Back to 15-puzzle

Back to 15-puzzle

- why we cannot exchange 14 and 15 in the puzzle?

Back to 15-puzzle

- why we cannot exchange 14 and 15 in the puzzle?
- each move is a transposition of an empty cell and some its neighbor

Back to 15-puzzle

- why we cannot exchange 14 and 15 in the puzzle?
- each move is a transposition of an empty cell and some its neighbor
- the required exchange requires an *odd* number of moves

Back to 15-puzzle

- why we cannot exchange 14 and 15 in the puzzle?
- each move is a transposition of an empty cell and some its neighbor
- the required exchange requires an *odd* number of moves
- but for other reasons it requires an *even* number of moves

Back to 15-puzzle

- why we cannot exchange 14 and 15 in the puzzle?
- each move is a transposition of an empty cell and some its neighbor
- the required exchange requires an *odd* number of moves
- but for other reasons it requires an *even* number of moves (why?)

Back to 15-puzzle


- why we cannot exchange 14 and 15 in the puzzle?
- each move is a transposition of an empty cell and some its neighbor
- the required exchange requires an *odd* number of moves
- but for other reasons it requires an *even* number of moves (why?)
- so the required exchange is impossible

Chessboard Helps Again

To bring back the empty cell we need an even number of moves. Why?


Chessboard Helps Again

To bring back the empty cell we need an even number of moves. Why?



Chessboard Helps Again


To bring back the empty cell we need an even number of moves. Why?



each move changes the color of the empty cell.

Chessboard Helps Again


To bring back the empty cell we need an even number of moves. Why?



each move changes the color of the empty cell.
bringing it back requires an even number of
moves.

Chessboard Helps Again

To bring back the empty cell we need an even number of moves. Why?



each move changes the color of the empty cell.
bringing it back requires an even number of
moves. **No risk to lose money!**

Outline

The Game

Permutations

Proof: The Difficult Part

Mission Impossible

Classify a Permutation

Programming Assignment

An array $a[1], \dots, a[n]$ is given, containing a permutation of numbers $1, \dots, n$.

Output 0 if the permutation is even (requires an even number of transpositions) and 1 otherwise.

A Possible Approach

A Possible Approach

- Do you remember a situation when a permutation is obtained by transpositions?

A Possible Approach

- Do you remember a situation when a permutation is obtained by transpositions?
- Sorting

A Possible Approach

- Do you remember a situation when a permutation is obtained by transpositions?
- Sorting (if done by exchanges)

A Possible Approach

- Do you remember a situation when a permutation is obtained by transpositions?
- Sorting (if done by exchanges)
- Hint: sort a and count the number of exchanges

A Possible Approach

- Do you remember a situation when a permutation is obtained by transpositions?
- Sorting (if done by exchanges)
- Hint: sort a and count the number of exchanges
- ...to see if this number is even or odd

Implementation?

```
// sorting a[1]..a[n]
sign=0 // sign = the number of transpositions mod 2
s=0 // first s elements are at the right places
while (s<n){
    u=s+1; t=u; //a[t] is minimal among a[s+1]..a[u];
    while (u<n){
        u=u+1;
        if a[u]<a[t] {t=u;}
    }
    // a[t] is minimal among a[s+1]..a[n]
    tmp=a[s+1]; a[s+1]=a[t]; a[t]=tmp; sign=1-sign;
}
```

Implementation?

```
// sorting a[1]..a[n]
sign=0 // sign = the number of transpositions mod 2
s=0 // first s elements are at the right places
while (s<n){
    u=s+1; t=u; //a[t] is minimal among a[s+1]..a[u];
    while (u<n){
        u=u+1;
        if a[u]<a[t] {t=u;}
    }
    // a[t] is minimal among a[s+1]..a[n]
    tmp=a[s+1]; a[s+1]=a[t]; a[t]=tmp; sign=1-sign;
}
```

What is wrong with this code?

Bonus Track

Bonus Track

- Running time (number of steps) for this approach?

Bonus Track

- Running time (number of steps) for this approach?
- $O(n^2)$ (two nested loops)

Bonus Track

- Running time (number of steps) for this approach?
- $O(n^2)$ (two nested loops)
- More efficient algorithm?

Bonus Track

- Running time (number of steps) for this approach?
- $O(n^2)$ (two nested loops)
- More efficient algorithm?
- Hint: use $O(n \log n)$ sorting to get $O(n \log n)$ algorithm

Bonus Track

- Running time (number of steps) for this approach?
- $O(n^2)$ (two nested loops)
- More efficient algorithm?
- Hint: use $O(n \log n)$ sorting to get $O(n \log n)$ algorithm
- Challenge: can you think of $O(n)$ algorithm?

Project

A position in 15-puzzle is given (the empty space is in the right corner, the other numbers are listed in the book order, so the standard position is represented as 1, 2, 3, 4, \dots , 14, 15). Find out whether the puzzle is solvable; if yes, give the sequence of moves (for each move, just write the number of a piece that is moving at that step).