

# Ramsey Numbers

---

Alexander Golovnev


# Outline

Balanced Graphs

Ramsey Numbers


Existence of Ramsey Numbers

# Balanced Graphs




# Balanced Graphs

The **Clique Number** and the **Independence Number** of  $C_5$  are 2.




# Balanced Graphs



# Balanced Graphs

Consider an arbitrary vertex  $v$  of  $G$




# Balanced Graphs

Consider an arbitrary vertex  $v$  of  $G$

Either  $v$  is connected to  $\geq 3$  vertices

Or  $v$  is not connected to  $\geq 3$  vertices




# Balanced Graphs

Consider an arbitrary vertex  $v$  of  $G$

Either  $v$  is connected to  $\geq 3$  vertices

Or  $v$  is not connected to  $\geq 3$  vertices






# Balanced Graphs

Consider an arbitrary vertex  $v$  of  $G$

Either  $v$  is connected to  $\geq 3$  vertices

Or  $v$  is not connected to  $\geq 3$  vertices




# Balanced Graphs

Consider an arbitrary vertex  $v$  of  $G$

Either  $v$  is connected to  $\geq 3$  vertices

Or  $v$  is not connected to  $\geq 3$  vertices




# Balanced Graphs

Consider an arbitrary vertex  $v$  of  $G$

Either  $v$  is connected to  $\geq 3$  vertices

Or  $v$  is not connected to  $\geq 3$  vertices




# Balanced Graphs

Consider an arbitrary vertex  $v$  of  $G$

Either  $v$  is connected to  $\geq 3$  vertices

Or  $v$  is not connected to  $\geq 3$  vertices




# Balanced Graphs

Consider an arbitrary vertex  $v$  of  $G$

Either  $v$  is connected to  $\geq 3$  vertices

Or  $v$  is not connected to  $\geq 3$  vertices




# Balanced Graphs

Consider an arbitrary vertex  $v$  of  $G$

Either  $v$  is connected to  $\geq 3$  vertices

Or  $v$  is not connected to  $\geq 3$  vertices



# Outline

Balanced Graphs

**Ramsey Numbers**

Existence of Ramsey Numbers

# Children Friendship

- 1950s, The Hungarian sociologist S. Szalai studies friendship between children



# Children Friendship

- 1950s, The Hungarian sociologist S. Szalai studies friendship between children
- Observes: in any group of 20 children, there are 4 mutual friends or 4 children s.t. no 2 are friends

# Children Friendship



# Children Friendship



# Children Friendship

- 1950s, A Hungarian sociologist S. Szalai studies friendship between children
- Observes: in any group of 20 children, there are 4 mutual friends or 4 children s.t. no 2 are friends

# Children Friendship

- 1950s, A Hungarian sociologist S. Szalai studies friendship between children
- Observes: in any group of 20 children, there are 4 mutual friends or 4 children s.t. no 2 are friends
- Consults Erdős, Turán and Sós

# Children Friendship

- 1950s, A Hungarian sociologist S. Szalai studies friendship between children
- Observes: in any group of 20 children, there are 4 mutual friends or 4 children s.t. no 2 are friends
- Consults Erdős, Turán and Sós
- This holds in any group of 18 and more people

# Ramsey Numbers

- For two integers  $k, \ell$ , the Ramsey Number  $R(k, \ell)$  is the minimum number, s.t. every graph with at least  $R(k, \ell)$  vertices must have

# Ramsey Numbers

- For two integers  $k, \ell$ , the Ramsey Number  $R(k, \ell)$  is the minimum number, s.t. every graph with at least  $R(k, \ell)$  vertices must have
  - either a clique of size  $k$



# Ramsey Numbers

- For two integers  $k, \ell$ , the Ramsey Number  $R(k, \ell)$  is the minimum number, s.t. every graph with at least  $R(k, \ell)$  vertices must have
  - either a clique of size  $k$
  - or an independent set of size  $\ell$

# Ramsey Numbers

- For two integers  $k, \ell$ , the **Ramsey Number**  $R(k, \ell)$  is the minimum number, s.t. every graph with at least  $R(k, \ell)$  vertices must have
  - either a clique of size  $k$
  - or an independent set of size  $\ell$
- $R(3, 3) = 6$

# Ramsey Numbers

- For two integers  $k, \ell$ , the **Ramsey Number**  $R(k, \ell)$  is the minimum number, s.t. every graph with at least  $R(k, \ell)$  vertices must have
  - either a clique of size  $k$
  - or an independent set of size  $\ell$
- $R(3, 3) = 6$
- $R(4, 4) = 18$

# Ramsey Numbers

- For two integers  $k, \ell$ , the **Ramsey Number**  $R(k, \ell)$  is the minimum number, s.t. every graph with at least  $R(k, \ell)$  vertices must have
  - either a clique of size  $k$
  - or an independent set of size  $\ell$
- $R(3, 3) = 6$
- $R(4, 4) = 18$
- $43 \leq R(5, 5) \leq 48$

# Outline

Balanced Graphs

Ramsey Numbers

Existence of Ramsey Numbers

## Does $R(k, \ell)$ make sense?

- Does  $R(k, \ell)$  **exist** for all values of  $k, \ell$ ?

## Does $R(k, \ell)$ make sense?

- Does  $R(k, \ell)$  **exist** for all values of  $k, \ell$ ?
- How do we know that **all** large graphs have either a large Clique or a large IS?

## Does $R(k, \ell)$ make sense?

- Does  $R(k, \ell)$  **exist** for all values of  $k, \ell$ ?
- How do we know that **all** large graphs have either a large Clique or a large IS?
- Ok,  $R(k, \ell)$  exists for  $k \leq 3$  and  $\ell \leq 3$



## Does $R(k, \ell)$ make sense?

- Does  $R(k, \ell)$  **exist** for all values of  $k, \ell$ ?
- How do we know that **all** large graphs have either a large Clique or a large IS?
- Ok,  $R(k, \ell)$  exists for  $k \leq 3$  and  $\ell \leq 3$
- We will prove  $R(k, \ell) \leq R(k-1, \ell) + R(k, \ell-1)$

## Does $R(k, \ell)$ make sense?

- Does  $R(k, \ell)$  **exist** for all values of  $k, \ell$ ?
- How do we know that **all** large graphs have either a large Clique or a large IS?
- Ok,  $R(k, \ell)$  exists for  $k \leq 3$  and  $\ell \leq 3$
- We will prove  $R(k, \ell) \leq R(k-1, \ell) + R(k, \ell-1)$
- This gives an upper bound on  $R(k, \ell)$  for all  $k, \ell$

## Does $R(k, \ell)$ make sense?

- Does  $R(k, \ell)$  **exist** for all values of  $k, \ell$ ?
- How do we know that **all** large graphs have either a large Clique or a large IS?
- Ok,  $R(k, \ell)$  exists for  $k \leq 3$  and  $\ell \leq 3$
- We will prove  $R(k, \ell) \leq R(k-1, \ell) + R(k, \ell-1)$
- This gives an upper bound on  $R(k, \ell)$  for all  $k, \ell$
- Therefore,  $R(k, \ell)$  **always exists!**

# Ramsey's Theorem

## Theorem (Ramsey's Theorem)

$$R(k, \ell) \leq R(k-1, \ell) + R(k, \ell-1).$$

# Ramsey's Theorem

## Theorem (Ramsey's Theorem)

$$R(k, \ell) \leq R(k-1, \ell) + R(k, \ell-1).$$

### Proof:

- Consider a graph  $G$  on  $R(k-1, \ell) + R(k, \ell-1)$  vertices

# Ramsey's Theorem

## Theorem (Ramsey's Theorem)

$$R(k, \ell) \leq R(k-1, \ell) + R(k, \ell-1).$$

### Proof:

- Consider a graph  $G$  on  $R(k-1, \ell) + R(k, \ell-1)$  vertices
- We'll prove  $G$  contains either a  $k$ -Clique or an  $\ell$ -Independent Set

# Ramsey's Theorem

## Theorem (Ramsey's Theorem)

$$R(k, \ell) \leq R(k - 1, \ell) + R(k, \ell - 1).$$

### Proof:

- Consider a graph  $G$  on  $R(k - 1, \ell) + R(k, \ell - 1)$  vertices
- We'll prove  $G$  contains either a  $k$ -Clique or an  $\ell$ -Independent Set
- Pick an arbitrary vertex  $v$ .  $A$  – set of neighbors of  $v$ ,  $B$  – the remaining vertices

# Ramsey's Theorem. Proof

- $A$  – neighbors of  $v$ ,  $B$  – not neighbors



# Ramsey's Theorem. Proof

- $A$  – neighbors of  $v$ ,  $B$  – not neighbors
- $|A| + |B| + 1 = R(k-1, \ell) + R(k, \ell-1)$

# Ramsey's Theorem. Proof

- $A$  – neighbors of  $v$ ,  $B$  – not neighbors
- $|A| + |B| + 1 = R(k-1, \ell) + R(k, \ell-1)$
- Either  $|A| \geq R(k-1, \ell)$  or  $|B| \geq R(k, \ell-1)$

# Ramsey's Theorem. Proof

- Either  $|A| \geq R(k-1, \ell)$  or  $|B| \geq R(k, \ell-1)$

# Ramsey's Theorem. Proof

- Either  $|A| \geq R(k-1, \ell)$  or  $|B| \geq R(k, \ell-1)$
- In the 1st case:

# Ramsey's Theorem. Proof

- Either  $|A| \geq R(k-1, \ell)$  or  $|B| \geq R(k, \ell-1)$
- In the 1st case:
  - Either  $A$  has an  $\ell$ -IS—done!

# Ramsey's Theorem. Proof

- Either  $|A| \geq R(k-1, \ell)$  or  $|B| \geq R(k, \ell-1)$
- In the 1st case:
  - Either  $A$  has an  $\ell$ -IS—done!
  - Or  $A \cup v$  has a  $k$ -Clique—done!

# Ramsey's Theorem. Proof

- Either  $|A| \geq R(k-1, \ell)$  or  $|B| \geq R(k, \ell-1)$
- In the 1st case:
  - Either  $A$  has an  $\ell$ -IS—done!
  - Or  $A \cup v$  has a  $k$ -Clique—done!
- In the 2nd case:

# Ramsey's Theorem. Proof


- Either  $|A| \geq R(k-1, \ell)$  or  $|B| \geq R(k, \ell-1)$
- In the 1st case:
  - Either  $A$  has an  $\ell$ -IS—done!
  - Or  $A \cup v$  has a  $k$ -Clique—done!
- In the 2nd case:
  - Either  $B$  has a  $k$ -clique—done!



# Ramsey's Theorem. Proof


- Either  $|A| \geq R(k-1, \ell)$  or  $|B| \geq R(k, \ell-1)$
- In the 1st case:
  - Either  $A$  has an  $\ell$ -IS—done!
  - Or  $A \cup v$  has a  $k$ -Clique—done!
- In the 2nd case:
  - Either  $B$  has a  $k$ -clique—done!
  - Or  $B \cup v$  has a  $\ell$ -IS—done!

# Ramsey's Theorem. Proof




# Ramsey's Theorem. Proof

If  $|A| \geq R(k-1, \ell)$ ,  
A contains a  
 $k-1$ -Clique



# Ramsey's Theorem. Proof


If  $|A| \geq R(k-1, \ell)$ ,  
A contains a  
 $k-1$ -Clique




# Ramsey's Theorem. Proof

If  $|A| \geq R(k-1, \ell)$ ,  
 $A$  contains a  
 $k-1$ -Clique

Then  $|A| \cup v$   
contains a  
 $k$ -Clique




# Ramsey's Theorem. Proof



If  $|B| \geq R(k, \ell - 1)$ ,  
 $B$  contains  
an  $\ell - 1$ -IS

# Ramsey's Theorem. Proof



If  $|B| \geq R(k, \ell - 1)$ ,  
 $B$  contains  
an  $\ell - 1$ -IS

Then  $|B| \cup v$   
contains an  
 $\ell$ -IS