

# Division by 2

---

Vladimir Podolskii

Computer Science Department, Higher School of Economics

# Outline

Division by 2

Binary System

# Division by 2

- Consider division of integers by 2

# Division by 2

- Consider division of integers by 2
- There are two possible remainders: 0 and 1

# Division by 2

- Consider division of integers by 2
- There are two possible remainders: 0 and 1
- If the remainder of  $a$  is 0, then  $a$  is divisible by 2

# Division by 2

- Consider division of integers by 2
- There are two possible remainders: 0 and 1
- If the remainder of  $a$  is 0, then  $a$  is divisible by 2
- These are even numbers

# Division by 2

- Consider division of integers by 2
- There are two possible remainders: 0 and 1
- If the remainder of  $a$  is 0, then  $a$  is divisible by 2
- These are **even numbers**
- If the remainder of  $a$  is 1,  $a$  is not divisible by 2

# Division by 2

- Consider division of integers by 2
- There are two possible remainders: 0 and 1
- If the remainder of  $a$  is 0, then  $a$  is divisible by 2
- These are **even numbers**
- If the remainder of  $a$  is 1,  $a$  is not divisible by 2
- These are **odd numbers**



# Sums of Even and Odd Numbers

## Splitting in pairs

Suppose there are two classes with  $a$  and  $b$  students respectively. We unite the classes and would like to split all students into pairs to work on a project. Is it possible if  $a$  is even and  $b$  is odd? What if both  $a$  and  $b$  are even? What if both  $a$  and  $b$  are odd?

# Sums of Even and Odd Numbers

## Splitting in pairs

Suppose there are two classes with  $a$  and  $b$  students respectively. We unite the classes and would like to split all students into pairs to work on a project. Is it possible if  $a$  is even and  $b$  is odd? What if both  $a$  and  $b$  are even? What if both  $a$  and  $b$  are odd?

- If  $a$  is even and  $b$  is odd we can split in pairs all students in the first class and all students except one in the second class

# Sums of Even and Odd Numbers

## Splitting in pairs

Suppose there are two classes with  $a$  and  $b$  students respectively. We unite the classes and would like to split all students into pairs to work on a project. Is it possible if  $a$  is even and  $b$  is odd? What if both  $a$  and  $b$  are even? What if both  $a$  and  $b$  are odd?

- If  $a$  is even and  $b$  is odd we can split in pairs all students in the first class and all students except one in the second class
- So there is one student left and the answer is no

# Sums of Even and Odd Numbers

## Splitting in pairs

Suppose there are two classes with  $a$  and  $b$  students respectively. We unite the classes and would like to split all students into pairs to work on a project. Is it possible if  $a$  is even and  $b$  is odd? What if both  $a$  and  $b$  are even? What if both  $a$  and  $b$  are odd?

- If both both  $a$  and  $b$  are even, we can just split students in pairs in both classes separately

# Sums of Even and Odd Numbers

## Splitting in pairs

Suppose there are two classes with  $a$  and  $b$  students respectively. We unite the classes and would like to split all students into pairs to work on a project. Is it possible if  $a$  is even and  $b$  is odd? What if both  $a$  and  $b$  are even? What if both  $a$  and  $b$  are odd?

- If both both  $a$  and  $b$  are even, we can just split students in pairs in both classes separately
- So the answer is yes

# Sums of Even and Odd Numbers

## Splitting in pairs

Suppose there are two classes with  $a$  and  $b$  students respectively. We unite the classes and would like to split all students into pairs to work on a project. Is it possible if  $a$  is even and  $b$  is odd? What if both  $a$  and  $b$  are even? What if both  $a$  and  $b$  are odd?

- If both  $a$  and  $b$  are odd, we can split into pairs all students except one in the first class and all student except one in the second class

# Sums of Even and Odd Numbers

## Splitting in pairs

Suppose there are two classes with  $a$  and  $b$  students respectively. We unite the classes and would like to split all students into pairs to work on a project. Is it possible if  $a$  is even and  $b$  is odd? What if both  $a$  and  $b$  are even? What if both  $a$  and  $b$  are odd?

- If both  $a$  and  $b$  are odd, we can split into pairs all students except one in the first class and all student except one in the second class
- Then we can pair two remaining students

# Sums of Even and Odd Numbers

## Splitting in pairs

Suppose there are two classes with  $a$  and  $b$  students respectively. We unite the classes and would like to split all students into pairs to work on a project. Is it possible if  $a$  is even and  $b$  is odd? What if both  $a$  and  $b$  are even? What if both  $a$  and  $b$  are odd?

- If both  $a$  and  $b$  are odd, we can split into pairs all students except one in the first class and all student except one in the second class
- Then we can pair two remaining students
- So the answer is yes



# Remainder of a Sum

## Remainder of a sum

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a + b$ ?

# Remainder of a Sum

## Remainder of a sum

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a + b$ ?

- It turns out that we can!

# Remainder of a Sum

## Remainder of a sum

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a + b$ ?

- It turns out that we can!
- If both  $a$  and  $b$  are even they have the form  $2 \times q_1$  and  $2 \times q_2$

# Remainder of a Sum

## Remainder of a sum

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a + b$ ?

- It turns out that we can!
- If both  $a$  and  $b$  are even they have the form  $2 \times q_1$  and  $2 \times q_2$
- So the sum is  $2 \times (q_1 + q_2)$  and is even

# Remainder of a Sum

## Remainder of a sum

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a + b$ ?

- If one number is even and the other is odd, they have the forms  $2 \times q_1$  and  $2 \times q_2 + 1$  respectively

# Remainder of a Sum

## Remainder of a sum

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a + b$ ?

- If one number is even and the other is odd, they have the forms  $2 \times q_1$  and  $2 \times q_2 + 1$  respectively
- So the sum is  $2 \times (q_1 + q_2) + 1$  and is odd

# Remainder of a Sum

## Remainder of a sum

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a + b$ ?

- If one number is even and the other is odd, they have the forms  $2 \times q_1$  and  $2 \times q_2 + 1$  respectively
- So the sum is  $2 \times (q_1 + q_2) + 1$  and is odd
- Finally if both  $a$  and  $b$  are odd they have the form  $2 \times q_1 + 1$  and  $2 \times q_2 + 1$

# Remainder of a Sum

## Remainder of a sum

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a + b$ ?

- If one number is even and the other is odd, they have the forms  $2 \times q_1$  and  $2 \times q_2 + 1$  respectively
- So the sum is  $2 \times (q_1 + q_2) + 1$  and is odd
- Finally if both  $a$  and  $b$  are odd they have the form  $2 \times q_1 + 1$  and  $2 \times q_2 + 1$
- So the sum is  $2 \times (q_1 + q_2 + 1)$  and is even



# Remainder of a Sum

- So we have the following table for the remainders of  $a + b$  when dividing by 2

+	0	1
0	0	1
1	1	0

# Remainder of a Sum

- So we have the following table for the remainders of  $a + b$  when dividing by 2
- Rows are remainders of  $a$  and columns are remainders of  $b$

+	0	1
0	0	1
1	1	0

# Remainder of a Sum

- So we have the following table for the remainders of  $a + b$  when dividing by 2
- Rows are remainders of  $a$  and columns are remainders of  $b$
- In the cells we have remainders of  $a + b$

+	0	1
0	0	1
1	1	0

# Remainder of an Opposite

## Remainder of an opposite

Suppose we know the remainder of  $a$  when divided by 2.  
Can we deduce the remainder of  $-a$ ?

# Remainder of an Opposite

## Remainder of an opposite

Suppose we know the remainder of  $a$  when divided by 2.  
Can we deduce the remainder of  $-a$ ?

- It is the same

# Remainder of an Opposite

## Remainder of an opposite

Suppose we know the remainder of  $a$  when divided by 2.  
Can we deduce the remainder of  $-a$ ?

- It is the same
- Indeed,  $a$  is divisible by 2 iff  $-a$  is divisible by 2

# Remainder of a Subtraction

## Remainder of a subtraction

The remainder of  $a - b$  is the same as the remainder of  $a + b$  when divided by 2

# Remainder of a Subtraction

## Remainder of a subtraction

The remainder of  $a - b$  is the same as the remainder of  $a + b$  when divided by 2

- So, for subtraction we have the same table

—	0	1
0	0	1
1	1	0



# Remainder of a Product

## Remainder of a product

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a \times b$ ?

# Remainder of a Product

## Remainder of a product

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a \times b$ ?

- Again, we can

# Remainder of a Product

## Remainder of a product

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a \times b$ ?

- Again, we can
- If at least one of the number is even, then it is divisible by 2

# Remainder of a Product

## Remainder of a product

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a \times b$ ?

- Again, we can
- If at least one of the number is even, then it is divisible by 2
- So the product  $a \times b$  is divisible by 2 and so is even

# Remainder of a Product

## Remainder of a product

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a \times b$ ?

- If both  $a$  and  $b$  are odd they have the form  $2 \times q_1 + 1$  and  $2 \times q_2 + 1$

# Remainder of a Product

## Remainder of a product

Suppose we know the remainders of  $a$  and  $b$  when divided by 2. Can we deduce the remainder of  $a \times b$ ?

- If both  $a$  and  $b$  are odd they have the form  $2 \times q_1 + 1$  and  $2 \times q_2 + 1$
- So the product is  $(q_1 \times 2 + 1) \times (q_2 \times 2 + 1)$   
 $= (q_1 \times q_2 \times 2 + q_1 + q_2) \times 2 + 1$  and is odd

# Remainder of a Product

- So we have the following table for the remainders of  $a \times b$  when dividing by 2

$\times$	0	1
0	0	0
1	0	1

# Remainder of a Product

- So we have the following table for the remainders of  $a \times b$  when dividing by 2
- Rows are remainders of  $a$  and columns are remainders of  $b$

$\times$	0	1
0	0	0
1	0	1



# Remainder of a Product

- So we have the following table for the remainders of  $a \times b$  when dividing by 2
- Rows are remainders of  $a$  and columns are remainders of  $b$
- In the cells we have remainders of  $a \times b$

$\times$	0	1
0	0	0
1	0	1

# Operations on Remainders

## Problem

What is the remainder of  $374 \times (419 + 267 \times 38) - 625$  when divided by 2?

# Operations on Remainders

## Problem

What is the remainder of  $374 \times (419 + 267 \times 38) - 625$  when divided by 2?

- Now we can do it without calculations!

# Operations on Remainders

## Problem

What is the remainder of  $374 \times (419 + 267 \times 38) - 625$  when divided by 2?

- Now we can do it without calculations!
- We can apply operations directly to remainders

# Operations on Remainders

## Problem

What is the remainder of  $374 \times (419 + 267 \times 38) - 625$  when divided by 2?

- Now we can do it without calculations!
- We can apply operations directly to remainders
- Substitute all numbers by remainders:

$$0 \times (1 + 1 \times 0) - 1$$

# Operations on Remainders

## Problem

What is the remainder of  $374 \times (419 + 267 \times 38) - 625$  when divided by 2?

- Now we can do it without calculations!
- We can apply operations directly to remainders
- Substitute all numbers by remainders:  
 $0 \times (1 + 1 \times 0) - 1$
- Computing by our rules gives us the answer 1

# Operations on Remainders

## Problem

What is the remainder of  $374 \times (419 + 267 \times 38) - 625$  when divided by 2?

- Now we can do it without calculations!
- We can apply operations directly to remainders
- Substitute all numbers by remainders:  
 $0 \times (1 + 1 \times 0) - 1$
- Computing by our rules gives us the answer 1
- So the expression in the problem has remainder 1

# Outline

Division by 2

Binary System



# Decimal System

- What is the decimal system?

# Decimal System

- What is the decimal system?
- We represent numbers by sequences of digits

# Decimal System

- What is the decimal system?
- We represent numbers by sequences of digits
- Which number is represented by 3926?

# Decimal System

- What is the decimal system?
- We represent numbers by sequences of digits
- Which number is represented by 3926?
- The number written here is

$$3926 = 3000 + 900 + 20 + 6 =$$

$$3 \times 10^3 + 9 \times 10^2 + 2 \times 10^1 + 6 \times 10^0$$

# Decimal System

- What is the decimal system?
- We represent numbers by sequences of digits
- Which number is represented by 3926?
- The number written here is
$$3926 = 3000 + 900 + 20 + 6 =$$
$$3 \times 10^3 + 9 \times 10^2 + 2 \times 10^1 + 6 \times 10^0$$
- In general, digits are multiplied by powers of 10

# Decimal System

- What is the decimal system?
- We represent numbers by sequences of digits
- Which number is represented by 3926?
- The number written here is
$$3926 = 3000 + 900 + 20 + 6 =$$
$$3 \times 10^3 + 9 \times 10^2 + 2 \times 10^1 + 6 \times 10^0$$
- In general, digits are multiplied by powers of 10
- Each non-negative integer number has a unique representation

# Binary System

- We can do the same with powers of 2

# Binary System

- We can do the same with powers of 2
- This is called **binary system**



# Binary System

- We can do the same with powers of 2
- This is called **binary system**
- There are only two bits 0 and 1 now instead of ten digits

# Binary System

- We can do the same with powers of 2
- This is called **binary system**
- There are only two bits 0 and 1 now instead of ten digits
- An example of a binary representation: 101101

# Binary System

- We can do the same with powers of 2
- This is called **binary system**
- There are only two bits 0 and 1 now instead of ten digits
- An example of a binary representation: 101101

- The number written here is

$$\begin{aligned} 101101 &= 1 \times 2^5 + 0 \times 2^4 + 1 \times 2^3 + 1 \times 2^2 + 0 \times 2^1 + \\ &1 \times 2^0 = 2^5 + 2^3 + 2^2 + 2^0 = 32 + 8 + 4 + 1 = 45 \end{aligned}$$

# Binary System

- We can do the same with powers of 2
- This is called **binary system**
- There are only two bits 0 and 1 now instead of ten digits
- An example of a binary representation: 101101
- The number written here is
$$101101 = 1 \times 2^5 + 0 \times 2^4 + 1 \times 2^3 + 1 \times 2^2 + 0 \times 2^1 + 1 \times 2^0 = 2^5 + 2^3 + 2^2 + 2^0 = 32 + 8 + 4 + 1 = 45$$
- Binary system is important since computers use it

# Binary System and Remainders

## Lemma

Suppose we divide  $a$  by 2 with a remainder. Then the remainder is the last bit of binary representation of  $a$  and the quotient is the number formed by all bits of the binary representation of  $a$  except the last one

# Binary System and Remainders

## Lemma

Suppose we divide  $a$  by 2 with a remainder. Then the remainder is the last bit of binary representation of  $a$  and the quotient is the number formed by all bits of the binary representation of  $a$  except the last one

- The proof is the same as for the decimal system

# Binary System and Remainders

## Lemma

Suppose we divide  $a$  by 2 with a remainder. Then the remainder is the last bit of binary representation of  $a$  and the quotient is the number formed by all bits of the binary representation of  $a$  except the last one

- The proof is the same as for the decimal system
- For simplicity consider some specific number, say, 11101 in binary

# Binary System and Remainders

## Lemma

Suppose we divide  $a$  by 2 with a remainder. Then the remainder is the last bit of binary representation of  $a$  and the quotient is the number formed by all bits of the binary representation of  $a$  except the last one

- The proof is the same as for the decimal system
- For simplicity consider some specific number, say, 11101 in binary
- This number is equal to
$$1 \times 2^4 + 1 \times 2^3 + 1 \times 2^2 + 0 \times 2^1 + 1 \times 2^0 = (2^3 + 2^2 + 2^1) \times 2 + 1$$



# Binary System and Remainders

## Lemma

Suppose we divide  $a$  by 2 with a remainder. Then the remainder is the last bit of binary representation of  $a$  and the quotient is the number formed by all bits of the binary representation of  $a$  except the last one

- The proof is the same as for the decimal system
- For simplicity consider some specific number, say, 11101 in binary
- This number is equal to
$$1 \times 2^4 + 1 \times 2^3 + 1 \times 2^2 + 0 \times 2^1 + 1 \times 2^0 = (2^3 + 2^2 + 2^1) \times 2 + 1$$
- So the remainder is 1 and the quotient is 1110 in binary

# Binary System

- Note that we are are not proving the correctness of binary system here

# Binary System

- Note that we are not proving the correctness of binary system here
- Why this system works? Why every integer is representable? Why the representation is unique?

# Binary System

- Note that we are not proving the correctness of binary system here
- Why this system works? Why every integer is representable? Why the representation is unique?
- We will omit the proof

# Binary System

- Note that we are not proving the correctness of binary system here
- Why this system works? Why every integer is representable? Why the representation is unique?
- We will omit the proof
- However, the proof is not hard; one can just combine our previous Lemma and induction

# Binary System

- Note that we are not proving the correctness of binary system here
- Why this system works? Why every integer is representable? Why the representation is unique?
- We will omit the proof
- However, the proof is not hard; one can just combine our previous Lemma and induction
- This is also a way to construct binary representation for a given number: apply previous lemma recursively

# Binary System and Logarithm

- How many bits are needed to represent a given number  $a$ ?

# Binary System and Logarithm

- How many bits are needed to represent a given number  $a$ ?
- Observe that to represent  $2^n$  we need  $n + 1$  bits:  
 $2^4$  is 10000



# Binary System and Logarithm

- How many bits are needed to represent a given number  $a$ ?
- Observe that to represent  $2^n$  we need  $n + 1$  bits:  
 $2^4$  is 10000
- And  $2^n$  is the smallest number that requires  $n + 1$  bits:  
 $2^4 - 1 = 2^3 + 2^2 + 2^1 + 1$  is 1111

# Binary System and Logarithm

- How many bits are needed to represent a given number  $a$ ?
- Observe that to represent  $2^n$  we need  $n + 1$  bits:  
 $2^4$  is 10000
- And  $2^n$  is the smallest number that requires  $n + 1$  bits:  
 $2^4 - 1 = 2^3 + 2^2 + 2^1 + 1$  is 1111
- The number of bits that is needed to represent  $a$  is determined by the largest  $n$  such that  $2^n \leq a$ : it is equal to  $n + 1$

# Binary Logarithm

- Recall that for real number  $x > 0$  by  $\log_2 x$  is denoted the number  $y$  such that  $2^y = x$

# Binary Logarithm

- Recall that for real number  $x > 0$  by  $\log_2 x$  is denoted the number  $y$  such that  $2^y = x$
- Logarithm is the inverse function to exponentiation:  
 $\log_2 2^y = y$

# Binary Logarithm

- Recall that for real number  $x > 0$  by  $\log_2 x$  is denoted the number  $y$  such that  $2^y = x$
- Logarithm is the inverse function to exponentiation:  
 $\log_2 2^y = y$
- In particular,  $\log_2 2^n = n$

# Binary Logarithm

- Recall that for real number  $x > 0$  by  $\log_2 x$  is denoted the number  $y$  such that  $2^y = x$
- Logarithm is the inverse function to exponentiation:  
 $\log_2 2^y = y$
- In particular,  $\log_2 2^n = n$
- Binary logarithm has a clear meaning in terms of binary representation

# Binary Logarithm

## Lemma

The number of bits needed to represent an integer  $a$  in binary representation is equal to  $\lfloor \log_2 a \rfloor + 1$

- By  $\lfloor x \rfloor$  we denote the largest integer that is less or equal to (possibly non-integer)  $x$

# Binary Logarithm

## Lemma

The number of bits needed to represent an integer  $a$  in binary representation is equal to  $\lfloor \log_2 a \rfloor + 1$

- By  $\lfloor x \rfloor$  we denote the largest integer that is less or equal to (possibly non-integer)  $x$
- Why the lemma is true?



# Binary Logarithm

## Lemma

The number of bits needed to represent an integer  $a$  in binary representation is equal to  $\lfloor \log_2 a \rfloor + 1$

- By  $\lfloor x \rfloor$  we denote the largest integer that is less or equal to (possibly non-integer)  $x$
- Why the lemma is true?
- Recall that the number of bits in binary representation of  $a$  is  $n + 1$  where  $n$  is the largest number such that  $2^n \leq a$

# Binary Logarithm

## Lemma

The number of bits needed to represent an integer  $a$  in binary representation is equal to  $\lfloor \log_2 a \rfloor + 1$

- By  $\lfloor x \rfloor$  we denote the largest integer that is less or equal to (possibly non-integer)  $x$
- Why the lemma is true?
- Recall that the number of bits in binary representation of  $a$  is  $n + 1$  where  $n$  is the largest number such that  $2^n \leq a$
- Applying logarithm to both sides of inequality we obtain that  $n$  is the largest number such that  $n \leq \log_2 a$