Markov's Inequality

Vladimir Podolskii

Computer Science Department, Higher School of Economics

Outline

From Expectation to Probability

Markov's Inequality

Application to Algorithms

Expectation vs. Probability

- In this week we have introduced random variables and their expected values

Expectation vs. Probability

- In this week we have introduced random variables and their expected values
- We will see now how these notions can help in studies of probabilities of events

Lottery Budget

Problem

A lottery ticket costs 10 dollars. A 40% of a lottery budget goes to prizes. Show that the chances to win 500 dollars or more are less than 1%

Lottery Budget

Problem

A lottery ticket costs 10 dollars. A 40% of a lottery budget goes to prizes. Show that the chances to win 500 dollars or more are less than 1%

- We will use proof by contradiction

Lottery Budget

Problem

A lottery ticket costs 10 dollars. A 40% of a lottery budget goes to prizes. Show that the chances to win 500 dollars or more are less than 1%

- We will use proof by contradiction
- Assume the contrary: the probability to win 500 dollars or more is at least 0.01

Lottery Budget

Problem

A lottery ticket costs 10 dollars. A 40% of a lottery budget goes to prizes. Show that the chances to win 500 dollars or more are less than 1%

- We will use proof by contradiction
- Assume the contrary: the probability to win 500 dollars or more is at least 0.01
- Denote the number of tickets sold by n

Lottery Budget

Problem

A lottery ticket costs 10 dollars. A 40% of a lottery budget goes to prizes. Show that the chances to win 500 dollars or more are less than 1%

- We will use proof by contradiction
- Assume the contrary: the probability to win 500 dollars or more is at least 0.01
- Denote the number of tickets sold by n
- Then the budget of the lottery is $10n$ dollars

Lottery Budget

Problem

A lottery ticket costs 10 dollars. A 40% of a lottery budget goes to prizes. Show that the chances to win 500 dollars or more are less than 1%

- We will use proof by contradiction
- Assume the contrary: the probability to win 500 dollars or more is at least 0.01
- Denote the number of tickets sold by n
- Then the budget of the lottery is $10n$ dollars
- $10n \times 0.4 = 4n$ dollars are spent on the prizes

Lottery Budget

Problem

A lottery ticket costs 10 dollars. A 40% of a lottery budget goes to prizes. Show that the chances to win 500 dollars or more are less than 1%

- By our assumption at least $\frac{n}{100}$ tickets win at least 500 dollars

Lottery Budget

Problem

A lottery ticket costs 10 dollars. A 40% of a lottery budget goes to prizes. Show that the chances to win 500 dollars or more are less than 1%

- By our assumption at least $\frac{n}{100}$ tickets win at least 500 dollars
- In total these tickets win $\frac{n}{100} \times 500 = 5n$ dollars

Lottery Budget

Problem

A lottery ticket costs 10 dollars. A 40% of a lottery budget goes to prizes. Show that the chances to win 500 dollars or more are less than 1%

- By our assumption at least $\frac{n}{100}$ tickets win at least 500 dollars
- In total these tickets win $\frac{n}{100} \times 500 = 5n$ dollars
- This exceeds the total prize budget of $4n$!

Lottery Budget

Problem

A lottery ticket costs 10 dollars. A 40% of a lottery budget goes to prizes. Show that the chances to win 500 dollars or more are less than 1%

- By our assumption at least $\frac{n}{100}$ tickets win at least 500 dollars
- In total these tickets win $\frac{n}{100} \times 500 = 5n$ dollars
- This exceeds the total prize budget of $4n$!
- We arrived into contradiction and the problem is solved

Outline

From Expectation to Probability

Markov's Inequality

Application to Algorithms

Markov's Inequality

Markov's Inequality

Suppose that f is a non-negative random variable. Then for any number $a > 0$ we have

$$\Pr[f \geq a] \leq \frac{\mathsf{E} f}{a}$$

Markov's Inequality

Markov's Inequality

Suppose that f is a non-negative random variable. Then for any number $a > 0$ we have

$$\Pr[f \geq a] \leq \frac{\mathsf{E} f}{a}$$

- The inequality allows to use expected value to bound probability of certain events

Markov's Inequality

Markov's Inequality

Suppose that f is a non-negative random variable. Then for any number $a > 0$ we have

$$\Pr[f \geq a] \leq \frac{\mathsf{E} f}{a}$$

- The inequality allows to use expected value to bound probability of certain events
- For the proof it is convenient to rewrite the inequality:

$$a \times \Pr[f \geq a] \leq \mathsf{E} f$$

Markov's Inequality

- We need to prove the inequality $a \times \Pr[f \geq a] \leq \mathsf{E} f$

Markov's Inequality

- We need to prove the inequality $a \times \Pr[f \geq a] \leq \mathsf{E} f$
- Consider the following random variable g on the same probability space: for an outcome such that $f = a_i$ we let $g = a$ on this outcome if $a \leq a_i$ and $g = 0$ otherwise; that is

$$g = \begin{cases} a & \text{if } f = a_i \geq a \\ 0 & \text{if } f = a_i < a \end{cases}$$

Markov's Inequality

- We need to prove the inequality $a \times \Pr[f \geq a] \leq \mathsf{E} f$
- Consider the following random variable g on the same probability space: for an outcome such that $f = a_i$ we let $g = a$ on this outcome if $a \leq a_i$ and $g = 0$ otherwise; that is

$$g = \begin{cases} a & \text{if } f = a_i \geq a \\ 0 & \text{if } f = a_i < a \end{cases}$$

- g is less or equal than f on each outcome

Markov's Inequality

- We need to prove the inequality $a \times \Pr[f \geq a] \leq \mathsf{E} f$
- Consider the following random variable g on the same probability space: for an outcome such that $f = a_i$ we let $g = a$ on this outcome if $a \leq a_i$ and $g = 0$ otherwise; that is

$$g = \begin{cases} a & \text{if } f = a_i \geq a \\ 0 & \text{if } f = a_i < a \end{cases}$$

- g is less or equal than f on each outcome
- So the average value $\mathsf{E} g$ of g is less or equal than the average value $\mathsf{E} f$ of f:

$$\mathsf{E} g \leq \mathsf{E} f$$

Markov's Inequality

$$g = \begin{cases} a & \text{if } f = a_i \geq a \\ 0 & \text{if } f = a_i < a \end{cases}$$

- What is the expectation of g?

Markov's Inequality

$$g = \begin{cases} a & \text{if } f = a_i \geq a \\ 0 & \text{if } f = a_i < a \end{cases}$$

- What is the expectation of g?
- g has only one nonzero value;

Markov's Inequality

$$g = \begin{cases} a & \text{if } f = a_i \geq a \\ 0 & \text{if } f = a_i < a \end{cases}$$

- What is the expectation of g?
- g has only one nonzero value;
- so Eg is this value multiplied by the sum of probabilities of all outcomes for this value

Markov's Inequality

$$g = \begin{cases} a & \text{if } f = a_i \geq a \\ 0 & \text{if } f = a_i < a \end{cases}$$

- What is the expectation of g?
- g has only one nonzero value;
- so Eg is this value multiplied by the sum of probabilities of all outcomes for this value
- But these outcomes are exactly the outcomes that form the event "$f \geq a$"

Markov's Inequality

$$g = \begin{cases} a & \text{if } f = a_i \geq a \\ 0 & \text{if } f = a_i < a \end{cases}$$

- What is the expectation of g?
- g has only one nonzero value;
- so Eg is this value multiplied by the sum of probabilities of all outcomes for this value
- But these outcomes are exactly the outcomes that form the event "$f \geq a$"
- Thus the sum of their probabilities is equal to $\Pr[f \geq a]$

Markov's Inequality

$$g = \begin{cases} a & \text{if } f = a_i \geq a \\ 0 & \text{if } f = a_i < a \end{cases}$$

- What is the expectation of g?
- g has only one nonzero value;
- so Eg is this value multiplied by the sum of probabilities of all outcomes for this value
- But these outcomes are exactly the outcomes that form the event "$f \geq a$"
- Thus the sum of their probabilities is equal to $\Pr[f \geq a]$
- Thus E$g = a \times \Pr[f \geq a]$

Markov's Inequality

$$g = \begin{cases} a & \text{if } f = a_i \geq a \\ 0 & \text{if } f = a_i < a \end{cases}$$

- Finally, we have $\mathsf{E}g \leq \mathsf{E}f$

Markov's Inequality

$$g = \begin{cases} a & \text{if } f = a_i \geq a \\ 0 & \text{if } f = a_i < a \end{cases}$$

- Finally, we have $\mathrm{E}g \leq \mathrm{E}f$
- And $\mathrm{E}g = a \times \Pr[f \geq a]$

Markov's Inequality

$$g = \begin{cases} a & \text{if } f = a_i \geq a \\ 0 & \text{if } f = a_i < a \end{cases}$$

- Finally, we have $\mathsf{E}g \leq \mathsf{E}f$
- And $\mathsf{E}g = a \times \Pr[f \geq a]$
- So

$$\mathsf{E}f \geq \mathsf{E}g = a \times \Pr[f \geq a]$$

Markov's Inequality

$$g = \begin{cases} a & \text{if } f = a_i \geq a \\ 0 & \text{if } f = a_i < a \end{cases}$$

- Finally, we have $\mathsf{E}g \leq \mathsf{E}f$
- And $\mathsf{E}g = a \times \Pr[f \geq a]$
- So

$$\mathsf{E}f \geq \mathsf{E}g = a \times \Pr[f \geq a]$$

- We have shown Markov's inequality

Geometric Interpretation

$$\mathsf{E} f \geq a \times \Pr[f \geq a]$$

Geometric Interpretation

$\mathsf{E}f \geq a \times \Pr[f \geq a]$

Suppose f obtains values a_1, a_2, a_3, a_4 with probabilities p_1, p_2, p_3, p_4

Geometric Interpretation

$\mathsf{E}f \geq a \times \Pr[f \geq a]$

Suppose f obtains values a_1, a_2, a_3, a_4 with probabilities p_1, p_2, p_3, p_4



Geometric Interpretation

$\mathsf{E}f \geq a \times \Pr[f \geq a]$

Suppose f obtains values a_1, a_2, a_3, a_4 with probabilities p_1, p_2, p_3, p_4



Geometric Interpretation

$\mathsf{E}f \geq a \times \Pr[f \geq a]$

Suppose f obtains values a_1, a_2, a_3, a_4 with probabilities p_1, p_2, p_3, p_4



Geometric Interpretation

$\mathsf{E}f \geq a \times \Pr[f \geq a]$

Suppose f obtains values a_1, a_2, a_3, a_4 with probabilities p_1, p_2, p_3, p_4



Geometric Interpretation

$\mathsf{E}f \geq a \times \Pr[f \geq a]$

Suppose f obtains values a_1, a_2, a_3, a_4 with probabilities p_1, p_2, p_3, p_4



- $\mathsf{E}f$ is the area of the gray region

Geometric Interpretation

$\mathrm{E}f \geq a \times \Pr[f \geq a]$

Suppose f obtains values a_1, a_2, a_3, a_4 with probabilities p_1, p_2, p_3, p_4



- $\mathrm{E}f$ is the area of the gray region
- $a \times \Pr[f \geq a]$ is the area of a red region

Geometric Interpretation

$\mathrm{E}f \geq a \times \Pr[f \geq a]$

Suppose f obtains values a_1, a_2, a_3, a_4 with probabilities p_1, p_2, p_3, p_4



- $\mathrm{E}f$ is the area of the gray region
- $a \times \Pr[f \geq a]$ is the area of a red region
- The gray region is large and the inequality follows

Outline

From Expectation to Probability

Markov's Inequality

Application to Algorithms

Application to Algorithms

Problem

Suppose there is a randomized algorithm that runs on average in time, say, n^2, where n is the size of input. The algorithm outputs the correct answer. Construct another randomized algorithm that always stops in time cn^2 for some constant c and makes a mistake with probability at most 10^{-3}

Application to Algorithms

Problem

Suppose there is a randomized algorithm that runs on average in time, say, n^2, where n is the size of input. The algorithm outputs the correct answer. Construct another randomized algorithm that always stops in time cn^2 for some constant c and makes a mistake with probability at most 10^{-3}

- We will apply Markov inequality

Application to Algorithms

- Running time of the algorithm is a random variable; denote it by f

Application to Algorithms

- Running time of the algorithm is a random variable; denote it by f
- We know that $\mathsf{E} f = n^2$

Application to Algorithms

- Running time of the algorithm is a random variable; denote it by f
- We know that $\mathsf{E} f = n^2$
- Here is a new algorithm

Application to Algorithms

- Running time of the algorithm is a random variable; denote it by f
- We know that $\mathsf{E} f = n^2$
- Here is a new algorithm
- Run the original algorithm for $10^3 n^2$ steps

Application to Algorithms

- Running time of the algorithm is a random variable; denote it by f
- We know that $\mathsf{E} f = n^2$
- Here is a new algorithm
- Run the original algorithm for $10^3 n^2$ steps
- If it stops, we also stop

Application to Algorithms

- Running time of the algorithm is a random variable; denote it by f
- We know that $\mathsf{E} f = n^2$
- Here is a new algorithm
- Run the original algorithm for $10^3 n^2$ steps
- If it stops, we also stop
- If not, stop and output, say, 0

Application to Algorithms

Claim

The probability that the original algorithm does not stop after $10^3 n^2$ number of steps is at most 10^{-3}

- Indeed, this probability is $\Pr[f \geq 10^3 n^2]$

Application to Algorithms

Claim

The probability that the original algorithm does not stop after $10^3 n^2$ number of steps is at most 10^{-3}

- Indeed, this probability is $\Pr[f \geq 10^3 n^2]$
- By Markov's inequality it is bounded by

$$\Pr[f \geq 10^3 n^2] \leq \frac{\mathsf{E} f}{10^3 n^2} = \frac{n^2}{10^3 n^2} = 10^{-3}$$

Conclusion

- We studied random variables

Conclusion

- We studied random variables
- Allow us to study quantitative aspects of randomness

Conclusion

- We studied random variables
- Allow us to study quantitative aspects of randomness
- Allow us to apply many analytic tools to study probability

Conclusion

- We studied random variables
- Allow us to study quantitative aspects of randomness
- Allow us to apply many analytic tools to study probability
- Expected value is one of the main characteristics of a random variable

Conclusion

- We studied random variables
- Allow us to study quantitative aspects of randomness
- Allow us to apply many analytic tools to study probability
- Expected value is one of the main characteristics of a random variable
- On one side, expectation bears a lot of information of a random variable

Conclusion

- We studied random variables
- Allow us to study quantitative aspects of randomness
- Allow us to apply many analytic tools to study probability
- Expected value is one of the main characteristics of a random variable
- On one side, expectation bears a lot of information of a random variable
- On the other side, expectation has very convenient mathematical properties