RESEARCH ARTICLE JOURNAL OF CHEMICAL THEORY AND COMPUTATION

# Scalable Contact-Gated Stochastic Operators and Universal Reduction Laws for Macromolecular Protein Dynamics, Allostery, and Kinetics

**Mehardeep Singh[a,*] and Gurarpit Singh[b]**

[a]Sri Chaitanya Educational Institutions, Chandigarh, India; [b]Vivekananda Institute of Professional Studies (VIPS), Guru Gobind Singh Indraprastha University (GGSIPU), Delhi, India

[*]Correspondence and requests for materials should be addressed to Mehardeep Singh (email: mehardeep.sim@gmail.com). Author emails: Mehardeep Singh (mehardeep.sim@gmail.com), Gurarpit Singh (gurarpit.sml@gmail.com).

**Abstract:** Constructing continuous, thermodynamically reversible reduced equations of motion from atomistic molecular dynamics (MD) that scale from fast-folding miniproteins to multi-domain allosteric enzymes in explicit water remains a foundational problem in computational biophysics. Across 15 molecular systems (890,000 SHA-256-verified frames, 0.89 $\mu$s cumulative sampling) spanning 13 structurally diverse proteins from Chignolin ($d = 18$) to explicit-solvent Adenylate Kinase ($d = 426$, 40,110 particles) and Maltose-Binding Protein ($d = 738$, 49,215 particles), plus two analytical verification peptides ($d = 2, 4$), we formulate a contact-gated, GENERIC-regularized stochastic operator on the backbone torus $\mathbb{T}^d$. We show that collective slow modes compress sub-linearly across two orders of magnitude in chain length—with spectral participation ratio scaling as $PR(d) = 1.061\, d^{0.494}$ (95% CI $[0.332, 0.666]$, $\Delta\mathrm{AIC}_c = -18.15$) and short-time diffusion rank as $r_{90}^{D}(d) = 0.799\, d^{0.930}$ ($\Delta\mathrm{AIC}_c = -14.39$)—whereas cumulative 90% equilibrium variance in large native-basin enzymes transitions to extensive proportionality (10.0% blind extrapolation error vs. 35.7% for small-protein power laws). Consequently, macromolecular tractability above $d > 200$ requires off-diagonal sparsity rather than global PCA truncation alone: universal peptide-plane crankshaft anti-correlation ($\bar{R}(\psi_i, \phi_{i+1}) = -0.603 \pm 0.054$) and sparse 3D tertiary contact gating ($\bar{k}_{\mathrm{eff}} = 5.97$, reaching 99.3% sparsity at $d = 738$) enable a contact-graph Cholesky estimator scaling empirically as $O(d^{1.13})$ in time and $O(d^{1.06})$ in memory (65.8× memory compression, 77.3× faster than Graphical LASSO, and higher blind held-out log-likelihood $+0.303/d$ vs. dense $+0.205/d$ on MBP). Biophysically, relative solvent-accessible surface area ($\mathrm{rSASA}_i$) unifies dry linear equipartition in protein cores ($\alpha_{\mathrm{core}} = 1.00$, 12.33% blind thermal error) with viscosity-coupled loops ($\alpha_{\mathrm{loop}} = 1.28$) and peptides ($\alpha_{\mathrm{peptide}} = 1.57$); parameter-free kinematic projection reproduces solution NMR Lipari–Szabo $S^2$ order parameters ($\bar{r}_s = +0.483$ across four BMRB enzymes, Ubiquitin Pearson $r = +0.921$) and X-ray $B$-factors ($\bar{r}_s = +0.276$); and 13-fold zero-shot Leave-One-Protein-Out transfer predicts unseen protein operators directly from static PDB coordinates ($r_s = +0.706$, $R^2 = +0.359$ ridge; $r_s = +0.721$, $R^2 = +0.410$ GBDT). Applied to functional discovery, explicit-solvent apo toroidal operators localize mechanical domain-closure hinges in AdK (PR-AUC = 0.348) and MBP (AUROC = 0.690 vs. 0.446 ANM), resolve distal Met20 ↔ FG catalytic loop coupling in *E. coli* DHFR (71.6% recovery at 93.2% sparsity) and $\beta_1$–$\beta_2$ ↔ Ile44 routing in Ubiquitin (7.1× amplification) on blind trajectories, and quantify sub-diffusive Mori–Zwanzig folding memory ($\tilde{\gamma}_M = 0.593 \pm 0.126$). A validation-tuned head-to-head benchmark against Markov State Models, TICA, Smoluchowski diffusion, Elastic Network Models, and unconstrained SDEs across all 13 proteins maps the exact physical regimes where continuous contact-gated GENERIC operators (lowest Frobenius error 0.288, exact 0.000 spurious entropy vs. 346.17 $k_B$/ns/$d$) and discrete or harmonic baselines each excel.

**Keywords:** macromolecular dynamics | stochastic differential operators | Mori–Zwanzig projection | allosteric hinges | NMR Lipari–Szabo $S^2$ | GENERIC thermodynamics

## INTRODUCTION

Macromolecular function—encompassing large-amplitude domain closure in Adenylate Kinase (AdK) and Maltose-Binding Protein (MBP), distal catalytic loop coordination in Dihydrofolate Reductase (DHFR), picosecond-to-nanosecond backbone amide bond vector fluctuations measured by solution NMR relaxation, and cooperative folding in miniproteins—emerges from stochastic trajectories evolving on rugged, high-dimensional free-energy landscapes (1–5). While explicit-solvent atomistic molecular dynamics (MD) integrates Newton's equations at femtosecond resolution across tens of thousands of solvent and solute atoms, extracting physically interpretable mechanisms and quantitative kinetic predictions requires projecting the microscopic Hamiltonian phase space $\dot{\Gamma} = \mathcal{L}_{\mathrm{micro}}\Gamma$ onto a reduced collective manifold $\mathcal{M}$ via the Mori–Zwanzig formalism (6–10).

Four established coarse-graining paradigms address distinct facets of this projection problem, yet face complementary theoretical and practical trade-offs:

1. *Discrete Markov State Models (MSMs) and TICA:* Partitioning continuous conformation space into $k$ Voronoi microstates at lag time $\tau$ (11, 12) or projecting onto slow linear time-lagged independent components (TICA) (13) effectively resolves multi-basin metastable populations when long trajectories sample abundant reversible transitions. However, discrete state boundaries discard continuous local restoring forces and position-dependent diffusion tensors $\mathbf{D}(z)$, incur sensitivity to $(k, \tau)$ selection under finite sampling, and cannot transfer operators zero-shot to unsimulated protein folds.
2. *Elastic Network Models (GNM and ANM):* Harmonic spring networks constructed around a single static crystal structure provide virtually instantaneous ($< 0.2$ s) trajectory-free estimates of isotropic crystallographic $B$-factors (14, 15). Yet $C_\alpha$-only harmonic networks lack dihedral steric exclusion, neglect compensatory peptide-plane crankshaft rotations ($\Delta\psi_{i-1} \approx -\Delta\phi_i$), and cannot model anharmonic barrier crossing or state-dependent diffusion surges.
3. *1D Reaction-Coordinate Smoluchowski Diffusion:* Projecting folding trajectories onto native contact fraction $Q$ via memory-calibrated 1D Fokker–Planck equations captures mean first-passage times (MFPTs) along dominant folding funnels (16, 17), but sacrifices residue-resolved $d$-dimensional off-diagonal allosteric couplings and breaks down in the low-friction Kramers energy-diffusion turnover regime (18, 19).
4. *Unconstrained Data-Driven and Neural SDE Surrogates:* Expressive basis-function and deep neural regressions parameterize nonlinear drift and diffusion fields (20, 21), yet finite-sample regression errors break Onsager reciprocal symmetry and microscopic detailed balance, inducing non-conservative steady-state probability currents ($J_{\mathrm{ss}} \neq 0$) and unphysical irreversible entropy production $\dot{S}_{\mathrm{irr}} > 0$ (22, 23).

Bridging these paradigms for genuine macromolecular proteins raises three open questions in statistical mechanics and biomolecular simulation: (i) *Scalability and Sparsity:* How do the spectral rank and off-diagonal coupling topology of continuous toroidal stochastic operators scale from miniproteins ($d = 18$) to solvated multi-domain enzymes ($d > 700$), and can physical locality break the $O(d^2)$–$d^3$) sample- and memory-complexity barrier on real proteins? (ii) *Experimental Fidelity and Allosteric Mechanism:* Can a parameter-free toroidal operator simultaneously unify internal vs. solvent friction, predict residue-resolved solution NMR Lipari–Szabo $S^2$ order parameters, transfer zero-shot across unseen PDB folds, and uncover mechanical hinges and distal allosteric loop pathways in enzymes such as AdK, MBP, and DHFR? (iii) *Fair Regime Mapping:* Under strict discovery/validation/blind-test governance (Replica 1 → Replica 2 → Replica 3), where do continuous contact-gated GENERIC operators outperform—and where do they tie or complement—validation-tuned MSM, TICA, Smoluchowski, and ENM baselines?

Here, we answer all three questions across a cryptographically verified 15-system atomistic atlas (890,000 frames, $d = 2 \rightarrow 738$; Table 1). Leading directly with 13 macromolecular proteins in explicit TIP3P Particle Mesh Ewald (PME) and Generalized Born (GBn2/OBC2) solvation, we establish six scaling and transfer laws (Figures 1–5), demonstrate concrete 3D allosteric hinge, distal loop routing, and folding-memory predictions across multi-domain enzymes and fast folders (Figures 1, 6–8), verify microscopic kinetic and memory closure on model peptides, and execute a validation-tuned 6-paradigm benchmark across all 13 proteins (Figure 9, Table 2).

## RESULTS

### *1. Macromolecular Scaling and Contact-Gated Sparsity Across 13 Proteins ($d = 18 \rightarrow 738$)*

**Toroidal Manifold Atlas and Spectral Convergence.** To evaluate stochastic operator reduction across genuine macromolecular architectures rather than toy coordinates, we assembled a 3-replica atomistic trajectory atlas spanning 13 structurally diverse proteins ($d = 2N_{\mathrm{res}} - 2 \in [18, 738]$) alongside two reference peptides (890,000 total frames saved at 1.0 ps stride; Table 1, Figure 1). Crucially, the largest catalytic and multi-domain allosteric enzymes—Hen Egg-White Lysozyme ($d = 256$, 17,110 atoms), *E. coli* DHFR ($d = 316$, 20,837 atoms), *E. coli* Adenylate Kinase ($d = 426$, 40,110 atoms), and Maltose-Binding Protein ($d = 738$, 49,215 atoms)—were simulated in explicit TIP3P-FB water with Particle Mesh Ewald (PME) electrostatics (0.15 M NaCl, 4.0 fs hydrogen mass repartitioning, three independent 10 ns replicas per enzyme), complementing multi-ten-nanosecond GBn2/OBC2 ensembles across Chignolin ($d = 18$), Trp-cage ($d = 38$), BBA5 ($d = 54$), Villin HP35 ($d = 64$), WW Domain FiP35 ($d = 68$), GB1 ($d = 110$), disulfide-stabilized BPTI ($d = 114$), Human Ubiquitin ($d = 150$), and Barnase ($d = 218$).

On the compact backbone torus $\mathbb{T}^d = [-\pi, \pi)^d$, naive Euclidean covariance es-

**Table 1. Master biophysical benchmark, spectral compression, contact-gated sparsity, experimental validation, and 13-fold zero-shot LOPO transfer across 15 molecular architectures ($d = 2 \to 738$, 890,000 verified frames)**

| System (PDB) | Fold / Functional Class | $N_{res}$ | Dim $d$ | Frames (Solvent) | Rank $r_{90}$ ($r_{90}^{cont}$) | $PR(d)$ | Crank $\bar{R}$ | Sparsity $S_{0.05}$ | Enrich. | Exp. NMR $S^2$ / Hinge / X-Ray | Zero-Shot LOPO Err. ($r_s$) |
|---|---|---|---|---|---|---|---|---|---|---|---|
| *Part I: Macromolecular Protein Atlas ($d = 18 \to 738$; Discovery Cohort $d \le 150$ & Blind Out-of-Scale Extrapolation Cohort $d \in \{218, 256, 316, 426, 738\}$)* | | | | | | | | | | | |
| **Chignolin (1UAO)** | $\beta$-hairpin fast folder | 10 | 18 | 150,000 (GBn2/OBC2) | 8 (7.81) | 5.08 | -0.512 | 71.2% | 3.01× | Memory $\gamma_M = 0.716$, $\tau^* = 50$ ps | 65.7% ($r_s = +0.761$) |
| **Trp-cage (1L2Y)** | $\alpha$/PPII miniprotein | 20 | 38 | 150,000 (GBn2/OBC2) | 14 (13.72) | 7.94 | -0.469 | 85.5% | 1.83× | $\Delta AIC_c = -11.39$ (power decay) | 57.3% ($r_s = +0.504$) |
| **BBA5 (1FME)** | $\beta\beta\alpha$ zinc-finger motif | 28 | 54 | 30,000 (GBn2) | 16 (15.23) | 6.26 | -0.605 | 91.1% | 1.88× | $\Delta AIC_c = -10.00$ (power decay) | 29.9% ($r_s = +0.728$) |
| **Villin HP35 (2F4K)** | 3-helix bundle folder | 33 | 64 | 30,000 (GBn2/OBC2) | 22 (21.47) | 6.03 | -0.566 | 92.5% | 1.07× | $\Delta AIC_c = -11.21$ (power decay) | 38.1% ($r_s = +0.475$) |
| **WW-FiP35 (2F21)** | 3-stranded $\beta$-sheet | 35 | 68 | 30,000 (GBn2) | 23 (22.64) | 9.60 | -0.616 | 89.8% | 2.51× | $\Delta AIC_c = -14.47$ (power decay) | 51.8% ($r_s = +0.587$) |
| **GB1 Domain (1PGA)** | $4\beta + 1\alpha$ IgG-binding | 56 | 110 | 7,500 (OBC2) | 52 (51.09) | 17.43 | -0.641 | 92.8% | 3.05× | NMR $S^2$: $r_s = +0.746$; $B$: +0.221 | 46.2% ($r_s = +0.693$) |
| **BPTI (5PTI)** | Disulfide Kunitz fold | 58 | 114 | 30,000 (GBn2) | 34 (33.18) | 8.79 | -0.623 | 94.4% | 2.22× | X-ray $B$-factor: $r_s = +0.346$ | 46.0% ($r_s = +0.496$) |
| **Ubiquitin (1UBQ/1D3Z)** | $\alpha + \beta$ signaling hub | 76 | 150 | 7,500 (OBC2) | 55 (54.34) | 11.53 | -0.613 | 94.7% | 2.68× | NMR $S^2$: $r_s = +0.512$ ($r = +0.921$) | 50.0% ($r_s = +0.551$) |
| **Barnase (1BNI)** | Microbial $\alpha + \beta$ RNase | 110 | 218 | 30,000 (GBn2) | 75 (74.15) | 10.74 | -0.614 | 96.9% | 1.79× | X-ray $B$-factor: $r_s = +0.322$ | 46.8% ($r_s = +0.529$) |
| **HEW Lysozyme (1AKI)** | Two-domain glycosidase | 129 | 256 | 30,000 (PME: 17,110) | 106 (105.41) | 13.29 | -0.642 | 97.9% | 2.38× | NMR $S^2$: $r_s = +0.491$; $B$: +0.467 | 37.4% ($r_s = +0.562$) |
| **E. coli DHFR (1RX2)** | Catalytic $\alpha/\beta$ Rossmann | 159 | 316 | 30,000 (PME: 20,837) | 161 (160.58) | 79.15 | -0.663 | 98.0% | 2.46× | Distal $r_s = +0.229$; $B$: +0.240 | 45.8% ($r_s = +0.480$) |
| **Adenylate Kinase (4AKE)** | 3-domain allosteric enzyme | 214 | 426 | 30,000 (PME: 40,110) | 178 (177.57) | 13.41 | -0.635 | 98.8% | 1.74× | Hinge PR-AUC: 0.348 (AUC 0.644) | 37.0% ($r_s = +0.630$) |
| **Maltose-Binding (1OMP)** | 2-lobe periplasmic receptor | 370 | 738 | 30,000 (PME: 49,215) | 312 (311.86) | 17.83 | -0.643 | 99.3% | 1.58× | Hinge AUROC: 0.690 (DynDom 0.619) | 39.4% ($r_s = +0.556$) |
| *Part II: Analytical Microscopic Verification Peptides ($d = 2, 4$; Exact Toroidal Grid & Extended Phase-Space Benchmarks)* | | | | | | | | | | | |
| **Dialanine (ACE-Ala-NME)** | Capped model dipeptide | 1† | 2 | 155,000 (GBn2/PME) | 2 (1.48) | 1.45 | — | 0.0% | — | Contact Frob: 0.079 ($\ell/d = -0.922$) | Tier-5 barrier MFPT err: +7.8% |
| **Trialanine (ACE-Ala$_2$-NME)** | Capped model tripeptide | 2† | 4 | 150,000 (GBn2/OBC2) | 3 (2.44) | 1.83 | +0.171 | 50.0% | — | Contact Frob: 0.092 ($\ell/d = -0.778$) | Top-1 solvent SVD: 100% (mode 1) |

*Notes:* All numerical entries are populated directly from SHA-256-verified trajectories summing to 890,000 frames (890 ns at 1.0 ps stride; Replica 1 discovery, Replica 2 validation, Replica 3 frozen blind test; inter-replica spectral convergence $\bar{R}^2 = 0.996$; covariance scaling on the four $\ge 150$ ns ensembles evaluated on balanced 60,000-frame windows). †Capped ACE/NME peptides carry $d = 2N_{Ala}$ torsions, whereas uncapped $N_{res}$-residue proteins satisfy $d = 2N_{res} - 2$. Explicit-PME entries list total solvated particle counts in rhombic dodecahedral 0.15 M NaCl TIP3P-FB boxes (4 fs HMR). Crank $\bar{R}$: signed peptide-bond increment correlation $R(\Delta\psi_i, \Delta\phi_{i+1})$; Sparsity $S_{0.05}$: fraction of off-diagonal couplings below $\varepsilon = 0.05$; Enrich.: non-local tertiary $C_\alpha$–$C_\alpha$ contact coupling enrichment ($|i-j| \ge 4$, $\langle r_{ij}^{C_\alpha} \rangle < 0.85$ nm).





**Figure 1. Three-dimensional macromolecular contact-gating topology, distal catalytic loop routing, and mechanical domain-closure hinges across the explicit/implicit protein atlas ($d = 18 \to 738$).** (*A*) Ray-traced atomic structure of Human Ubiquitin (1UBQ, $N_{res} = 76$, $d = 150$) illustrating non-local tertiary $C_\alpha$ contact-gating chords $\mathcal{E}_C$ ($\bar{k}_{eff} = 5.97$, 94.7% off-diagonal sparsity), rigid $\beta$-grasp core amides reproduced parameter-free against BMRB 6457 solution NMR Lipari–Szabo $S^2$ order parameters ($r = +0.921$, $r_S = +0.512$), and the $\beta_1$–$\beta_2$ pincer (res 6–12) ↔ Ile44 hydrophobic recognition patch allosteric relay (7.1× blind amplification over sequence-banded operators). (*B*) Explicit-PME *E. coli* DHFR (1RX2, $d = 316$, 20,837 atoms) showing 3D heavy-atom contact bridges ($r_c^{heavy} < 0.50$ nm) connecting the active-site Met20 loop (res 9–24) to the distal $\beta$F–$\beta$G loop (res 116–132, Gly121 mutational hotspot > 15 Å away), recovering 71.6% of held-out blind MD coupling at 93.2% sparsity. (*C*) Explicit-PME apo Maltose-Binding Protein (1OMP open reference aligned on its N-terminal lobe against 1ANF closed holo, 35° C-lobe swing, $d = 738$, 49,215 atoms), highlighting the central $\beta/\alpha$ mechanical hinge linkers (res 110–114, 258–264, 311–318) localized directly from apo toroidal slow modes (AUROC = 0.690 vs. 0.446 Cartesian ANM).

timation suffers severe $2\pi$ jump discontinuities across the $\pm\pi$ branch cut. Applying maximal-gap circular chart shifting (dPCA+) (24) centers each angle's branch cut inside its widest zero-density steric barrier arc $\tilde{\theta}_a = (\theta_a - b_a + \pi) \bmod 2\pi - \pi$, preserving an exact $d$-dimensional intrinsic chart without doubling dimensionality to $2d$ $(\cos\theta, \sin\theta)$ coordinates. Across all 15 systems, independent replica spectral profiles exhibit inter-replica eigenvalue $R^2 = 0.996$, and label-shift-corrected variational mutual information lower bounds (25, 26) on Trp-cage confirm that backbone torsions span the slow collective folding subspace (1.000 principal subspace overlap with rotamer-augmented embeddings, while core $\chi_1/\chi_2$ sidechains contribute localized +12.8% state-discrimination information within Shannon entropy bounds).

**Law 1: Sub-Linear Slow-Mode Compression vs. Extensive Native-Basin Tail Variance.** Analyzing spectral dimensionality across two orders of magnitude ($d = 2 \to 738$; Figure 2) resolves a critical physical distinction between *collective dominant modes* and *cumulative native-basin thermal fluctuations*. Across the full 15-system atlas, integer 90% variance rank scales as $r_{90}(d) = 0.735\, d^{0.874}$ ($R^2 = 0.969$, $\Delta AIC_c = -4.8$; continuous rank $r_{90}^{cont}(d) = 0.577\, d^{0.919}$, $R^2 = 0.978$), compressing MBP ($d = 738$) by 2.37× to $r_{90} = 312$ (1.62× below its uncoupled marginal-variance null). Even more strikingly, the spectral participation ratio $PR(d) = (\sum_i \lambda_i)^2 / \sum_i \lambda_i^2$ and short-time ($\tau = 1$ ps) Kramers–Moyal diffusion operator rank $r_{90}^D(d)$ obey decisive sub-linear scaling laws across the entire macromolecular range (Figure 2B):

$$PR(d) = 1.061\, d^{0.494} \quad (\Delta AIC_c = -18.15), \qquad r_{90}^D(d) = 0.799\, d^{0.930} \quad (\Delta AIC_c = -14.39)$$

with 95% bootstrap confidence intervals $\alpha_{PR} \in [0.332, 0.666]$ and $\alpha_D \in [0.904, 0.998]$ strictly below unity (e.g., $PR = 13.41$ in AdK and 17.83 in MBP despite $d = 426$ and 738).

However, prospective out-of-scale extrapolation—fitting exclusively on the 10 discovery systems with $d \le 150$ ($r_{90}^{train}(d) = 0.762\, d^{0.819}$ vs. proportional linear $r_{90}^{lin}(d) = 0.402\, d$) and predicting zero-shot on the 5 unseen large proteins ($d \in \{218, 256, 316, 426, 738\}$)—reveals why naive global PCA truncation fails on large enzymes (Figure 2A). Whereas reversible fast folders ($d \le 68$) concentrate variance into low-dimensional folding funnels, 300 K explicit-solvent multi-domain enzymes populate hundreds of independent, locally rattling peptide planes across separate lobes, causing cumulative 90% variance across proteins ($d = 18 \to 738$) to transition to extensive proportionality ($r_{90} \approx 0.376\, d$, $R^2 = 0.975$, yielding 10.0% blind extrapolation MAPE vs. 35.7% for the small-protein power law). Thus, while collective allosteric modes remain low-dimensional ($PR \propto d^{0.494}$), preserving residue-resolved mechanics at $d > 200$ requires exploiting *spatial off-diagonal operator sparsity* rather than discarding $d - r$ tail modes.

**Laws 2 & 3: Crankshaft Parity, Contact Gating, and Real-Protein $O(d^{1.13})$ Sparse Chordal Estimation.** Decomposing off-diagonal correlation and diffusion matrices across all 13 proteins uncovers the microscopic origin of operator locality (Figure 3, Left). Along the polypeptide chain, off-diagonal couplings decay algebraically as $C(\Delta) = A\Delta^{-\beta} + c_0$, outperforming exponential decay across every protein ($\Delta AIC_c = -1.85$ in Chignolin, $-14.26$ in Ubiquitin, $-21.34$ in Barnase, $-9.62$ in AdK, and $-11.43$ in MBP). Crucially, inspecting signed increment correlations across adjacent torsions reveals a universal mechanical invariant across all 13 folded architectures (Figure 3, Left B): flanking torsions across each planar C′–N peptide bond exhibit strong anti-correlated *crankshaft counter-rotation*,

$$\bar{R}(\Delta\psi_i, \Delta\phi_{i+1}) = -0.603 \pm 0.054 \quad (\text{vs. intra-residue } \bar{R}(\Delta\phi_i, \Delta\psi_i) = -0.211)$$

reaching $-0.642$ in Lysozyme, $-0.663$ in DHFR, $-0.635$ in AdK, and $-0.643$ in MBP. Compensatory ($\Delta\psi_i \approx -\Delta\phi_{i+1}$) rotations rock individual peptide planes while localizing steric strain so distal segments are not swept through solvent. Beyond primary sequence separation ($|i-j| \ge 4$), 3D spatial proximity ($\langle r_{ij}^{C_\alpha} \rangle < 0.85$ nm) selectively gates non-local dynamical coupling with a 1.07×–3.05× enrichment (mean 2.17×; positive partial Spearman controlling for $|i-j|$ across all 13 folds; Figure 3, Left C). Because 3D steric packing bounds tertiary coordination, each dihedral couples above $\varepsilon = 0.05$ to only $\bar{k}_{eff} = 5.97$ partners ($k_{eff} \in [4.70, 7.95]$), driving





**Figure 2. Macromolecular manifold rank scaling, slow-mode compression, and blind out-of-scale extrapolation across $d = 2 \to 738$.** (*A*) Intrinsic 90% variance rank $r_{90}(d)$ across the 10-system discovery cohort ($d \le 150$, circles) and 5 unseen out-of-scale macromolecular proteins ($d \in \{218, 256, 316, 426, 738\}$, stars). Across all 15 systems, $r_{90}(d) = 0.735\, d^{0.874}$ ($R^2 = 0.969$, $\Delta\mathrm{AIC}_c = -4.8$), while within 300 K folded enzymes cumulative 90% variance transitions toward extensive proportionality ($r_{90} \approx 0.376\, d$, achieving 10.0% blind extrapolation MAPE vs. 35.7% for small-protein folding extrapolation). (*B*) Spectral participation ratio $PR(d) = (\sum \lambda_i)^2 / \sum \lambda_i^2$ compresses strongly sub-linearly as $PR(d) = 1.061\, d^{0.494}$ (95% CI $[0.332, 0.666]$, $\Delta\mathrm{AIC}_c = -18.15$), alongside short-time diffusion operator rank $r_{90}^{\mathbf{D}}(d) = 0.799\, d^{0.930}$ ($R^2 = 0.997$, $\Delta\mathrm{AIC}_c = -14.39$).





**Figure 3. Universal crankshaft locality, 3D contact gating, and $O(d^{1.13})$ sparse contact-graph operator estimation across real proteins ($d = 18 \to 738$).** (*Left, A–D*) Covalent sequence correlation decay $|R(\Delta)|$ favors algebraic power laws over exponentials ($\Delta\mathrm{AIC}_c = -1.85$ to $-21.34$), driven by universal anti-correlated peptide-bond crankshaft counter-rotation $\bar{R}(\psi_i, \phi_{i+1}) = -0.603 \pm 0.054$ (*B*) and positive partial Spearman tertiary contact gating (*C*), bounding effective degree to $\bar{k}_{\mathrm{eff}} = 5.97$ and driving sparsity $S_{0.05}(d)$ from 71.2% to 99.3% (*D*). (*Right, A–D*) Evaluated on blind Replica 3 trajectories, the sparse contact-graph Cholesky estimator achieves higher held-out Gaussian log-likelihood per DOF than dense Tikhonov and Graphical LASSO at $d \ge 218$ (*A*), avoids the high-dimensional sample-complexity crossover of dense covariance (*B*), preserves slow-subspace overlap (*C*), and scales in memory as $O(d^{1.06})$ vs. dense $O(d^{2.00})$ (65.8× compression on MBP, *D*).

operator sparsity $S_{0.05}(d) = 1 - k_{\mathrm{eff}}/(d-1)$ asymptotically to unity—from 71.2% in Chignolin ($d = 18$) to 96.9% in Barnase ($d = 218$), 98.8% in AdK ($d = 426$), and 99.3% in MBP ($d = 738$; Figure 3, Left D).

Exploiting this graph structure directly on real protein trajectories ($d = 18 \to 738$), we benchmarked a sparse contact-graph Cholesky estimator (Replica 1 graph/covariance, Replica 2 cutoff $r_c^{C_\alpha} \in \{0.65, 0.80\}$ nm and shrinkage validation, Replica 3 blind evaluation) against Diagonal-Only, Chordal Sequence-Banded, Tikhonov-tuned Full Dense, Ledoit–Wolf Linear Shrinkage, and $L_1$-penalized Graphical LASSO (Figure 3, Right). Empirically across the 13 proteins, contact-graph operator memory scales as $O(d^{1.06})$ versus $O(d^{2.00})$ for dense storage, and wall-clock estimation time scales as $O(d^{1.13})$ versus $O(d^{1.27})$ for Graphical LASSO. On MBP ($d = 738$), our sparse contact-graph estimator compresses operator memory by 65.8× (64.7 KB vs. 4255.0 KB dense) and accelerates estimation by 77.3× over Graphical LASSO (26.7 ms vs. 2062.7 ms, and 10.3× faster than dense 275.4 ms). Most importantly, above $d \ge 218$, eliminating $\sim 2.7 \times 10^5$ noisy non-contact off-diagonal entries prevents finite-sample variance inflation (Figure 3, Right A–B), enabling the contact-graph operator to achieve *higher* blind Replica 3 Gaussian log-likelihood per degree of freedom (+0.303/$d$ vs. dense +0.205/$d$ and Graphical LASSO +0.247/$d$) and lower relative Frobenius error (0.501 vs. dense 0.601).

### *2. SASA-Shielded Internal Friction, Solution NMR $S^2$ Order Parameters, and 13-Fold Zero-Shot PDB Transfer*

**Law 4: Relative SASA Reconciles Dry Protein-Core Equipartition with Peptide Viscosity Coupling.** A classic paradox in biomolecular dynamics is why short peptides exhibit super-linear thermal diffusivity scaling ($a(T) \propto (T/T_0)^{1.57}$) tracking bulk water viscosity reduction $\eta(T)^{-1}$, whereas folded protein interiors appear nearly insensitive to external solvent viscosity. Evaluating nested candidate laws across temperature (280–360 K) and Langevin bath collision sweeps ($\gamma \in [0.5, 5.0]$ ps$^{-1}$ conditioned on folded frames $Q_{\mathrm{BHE}} \ge 0.70$; Figure 4, Left) resolves this dichotomy via residue relative solvent-accessible surface area ($\mathrm{rSASA}_i$). Inside buried cores ($\mathrm{rSASA}_i < 0.20$), zero-parameter linear Einstein equipartition ($D(T) = D(T_0)(T/T_0)$) wins held-out parsimony selection with 12.33% blind prediction MAPE (beating 1-parameter power-law scaling at 12.79%; Figure 4, Left B) and exhibits high internal steric friction ($\gamma_{\mathrm{int}} \approx 250$ ps$^{-1}$) that shields core torsions from bath friction $\gamma$ (Figure 4, Left C). As $\mathrm{rSASA}_i$ increases from buried protein cores (rSASA $\approx 0.07$ in MBP, Ubiquitin, and Lysozyme) to exposed loops (rSASA $> 0.50$, $\gamma_{\mathrm{int}} = 14.0$ ps$^{-1}$) and fully solvated peptides (rSASA $\ge 0.90$), the effective thermal exponent $\alpha_{\mathrm{eff}}(\mathrm{rSASA}_i)$ interpolates monotonically from 1.00 to 1.28 and 1.57 (Figure 4, Left A).

**Law 5: Parameter-Free Solution NMR Lipari–Szabo $S^2$ Order Parameters and X-Ray $B$-Factors.** To test whether toroidal operators predict residue-resolved spectroscopic observables without trajectory-fitted scaling parameters, we derived the exact second-order kinematic Jacobian $\mathbf{J}_i \in \mathbb{R}^{3\times4}$ mapping local 4×4 flanking toroidal covariances $\hat{\boldsymbol{\Sigma}}_{\mathrm{local},i}$ on $(\psi_{i-1}, \phi_i, \psi_i, \phi_{i+1})$ onto unit amide N–H$_i$ bond-vector fluctuations (27–30):

$$\hat{S}_i^2 = 1 - \frac{3}{2}\,\mathrm{Tr}\left(\mathbf{J}_i \hat{\boldsymbol{\Sigma}}_{\mathrm{local},i} \mathbf{J}_i^{\mathsf{T}}\right)$$

Evaluated across experimental Biological Magnetic Resonance Data Bank (BMRB)





**Figure 4. SASA-shielded internal vs. solvent friction (Law 4) and parameter-free solution NMR Lipari–Szabo $S^2$ validation (Law 5).** (*Left, A–D*) Effective thermal mobility exponent $\alpha_{\text{eff}} = \partial \ln D / \partial \ln T$ transitions smoothly with relative solvent-accessible surface area (rSASA$_i$) from dry Einstein equipartition in buried protein cores ($\alpha_{\text{core}} = 1.00$) through surface loops ($\alpha_{\text{loop}} = 1.28$, $\gamma_{\text{int}} = 14.0\ \text{ps}^{-1}$) to bulk-viscosity-coupled short peptides ($\alpha_{\text{peptide}} = 1.57$), with zero-parameter linear Einstein equipartition achieving 12.33% blind thermal MAPE (*B*) and rank-1 cross-solvent SVD dominance (*D*). (*Right, A–D*) Kinematic Jacobian projection of $4 \times 4$ toroidal crankshaft blocks predicts BMRB solution NMR $S^2$ order parameters across 387 internal flanking-quartet residues (*A*; Ubiquitin profile in *B*: $r_s = +0.512$, $r = +0.921$, RMSE = 0.110), compares favorably against baseline models (*C*), and resolves peptide-plane rocking excess X-ray $B$-factors driven by $|\Sigma_{\psi_{i-1},\phi_i}|$ (*D*).

solution NMR $S^2$ datasets ($n = 387$ internal residues where all four flanking torsions are defined) for four benchmark enzymes—Human Ubiquitin (1D3Z/BMRB 6457: $r_s = +0.512$, Pearson $r = +0.921$, RMSE = 0.110; Figure 4, Right B), GB1 (2QMT/BMRB 7280: $r_s = +0.746$, RMSE = 0.153), HEW Lysozyme (1AKI/BMRB 4562: $r_s = +0.491$, RMSE = 0.137), and *E. coli* DHFR (1RX2/BMRB 4380: $r_s = +0.182$, where microsecond cofactor-loop exchange contributes beyond sub-nanosecond librations)—the $4\times4$ crankshaft-coupled operator achieves a higher mean within-protein Spearman rank correlation ($\bar{r}_s = +0.483$) than both 1D flanking diagonal variance ($\bar{r}_s = +0.462$) and validation-tuned isotropic GNM ($\bar{r}_s = +0.408$), as well as the highest pooled Pearson correlation ($r = +0.522$, $p = 3.50\text{e-}07$, RMSE = 0.138; Figure 4, Right A). Notably, when pooling un-normalized $S^2$ values across all four proteins measured at heterogeneous NMR temperatures (285–308 K), LOPO-linearly calibrated GNM achieves a higher pooled rank correlation ($r_s = +0.378$ vs. parameter-free crankshaft +0.255 and direct MD +0.311; Figure 4, Right C) because its fitted linear slope/intercept absorbs baseline offset differences across proteins, whereas within any single protein fold covalent crankshaft cancellation ($\Sigma_{\psi_{i-1},\phi_i} < 0$) is essential to capture rigid peptide-plane rocking and crystallographic $B$-factor profiles across 8 X-ray structures ($\bar{r}_s = +0.276$: Lysozyme +0.467, BPTI +0.346, Barnase +0.322, DHFR +0.240, Ubiquitin +0.221, GB1 +0.221, AdK +0.206, and MBP +0.188; Figure 4, Right D).

**Law 6: 13-Fold Zero-Shot Operator Prediction From Static PDB Coordinates.** Across all 13 proteins (2,570 dihedrals), folded interior backbone mobilities cluster around secondary-structure plateaus of 0.0320 rad$^2$/ps in $\alpha$-helices, 0.0407 rad$^2$/ps in $\beta$-strands, and 0.0485 rad$^2$/ps in loops ($\bar{D}_{\text{interior}} = 0.0419$ rad$^2$/ps, CV = 24.7%; Figure 5D). Training strictly on 12 proteins to predict $\ln D_{aa}$ and contact-gated off-diagonal couplings on the held-out 13th protein from native PDB features (12-feature structural ridge: secondary structure, Gly/Pro/pre-Pro identity, chain-terminus depth, $C_\alpha$ contact burial, rSASA$_i$, and hydrogen-bond geometry) yields pooled zero-shot Spearman $r_s = +0.706$ ($p < 10^{-100}$, Pearson $r = +0.735$, mean out-of-sample $R^2 = +0.359$ vs. 9-feature local ridge baseline $R^2 = +0.350$) and lowers full-matrix Frobenius error from 60.0% (constant null) to 45.5% (Figure 5A–C; e.g., BBA5 29.9%, AdK 37.0%, Lysozyme 37.4%, MBP 39.4%). Nonlinear histogram gradient-boosted decision trees (GBDT) further improve held-out accuracy to $r_s = +0.721$, $R^2 = +0.410$, and 44.8% Frobenius error toward the 16.9% inter-replica floor.

### 3. Mechanistic Predictions in Multi-Domain Allosteric Enzymes and Fast-Folding Proteins

**Mechanical Hinge Localization in Explicit-Solvent Apo AdK ($d = 426$) and MBP ($d = 738$).** To test whether reduced toroidal operators extract functional conformational mechanisms in large multi-domain enzymes without access to ligand-bound holo crystal structures, we analyzed 30,000 explicit-PME apo open-state trajectory frames ($3 \times 10$ ns) alongside static open PDB structures for *E. coli* Adenylate Kinase (4AKE open → 1AKE closed; CORE, LID, and NMP domains, 214 residues) (31, 32) and Maltose-Binding Protein (1OMP open → 1ANF closed; N- and C-terminal periplasmic lobes connected by a three-segment $\beta/\alpha$ linker, 370 residues) (33) (Figures 1C, 6). Computing residue hinge scores from the top slow collective eigenmodes of the apo toroidal covariance weighted by local tertiary contact torque ($\log(1 + n_{\text{contacts},i})$) reveals sharp mechanical peaks aligned with the crystallographic NMP/LID hinges in AdK and the inter-lobe linker strands/helix in MBP (Figure 6A–B).

Quantitatively, in 370-residue MBP ($d = 738$; Figure 6D), the explicit-PME apo toroidal operator predicts crystallographic open-to-closed strain hinges with AUROC = 0.690 and continuous Spearman $\rho = +0.315$ ($p = 6.7 \times 10^{-10}$), substantially outperforming both Cartesian ANM (AUROC = 0.446) and Kabsch-superposed Cartesian apo-MD PCA (AUROC = 0.635). Evaluated against literature DynDom inter-lobe consensus hinges, static contact-gated toroidal modes achieve AUROC = 0.619 (and apo-MD toroidal modes reach 0.693) compared to 0.306 for Cartesian ANM, which misassigns maximum displacement to peripheral solvent-exposed lobe tips rather than the central mechanical pivot. In AdK ($d = 426$; Figure 6C), where peripheral LID/NMP translations correlate more closely with harmonic lever-arm amplitude, static and apo-MD toroidal operators match Cartesian ANM in global ROC-AUC (0.644 and 0.630 vs. 0.647 ANM, beating Cartesian apo PCA 0.580), while achieving 51% higher precision-recall (PR-AUC = 0.348, Precision@15 = 0.400 vs. ANM PR-AUC = 0.230, Precision@15 = 0.267), continuous Spearman $\rho = +0.312$ ($p = 3.7 \times 10^{-6}$), and higher DynDom consensus accuracy (AUROC = 0.529, Precision@15 = 0.267 vs. 0.000 ANM).

**Distal Allosteric Loop Routing in DHFR ($d = 316$), Lysozyme ($d = 256$), and Ubiquitin ($d = 150$).** Beyond localized hinges, enzymatic catalysis in *E. coli* DHFR (1RX2) relies on distal allosteric communication between the active-site Met20 loop (res 9–24) and the $\beta$F–$\beta$G (FG) loop (res 116–132, housing the classic Gly121 mutational hotspot > 15 Å from the catalytic center) (34). Evaluating the matrix-exponential Markov propagator $\mathcal{P}(\tau) = \exp(-\tau \hat{\mathbf{D}} \hat{\mathbf{\Sigma}}^{-1})$ at $\tau = 5$ ps (heavy-atom tertiary contact gating $r_c^{\text{heavy}} = 0.50$ nm) strictly on blind Replica 3 trajectories for sequence-distant pairs ($|i-j| \geq 10$; Figures 1A–B, 7) demonstrates that contact-gated toroidal operators achieve the lowest distal coupling RMSE across all three benchmark enzymes: 0.0441 in Ubiquitin (85.9% sparsity), 0.0335 in Lysozyme (91.1% sparsity, $r_s = +0.446$, Pearson $r = +0.448$), and 0.0441 in DHFR (93.2% sparsity, $r_s = +0.229$, Pearson $r = +0.153$), compared to $3.0\times$–$3.6\times$ larger errors for Kabsch-aligned Cartesian dynamic cross-correlation matrices (RMSE = 0.1580, 0.1182, and 0.1289; DHFR DCCM $r_s = +0.021$; Figure 7B), which are contaminated by global rigid-body domain rocking. Crucially, compared to purely sequence-banded operators ($w = 2$), non-local 3D contact edges increase blind Replica 3 Met20 ↔ FG loop relay recovery in DHFR from 0.0238 to 0.0327 (71.6% of ground-truth MD 0.0458; Figure 7C–D) and amplify $\beta_1$–$\beta_2$ pincer (res 6–11) ↔ Ile44 hydrophobic recognition-patch coupling in Ubiquitin by $7.1\times$ (0.0116 vs. 0.0016 sequence-banded; MD 0.0300).

**Multi-Protein Fast-Folding Kinetics and Mori–Zwanzig Memory Scaling.** Across 7 proteins spanning $d = 18 \rightarrow 150$ (Chignolin, Trp-cage, BBA5, Villin HP35, WW-FiP35, GB1, and Ubiquitin; Figure 8), projecting $d$ backbone torsions onto the smooth Best–Hummer–Eaton native contact coordinate $Q_{\text{BHE}} \in [0,1]$ (17) reveals a universal sub-diffusive Mori–Zwanzig memory regime: finite-lag diffusivity decays across $\tau \in [1, 400]$ ps as $\bar{D}(\tau) \propto \tau^{-\gamma_M}$ with mean exponent $\bar{\gamma}_M = 0.593 \pm 0.126$ (Chignolin 0.716 at $\tau^* = 50$ ps; Figure 8B, F), trending downward with chain dimension ($r_s = -0.714$, exact permutation $p = 0.088$) as larger native cores damp short-time ballistic recrossings. Enforcing cryptographic SHA-256 prospective





**Figure 5. Law 6: 13-fold zero-shot Leave-One-Protein-Out (LOPO) operator prediction from static PDB coordinates ($d = 18 \to 738$).** (*A*) Parity plot across all $n = 2,570$ held-out backbone dihedrals in 13-fold LOPO cross-validation using 12-feature structural ridge regression (pooled $r_S = +0.706$, $r = +0.735$, $\bar{R}^2 = +0.359$). (*B–C*) Out-of-sample $R^2$ (*B*) and full $d \times d$ operator Frobenius error (*C*) across nested structural models relative to the constant baseline (60.0%) and inter-replica MD noise floor (16.9%); histogram gradient-boosted decision trees (GBDT) reach $r_S = +0.721$, $\bar{R}^2 = +0.410$, Frob = 44.8%. (*D*) Conserved interior mobility plateaus across secondary structure classes (α-helix 0.0320, β-strand 0.0407, coil 0.0485, all interior 0.0419 rad$^2$/ps, CV = 24.7%).





**Figure 6. Mechanical hinge localization in explicit-solvent apo Adenylate Kinase ($d = 426$) and Maltose-Binding Protein ($d = 738$).** (*A–B*) Residue-resolved crystallographic open-to-closed dihedral strain $\Delta\theta_i^{xtal}$ (solid black: 4AKE→1AKE in AdK; 1OMP→1ANF in MBP) overlaid with predicted toroidal hinge operator profiles (dashed) and canonical domain-closure hinge segments (shaded). (*C*) In AdK, toroidal operators match Cartesian ANM in overall ROC-AUC (0.644 static, 0.630 apo MD vs. 0.647 ANM and 0.580 Cartesian apo PCA) while achieving 1.5× higher precision-recall (PR-AUC = 0.348 vs. 0.230) and Spearman $\rho = +0.312$. (*D*) In 370-residue MBP, the explicit-PME apo toroidal operator achieves AUROC = 0.690 ($\rho = +0.315$), decisively outperforming Cartesian ANM (0.446) and Cartesian apo PCA (0.635), while static contact-gated toroidal modes beat ANM on DynDom inter-lobe hinges (0.619 vs. 0.306).

model freezing prior to Replica 3 inspection (Supporting Information Section S4) confirms exact detailed balance to machine precision ($< 10^{-15}$ across all 7 systems) and reduces blind Replica 3 committed MFPT log-error from 1.575 (uncalibrated $\tau = 1$ ps Kramers–Moyal $\mathcal{K}_0$) to 1.354 under memory-calibrated $\mathcal{K}_2$ (Figure 8C–E; strict Pois-





**Figure 7. Blind Replica 3 distal allosteric loop routing via sparse contact-gated Markov propagators in DHFR ($d = 316$), Lysozyme ($d = 256$), and Ubiquitin ($d = 150$).** (*A*) Distal propagator parity ($|i - j| \geq 10$ residues, $\tau = 5$ ps) on held-out Replica 3 trajectories in *E. coli* DHFR comparing contact-gated vs. sequence-banded operators. (*B*) Distal coupling Spearman rank correlation on DHFR (+0.229 contact-gated vs. +0.216 sequence-banded and +0.021 Cartesian DCCM). (*C*) Catalytic Met20 (res 9–24) ↔ distal FG loop (res 116–132, Gly121) allosteric relay strength on blind Replica 3: contact-gated operators recover 71.6% (0.0327) of ground-truth MD coupling (0.0458) vs. 0.0238 for sequence-banded operators at 93.2% matrix sparsity. (*D*) Full 159 × 159 residue-resolved contact-gated propagator $\mathcal{P}_{ij}$ (5 ps) of DHFR.





**Figure 8. Multi-protein folding committors, sub-diffusive Mori–Zwanzig memory scaling, and blind kinetic validation across 7 proteins ($d = 18 \rightarrow 150$).** (*A*) Exact adjoint committor profiles $q_+(Q_{\mathrm{BHE}})$ and transition-state separatrices ($q_+ = 0.5$). (*B*) Universal power-law memory attenuation $\bar{D}(\tau) \propto \tau^{-\gamma_M}$ across coarse-graining lag times $\tau \in [1, 400]$ ps ($\bar{\gamma}_M = 0.593 \pm 0.126$). (*C*) Prospective blind Replica 3 MFPT parity comparing uncalibrated $\mathcal{K}_0$ ($\tau = 1$ ps, crosses, mean log-error 1.575) against forward-calibrated $\mathcal{K}_2$ ($\tau^*$, circles, mean log-error 1.354). (*D*) Hummer transition-path durations $\tau_{\mathrm{TP}}$. (*E*) Empirical 2D free-energy surface $\Delta F(Q_{\mathrm{BHE}}, \psi_1)$ and $q_+ = 0.5$ separatrix on Chignolin. (*F*) Scaling of memory attenuation exponent $\gamma_M$ with chain dimension $d$ ($r_S = -0.714$, exact permutation $p = 0.088$).

son 95% CI hit rate 1/6 on ultra-rare unfolded excursions due to replica-to-replica sampling variance, while capturing Hummer transition-path times $\tau_{\mathrm{TP}}$ within 1–2 ps on Trp-cage through GB1; Figure 8D).

#### 4. Analytical Microscopic Verification and Fair 6-Paradigm SOTA Benchmark

**Microscopic Verification on Model Peptides ($d = 2, 4$).** Capped alanine dipeptide ($d = 2,155,000$ frames) and trialanine ($d = 4,150,000$ frames) provide exact low-dimensional analytical controls where dense $6^2$–$6^4$ Fourier/grid operators can be solved without spatial truncation: (i) a 5-tier tensor ablation on dialanine confirms that coupled state-dependent diagonal diffusion anisotropy (10.3× saddle mobility surge along $\phi$ at $-1.2$ rad) reduces barrier-crossing MFPT error from +145.8% (constant isotropic $D_0\mathbf{I}$) to +18.0% (state-dependent diagonal) and +7.8% (full positive-definite tensor, KS $p = 0.612$); (ii) under low solvent friction ($\gamma = 0.5$ ps$^{-1}$, Kramers turnover), embedding torsions into an auxiliary Markovian phase space $(z, v, w) \in \mathbb{T}^2 \times \mathbb{R}^2 \times \mathbb{R}^2$ parameterized directly from the second-order analytical velocity autocorrelation function ($R^2 \geq 0.9948$) reduces MFPT error from 47.0% (overdamped) to 25.8% (KS $p = 0.210$) and compresses multi-step non-Markovian memory $M_{32}$ from 0.330 to 0.026 nats; and (iii) across implicit/explicit solvent perturbations, residual diffusion shifts collapse onto a dominant rank-1 singular mode (Figure 4, Left D).

**Fair 6-Paradigm Validation-Tuned SOTA Benchmark Across All 13 Proteins.** Finally, to establish precisely where continuous contact-gated GENERIC operators advance beyond existing methods—and where classical paradigms remain preferable—we executed a controlled benchmark across all 13 proteins ($d = 18 \rightarrow 738$; Figure 9, Table 2). Every baseline (VAMP-2 Reversible MSM, TICA Linear Langevin, Calibrated 1D Smoluchowski, Unconstrained Basis/Polynomial SDE, our Contact-Gated GENERIC Operator, and Calibrated GNM/ENM) underwent a full hyperparameter grid search on Replica 2 (Figure 9A) before single-shot blind evaluation on Replica 3:

1. *Full-Dimensional Operator Reconstruction (13/13 Wins):* Across every protein from Chignolin (0.095) to AdK (0.297) and MBP (0.365), the Contact-Gated GENERIC operator achieves the lowest blind Replica 3 diffusion operator relative Frobenius error (mean 0.288, beating unconstrained SDE 0.497, diagonal Smoluchowski 0.754, and low-rank TICA Langevin 1.759; Figure 9D).
2. *Thermodynamic Consistency and Lyapunov Stability:* Unconstrained regression SDEs violate microscopic reversibility ($J_{\text{ss}} \neq 0$), producing a mean spurious entropy rate of 346.17 $k_B$/ns/$d$ (102.25 in Ubiquitin, 104.66 in AdK, 73.51 in MBP; Figure 9C). Enforcing the GENERIC Helmholtz–Hodge identity $f_{\text{GENERIC}}(z) = -\mathbf{D}(z)\nabla[\beta F(z)] + \operatorname{div}\mathbf{D}(z)$ suppresses 100.0% of spurious cycle flux ($\dot{S}_{\text{irr}} \equiv 0.000$) and guarantees strict Hurwitz Lyapunov stability ($\max \operatorname{Re}\lambda(\mathbf{A}) < 0$ across 13/13 proteins).
3. *Zero-Shot Structural Generalization and Direct Disorder:* Unlike trajectory-only estimators (MSM, TICA, Smoluchowski, and unconstrained SDEs), our 13-fold LOPO structural prior predicts unseen protein operators from PDB coordinates at $r_s = +0.706$ vs. +0.328 for calibrated ENM (Figure 9E), and achieves the highest uncalibrated Replica 3 residue-disorder rank correlation across all 398 BMRB-matched NMR residues ($r_s = +0.348$, $p = 9.2\times10^{-13}$, including 11 chain-border positions excluded from the 387-residue $4\times4$ flanking-quartet Jacobian in Law 5; vs. MSM +0.307, SDE +0.250, and uncalibrated ENM +0.169).
4. *Candid Ties and Baseline Strengths:* Along a pre-defined 1D folding coordinate $Q_{\text{BHE}}$, Contact-Gated GENERIC (1.76) and calibrated 1D Smoluchowski (1.76) achieve a statistical tie on blind Replica 3 MFPT log-error (both beating unconstrained SDE 2.19, TICA Langevin 3.34, and discrete MSM 3.68; Figure 9B), confirming that 1D Smoluchowski suffices when only scalar folding rates are sought. Conversely, when linear slope/intercept LOPO rescaling is applied across pooled multi-temperature NMR $S^2$ datasets, calibrated GNM achieves the highest pooled rank correlation (+0.378; Figure 9F) at $< 0.2$ s trajectory-free cost, whereas our crankshaft operator wins within-protein parameter-free $S^2$ (+0.483 vs. +0.408) and Pearson linearity (+0.522).

## DISCUSSION

Extending continuous stochastic operator discovery from short peptides to multi-domain explicit-solvent proteins up to $d = 738$ (49,215 particles) yields three broad insights for biomolecular theory and multiscale simulation.

First, **macromolecular dimensional reduction is dual-natured: slow functional modes compress sub-linearly, whereas native-basin thermal variance is extensive.** In small fast-folding miniproteins ($d \leq 68$), cooperative folding funnels compress both $PR(d)$ and $r_{90}(d)$. In large multi-domain enzymes at 300 K (`Lysozyme`, `DHFR`, `AdK`, `MBP`), however, dominant allosteric and hinge modes still concentrate into a compact sub-linear subspace ($PR(d) = 1.061\, d^{0.494}$), yet cumulative 90% variance grows proportionally with chain length ($r_{90} \approx 0.376\, d$). Consequently, discarding $d - r$ modes via low-rank TICA/PCA loses local loop and sidechain-coupled backbone mechanics (Frob = 1.759), whereas dense $d \times d$ estimation suffers sample-complexity degradation above $d > 200$. Our sparse contact-graph Cholesky architecture resolves this dilemma by retaining all $d$ physical torsions while zeroing non-contact off-diagonal noise (99.3% sparsity at $d = 738$), simultaneously delivering $O(d^{1.13})$ runtime, 65.8× memory compression, and superior held-out log-likelihood.

Second, **covalent peptide-plane crankshaft geometry and 3D contact topology directly bridge MD trajectories, NMR relaxation, and allostery.** The universal $\bar{R}(\psi_i, \phi_{i+1}) = -0.603$ anti-correlation across all 13 folds demonstrates that the planar C′–N peptide group acts as a local mechanical rock-and-cancel linkage: accounting for $\Sigma_{\psi_{i-1},\phi_i} < 0$ is essential to predict solution NMR Lipari–Szabo $S^2$ order parameters parameter-free from dihedral fluctuations, while augmenting covalent bands with sparse 3D tertiary contacts captures distal Met20 ↔ FG loop coupling in DHFR (34) and inter-lobe mechanical hinges in MBP (33) and AdK (31, 32) without rigid-body Kabsch alignment artifacts.

Third, **structure-preserving GENERIC inductive biases complement discrete MSMs and generative machine learning.** In Riemannian flow-matching and graph neural network (GNN) coarse-grained force fields on $\mathbb{T}^d$ (20, 23), replacing dense $O(d^2)$ all-to-all attention and unconstrained drift networks with our sparse band-plus-contact graph ($\bar{k}_{\text{eff}} = 5.97$) and reversible GENERIC projection ($J_{\text{ss}} \equiv 0$) guarantees infinite-horizon Boltzmann stability while preserving barrier diffusivity surges and rSASA$_i$-interpolated internal friction. Future work integrating state-dependent contact-graph switching $\mathcal{E}(z)$ will extend these operators to intrinsically disordered proteins (IDPs) and millisecond domain-swapping assemblies.

## MATERIALS AND METHODS

**Atomistic MD Atlas and Cryptographic Governance.** Explicit-solvent simulations (`Lysozyme 1AKI`, `DHFR 1RX2`, `AdK 4AKE`, `MBP 1OMP`, plus peptide solvent controls) were performed in OpenMM 8.4/8.6 (35) using AMBER `ff99SB-ILDN + TIP3P-FB` water in rhombic dodecahedral boxes (1.2 nm padding, 0.15 M NaCl, PME 1.0 nm real-space cutoff, Monte Carlo barostat at 1.0 atm, `LangevinMiddleIntegrator` BAOAB at 300 K, $\gamma = 1.0\,\text{ps}^{-1}$, 4.0 fs timestep with 1.5 amu hydrogen mass repartitioning, saving every 1.0 ps). Matched implicit-solvent ensembles (`GBn2` and `OBC2` with AMBER `ff14SB/ff99SB-ILDN`) were integrated at 2.0 fs timestep (1.0 ps stride). Every trajectory chunk was locked with SHA-256 checksums (Supporting Information Table S3) and partitioned into Replica 1 (discovery), Replica 2 (hyperparameter validation), and Replica 3 (single-shot blind test).

**Toroidal dPCA+, Contact-Graph Cholesky, and GENERIC Operator.** Backbone torsions $\boldsymbol{\theta} \in [-\pi, \pi)^d$ were unwrapped via maximal-gap dPCA+ shifts computed strictly on Replica 1 (24). Short-time diffusion tensors $\hat{\mathbf{D}} = (2\tau)^{-1}\langle \Delta_{\mathbb{T}^d}\boldsymbol{\theta}(\tau)\Delta_{\mathbb{T}^d}\boldsymbol{\theta}(\tau)^\top\rangle$ ($\tau = 1$ ps) and stationary covariances $\hat{\boldsymbol{\Sigma}}$ were projected onto the sparse graph $\mathcal{G} = \{(a,b) : |\text{res}(a) - \text{res}(b)| \leq w \text{ or } \langle r_{\text{res}(a),\text{res}(b)}\rangle_{\text{disc}} < r_c\}$ via sparse chordal Cholesky completion and Replica 2-validated shrinkage $\beta_c$. Reversible GENERIC drift fields were enforced via $f_{\text{GENERIC}}(z) = -\hat{\mathbf{D}}(z)\nabla[\beta F(z)] + \operatorname{div}\hat{\mathbf{D}}(z)$, guaranteeing $\dot{S}_{\text{irr}} \equiv 0$.

**NMR $S^2$ Jacobian, Allosteric Propagators, and Benchmark Baselines.** Amide N–H$_i$ order parameters $\hat{S}_i^2$ were evaluated analytically from $\mathbf{J}_i = \partial\mathbf{u}_{\text{NH},i}/\partial(\psi_{i-1}, \phi_i, \psi_i, \phi_{i+1})$ at native reference geometry without free parameters against experimental BMRB entries `6457` (28), `7280` (29), `4562` (30), and `4380` (36). Allosteric Markov propagators $\mathcal{P}(\tau) = \exp(-\tau\hat{\mathbf{D}}\hat{\boldsymbol{\Sigma}}^{-1})$ and AdK/MBP hinge scores were evaluated against crystallographic `4AKE/1AKE` and `1OMP/1ANF` strain and DynDom consensus annotations (31–33). All six modeling paradigms in Table 2 were tuned on Replica 2 and evaluated once on Replica 3; complete derivations and provenance digests are provided in the Supporting Information.

## DATA AND CODE AVAILABILITY

All simulation scripts, SHA-256 trajectory manifests, frozen model locks, and automated analysis pipelines are available at `https://github.com/Mehardeep-Singh-GG/discovering-reduced-dynamical-laws-md`. Complete mathematical proofs, per-protein tables, and validation sweep logs are provided in the Supporting Information.

## ACKNOWLEDGMENTS

We thank the developers of OpenMM, MDTraj, deeptime, SciPy, and the Biological Magnetic Resonance Data Bank (BMRB). **Funding:** No external funding was received. **Competing Interests:** None declared.

## REFERENCES

1. Karplus, M. & Kuriyan, J. Molecular dynamics and protein function. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 6679–6685 (2005).
2. Lindorff-Larsen, K., Piana, S., Dror, R. O. & Shaw, D. E. How fast-folding proteins fold. *Science* **334**, 517–520 (2011).
3. Henzler-Wildman, K. & Kern, D. Dynamic personalities of proteins. *Nature* **450**, 964–972 (2007).
4. Dill, K. A. & MacCallum, J. L. The protein-folding problem, 50 years on. *Science* **338**, 1042–1046 (2012).
5. Boehr, D. D., Nussinov, R. & Wright, P. E. The role of dynamic conformational ensembles in biomolecular recognition. *Nat. Chem. Biol.* **5**, 789–796 (2009).
6. Kramers, H. A. Brownian motion in a field of force and the diffusion model of chemical reactions. *Physica* **7**, 284–304 (1940).
7. Onsager, L. Reciprocal relations in irreversible processes. I. *Phys. Rev.* **37**, 405–426 (1931).
8. Zwanzig, R. Memory effects in irreversible thermodynamics. *Phys. Rev.* **124**, 983–992 (1961).
9. Mori, H. Transport, collective motion, and Brownian motion. *Prog. Theor. Phys.* **33**, 423–455 (1965).

**Table 2. Fair validation-tuned head-to-head benchmark across six dynamical modeling paradigms on frozen blind test trajectories (Replica 3) across all 13 macromolecular proteins ($d = 18 \rightarrow 738$)**

| Method Family | Validation (Replica 2) Sweep Grid | A: Blind MFPT Log-Err | B: Spurious $\dot{S}_{\mathrm{irr}}$ ($k_B$/ns/$d$) | C: Blind D Frob. Err | D: 13-Fold LOPO ($r_s$) | E: Direct Disorder / Cal. $S^2$ | Optimal Physical Regime & Candid Trade-Offs |
|---|---|---|---|---|---|---|---|
| **Contact-Gated GENERIC (Ours)** | $r_c = 0.65$ nm, $\beta_c \in [0, 0.5]$, $\tau^*$ | **1.76 (Tie #1)** | **0.000 (Exact, 13/13)** | **0.288 (#1 on 13/13)** | **+0.706 (#1)** | **+0.348 (#1)** / +0.255 | Full $d$-dim reversible operator, hinges, distal allostery, PDB transfer |
| **VAMP-2 Reversible MSM** | $k \in [20, 200]$, $\tau \in [2, 50]$ ps | 3.68 | **0.000** (0.79 non-rev) | N/A (discrete $k \times k$) | N/A (traj-only) | +0.307 / +0.307 | Coarse multi-basin populations & flux networks with long sampling |
| **TICA Linear Langevin** | $r \in [3, 12]$, $\tau \in [2, 20]$ ps | 3.34 | **0.000** (Gaussian OU) | 1.759 | N/A (traj-only) | +0.145 / +0.145 | Optimal linear slow-subspace basis; underfits anharmonic barriers |
| **Calibrated 1D Smoluchowski** | $M \in [20, 40]$, $\tau^* \in [1, 100]$ ps | **1.76 (Tie #1)** | **0.000** (1D exact) | 0.754 (diag null) | N/A (1D $Q$ only) | -0.110 / -0.110 | Fast, exact 1D barrier kinetics once reaction coordinate $Q$ is known |
| **Unconstrained Basis/Poly SDE** | $B \in [1, 3]$, $\lambda \in [10^{-4}, 10^{-1}]$ | 2.19 | **346.17** ($J_{\mathrm{ss}} \neq 0$) | 0.497 | N/A (overfits fold) | +0.250 / +0.250 | Flexible drift regression, but violates detailed balance ($\dot{S}_{\mathrm{irr}} \gg 0$) |
| **Calibrated GNM / ENM** | $r_c \in [0.7, 1.5]$ nm | N/A (harmonic) | **0.000** (static eq.) | N/A (unscaled $C_\alpha$) | +0.328 | +0.169 / **+0.378 (#1 cal.)** | $< 0.2$ s zero-MD crystal $B$-factors & linearly rescaled pooled $S^2$ |





**Figure 9. Fair 6-paradigm validation-tuned SOTA benchmark across all 13 proteins (Replica 1 discovery → Replica 2 hyperparameter sweep → Replica 3 single-shot blind test).** (*A*) Representative Replica 2 cross-validated VAMP-2 tuning curves selecting optimal MSM cluster counts $k^*$. (*B*) Metric A: mean blind Replica 3 committed MFPT log-error, showing a statistical tie between calibrated 1D Smoluchowski (1.76) and Contact-Gated GENERIC (1.76). (*C*) Metric B: steady-state spurious entropy production $\dot{S}_{\mathrm{irr}}$, where unconstrained SDEs generate 346.17 $k_B$/ns/$d$ vs. exact 0.000 for GENERIC projection. (*D*) Metric C: blind Replica 3 diffusion tensor Frobenius error across all 13 proteins (Contact-Gated GENERIC wins 13/13, mean 0.288). (*E*) Metric D: 13-fold zero-shot PDB-to-mobility correlation (+0.706 toroidal LOPO vs. +0.328 calibrated ENM). (*F*) Metric E: pooled cross-temperature BMRB NMR $S^2$ correlation after linear LOPO calibration.


10. Zwanzig, R. *Nonequilibrium Statistical Mechanics* (Oxford Univ. Press, 2001).
11. Prinz, J.-H. et al. Markov models of molecular kinetics: Generation and validation. *J. Chem. Phys.* **134**, 174105 (2011).
12. Chodera, J. D. & Noé, F. Markov state models of biomolecular kinetics. *Curr. Opin. Struct. Biol.* **25**, 135–144 (2014).
13. Perez-Hernandez, G., Paul, F., Giorgino, T., De Fabritiis, G. & Noé, F. Identification of slow molecular order parameters for Markov model construction. *J. Chem. Phys.* **139**, 015102 (2013).
14. Bahar, I., Atilgan, A. R. & Erman, B. Direct evaluation of thermal fluctuations in proteins using a single-parameter harmonic potential. *Fold. Des.* **2**, 173–181 (1997).
15. Atilgan, A. R. et al. Anisotropy of fluctuation dynamics of proteins with an elastic network model. *Biophys. J.* **80**, 505–515 (2001).
16. Hummer, G. Position-dependent diffusion coefficients and free energy profiles from slow, equilibrium molecular dynamics simulations. *New J. Phys.* **7**, 34 (2005).
17. Best, R. B., Hummer, G. & Eaton, W. A. Native contacts determine protein folding mechanisms in atomistic simulations. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 17874–17879 (2013).
18. Hänggi, P., Talkner, P. & Borkovec, M. Reaction-rate theory: fifty years after Kramers. *Rev. Mod. Phys.* **62**, 251–341 (1990).
19. Best, R. B. & Hummer, G. Coordinate-dependent diffusion in protein folding. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 1088–1093 (2010).
20. Noé, F., Olsson, S., Köhler, J. & Wu, H. Boltzmann generators: Sampling equilibrium states of many-body systems with deep learning. *Science* **365**, eaaw1147 (2019).
21. Klus, S. et al. Data-driven approximation of the Koopman generator. *J. Nonlinear Sci.* **30**, 295–336 (2018).
22. Grmela, M. & Öttinger, H. C. Dynamics and thermodynamics of complex fluids. I. *Phys. Rev. E* **56**, 6620–6632 (1997).
23. Onsager, L. & Machlup, S. Fluctuations and irreversible processes. *Phys. Rev.* **91**, 1505–1512 (1953).
24. Sittel, F., Jain, A. & Stock, G. Principal component analysis of molecular dynamics: On the use of cartesian vs. internal coordinates. *J. Chem. Phys.* **141**, 014111 (2014).
25. Barber, D. & Agakov, F. The IM algorithm: a variational approach to information maximization. *Adv. Neural Inf. Process. Syst.* **16**, 201–208 (2003).
26. Lipton, Z., Wang, Y.-X. & Smola, A. Detecting and correcting for label shift with black box predictors. *Proc. Mach. Learn. Res.* **80**, 3122–3130 (2018).
27. Lipari, G. & Szabo, A. Model-free approach to the interpretation of nuclear magnetic resonance relaxation in macromolecules. 1. *J. Am. Chem. Soc.* **104**, 4546–4559 (1982).
28. Tjandra, N., Feller, S. E., Pastor, R. W. & Bax, A. Rotational diffusion anisotropy of human ubiquitin from $^{15}$N NMR relaxation. *J. Am. Chem. Soc.* **117**, 12562–12566 (1995).
29. Hall, J. B. & Fushman, D. Characterization of the overall and local dynamics of a protein with intermediate rotational anisotropy: Differentiating between conformational exchange and anisotropic diffusion in the B3 domain of protein G. *J. Biomol. NMR* **27**, 261–275 (2003).
30. Buck, M., Boyd, J., Redfield, C., MacKenzie, D. A., Jeenes, D. J., Archer, D. B. & Dobson, C. M. Structural determinants of protein dynamics: Analysis of $^{15}$N NMR relaxation measurements for main-chain and side-chain nuclei of hen egg white lysozyme. *Biochemistry* **34**, 4041–4055 (1995).
31. Müller, C. W., Schlauderer, G. J., Reinstein, J. & Schulz, G. E. Adenylate kinase motions during catalysis: An energetic counterweight balancing substrate binding. *Structure* **4**, 147–156 (1996).
32. Henzler-Wildman, K. A. et al. Intrinsic motions along an enzymatic reaction trajectory. *Nature* **450**, 838–844 (2007).
33. Sharff, A. J., Rodseth, L. E., Spurlino, J. C. & Quiocho, F. A. Crystallographic evidence of a large ligand-induced hinge-twist motion between the two domains of the maltodextrin binding protein involved in active transport and chemotaxis. *Biochemistry* **31**, 10657–10663 (1992).
34. Boehr, D. D., McElheny, D., Dyson, H. J. & Wright, P. E. The dynamic energy landscape of dihydrofolate reductase catalysis. *Science* **313**, 1638–1642 (2006).
35. Eastman, P. et al. OpenMM 8: Modern materials and biophysical simulation. *PLoS Comput. Biol.* **20**, e1011807 (2024).
36. Epstein, D. M., Benkovic, S. J. & Wright, P. E. Dynamics of the dihydrofolate reductase–folate complex: Catalytic sites and regions known to undergo conformational change exhibit diverse dynamical features. *Biochemistry* **34**, 11037–11048 (1995).
37. Lindorff-Larsen, K. et al. Improved side-chain torsion potentials for the Amber ff99SB protein force field. *Proteins* **78**, 1950–1958 (2010).
38. Wang, L.-P., Martinez, T. J. & Pande, V. S. Building force fields: An automatic, systematic, and reproducible approach. *J. Phys. Chem. Lett.* **5**, 1885–1891 (2014).
39. Nguyen, H., Roe, D. R. & Simmerling, C. Improved Generalized Born solvent model parameters for protein simulations. *J. Chem. Theory Comput.* **9**, 2020–2034 (2013).
40. Honda, S., Yamasaki, K., Sawada, Y. & Morii, H. 10 residue folded peptide designed by segment statistics. *Structure* **12**, 1507–1518 (2004).
41. Neidigh, J. W., Fesinmeyer, R. M. & Andersen, N. H. Designing a 20-residue protein. *Nat. Struct. Biol.* **9**, 425–430 (2002).
42. Wlodawer, A., Walter, J., Huber, R. & Sjölin, L. Structure of bovine pancreatic trypsin inhibitor: Results of joint neutron and X-ray refinement of crystal form II. *J. Mol. Biol.* **180**, 301–329 (1984).
43. Buckle, A. M., Henrick, K. & Fersht, A. R. Crystal structural analysis of mutations in the hydrophobic core of barnase. *J. Mol. Biol.* **234**, 847–860 (1993).
44. Vijay-Kumar, S., Bugg, C. E. & Cook, W. J. Structure of ubiquitin refined at 1.8 Å resolution. *J. Mol. Biol.* **194**, 531–544 (1987).
45. Hoffmann, M. et al. Deeptime: a Python library for machine learning dynamical models from time series data. *Mach. Learn.: Sci. Technol.* **3**, 015009 (2022).
46. Ledoit, O. & Wolf, M. A well-conditioned estimator for large-dimensional covariance matrices. *J. Multivar. Anal.* **88**, 365–411 (2004).
47. Friedman, J., Hastie, T. & Tibshirani, R. Sparse inverse covariance estimation with the graphical lasso. *Biostatistics* **9**, 432–441 (2008).