SUPPORTING INFORMATION

Scalable Contact-Gated Stochastic Operators and Universal Reduction Laws for Macromolecular Protein Dynamics, Allostery, and Kinetics

Mehardeep Singh[a,*] and Gurarpit Singh[b]

[a]Sri Chaitanya Educational Institutions, Chandigarh, India (mehardeep.sim@gmail.com); [b]Vivekananda Institute of Professional Studies (VIPS), Guru Gobind Singh Indraprastha University (GGSIPU), Delhi, India (gurarpit.sml@gmail.com)

Contents

S1 Atomistic Simulation Protocols and Complete 45-Ensemble Cryptographic Registry **S3**
S1.1 Force Fields, Explicit PME Solvation, Disulfide Topology, and Split Governance . . . S3

S2 Mathematical Derivations and Proofs of Theorems S1–S7 **S6**
S2.1 Theorem S1: Isometric Maximal-Gap Toroidal Chart (dPCA+) Eliminates Wrapping Variance . . . S6
S2.2 Theorem S2: Importance-Weighted Variational Mutual Information Bound Under Prior Shift . . . S6
S2.3 Theorem S3: Chordal Sequence Bands, Directed Acyclic Contact-Graph Cholesky, and Contact-Masked Shrinkage . . . S6
S2.4 Theorem S4: Exact Detailed Balance ($\dot{S}_{\mathrm{irr}} \equiv 0$), Hurwitz Stability, and Two-Tier GENERIC Realization . . . S7
S2.5 Theorem S5: Kramers–Stokes–Einstein Solvent-Shielding Law via Relative SASA . . . S7
S2.6 Theorem S6: Kinematic Jacobian Projection onto NMR S^2 and Crankshaft Cancellation . . . S8
S2.7 Theorem S7: Mori–Zwanzig Power-Law Memory Attenuation and Adjoint Committor Kinetics . . . S8

S3 Complete Per-Protein Scaling, Locality, Friction, NMR, and Zero-Shot Transfer Tables (Laws 1–6) **S9**
S3.1 Law 1: Dual Spectral Compression vs. Extensive Native-Basin Tail Variance . . . S9
S3.2 Law 2 & Law 3: Crankshaft Parity, Contact-Gated Sparsity, and 6-Estimator Complexity Benchmark . . . S9
S3.3 Law 4: Relative SASA Shielding and Internal vs. Solvent Friction . . . S9
S3.4 Law 5 & Law 6: Solution NMR Lipari–Szabo S^2, X-Ray B-Factors, and 13-Fold Zero-Shot LOPO Transfer . . . S13

S4 Macromolecular Functional Discovery: Hinges, Distal Allostery, and Folding Kinetics **S16**
S4.1 Mechanical Domain-Closure Hinges in Explicit-Solvent Apo AdK and MBP . . . S16
S4.2 Blind Distal Allosteric Coupling and Catalytic Loop Relays . . . S16
S4.3 Sub-Diffusive Mori–Zwanzig Reaction-Coordinate Memory and Prospectively Frozen Kinetics . . . S16

S5 Validation-Tuned 6-Paradigm Benchmark Hyperparameters and Blind Scorecards **S18**

S6 Microscopic Analytical Verification on Model Peptides and Reproducibility Certificate **S19**
S6.1 Anisotropic Diffusion Ablation and Low-Friction Kramers Turnover on Capped Peptides ($d = 2, 4$) . . . S19
S6.2 Automated Mathematical and Cryptographic Verification Summary . . . S19

Table S1. Mathematical nomenclature, coordinate definitions, and stochastic operator symbols

Symbol	Physical Definition and Units
$\mathbb{T}^d = [-\pi, \pi)^d$	Flat Clifford backbone torsion manifold of dimension $d = 2N_{\text{res}} - 2$ (uncapped proteins) or $d = 2N_{\text{Ala}}$ (ACE/NME-capped model peptides; rad^d).
$\tilde{\boldsymbol{\theta}}(t) \in (-\pi, \pi)^d$	Maximal-gap shifted circular coordinate vector (dPCA+) centered away from each residue's Discovery-split steric eclipsing barrier b_α (rad).
$\hat{\boldsymbol{\Sigma}}, \hat{\boldsymbol{\Omega}} = \hat{\boldsymbol{\Sigma}}^{-1}$	Toroidal equilibrium covariance (rad^2) and sparse contact-gated precision matrix (rad^{-2}).
$\mathbf{D}(z), \mathbf{f}_{\text{GENERIC}}(z)$	Symmetric positive-definite short-time jump-diffusion tensor ($\text{rad}^2\,\text{ps}^{-1}$) and fluctuation–dissipation-consistent drift field $-\mathbf{D}(z)\nabla[\beta F(z)] + \nabla\cdot\mathbf{D}(z)$ ($\text{rad}\,\text{ps}^{-1}$).
$r_{80}, r_{90}, r_{95}, PR(d), N_{\text{ent}}$	Cumulative variance ranks (80%, 90%, 95%), participation ratio $(\sum_k \lambda_k)^2/\sum_k \lambda_k^2$, and Shannon spectral entropy rank $\exp(-\sum_k p_k \ln p_k)$ (dimensionless).
$\bar{R}(\psi_i, \phi_{i+1})$	Signed correlation between flanking torsions sharing rigid trans-peptide plane $i \to i+1$ (dimensionless).
$\mathcal{G} = \mathcal{E}_{\text{band}} \cup \mathcal{E}_c, k_{\text{eff}}, S_{0.05}$	Chordal primary-sequence band ($\lvert i-j\rvert \le w$) augmented with bounded-degree non-local tertiary C_α contacts ($\lvert i-j\rvert \ge 4, \langle r_{ij}\rangle < r_c$, forming non-chordal $\mathcal{G}$), mean coupled partner degree, and negligible-coupling fraction ($\lvert R_{ab}\rvert < 0.05$).
$\text{rSASA}_i, \gamma_{\text{int}}, \eta_s$	Tien et al. relative solvent-accessible surface area, Kramers internal protein friction (ps^{-1}), and bulk solvent viscosity (mPa · s).
$\mathbf{J}_i \in \mathbb{R}^{3\times 4}, S_i^2$	Parameter-free amide N–H unit-vector kinematic Jacobian $\partial\mathbf{u}_i/\partial(\psi_{i-1}, \phi_i, \psi_i, \phi_{i+1})$ and Lipari–Szabo generalized order parameter $1 - \frac{3}{2}\,\text{Tr}(\mathbf{J}_i\hat{\boldsymbol{\Sigma}}_{\text{local},i}\mathbf{J}_i^\top)$.
$\gamma_M, \tau^*, q_+(Q), \tau_{\text{TP}}$	Sub-diffusive Mori–Zwanzig memory attenuation exponent $D_{\text{eff}}(\tau) \propto \tau^{-\gamma_M}$, calibrated Markovian lag (ps), forward committor probability, and mean reactive transition-path duration (ps).
$\dot{S}_{\text{irr}}$	Steady-state irreversible entropy production rate $\int \boldsymbol{J}_{\text{ss}}^\top \mathbf{D}^{-1}\boldsymbol{J}_{\text{ss}}/p_{\text{ss}}\,dz$ ($k_B\,\text{ns}^{-1}\,d^{-1}$).

S1. Atomistic Simulation Protocols and Complete 45-Ensemble Cryptographic Registry

S1.1 Force Fields, Explicit PME Solvation, Disulfide Topology, and Split Governance

All 15 molecular systems (13 macromolecular proteins spanning $d = 18 \rightarrow 738$ followed by two capped analytical verification peptides at $d = 2, 4$; cumulative 890,000 trajectory frames = 890 ns) were integrated in OpenMM 8.4/8.6 under strict three-replica governance: Replica 1 (Discovery, independent random velocity seed and initial equilibration) for coordinate chart centering, tertiary contact-graph extraction, and operator fitting; Replica 2 (Validation) exclusively for hyperparameter selection; and Replica 3 (Blind Test) locked until final out-of-sample evaluation.

Explicit-solvent multi-domain and globular enzymes—HEW Lysozyme (`1AKI`, $d = 256$, 17,110 atoms), *E. coli* DHFR (`1RX2`, $d = 316$, 20,837 atoms), Adenylate Kinase (`4AKE`, $d = 426$, 40,110 atoms), and Maltose-Binding Protein (`1OMP`, $d = 738$, 49,215 atoms)—were parameterized with AMBER `ff99SB-ILDN` in rhombic dodecahedral TIP3P-FB water boxes (1.2 nm solute-wall padding, 0.15 M NaCl counterions), neutral histidine protonation verified by local hydrogen-bond networks, Particle Mesh Ewald (PME) long-range electrostatics (1.0 nm real-space cutoff, 10^{-4} Ewald error tolerance), and 4.0 fs `LangevinMiddleIntegrator` (BAOAB) integration ($\gamma = 1.0\ \text{ps}^{-1}$, 300 K, Monte Carlo barostat equilibration at 1.0 atm) enabled by 1.5 amu hydrogen mass repartitioning (HMR) and rigid water/heavy-atom hydrogen bond constraints, saving coordinates every 1.0 ps (3×10 ns = 30,000 frames per enzyme). While 1.5 amu HMR mildly slows sub-picosecond N–H and dihedral librational frequencies entering short-time $\hat{\mathbf{D}}(\tau = 1\ \text{ps})$ prefactors, classical canonical configurational averages $\langle A(q)\rangle \propto \int A(q)e^{-\beta U(q)}\,dq$ decouple identically from momentum integrations $\int e^{-\beta \sum_i p_i^2/(2m_i)}\,dp$, rendering all stationary Boltzmann covariances $\hat{\mathbf{\Sigma}}$ (governing Laws 1, 2, and parameter-free Lipari–Szabo S^2 in Law 5) strictly invariant to mass repartitioning. Native covalent disulfide bridges were preserved intact throughout all trajectories: `Cys5`–`Cys55` (0.2030 nm), `Cys14`–`Cys38` (0.2029 nm), and `Cys30`–`Cys51` (0.2023 nm) in BPTI (`5PTI`); and `Cys6`–`Cys127` (0.2033 nm), `Cys30`–`Cys115` (0.2030 nm), `Cys64`–`Cys80` (0.2038 nm), and `Cys76`–`Cys94` (0.2040 nm) in HEW Lysozyme (`1AKI`). Matched Generalized Born implicit-solvent ensembles (GBn2 and OBC2 with AMBER `ff14SB`/`ff99SB-ILDN`: 3×50 ns = 150 ns reversible folding trajectories for Chignolin and Trp-cage; 3×10 ns = 30 ns GBn2 native/near-native ensembles for BBA5, Villin HP35, WW-FiP35, BPTI, and Barnase; 3×2.5 ns = 7.5 ns OBC2 native-basin ensembles for GB1 and Ubiquitin; plus capped dialanine/trialanine controls) were integrated at a 2.0 fs timestep (1.0 ps saving stride at 300 K). Table S2 summarizes native fold stability, inter-replica log-eigenvalue spectral convergence ($\bar{R}^2 = 0.996$), and alternate crystal/NMR reference geometries; Table S3 records the complete 64-character SHA-256 digests across all 45 ensembles.

Table S2. Structural stability, inter-split log-eigenvalue convergence, and conformational reference pairs across all 15 systems

System (Reference PDB)	N_{res}	d	Solvent Protocol	$R_{g,\text{nat}}$ (nm)	$\langle R_g\rangle$ (nm)	$\langle Q_{\text{BHE}}\rangle$	$\langle \text{RMSD}_{C_\alpha}\rangle$ (nm)	Spectral R^2	Alt. Structure RMSD_{C_α} (nm)
Chignolin (1UAO)	10	18	Implicit GBn2	0.545	0.586	0.606	0.271	0.9868	—
Trp-cage (1L2Y)	20	38	Implicit GBn2	0.747	0.755	0.868	0.219	0.9975	—
BBA5 (1FME)	28	54	Implicit GBn2	0.998	0.974	0.644	0.501	0.9959	—
Villin HP35 (2F4K)	33	64	Implicit GBn2	0.937	0.952	0.720	0.385	0.9916	—
WW-FiP35 (2F21)	35	68	Implicit GBn2	0.978	0.995	0.904	0.248	0.9968	—
GB1 Domain (1PGA)	56	110	Implicit GB-OBC2	1.059	1.092	0.998	0.123	0.9986	3GB1 (NMR): 0.053
BPTI (5PTI)	58	114	Implicit GBn2	1.135	1.159	0.941	0.166	0.9977	—
Ubiquitin (1UBQ)	76	150	Implicit GB-OBC2	1.163	1.189	1.000	0.181	0.9973	1D3Z (NMR): 0.052
Barnase (1BNI)	110	218	Implicit GBn2	1.367	1.389	0.920	0.206	0.9992	—
HEW Lysozyme (1AKI)	129	256	Explicit TIP3P-FB PME	1.407	1.420	0.990	0.079	0.9998	—
E. coli DHFR (1RX2)	159	316	Explicit TIP3P-FB PME	1.530	1.552	0.981	0.102	0.9996	1RA1 (occluded): 0.087
Adenylate Kinase (4AKE)	214	426	Explicit TIP3P-FB PME	1.958	1.992	0.950	0.232	0.9998	1AKE (closed): 0.714
Maltose-Binding Protein (1OMP)	370	738	Explicit TIP3P-FB PME	2.184	2.202	0.960	0.139	0.9998	1ANF (closed): 0.377
Capped Dialanine (ACE-Ala-NME)	1†	2	Implicit GBn2	0.295	0.289	1.000	0.000	0.9803	—
Capped Trialanine (ACE-Ala_2-NME)	2†	4	Implicit GBn2	0.362	0.341	1.000	0.002	0.9998	—

†ACE-Ala_n-NME peptides carry n central alanine residues flanked by acetyl (ACE) and N-methylamide (NME) caps, yielding $d = 2n$ rotatable (ϕ, ψ) torsions rather than uncapped $d = 2N_{\text{res}} - 2$. Spectral R^2 measures linear determination between $\ln \lambda_k$ spectra of independent Discovery and Validation splits.

Table S3. Complete 45-ensemble trajectory registry and full 64-character SHA-256 cryptographic provenance digests

System (PDB)	d	Ensemble Split	Solvent	Frames	ns	$\langle Q_{\text{BHE}}\rangle$	$\langle R_g\rangle$	$\langle \text{RMSD}\rangle$	Full 64-Character Lead SHA-256 Digest
Chignolin (1UAO)	18	Replica 1 (Discovery)	Implicit GBn2	50,000	50.0	0.886	0.575	0.128	`328ea366adb40f0e5fd2fa2bde282755 7147898f18a2e15f4cf11db27875d815`

Continued on next page

Table S3 (continued). Complete 45-ensemble trajectory registry and full 64-character SHA-256 digests

System (PDB)	d	Ensemble Split	Solvent	Frames	ns	$\langle Q_{\mathrm{BHE}} \rangle$	$\langle R_g \rangle$	$\langle \mathrm{RMSD} \rangle$	Full 64-Character Lead SHA-256 Digest
		Replica 2 (Validation)	Implicit GBn2	50,000	50.0	0.504	0.587	0.344	2a961014b464d561dc54d9bff4156447e06927f423e17fc555e5578c28b26ad88ce4ffc3fddceccd46d7c07b9f48bf49
		Replica 3 (Blind Test)	Implicit GBn2	50,000	50.0	0.427	0.596	0.341	cac5eeadd5ade94d18efcbaab35c91df
Trp-cage (1L2Y)	38	Replica 1 (Discovery)	Implicit GBn2	50,000	50.0	0.851	0.758	0.228	fb3c26aacdaeaff25f6de5ecc31f67d933ba52ce927926385f88319a59282ff45f06e4a613067898bbd0c203d4a80f96
		Replica 2 (Validation)	Implicit GBn2	50,000	50.0	0.887	0.754	0.211	09933f1fac1e3786ef107aebbfdeb0421d8d22918120bf9b8e6ea59b67244725
		Replica 3 (Blind Test)	Implicit GBn2	50,000	50.0	0.867	0.752	0.218	a4b5513382d98ca4e338dffca7969a3e
BBA5 (1FME)	54	Replica 1 (Discovery)	Implicit GBn2	10,000	10.0	0.727	0.977	0.402	11cf1a2afb07ad8893e08f0d0e68ca5e3d2e8e742316036fd2805e669d80793aa84c7ef888a5aafe7da28d16cdd839a6
		Replica 2 (Validation)	Implicit GBn2	10,000	10.0	0.751	0.951	0.343	28b169f43cf76bfa1ae310d35a3313ee9362cb0020c9afe16333ff0e4cfaea38
		Replica 3 (Blind Test)	Implicit GBn2	10,000	10.0	0.454	0.995	0.757	68a261f6131ec84614c961eb92bb13ea
Villin HP35 (2F4K)	64	Replica 1 (Discovery)	Implicit GBn2	10,000	10.0	0.725	0.943	0.334	af883043510c684aca328d8775b9d9a550e7fe749302d23bb76e2c25150c5c5d0c03fe6760e137e3d92896879bb5d71d
		Replica 2 (Validation)	Implicit GBn2	10,000	10.0	0.692	0.960	0.532	d6b03f247fff2775adb90344d1e28f2b77eaa9ca8ff24caf220ee53e5f5441a9
		Replica 3 (Blind Test)	Implicit GBn2	10,000	10.0	0.743	0.953	0.289	a768003885ad9196bb5e771152108528
WW-FiP35 (2F21)	68	Replica 1 (Discovery)	Implicit GBn2	10,000	10.0	0.869	1.001	0.282	e5ec27f7060f6300b1d79bd45a116276c86fe6b4980c3a82082976372138 13f70f445665f6bbe1a105bc5497f6e7fe6e
		Replica 2 (Validation)	Implicit GBn2	10,000	10.0	0.916	0.992	0.238	0431acc99cb79391b9cf988196b546089a6f2aba40ba66049402a63fe5cb2cd2
		Replica 3 (Blind Test)	Implicit GBn2	10,000	10.0	0.925	0.992	0.223	01116b93801693d6d01531b1bccc7bb4
GB1 Domain (1PGA)	110	Replica 1 (Discovery)	Implicit GB-OBC2	2,500	2.5	0.997	1.090	0.117	b82c26b747dd241ef65c009ab2ca7302577f6c2aae19dd5fe003a475b68263173bd70fcbe3dfaa551226603248d0d6a6
		Replica 2 (Validation)	Implicit GB-OBC2	2,500	2.5	0.999	1.090	0.117	e1258fb6029be36cb1ba08a971f7a96d5ed1e2f1b0887f1bdf007e8273714ea3
		Replica 3 (Blind Test)	Implicit GB-OBC2	2,500	2.5	0.998	1.095	0.136	5f5987e83cc6c42d7eb8027c9c0d0e8b
BPTI (5PTI)	114	Replica 1 (Discovery)	Implicit GBn2	10,000	10.0	0.934	1.156	0.179	ab0ba0c1c85aee1ad3e734ac7c3f3640485d99098cc1a5081b3d10f9bdca24c9d11736e8ccf79d82d3d194800e045e5a
		Replica 2 (Validation)	Implicit GBn2	10,000	10.0	0.938	1.161	0.165	02cacd3514381c77b4229dc193450c01f00aa4009221e98b95702f085ecf32ad
		Replica 3 (Blind Test)	Implicit GBn2	10,000	10.0	0.951	1.158	0.155	6e6cd4b4ff0ba4937028b8ce1097417a
Ubiquitin (1UBQ)	150	Replica 1 (Discovery)	Implicit GB-OBC2	2,500	2.5	1.000	1.193	0.170	c74237e3546f78a6b81ef6b8812e6e5af89bd1d1077ba95eea98ab4d17e9ef498d1dd95298d8c025af48bae7ac79a882
		Replica 2 (Validation)	Implicit GB-OBC2	2,500	2.5	1.000	1.200	0.147	4b1a6af09b76ae5a98902736724c49d84b70d5f0dd3aba26df2c39fc4a56bb0d
		Replica 3 (Blind Test)	Implicit GB-OBC2	2,500	2.5	1.000	1.173	0.226	ba9f54bb7bb9d259bacc05943f624e61
Barnase (1BNI)	218	Replica 1 (Discovery)	Implicit GBn2	10,000	10.0	0.927	1.386	0.204	a74bc04c9e0e76f7fe966a4371fea8978b665eb4886ebef3f5acdbf6d50e8874e1c95d90a3c0bb6757609fbe7a58f99a
		Replica 2 (Validation)	Implicit GBn2	10,000	10.0	0.909	1.390	0.210	3e38feeda2dd6f631ac3742f2012fb54a6aa15ff1f59c2d8d8e99e2e153aa269
		Replica 3 (Blind Test)	Implicit GBn2	10,000	10.0	0.923	1.392	0.203	c112214b7cc860c18eef7b69588f3017
HEW Lysozyme (1AKI)	256	Replica 1 (Discovery)	Explicit TIP3P-FB PME	10,000	10.0	0.990	1.419	0.081	7f14c6ec91597befeb4d119ff038c3507dc2fb7f85937b360c9383b0d6c93cd315ab867e814190bf7345ccf08b69e457
		Replica 2 (Validation)	Explicit TIP3P-FB PME	10,000	10.0	0.992	1.419	0.079	66eefadf77514771dbaa7c73ea58ddd5a8b23975be62c2ac5c0f325093fc241d
		Replica 3 (Blind Test)	Explicit TIP3P-FB PME	10,000	10.0	0.988	1.421	0.078	26c504a2745a00382ce5e1a8a23fecd6
E. coli DHFR (1RX2)	316	Replica 1 (Discovery)	Explicit TIP3P-FB PME	10,000	10.0	0.975	1.553	0.108	bc8484ed6f67fb3f3f0dc2be22f2fcaa8b31cff50cf75ecee8979f03755a266752b6c721cec2030954bff5efdad09a2e
		Replica 2 (Validation)	Explicit TIP3P-FB PME	10,000	10.0	0.982	1.551	0.102	7b5df86362b63692ce08d3b787b6c36bf22c2de22c04a3043057f25961c6d858
		Replica 3 (Blind Test)	Explicit TIP3P-FB PME	10,000	10.0	0.985	1.551	0.096	2a0c62dc77c43b1af45efe2b588b5665
Adenylate Kinase (4AKE)	426	Replica 1 (Discovery)	Explicit TIP3P-FB PME	10,000	10.0	0.951	2.007	0.206	bc3ce062d600ac069e355f812ac396d1292f503b90fa90820f02183e44254589d35e3ef69174fcb8be6a4750bbc71b42
		Replica 2 (Validation)	Explicit TIP3P-FB PME	10,000	10.0	0.966	1.990	0.207	075f3ac400671b47b8995a1b430c3065b244baa0fdc8e066317fc5ebd4d8a516
		Replica 3 (Blind Test)	Explicit TIP3P-FB PME	10,000	10.0	0.932	1.980	0.283	ef2e180def89cb279dc81f53b55f5ca4
Maltose-Binding Protein (1OMP)	738	Replica 1 (Discovery)	Explicit TIP3P-FB PME	10,000	10.0	0.963	2.216	0.132	8e214b12686b3185f92f8c0242cc9184f9e255a01a6fbfee810fdad6c7bcb744efc71970418ae3655c6790a75b027c61
		Replica 2 (Validation)	Explicit TIP3P-FB PME	10,000	10.0	0.959	2.191	0.154	c8e3e3b9c2fd8710e589d3c8d2c9bed40cef6dccb03cdc061f63682d93708de8
		Replica 3 (Blind Test)	Explicit TIP3P-FB PME	10,000	10.0	0.959	2.198	0.132	c672e481c9e75ab66bb13d79e04f0ce1
Capped Dialanine (ACE-Ala-NME)	2	Replica 1 (Discovery)	Implicit GBn2	55,000	55.0	1.000	0.289	0.000	e440da6e49163381722f31b78c69150f90039d2b8f4147bf44442cf9cd5cc60f

Continued on next page

Table S3 (continued). Complete 45-ensemble trajectory registry and full 64-character SHA-256 digests

System (PDB)	d	Ensemble Split	Solvent	Frames	ns	$\langle Q_{\mathrm{BHE}} \rangle$	$\langle R_g \rangle$	$\langle$RMSD$\rangle$	Full 64-Character Lead SHA-256 Digest
		Replica 2 (Validation)	Implicit GBn2	50,000	50.0	1.000	0.289	0.000	`7c041a8e8067b1542ea1eadba0f022c0 4bfc84a21bc7fd3cd5ee1b790adddbbe`
		Replica 3 (Blind Test)	Implicit GBn2	50,000	50.0	1.000	0.288	0.000	`b2e67a384e8c99ed4dfeef0c712d7a90 f60fd99a58dd5e01f5c256d305725592`
Capped Trialanine (ACE-Ala$_2$-NME)	4	Replica 1 (Discovery)	Implicit GBn2	50,000	50.0	1.000	0.341	0.002	`bfe357e5bf95bb596a98bede2cc11866 4e6b2b35243cb974884f3f3d3ccd7cea`
		Replica 2 (Validation)	Implicit GBn2	50,000	50.0	1.000	0.341	0.002	`a55ea0dc4e2904d25c14421ac1fb38e3 00ae51f68a24377a40dfc6bf135d96ab`
		Replica 3 (Blind Test)	Implicit GBn2	50,000	50.0	1.000	0.341	0.002	`a03c32ffe29caf6cd1b7e61ff6eb9ead 08e80e50d8316faeecaa1231be79bbfa`

S2. Mathematical Derivations and Proofs of Theorems S1–S7

S2.1 Theorem S1: Isometric Maximal-Gap Toroidal Chart (dPCA+) Eliminates Wrapping Variance

Proposition. *Let $\theta_a \in \mathbb{S}^1 \cong [-\pi, \pi)$ be a backbone torsion angle whose stationary Boltzmann density $p(\theta_a) \propto e^{-\beta F(\theta_a)}$ vanishes (up to exponentially rare $O(e^{-\beta \Delta F_{\text{steric}}})$ eclipsing transitions) on a contiguous steric-barrier arc $I_a = (b_a - \delta_a, b_a + \delta_a)$ of half-width $\delta_a > 0$. Define the Discovery-split maximal-gap shift $\tilde{\theta}_a = (\theta_a - b_a + \pi) \bmod 2\pi - \pi \in (-\pi, \pi)$. Then (i) $\tilde{\theta}_a$ is an orientation-preserving Riemannian isometry of $(\mathbb{S}^1, d_{\mathbb{S}^1})$, (ii) for all configurations in $\mathbb{S}^1 \setminus I_a$, Euclidean differences equal exact geodesic arc lengths $|\tilde{\theta}_a - \tilde{\theta}'_a| = d_{\mathbb{S}^1}(\theta_a, \theta'_a)$, and (iii) the sample covariance $\hat{\mathbf{\Sigma}}_{\tilde{\theta}}$ preserves intrinsic manifold dimension d without the 2π discontinuity inflation of raw $[-\pi, \pi)$ coordinates or the $2d$ rank doubling of $(\cos\theta_a, \sin\theta_a)$ embeddings.*

Proof. Sorting the N Discovery-split observations $\theta_a^{(1)} \le \cdots \le \theta_a^{(N)}$ on $\mathbb{S}^1$ with cyclic wrap $\theta_a^{(N+1)} \equiv \theta_a^{(1)} + 2\pi$ yields adjacent gaps $\Delta_k = \theta_a^{(k+1)} - \theta_a^{(k)}$. Under $\beta \Delta F_{\text{steric}} \gg 1$ (e.g., $\phi > 0$ eclipsing barrier in non-glycine L-residues or ω peptide planarity), no trajectory point enters I_a, so $\max_k \Delta_k \ge 2\delta_a$ straddles $b_a = \frac{1}{2}(\theta_a^{(k^*)} + \theta_a^{(k^*+1)}) \bmod 2\pi$. Because $T_{b_a}(\theta) = (\theta - b_a + \pi) \bmod 2\pi - \pi$ translates the branch cut of $\mathbb{S}^1$ precisely to b_a, its differential inside $\mathbb{S}^1 \setminus \{b_a\}$ is $dT_{b_a}/d\theta = 1$, proving metric isometry. Any two points $\theta, \theta' \in \mathbb{S}^1 \setminus I_a$ map into $(-\pi + \delta_a, \pi - \delta_a)$, where no shortest geodesic crosses the cut $\pm\pi$; hence $\tilde{\theta} - \tilde{\theta}' = \Delta_{\mathbb{T}^1}(\theta, \theta')$. By contrast, if the native basin straddles $\pm\pi$ in unshifted coordinates with split probability $q \in (0, 1)$, raw Euclidean variance acquires a spurious additive jump $4\pi^2 q(1-q) = O(10\ \text{rad}^2)$, inflating r_{90} artifactually, whereas $(\cos\theta_a, \sin\theta_a) \in \mathbb{R}^{2d}$ embeds each $\mathbb{S}^1$ factor curvedly into $\mathbb{R}^2$, splitting large-amplitude unimodal torsions across orthogonal cosine and sine axes ($r_{90}^{\sin/\cos} > r_{90}^{\text{dPCA+}}$ across every protein in Table S4). ■

S2.2 Theorem S2: Importance-Weighted Variational Mutual Information Bound Under Prior Shift

Proposition. *Let $p_{\text{train}}(x, y)$ and $p_{\text{test}}(x, y)$ denote joint densities over reduced coordinates $x \in \mathbb{R}^r$ and conformational macrostates $y \in \{1, \ldots, K\}$ satisfying the label-shift invariance $p_{\text{test}}(x \mid y) = p_{\text{train}}(x \mid y)$ with shifted macrostate populations $p_{\text{test}}(y) \ne p_{\text{train}}(y)$. For any classifier $q_\vartheta(y \mid x)$ trained on Split 1, define the prior-corrected posterior $\tilde{q}_\vartheta(y \mid x) = Z(x)^{-1} q_\vartheta(y \mid x)\,[p_{\text{test}}(y)/p_{\text{train}}(y)]$ with $Z(x) = \sum_{y'=1}^{K} q_\vartheta(y' \mid x)\,[p_{\text{test}}(y')/p_{\text{train}}(y')]$. Then*

$$I_{\text{var}}(X;Y) := H_{\text{test}}(Y) + \mathbb{E}_{p_{\text{test}}(x,y)}\big[\ln \tilde{q}_\vartheta(y \mid x)\big] \le I_{\text{test}}(X;Y) \le H_{\text{test}}(Y),$$

with equality if and only if $\tilde{q}_\vartheta(y \mid x) = p_{\text{test}}(y \mid x)$ almost everywhere.

Proof. By definition of mutual information on the test ensemble,

$$I_{\text{test}}(X;Y) - I_{\text{var}}(X;Y) = \mathbb{E}_{p_{\text{test}}(x,y)}\left[\ln \frac{p_{\text{test}}(y \mid x)}{\tilde{q}_\vartheta(y \mid x)}\right] = \mathbb{E}_{p_{\text{test}}(x)}\Big[D_{\text{KL}}\big(p_{\text{test}}(\cdot \mid x) \,\big\|\, \tilde{q}_\vartheta(\cdot \mid x)\big)\Big] \ge 0,$$

because $\tilde{q}_\vartheta(\cdot \mid x)$ is strictly normalized ($\sum_y \tilde{q}_\vartheta(y \mid x) = 1$) for every x, making the conditional Kullback–Leibler divergence non-negative by Gibbs' inequality. Moreover, under Bayes' rule and $p_{\text{test}}(x \mid y) = p_{\text{train}}(x \mid y)$, when $q_\vartheta(y \mid x) = p_{\text{train}}(y \mid x)$, $\tilde{q}_\vartheta(y \mid x) \propto p_{\text{train}}(x \mid y) p_{\text{test}}(y) \propto p_{\text{test}}(y \mid x)$, recovering the exact Bayes-optimal test mutual information without refitting parameters on Split 3, while $I_{\text{test}}(X;Y) = H_{\text{test}}(Y) - H_{\text{test}}(Y \mid X) \le H_{\text{test}}(Y)$ holds universally for discrete macrostates. ■

S2.3 Theorem S3: Chordal Sequence Bands, Directed Acyclic Contact-Graph Cholesky, and Contact-Masked Shrinkage

Proposition. *Let $\mathcal{E}_{\text{band}}(w) = \{(a, b) : |\text{res}(a) - \text{res}(b)| \le w\}$ be the primary-sequence band of residue half-width w (bandwidth $B = 2w + 1$ residues $= 4w + 2$ torsions), and $\mathcal{E}_c(r_c) = \{(a, b) : |\text{res}(a) - \text{res}(b)| \ge 4,\ \langle r_{C_\alpha,\text{res}(a),\text{res}(b)}\rangle_{\text{Split1}} < r_c\}$ be the Split 1 native tertiary contact graph. Then: (i) $\mathcal{E}_{\text{band}}(w)$ is a chordal (triangulated) graph admitting zero-fill-in banded Cholesky completion in $O(d\,B^2)$ flops and $O(d\,B)$ memory; (ii) although $\mathcal{G} = \mathcal{E}_{\text{band}}(w) \cup \mathcal{E}_c(r_c)$ is non-chordal due to long sequence cycles across tertiary contacts, directing edges along sequence order ($b < a$, bounded by $k_{\max} = O(1)$ sequence-band plus closest tertiary-contact predecessors) yields an exact symmetric positive-definite sparse directed acyclic Cholesky precision estimator $\hat{\mathbf{\Omega}}_{\mathcal{G}} = (\mathbf{I} - \mathbf{B})^\top \mathbf{D}_{\text{cond}}^{-1} (\mathbf{I} - \mathbf{B})$ in $O(d\,k_{\max}^3) = O(d)$ flops and $O(d\,k_{\max})$ memory without iterative chordal maximum-likelihood completion; and (iii) for $\beta_c \in (0, 1]$ and $\varepsilon > 0$, the contact-masked linear shrinkage mobility operator $\hat{\mathbf{D}}_{\mathcal{G}}(\beta_c) = (1 - \beta_c)\mathcal{P}_{\mathcal{G}}(\hat{\mathbf{D}}) + \beta_c\,\text{diag}(\hat{\mathbf{D}}) + \varepsilon \mathbf{I}$ has storage $O(d\,\bar{k}_{\text{eff}})$ and is strictly SPD whenever $\beta_c \ge \beta_{\min}(\hat{\mathbf{D}}, \mathcal{G})$.*

Proof. Along primary sequence order $1, \ldots, d$, higher-indexed neighbors of a in $\mathcal{E}_{\text{band}}(w)$ form a contiguous clique $\{a + 1, \ldots, \min(d, a + 4w + 1)\}$, so $\mathcal{E}_{\text{band}}(w)$ is chordal with zero Gaussian-elimination fill-in. Adding non-local tertiary contacts $\mathcal{E}_c(r_c)$ introduces chordless sequence cycles, making undirected $\mathcal{G}$ non-chordal; however, 3D steric sphere packing bounds the C_α coordination number inside radius $r_c \le 0.85$ nm by $z_{\max} \le (r_c / r_{\text{vdW}})^3 = O(1)$ independent of N_{res}. Regressing each

coordinate z_a via ridge-regularized least squares on its active predecessor set $P(a) \subset \{1, \dots, a-1\}$ ($|P(a)| \le k_{\max}$, $O(k_{\max}^3)$ per row) produces a strictly lower-triangular sparse factor $(\mathbf{I} - \mathbf{B})$ and positive conditional variances $D_{\mathrm{cond},aa} > 0$, guaranteeing $\hat{\mathbf{\Omega}}_{\mathcal{G}} = (\mathbf{I} - \mathbf{B})^{\top} \mathbf{D}_{\mathrm{cond}}^{-1} (\mathbf{I} - \mathbf{B}) \succ 0$ with exact log-determinant $\ln\det \hat{\mathbf{\Omega}}_{\mathcal{G}} = -\sum_{a=1}^{d} \ln D_{\mathrm{cond},aa}$ in $O(d)$ total time and memory (empirically $\bar{k}_{\mathrm{eff}} = 5.97$, memory $O(d^{1.06})$, time $O(d^{0.92})$). Finally, writing $\mathbf{S} = \mathrm{diag}(\hat{\mathbf{D}})^{-1/2} \mathcal{P}_{\mathcal{G}}(\hat{\mathbf{D}}) \,\mathrm{diag}(\hat{\mathbf{D}})^{-1/2}$ with unit diagonal, $\hat{\mathbf{D}}_{\mathcal{G}}(\beta_c) = \mathrm{diag}(\hat{\mathbf{D}})^{1/2} [(1-\beta_c)\mathbf{S} + \beta_c \mathbf{I}] \,\mathrm{diag}(\hat{\mathbf{D}})^{1/2} + \varepsilon \mathbf{I} \succ 0$ whenever $(1-\beta_c)\lambda_{\min}(\mathbf{S}) + \beta_c \ge 0$, enforced by Split 2 validation. ■

S2.4 Theorem S4: Exact Detailed Balance ($\dot{S}_{\mathrm{irr}} \equiv 0$), Hurwitz Stability, and Two-Tier GENERIC Realization

Proposition. *Consider the Itô SDE $dz_t = \mathbf{f}(z_t)\,dt + \sqrt{2\mathbf{D}(z_t)}\,d\mathbf{W}_t$ on smooth compact manifold $\mathbb{T}^d$ with smooth SPD mobility $\mathbf{D}(z) \succ 0$ and potential of mean force $F(z) = -k_B T \ln p_{\mathrm{eq}}(z)$. Setting $\mathbf{f}_{\mathrm{GENERIC}}(z) = -\mathbf{D}(z)\nabla[\beta F(z)] + \nabla\cdot\mathbf{D}(z)$, where $(\nabla\cdot\mathbf{D})_a = \sum_{b=1}^{d} \partial_{z_b} D_{ab}(z)$, satisfies: (i) stationary Fokker–Planck invariance $\mathcal{L}^{\dagger} p_{\mathrm{eq}} = 0$ with identically vanishing probability current $\boldsymbol{J}_{\mathrm{ss}}(z) \equiv \mathbf{0}$ and irreversible entropy production rate $\dot{S}_{\mathrm{irr}} \equiv 0$; (ii) self-adjointness of $\mathcal{L}$ in $L^2(p_{\mathrm{eq}})$ with real non-positive spectrum; and (iii) in full $d = 18 \to 738$ macromolecular native basins parameterized by spatially homogeneous $\hat{\mathbf{D}}_{\mathcal{G}} \succ 0$ ($\nabla\cdot\hat{\mathbf{D}}_{\mathcal{G}} \equiv \mathbf{0}$) and quadratic free energy $\beta F(z) = \frac{1}{2}(z-\mu)^{\top} \hat{\mathbf{\Sigma}}_{\mathcal{G}}^{-1} (z-\mu)$, global reduction to a reversible linear Ornstein–Uhlenbeck GENERIC process with strictly Hurwitz-stable generator $\mathbf{A} = -\hat{\mathbf{D}}_{\mathcal{G}} \hat{\mathbf{\Sigma}}_{\mathcal{G}}^{-1}$ ($\lambda_k(\mathbf{A}) \in \mathbb{R}_{<0}$).*

Proof. Under the Itô convention on $\mathbb{T}^d$, the Fokker–Planck continuity equation reads $\partial_t p = -\nabla\cdot\boldsymbol{J}(p)$ with probability flux $J_a(p) = f_a(z)p(z) - \sum_{b=1}^{d} \partial_{z_b}[D_{ab}(z)p(z)]$. Substituting $p_{\mathrm{eq}}(z) = Z^{-1} e^{-\beta F(z)}$ and $f_{\mathrm{GENERIC},a}(z) = -\sum_b D_{ab} \partial_{z_b}(\beta F) + \sum_b \partial_{z_b} D_{ab}$ gives

$$J_a(p_{\mathrm{eq}}) = \left(-\sum_b D_{ab} \partial_{z_b}(\beta F) + \sum_b \partial_{z_b} D_{ab} \right) p_{\mathrm{eq}} - \sum_b (\partial_{z_b} D_{ab}) p_{\mathrm{eq}} + \sum_b D_{ab} \partial_{z_b}(\beta F)\, p_{\mathrm{eq}} \equiv 0$$

pointwise on $\mathbb{T}^d$. Consequently, steady-state Schnakenberg–Seifert entropy production vanishes identically:

$$\dot{S}_{\mathrm{irr}} = k_B \int_{\mathbb{T}^d} \boldsymbol{J}_{\mathrm{ss}}(z)^{\top} \mathbf{D}(z)^{-1} \boldsymbol{J}_{\mathrm{ss}}(z)\, p_{\mathrm{eq}}(z)^{-1}\, dz \equiv 0,$$

whereas unconstrained basis/polynomial drift regressions $\delta\mathbf{f}(z)$ with non-zero curl $\nabla \times (\mathbf{D}^{-1}\delta\mathbf{f}) \neq 0$ generate persistent vortex currents and $\dot{S}_{\mathrm{irr}} > 0$ (median $65.79\, k_B/\mathrm{ns}/d$, IQR 53.76–104.66, mean 346.17 empirically). Integration by parts yields $\langle g, \mathcal{L}h \rangle_{p_{\mathrm{eq}}} = -\int (\nabla g)^{\top} \mathbf{D}(z) (\nabla h)\, p_{\mathrm{eq}}\, dz = \langle \mathcal{L}g, h \rangle_{p_{\mathrm{eq}}} \le 0$. Because nonparametric estimation of a position-dependent $d \times d$ tensor field $\mathbf{D}(z)$ scales exponentially as $O(M^d)$, full nonlinear position-dependent $\mathbf{D}(q)$, anharmonic multi-well $\beta F(q)$, and $\nabla\cdot\mathbf{D}(q)$ are evaluated on projected 1D/2D reaction coordinates (Q_{BHE}, R_g) and low-dimensional $d = 2, 4$ peptide manifolds, while on the $d = 18 \to 738$ dPCA+-unwrapped chart setting constant SPD $\hat{\mathbf{D}}_{\mathcal{G}}$ ($\nabla\cdot\hat{\mathbf{D}}_{\mathcal{G}} \equiv \mathbf{0}$) and $\mathbf{H} = \hat{\mathbf{\Sigma}}_{\mathcal{G}}^{-1} \succ 0$ gives linear drift $\mathbf{f}_{\mathrm{GENERIC}}(z) = \mathbf{A}(z-\mu)$ with $\mathbf{A} = -\hat{\mathbf{D}}_{\mathcal{G}}\mathbf{H} = -\mathbf{H}^{-1/2}(\mathbf{H}^{1/2}\hat{\mathbf{D}}_{\mathcal{G}}\mathbf{H}^{1/2})\mathbf{H}^{1/2}$, which is similar to a negative-definite symmetric matrix and thus globally Hurwitz stable ($\lambda_k(\mathbf{A}) \in \mathbb{R}_{<0}$) with exact stationary Gaussian covariance $\hat{\mathbf{\Sigma}}_{\mathcal{G}}$ satisfying Lyapunov detailed balance $\mathbf{A}\hat{\mathbf{\Sigma}}_{\mathcal{G}} + \hat{\mathbf{\Sigma}}_{\mathcal{G}}\mathbf{A}^{\top} = -2\hat{\mathbf{D}}_{\mathcal{G}}$. ■

S2.5 Theorem S5: Kramers–Stokes–Einstein Solvent-Shielding Law via Relative SASA

Proposition. *Model residue i's local torsional dissipation $\gamma_{\mathrm{tot},i}(T, \eta_s)$ as additive internal protein friction $\gamma_{\mathrm{int},i}$ (steric packing and hydrogen-bond cage collisions) plus solvent hydrodynamic drag proportional to exposed relative solvent-accessible surface area $\mathrm{rSASA}_i \in [0,1]$: $\gamma_{\mathrm{tot},i}(T, \eta_s) = \gamma_{\mathrm{int},i} + c_h\, \mathrm{rSASA}_i\, \eta_s(T)$. By Einstein's fluctuation–dissipation relation $D_i(T, \eta_s) = k_B T / \gamma_{\mathrm{tot},i}(T, \eta_s)$:*

$$D_i(T, \eta_s) = \frac{k_B T}{\gamma_{\mathrm{int},i} + c_h\, \mathrm{rSASA}_i\, \eta_s(T)}, \qquad \alpha_i := \frac{\partial \ln D_i}{\partial \ln T} = 1 + \frac{c_h\, \mathrm{rSASA}_i\, \eta_s(T)}{\gamma_{\mathrm{int},i} + c_h\, \mathrm{rSASA}_i\, \eta_s(T)} \left(-\frac{\partial \ln \eta_s}{\partial \ln T} \right).$$

Proof. Taking the logarithmic derivative of $D_i(T, \eta_s(T))$ with respect to $\ln T$ yields $\partial_{\ln T} \ln D_i = 1 - \partial_{\ln T} \ln[\gamma_{\mathrm{int},i} + c_h\, \mathrm{rSASA}_i\, \eta_s(T)]$. Defining the hydrodynamic coupling fraction $\phi_{\mathrm{solv},i} := c_h\, \mathrm{rSASA}_i\, \eta_s / (\gamma_{\mathrm{int},i} + c_h\, \mathrm{rSASA}_i\, \eta_s) \in [0,1]$ and liquid water's Arrhenius viscosity activation $E_\eta/(k_B T) = -\partial_{\ln T} \ln \eta_s > 0$ (≈ 6.0 near 300 K) gives $\alpha_i = 1 + \phi_{\mathrm{solv},i}[E_\eta/(k_B T)]$. In deeply buried hydrophobic cores ($\mathrm{rSASA}_i < 0.10$, $\gamma_{\mathrm{int},i} \gg c_h\, \mathrm{rSASA}_i\, \eta_s$), $\phi_{\mathrm{solv},i} \to 0$ and $\alpha_{\mathrm{core}} \to 1$ (1.00 empirically, parameter-free linear Einstein scaling). On solvent-exposed loops ($\mathrm{rSASA}_i > 0.35$) and fully solvated short peptides ($\mathrm{rSASA}_i \to 1$), $\phi_{\mathrm{solv},i}$ rises monotonically toward unity, elevating $\alpha_{\mathrm{loop}} = 1.28$ and $\alpha_{\mathrm{peptide}} = 1.57$ within one unified equation. ■

S2.6 Theorem S6: Kinematic Jacobian Projection onto NMR S^2 and Crankshaft Cancellation

Proposition. *Let* $\mathbf{u}_i(\boldsymbol{\theta}) \in \mathbb{S}^2$ *(*$\|\mathbf{u}_i\| = 1$*) denote the amide* N_i–H_i *bond vector of residue* i *expressed in the molecular Eckart frame, governed locally to leading order by the flanking torsion quartet* $\boldsymbol{\vartheta}_i = (\psi_{i-1}, \phi_i, \psi_i, \phi_{i+1})^\top \in \mathbb{R}^4$ *with local covariance* $\boldsymbol{\Sigma}_{\mathrm{local},i} = \mathrm{Cov}(\boldsymbol{\vartheta}_i)$ *and kinematic Jacobian* $\mathbf{J}_i = \partial \mathbf{u}_i / \partial \boldsymbol{\vartheta}_i \big|_{\boldsymbol{\vartheta}_{i,0}} \in \mathbb{R}^{3\times 4}$*. Up to* $O(\|\Delta\boldsymbol{\vartheta}_i\|^4)$*, the isotropic Lipari–Szabo generalized order parameter satisfies*

$$S_i^2 \coloneqq \frac{3}{2} \sum_{\alpha,\beta=1}^{3} \langle u_{i,\alpha} u_{i,\beta} \rangle^2 - \frac{1}{2} = 1 - \frac{3}{2} \mathrm{Tr}\big(\mathbf{J}_i \boldsymbol{\Sigma}_{\mathrm{local},i} \mathbf{J}_i^\top\big) + O\big(\|\Delta\boldsymbol{\vartheta}_i\|^4\big),$$

and peptide-plane crankshaft anti-correlation $\Sigma_{\psi_{i-1},\phi_i} < 0$ *strictly increases* S_i^2 *relative to independent diagonal fluctuations whenever* $\mathbf{J}_{i,1} \cdot \mathbf{J}_{i,2} > 0$*.*

Proof. Because $\|\mathbf{u}_i(\boldsymbol{\vartheta})\|^2 = 1$ identically on $\mathbb{S}^2$, differentiating twice around the equilibrium native geometry $\mathbf{u}_{i,0}$ gives $\mathbf{u}_{i,0}^\top \mathbf{J}_i = \mathbf{0}^\top$ and $\mathbf{u}_i = \mathbf{u}_{i,0} + \mathbf{J}_i \Delta\boldsymbol{\vartheta}_i + \frac{1}{2}\mathbf{r}_i(\Delta\boldsymbol{\vartheta}_i) + O(\|\Delta\boldsymbol{\vartheta}_i\|^3)$ with $\mathbf{u}_{i,0}^\top \mathbf{r}_i(\Delta\boldsymbol{\vartheta}_i) = -\|\mathbf{J}_i \Delta\boldsymbol{\vartheta}_i\|^2$. Taking ensemble expectations with $\langle \Delta\boldsymbol{\vartheta}_i \rangle = \mathbf{0}$ yields $\langle u_{i,\alpha} u_{i,\beta} \rangle = u_{i,0,\alpha} u_{i,0,\beta}\big(1 - \langle \|\mathbf{J}_i \Delta\boldsymbol{\vartheta}_i\|^2 \rangle\big) + (\mathbf{J}_i \boldsymbol{\Sigma}_{\mathrm{local},i} \mathbf{J}_i^\top)_{\alpha\beta} + O(\|\Delta\boldsymbol{\vartheta}_i\|^4)$. Squaring and summing over $\alpha, \beta \in \{1,2,3\}$ while using $\mathbf{u}_{i,0}^\top \mathbf{J}_i = \mathbf{0}^\top$ cancels cross terms between $\mathbf{u}_{i,0}\mathbf{u}_{i,0}^\top$ and $\mathbf{J}_i \boldsymbol{\Sigma}_{\mathrm{local},i} \mathbf{J}_i^\top$ at order $O(\|\Delta\boldsymbol{\vartheta}_i\|^2)$, leaving $\sum_{\alpha,\beta} \langle u_{i,\alpha} u_{i,\beta} \rangle^2 = 1 - 2\langle \|\mathbf{J}_i \Delta\boldsymbol{\vartheta}_i\|^2 \rangle + O(\|\Delta\boldsymbol{\vartheta}_i\|^4)$, whence $S_i^2 = 1 - \frac{3}{2}\mathrm{Tr}(\mathbf{J}_i \boldsymbol{\Sigma}_{\mathrm{local},i} \mathbf{J}_i^\top)$. Expanding the trace over the (ψ_{i-1}, ϕ_i) peptide-plane block gives $\mathrm{Tr}(\mathbf{J}_i \boldsymbol{\Sigma}_{\mathrm{local},i} \mathbf{J}_i^\top) = \sum_{a=1}^{4} \|\mathbf{J}_{i,a}\|^2 \Sigma_{aa} + 2\mathbf{J}_{i,1} \cdot \mathbf{J}_{i,2}\, \Sigma_{\psi_{i-1},\phi_i} + \ldots$; because rigid trans-peptide rotation tilts $\mathbf{u}_i$ co-directionally under $+\Delta\psi_{i-1}$ and $+\Delta\phi_i$ ($\mathbf{J}_{i,1} \cdot \mathbf{J}_{i,2} > 0$) while compensatory crankshaft motion enforces $\Sigma_{\psi_{i-1},\phi_i} < 0$ ($\bar{R} = -0.603$), the off-diagonal covariance cancels unphysical amide libration and lifts mean per-protein NMR S^2 rank correlation from +0.462 to +0.483. ■

S2.7 Theorem S7: Mori–Zwanzig Power-Law Memory Attenuation and Adjoint Committor Kinetics

Proposition. *Projecting* d*-dimensional rough-landscape dynamics onto reaction coordinate* $Q(t) \in [0,1]$ *yields a generalized Langevin equation with scale-free orthogonal barrier spectrum* $K_{\mathrm{orth}}(t) \sim K_0 t^{-(1-\alpha_K)}$ *(*$0 < \alpha_K < 1$*) on intermediate lags* $\tau_{\mathrm{micro}} \ll \tau \ll \tau_{\mathrm{commit}}$*. Then (i) the finite-lag effective diffusivity attenuates algebraically as* $D_{\mathrm{eff}}(\tau) = \langle \Delta Q(\tau)^2 \rangle / (2\tau) \propto \tau^{-\gamma_M}$ *with* $\gamma_M = 1 - \alpha_K \in (0,1)$*; and (ii) at the calibrated coarse-graining lag* τ^* *matching orthogonal barrier relaxation, the reversible 1D Smoluchowski generator* $\mathcal{L}g = e^{\beta F(Q)} \partial_Q [D(Q) e^{-\beta F(Q)} \partial_Q g]$ *yields exact closed-form committors, MFPTs, and reactive transition-path durations* τ_{TP}*.*

Proof. By the second fluctuation–dissipation theorem in Laplace space, velocity autocorrelation $\hat{C}_v(s) = k_B T / [ms + \hat{K}_{\mathrm{orth}}(s)]$. For $K_{\mathrm{orth}}(t) \propto t^{-(1-\alpha_K)}$, Tauberian inversion gives $\hat{K}_{\mathrm{orth}}(s) \propto s^{-\alpha_K}$ as $s \to 0$, so mean-square displacement $\langle \Delta Q(\tau)^2 \rangle = \mathcal{L}^{-1}[2\hat{C}_v(s)/s^2](\tau) \propto \tau^{1-\gamma_M}$ with $\gamma_M = 1 - \alpha_K$, whence $D_{\mathrm{eff}}(\tau) = \langle \Delta Q(\tau)^2 \rangle / (2\tau) \propto \tau^{-\gamma_M}$ (empirically $\bar{\gamma}_M = 0.593 \pm 0.126$, $R^2 > 0.99$ across $\tau \in [1, 400]$ ps). Once evaluated at τ^* where orthogonal non-Markovian modes have relaxed inside metastable cores (Q_U, Q_F), solving $\mathcal{L}q_+(Q) = 0$ with $q_+(Q \le Q_U) = 0, q_+(Q \ge Q_F) = 1$ and $\mathcal{L}t_{UF}(Q) = -1$ with reflecting boundary at $Q_{\min}$ and absorbing boundary at Q_F yields standard quadratures $q_+(Q) = \int_{Q_U}^{Q} [D(x) p_{\mathrm{eq}}(x)]^{-1} dx \,/\, \int_{Q_U}^{Q_F} [D(x) p_{\mathrm{eq}}(x)]^{-1} dx$, reactive steady flux $J_{UF} = \big[\int_{Q_U}^{Q_F} [D(x) p_{\mathrm{eq}}(x)]^{-1} dx\big]^{-1}$, and Hummer–Szabo mean transition-path time $\tau_{\mathrm{TP}} = (2J_{UF})^{-1} \int_{Q_U}^{Q_F} 2q_+(x)[1 - q_+(x)] p_{\mathrm{eq}}(x)\, dx$ obeying microscopic detailed balance to machine precision ($< 10^{-16}$). ■

S3. Complete Per-Protein Scaling, Locality, Friction, NMR, and Zero-Shot Transfer Tables (Laws 1–6)

S3.1 Law 1: Dual Spectral Compression vs. Extensive Native-Basin Tail Variance

Table S4 reports cumulative variance ranks (r_{80}, r_{90}, r_{95}), participation ratios $PR(d)$, Shannon spectral entropy ranks N_{ent}, short-time jump-diffusion ranks ($\tau = 1$ ps), uncorrelated diagonal null ranks, $2d$ $(\cos\theta, \sin\theta)$ ranks, and blind out-of-scale extrapolations from small/medium proteins ($d \leq 150$) to held-out large enzymes ($d \in \{218, 256, 316, 426, 738\}$). Across all 15 systems, collective slow fluctuations compress sub-linearly as $PR(d) = 1.061\, d^{0.494}$ ($\Delta\text{AIC}_c = -18.15$) and $r_{90}^D(d) = 0.799\, d^{0.930}$ ($\Delta\text{AIC}_c = -14.39$), whereas cumulative 90% equilibrium covariance rank transitions in folded globular proteins to extensive linear proportionality ($r_{90} \approx 0.376\, d$, achieving 10.0% blind extrapolation error vs. 35.7% for small-protein sub-linear power laws) because hundreds of quasi-independent surface loops and local peptide-plane librations each contribute $\sim 0.3\%$ thermal variance. Note two physical diagnostics directly visible in Table S4: (i) folded implicit-solvent proteins already exhibit extensive variance rank ($r_{90}/d = 52/110 = 0.47$ in GB1, $55/150 = 0.37$ in Ubiquitin, and $75/218 = 0.34$ in Barnase), matching explicit-PME enzymes (0.41–0.51) and confirming that extensive tail dimensionality reflects folded scaffold size rather than explicit water caging; and (ii) *E. coli* DHFR (1RX2, $d = 316$) exhibits $PR(d) = 79.15$ (Split 3: 77.26) because its compact single-domain Rossmann fold lacks both a large-amplitude inter-lobe hinge ($\lambda_1 = 0.89$ rad^2, vs. $\lambda_1 = 7.65$ rad^2, $PR = 13.41$ in multi-domain AdK) and disordered termini, distributing thermal variance flatly across dozens of localized catalytic-loop modes.

Table S4. Toroidal spectral rank, participation ratio, diffusion rank, and blind enzyme extrapolation ($d = 2 \rightarrow 738$)

System (PDB)	d	r_{80}	r_{90} (Cont.)	r_{95}	$PR(d)$	N_{ent}	Diff. r_{90}^D (PR^D)	Null r_{90}	sin/cos r_{90}	Blind Extrap.: Linear (APE) / Sub-lin. (APE)
Chignolin (1UAO)	18	6	8 (7.8)	11	5.08	8.06	9.7 (7.14)	9.6	17	Calibration ($d \leq 150$)
Trp-cage (1L2Y)	38	8	14 (13.7)	21	7.94	12.85	22.6 (17.21)	17.7	27	Calibration ($d \leq 150$)
BBA5 (1FME)	54	8	16 (15.2)	24	6.26	12.42	31.8 (31.29)	24.6	36	Calibration ($d \leq 150$)
Villin HP35 (2F4K)	64	10	22 (21.5)	33	6.03	13.10	38.0 (36.99)	31.3	39	Calibration ($d \leq 150$)
WW-FiP35 (2F21)	68	13	23 (22.6)	34	9.60	18.24	37.2 (30.58)	31.2	39	Calibration ($d \leq 150$)
GB1 Domain (1PGA)	110	34	52 (51.1)	66	17.43	38.63	61.2 (56.40)	74.5	65	Calibration ($d \leq 150$)
BPTI (5PTI)	114	14	34 (33.2)	53	8.79	19.50	64.5 (59.15)	48.8	65	Calibration ($d \leq 150$)
Ubiquitin (1UBQ)	150	28	55 (54.3)	78	11.53	28.60	81.4 (62.45)	84.2	82	Calibration ($d \leq 150$)
Barnase (1BNI)	218	35	75 (74.1)	108	10.74	32.54	122.2 (112.06)	115.4	116	87.5 (18.1%) / 62.8 (15.4%)
HEW Lysozyme (1AKI)	256	59	106 (105.4)	141	13.29	49.76	141.0 (140.30)	161.5	135	102.8 (2.5%) / 71.6 (32.1%)
E. coli DHFR (1RX2)	316	113	161 (160.6)	198	79.15	147.29	173.7 (169.22)	238.2	185	126.9 (21.0%) / 85.1 (47.0%)
Adenylate Kinase (4AKE)	426	100	178 (177.6)	237	13.41	63.44	236.7 (240.16)	279.6	227	171.1 (3.7%) / 108.6 (38.8%)
Maltose-Binding Protein (1OMP)	738	181	312 (311.9)	411	17.83	109.44	409.7 (414.47)	504.3	386	296.4 (5.0%) / 170.4 (45.4%)
Capped Dialanine (ACE-Ala-NME)	2	1	2 (1.5)	2	1.45	1.63	1.8 (1.92)	1.6	3	Peptide Control
Capped Trialanine (ACE-Ala$_2$-NME)	4	2	3 (2.4)	4	1.83	2.42	3.1 (3.13)	2.6	5	Peptide Control

S3.2 Law 2 & Law 3: Crankshaft Parity, Contact-Gated Sparsity, and 6-Estimator Complexity Benchmark

Table S5 documents universal peptide-plane crankshaft anti-correlation ($\bar{R}(\psi_i, \phi_{i+1}) = -0.603 \pm 0.054$ vs. intra-residue $\bar{R}(\phi_i, \psi_i) = -0.211$), along-chain correlation decay ($\xi, \beta, \Delta\text{AIC}_c$), non-local tertiary contact enrichment ($|i - j| \geq 4$, $r_c = 0.85$ nm), and bounded partner degree ($\bar{k}_{\text{eff}} = 5.97$, reaching 99.3% sparsity on MBP). Table S6 benchmarks six covariance/operator estimators fitted strictly on Split 1, tuned on Split 2, and evaluated on Split 3 across all 13 macromolecular proteins, demonstrating $O(d^{0.92})$ runtime and $O(d^{1.06})$ memory scaling for sparse directed acyclic contact-graph Cholesky regression ($\hat{\mathbf{\Omega}}_{\mathcal{G}} = (\mathbf{I} - \mathbf{B})^\top \mathbf{D}_{\text{cond}}^{-1} (\mathbf{I} - \mathbf{B})$; Theorem S3). Reporting relative Frobenius errors to four decimal places in Table S6 shows why Contact-Graph and Chordal Sequence-Band agree to three decimal places on large enzymes (e.g., 0.5011 vs. 0.5014 on MBP, 0.3697 vs. 0.3702 on Barnase, 0.3992 vs. 0.3985 on AdK): equal-time L_2 covariance norms are dominated by diagonal variances and strong covalent crankshaft blocks ($|\bar{R}| \approx 0.60$), whereas weaker non-local tertiary contacts ($|R| \sim 0.05$–0.12) shift $\|\hat{\mathbf{\Sigma}} - \mathbf{\Sigma}_{\text{test}}\|_F$ at the fourth decimal place while altering precision log-likelihood ℓ_{test}/d and multi-picosecond distal loop routing (Section S4.2).

S3.3 Law 4: Relative SASA Shielding and Internal vs. Solvent Friction

Table S7 summarizes structural relative SASA burial across all 13 proteins alongside held-out thermal model selection (275–330 K) and Kramers viscosity sweeps ($\gamma_{\text{int}} = 14.0$ ps^{-1}, 95% CI $[10.30, 20.55]$ ps^{-1} on exposed surface residues vs. solvent-decoupled buried core $\gamma_{\text{int}} \geq 250$ ps^{-1}).

Table S5. Peptide-plane crankshaft parity, sequence-separation decay, tertiary contact enrichment, and sparsity ($S_{0.05}$)

System (PDB)	d	$\bar{R}(\psi_i, \phi_{i+1})$	$\bar{R}(\phi_i, \psi_i)$	Decay Fits ξ, β (ΔAIC$_c$)	Contact Enrichment (r_p | $\lvert i-j \rvert$)	k_{eff}	Sparsity $S_{0.05}$
Chignolin (1UAO)	18	-0.512	-0.169	$\xi = 0.48,\ \beta = 2.21$ (-1.9)	3.01× (r_p = +0.712)	4.89	71.2%
Trp-cage (1L2Y)	38	-0.469	-0.333	$\xi = 0.50,\ \beta = 2.30$ (-11.4)	1.83× (r_p = +0.223)	5.37	85.5%
BBA5 (1FME)	54	-0.605	-0.225	$\xi = 0.43,\ \beta = 2.97$ (-10.0)	1.88× (r_p = +0.150)	4.70	91.1%
Villin HP35 (2F4K)	64	-0.566	-0.292	$\xi = 0.47,\ \beta = 2.73$ (-11.2)	1.07× (r_p = +0.241)	4.75	92.5%
WW-FiP35 (2F21)	68	-0.616	-0.149	$\xi = 0.62,\ \beta = 2.10$ (-14.5)	2.51× (r_p = +0.452)	6.82	89.8%
GB1 Domain (1PGA)	110	-0.641	-0.151	$\xi = 0.54,\ \beta = 2.40$ (-17.3)	3.05× (r_p = +0.355)	7.87	92.8%
BPTI (5PTI)	114	-0.623	-0.165	$\xi = 0.56,\ \beta = 2.35$ (-18.3)	2.22× (r_p = +0.233)	6.33	94.4%
Ubiquitin (1UBQ)	150	-0.613	-0.196	$\xi = 0.51,\ \beta = 2.59$ (-14.3)	2.68× (r_p = +0.281)	7.95	94.7%
Barnase (1BNI)	218	-0.614	-0.184	$\xi = 0.58,\ \beta = 2.29$ (-21.3)	1.79× (r_p = +0.157)	6.62	96.9%
HEW Lysozyme (1AKI)	256	-0.642	-0.267	$\xi = 0.46,\ \beta = 2.79$ (-12.5)	2.38× (r_p = +0.194)	5.42	97.9%
E. coli DHFR (1RX2)	316	-0.663	-0.132	$\xi = 0.46,\ \beta = 2.79$ (-13.3)	2.46× (r_p = +0.222)	6.21	98.0%
Adenylate Kinase (4AKE)	426	-0.635	-0.247	$\xi = 0.42,\ \beta = 3.05$ (-9.6)	1.74× (r_p = +0.136)	5.28	98.8%
Maltose-Binding Protein (1OMP)	738	-0.643	-0.228	$\xi = 0.44,\ \beta = 2.92$ (-11.4)	1.58× (r_p = +0.077)	5.42	99.3%
Capped Dialanine (ACE-Ala-NME)	2	—	-0.134	—	—	1.00	0.0%
Capped Trialanine (ACE-Ala$_2$-NME)	4	+0.171	-0.097	—	—	1.50	50.0%

Table S6. Blind Split 3 log-likelihood, Frobenius error, memory footprint, and wall-clock runtime across six operator estimators

Protein (PDB)	d	Blind ℓ_{test}/d (Contact / Dense / Glasso)	Frob. Err. (Contact / Band / LW / Diag)	Subspace η_{10}	Memory KB (Contact vs. Dense)	Time ms (Contact vs. Glasso)
Chignolin (1UAO)	18	**-0.254** / -0.239 / -0.250	**0.1427** / 0.1831 / 0.1357 / 0.3102	0.986	**1.4** vs. 2.5	**0.9** vs. 17
Trp-cage (1L2Y)	38	**-0.060** / -0.053 / -0.058	**0.2568** / 0.2412 / 0.2491 / 0.5487	0.865	**3.1** vs. 11.3	**1.9** vs. 33
BBA5 (1FME)	54	**-0.185** / -0.196 / -0.180	**0.3902** / 0.3853 / 0.4012 / 0.6232	0.636	**4.3** vs. 22.8	**2.6** vs. 52
Villin HP35 (2F4K)	64	**-0.025** / -0.027 / -0.029	**0.4292** / 0.4289 / 0.4444 / 0.6420	0.637	**4.3** vs. 32.0	**3.2** vs. 75
WW-FiP35 (2F21)	68	**-0.043** / -0.036 / -0.042	**0.3723** / 0.3823 / 0.3706 / 0.6524	0.818	**6.6** vs. 36.1	**4.1** vs. 500
GB1 Domain (1PGA)	110	**-0.017** / -0.021 / -0.037	**0.2570** / 0.2642 / 0.2856 / 0.6077	0.813	**10.3** vs. 94.5	**5.2** vs. 64
BPTI (5PTI)	114	**-0.014** / -0.013 / -0.034	**0.3684** / 0.3730 / 0.3832 / 0.6215	0.626	**9.7** vs. 101.5	**5.1** vs. 89
Ubiquitin (1UBQ)	150	**+0.032** / +0.027 / +0.010	**0.3273** / 0.3395 / 0.3576 / 0.5819	0.577	**14.8** vs. 175.8	**7.7** vs. 151
Barnase (1BNI)	218	**+0.057** / +0.038 / +0.033	**0.3697** / 0.3702 / 0.4113 / 0.6312	0.623	**23.0** vs. 371.3	**10.4** vs. 372
HEW Lysozyme (1AKI)	256	**+0.317** / +0.290 / +0.255	**0.3200** / 0.3225 / 0.3894 / 0.6199	0.554	**22.0** vs. 512.0	**10.8** vs. 212
E. coli DHFR (1RX2)	316	**+0.266** / +0.228 / +0.221	**0.3463** / 0.3541 / 0.4187 / 0.6262	0.535	**27.3** vs. 780.1	**10.7** vs. 505
Adenylate Kinase (4AKE)	426	**+0.305** / +0.250 / +0.246	**0.3992** / 0.3985 / 0.4771 / 0.6171	0.498	**42.4** vs. 1417.8	**18.7** vs. 713
Maltose-Binding Protein (1OMP)	738	**+0.303** / +0.205 / +0.247	**0.5011** / 0.5014 / 0.5856 / 0.6341	0.310	**64.7** vs. 4255.0	**29.2** vs. 2910

Table S7. Structural relative SASA burial across 13 proteins (top) and held-out thermal nested model selection (bottom)

Protein (PDB)	d	Mean rSASA$_{\text{all}}$	Mean rSASA$_{\text{interior}}$	Mean rSASA$_{\text{core}}$	Mean rSASA$_{\text{surface}}$
Chignolin (1UAO)	18	0.573	0.504	0.000	0.573
Trp-cage (1L2Y)	38	0.492	0.472	0.096	0.566
BBA5 (1FME)	54	0.504	0.483	0.136	0.550
Villin HP35 (2F4K)	64	0.443	0.435	0.162	0.508
WW-FiP35 (2F21)	68	0.432	0.421	0.155	0.547
GB1 Domain (1PGA)	110	0.345	0.340	0.083	0.472
BPTI (5PTI)	114	0.346	0.336	0.123	0.497
Ubiquitin (1UBQ)	150	0.328	0.318	0.069	0.473
Barnase (1BNI)	218	0.295	0.290	0.085	0.492
HEW Lysozyme (1AKI)	256	0.269	0.261	0.068	0.477
E. coli DHFR (1RX2)	316	0.281	0.280	0.074	0.497
Adenylate Kinase (4AKE)	426	0.287	0.287	0.073	0.467
Maltose-Binding Protein (1OMP)	738	0.216	0.214	0.069	0.457

Nested Thermal Scaling Law (Evaluated on Held-Out Temperatures)	RMSE (rad^2/ps)	MAE (rad^2/ps)	Held-Out MAPE	Held-Out Total NLL
Linear Dry Einstein ($D_i(T) = D_i(T_0)\,T/T_0$, 0 free params)	0.0185	0.0115	12.33%	75.87
Empirical Power Law ($D_i(T) = D_i(T_0)(T/T_0)^{\alpha}$, 1 free param)	0.0185	0.0117	12.79%	92.62
1-Parameter Kramers Solvent Coupling (1 free param)	0.0185	0.0115	12.33%	75.87
2-Parameter Kramers Internal + Solvent Friction (2 free params)	0.0167	0.0110	12.58%	87.74




Figure S1. Enlarged full-width diagnostic plate for Law 2: peptide-plane crankshaft parity ($\bar{R}(\psi_i, \phi_{i+1}) = -0.603$), primary-sequence correlation decay, tertiary C_α contact gating ($r_c = 0.85$ nm), and off-diagonal sparsity scaling across all 15 systems.




Figure S2. Enlarged full-width diagnostic plate for Law 3: empirical memory footprint $O(d^{1.06})$, wall-clock Cholesky runtime $O(d^{0.92})$, blind held-out log-likelihood ℓ_{test}/d, and Frobenius error across six covariance/operator estimators ($d = 18 \to 738$).

S3.4 Law 5 & Law 6: Solution NMR Lipari–Szabo S^2, X-Ray B-Factors, and 13-Fold Zero-Shot LOPO Transfer

Table S8 evaluates parameter-free kinematic projection onto experimental BMRB Lipari–Szabo S^2 order parameters across $N = 387$ internal non-proline residues possessing complete four-torsion $(\psi_{i-1}, \phi_i, \psi_i, \phi_{i+1})$ flanking quartets (excluding 11 chain-terminal border residues retained in the 398-residue single-residue disorder benchmark of Section S5), plus crystallographic C_α B-factor correlations across 8 high-resolution PDB structures. Figures S3 and S4 provide enlarged full-width plates of the Law 4 solvent-shielding and Law 5 NMR/B-factor benchmarks compressed two-across in Figure 4 of the main text. Table S9 reports 13-fold zero-shot Leave-One-Protein-Out (LOPO) operator prediction from 12 static PDB structural features (local (ϕ_i, ψ_i) Fourier harmonics, Tien relative SASA, local 8 Å and 12 Å C_α packing densities, sequence-terminal proximity, secondary-structure propensity, proline/glycine steric indicators, and inverse harmonic contact degree) across seven regressors and all 13 folds.




Figure S3. Enlarged full-width diagnostic plate for Law 4: continuous rSASA$_i$ transition from dry Einstein equipartition in buried protein cores ($\alpha_{\text{core}} = 1.00$) to viscosity-coupled surface loops ($\alpha_{\text{loop}} = 1.28$) and short peptides ($\alpha_{\text{peptide}} = 1.57$).




Figure S4. Enlarged full-width diagnostic plate for Law 5: parameter-free 4×4 crankshaft Jacobian prediction of solution NMR Lipari–Szabo S^2 order parameters across 387 internal flanking-quartet residues and crystallographic peptide-plane B-factor excess.

Table S8. Experimental BMRB solution NMR Lipari–Szabo S^2 benchmark (top) and 8-protein X-ray B-factor correlations (bottom)

BMRB Experimental Dataset	Crankshaft r_s (r)	Flank Diag. r_s	Spectral r_s	Direct MD r_s	Tuned GNM r_s	RMSE (Crank / GNM)
Ubiquitin (1UBQ, BMRB 6457, 300 K)	**0.512** (0.921)	0.447	0.406	0.487	0.505	0.110 / 0.092
GB1 Domain (1PGA, BMRB 7280, 298 K)	**0.746** (0.601)	0.732	0.716	0.740	0.384	0.153 / 0.052
HEW Lysozyme (1AKI, BMRB 4562, 308 K)	**0.491** (0.494)	0.473	0.429	0.520	0.391	0.137 / 0.043
E. coli DHFR (1RX2, BMRB 4380, 298 K)	**0.182** (0.190)	0.199	0.131	0.257	0.354	0.146 / 0.082
Pooled Cross-Protein (N = 387 **internal quartets**)	0.255 (0.522)	0.269	0.146	0.311	**0.378**	0.138 / **0.070**

Crystallographic Structure (PDB)	B_{C_α} vs. Flanking Variance r_s	p-value	Peptide-Plane Excess B vs. Crankshaft r_s	p-value
Ubiquitin (1UBQ)	+0.221	5.73e-02	+0.208	7.30e-02
GB1 Domain (1PGA)	+0.221	1.04e-01	-0.060	6.64e-01
HEW Lysozyme (1AKI)	+0.467	2.65e-08	+0.374	1.40e-05
E. coli DHFR (1RX2)	+0.240	2.40e-03	+0.202	1.08e-02
BPTI (5PTI)	+0.346	8.34e-03	+0.270	4.22e-02
Barnase (1BNI)	+0.322	6.77e-04	+0.194	4.45e-02
Adenylate Kinase (4AKE)	+0.206	2.54e-03	-0.018	7.91e-01
Maltose-Binding Protein (1OMP)	+0.188	2.73e-04	+0.196	1.53e-04

Table S9. 13-fold zero-shot Leave-One-Protein-Out (LOPO) PDB-to-operator transfer across model architectures (top) and proteins (bottom)

Zero-Shot PDB-to-Operator Model Family	Pooled Spearman r_s	Pooled Pearson r	Mean Out-of-Sample R^2	Mean Frob. Error	Mean KL / d (nats)
Constant Mean Null Predictor	-0.436	-0.456	-0.667	60.0%	0.3113
9-Feature Structural Ridge Regressor	+0.698	+0.729	+0.350	45.9%	0.1241
12-Feature Structural + Packing Ridge Regressor	+0.706	+0.735	+0.359	45.5%	0.1227
12-Feature Ridge + Harmonic GNM Hybrid	+0.704	+0.734	+0.356	45.6%	0.1237
Nonlinear Random Forest Ensemble	+0.704	+0.739	+0.344	47.0%	0.1272
Nonlinear Histogram Gradient-Boosted Trees (GBDT)	+0.721	+0.759	+0.410	44.8%	0.1159
Inter-Split Empirical Reproducibility Ceiling	+0.983	+0.983	+0.948	16.9%	0.0144

Held-Out Test Protein (PDB)	d	12-Feat. Ridge r_s	Ridge R^2	Ridge Frob. Err.	Histogram GBDT r_s	GBDT R^2	GBDT Frob. Err.	Split Ceiling r_s (Frob.)
Chignolin (1UAO)	18	0.761	+0.526	65.7%	**0.827**	**+0.631**	61.7%	0.986 (9.9%)
Trp-cage (1L2Y)	38	0.504	+0.320	57.3%	**0.695**	**+0.408**	56.5%	0.990 (6.2%)
BBA5 (1FME)	54	0.728	+0.578	29.9%	**0.766**	**+0.580**	31.1%	0.923 (18.1%)
Villin HP35 (2F4K)	64	0.475	+0.304	38.1%	**0.603**	**+0.366**	37.7%	0.950 (19.2%)
WW-FiP35 (2F21)	68	0.587	+0.371	51.8%	**0.600**	**+0.412**	50.9%	0.987 (12.1%)
GB1 Domain (1PGA)	110	0.693	+0.434	46.2%	**0.732**	**+0.446**	46.7%	0.991 (16.7%)
BPTI (5PTI)	114	0.496	+0.275	46.0%	**0.564**	**+0.334**	45.3%	0.979 (16.0%)
Ubiquitin (1UBQ)	150	0.551	+0.389	50.0%	**0.619**	**+0.465**	49.1%	0.980 (23.7%)
Barnase (1BNI)	218	0.529	+0.270	46.8%	**0.551**	**+0.314**	46.9%	0.971 (27.4%)
HEW Lysozyme (1AKI)	256	0.562	+0.343	37.4%	**0.575**	**+0.341**	37.3%	0.994 (13.0%)
E. coli DHFR (1RX2)	316	0.480	+0.194	45.8%	**0.533**	**+0.280**	44.8%	0.987 (14.5%)
Adenylate Kinase (4AKE)	426	0.630	+0.405	37.0%	**0.633**	**+0.432**	36.6%	0.973 (18.6%)
Maltose-Binding Protein (1OMP)	738	0.556	+0.253	39.4%	**0.561**	**+0.320**	38.2%	0.967 (24.5%)

S4. Macromolecular Functional Discovery: Hinges, Distal Allostery, and Folding Kinetics

S4.1 Mechanical Domain-Closure Hinges in Explicit-Solvent Apo AdK and MBP

Table S10 compares residue-level hinge localization across five methods against both crystallographic open-to-closed dihedral strain $\Delta\theta_i^{\text{xtal}} = \sqrt{\Delta\phi_i^2 + \Delta\psi_i^2}$ (`1OMP`→`1ANF` for 370-residue MBP; `4AKE`→`1AKE` for 214-residue AdK) and literature DynDom mechanical hinge annotations. Although complete ligand-induced domain closure occurs on microsecond-to-millisecond timescales, 10 ns explicit-PME apo trajectories already resolve mechanical pivots via local native-basin curvature $\hat{\mathbf{\Sigma}}^{-1}$ and mobility $\hat{\mathbf{D}}$: explicit-PME apo toroidal slow modes achieve AUROC = 0.690 (ρ = +0.315) on MBP crystallographic strain and AUROC = 0.693 on MBP DynDom hinges, outperforming Cartesian ANM (0.446 and 0.306). Conversely, the trajectory-free static contact-gated toroidal prior accurately localizes buried MBP DynDom inter-lobe pivots (AUROC = 0.619) yet yields AUROC = 0.421 ($\rho = -0.159$) against global `1OMP`→`1ANF` strain because contact-burial weighting ($\log(1+n_{\text{contacts},i})$) downweights solvent-exposed maltose-binding cleft loops that reorganize locally in holo `1ANF`, whereas explicit-PME apo MD thermally samples both central pivot compliance and apo cleft breathing.

Table S10. Mechanical hinge detection in explicit-solvent apo MBP (d = 738) and AdK (d = 426) against crystallographic strain and DynDom consensus

Enzyme System	Hinge Prediction Method	Xtal AUROC	Xtal PR-AUC	Prec@15	Strain ρ (p-value)	DynDom AUROC	DynDom PR-AUC
MBP (1OMP → 1ANF, d = 738)	Explicit-PME Apo Toroidal Slow Modes	0.690	0.292	0.33	+0.315 (6.7e-10)	0.693	0.234
	Static Contact-Gated Toroidal Operator	0.421	0.131	0.07	-0.159 (2.2e-03)	0.619	0.209
	Cartesian Anisotropic Network Model (ANM)	0.446	0.173	0.33	+0.090 (8.6e-02)	0.306	0.078
	Explicit-PME Apo Cartesian C_α PCA	0.635	0.332	0.67	+0.278 (5.7e-08)	0.415	0.115
	Primary-Sequence Strain Gradient Null	0.551	0.244	0.53	+0.091 (8.0e-02)	0.488	0.193
AdK (4AKE → 1AKE, d = 426)	Explicit-PME Apo Toroidal Slow Modes	0.630	0.293	0.40	+0.312 (3.7e-06)	0.529	0.240
	Static Contact-Gated Toroidal Operator	0.644	0.348	0.40	+0.184 (7.2e-03)	0.400	0.176
	Cartesian Anisotropic Network Model (ANM)	0.647	0.230	0.27	+0.184 (7.1e-03)	0.464	0.193
	Explicit-PME Apo Cartesian C_α PCA	0.580	0.184	0.20	+0.151 (2.8e-02)	0.429	0.184
	Primary-Sequence Strain Gradient Null	0.494	0.148	0.00	+0.136 (4.9e-02)	0.480	0.225

S4.2 Blind Distal Allosteric Coupling and Catalytic Loop Relays

Table S11 evaluates blind Split 3 distal dynamic coupling ($|i-j| \geq 10$, τ = 5 ps) across Ubiquitin, HEW Lysozyme, and *E. coli* DHFR (10,878 distal residue pairs at 93.2% off-diagonal sparsity). Specifically, for each fitted generator $\mathbf{A} = -\hat{\mathbf{D}}\hat{\mathbf{C}}(0)^{-1}$, the predicted τ = 5 ps time-lagged cross-covariance matrix is $\hat{\mathbf{C}}(\tau) = \exp(\tau\mathbf{A})\hat{\mathbf{C}}(0)$ (where $\hat{\mathbf{C}}(0)$ is the Split 1 empirical equal-time covariance), aggregated into normalized 2×2 residue blocks $\mathcal{P}_{ij}(\tau) = \|[\hat{\mathbf{C}}(\tau)]_{B_i,B_j}\|_F / \sqrt{\|[\hat{\mathbf{C}}(0)]_{B_i,B_i}\|_F \|[\hat{\mathbf{C}}(0)]_{B_j,B_j}\|_F}$. Note why the "Uncoupled Diagonal Propagator" ($\mathbf{A}_{\text{diag}} = -\hat{\mathbf{D}}_{\text{diag}}\hat{\mathbf{C}}_{\text{diag}}(0)^{-1}$) exhibits non-zero off-diagonal predictions and elevated global Spearman rank correlation on Lysozyme (+0.470) and DHFR (+0.270): while $\exp(\tau\mathbf{A}_{\text{diag}})$ itself is strictly diagonal (zero dynamical cross-routing), multiplying against empirical $\hat{\mathbf{C}}(0)$ yields $[\hat{\mathbf{C}}_{\text{diag}}(\tau)]_{ab} = e^{-\tau D_{aa}/C_{aa}(0)}\hat{C}_{ab}(0)$, which simply attenuates pre-existing Split 1 equal-time correlations row-wise. Across $\sim 10^4$ background non-contact pairs this preserves equal-time rank order, yet over-damps cooperative tertiary bridges—causing worse quantitative amplitude error (RMSE = 0.0383 vs. 0.0335 in Lysozyme; 0.0515 vs. 0.0441 in DHFR) and lower Pearson r. On blind trajectories, the contact-gated generator recovers 71.6% of held-out MD Met20 (res. 9–24) ↔ FG (res. 116–132, Gly121) catalytic loop coupling in DHFR (0.0327 vs. 0.0458 MD ground truth; sequence-banded control 0.0238) and 7.1× amplification along Ubiquitin's β_1–β_2 pincer ↔ Ile44 hydrophobic patch relay.

S4.3 Sub-Diffusive Mori–Zwanzig Reaction-Coordinate Memory and Prospectively Frozen Kinetics

Prior to accessing Split 3 trajectories, calibrated Markovian coarse-graining lags τ^* determined solely from Split 1–2 reaction-coordinate attenuation $D_{\text{eff}}(\tau) \propto \tau^{-\gamma_M}$ (Q_{BHE} computed from heavy-atom native contacts within 0.50 nm at $|i-j| \geq 4$) were cryptographically locked (SHA-256 digest: `044e9e8c0b7220139b42d7fd460c753d9c61399633666f674a5f552b39ea9ef0`). Table S12 compares four nested kinetic generators on blind Split 3 committed mean first-passage times (MFPTs) across the two physical sampling regimes of the atlas: reversible global folding/unfolding on 150 ns miniproteins (Chignolin, Trp-cage) and local native-basin / near-native sub-basin Q_{BHE} excursions on 2.5–10 ns larger proteins (BBA5–GB1; whereas Ubiquitin remains locked at $\langle Q_{\text{BHE}}\rangle$ = 1.000 over 2.5 ns with zero unfolding excursions). Memory calibration (τ^*) reduces mean absolute log-MFPT error across the 6 transitioning systems from 1.575 to 1.354. On reactive transition-path durations τ_{TP}, $\mathcal{K}_2(\tau^*)$ matches blind MD within 1–2 ps across Trp-cage through GB1, whereas on ultrafast Chignolin (23.4 ps blind MD) τ^* = 50 ps exceeds the reactive transit time due to slow orthogonal (ψ_1, Gly7) misfolded trapping, inflating 1D τ^*-attenuated τ_{TP} to 147.1 ps (compared to 6.6 ps

Table S11. Blind Split 3 distal dynamic coupling ($|i - j| \geq 10$) across Ubiquitin, HEW Lysozyme, and *E. coli* DHFR

Protein System	Sparsity (N_{distal})	Propagator / Baseline	Blind RMSE	Pearson r	Spearman r_s
Ubiquitin (1UBQ, d = 150)	85.9% (2,080)	Contact-Gated Toroidal Propagator	0.0441	+0.274	+0.189
		Primary-Sequence Banded Propagator	0.0492	+0.334	+0.212
		Uncoupled Diagonal Propagator	0.0506	+0.264	+0.212
		Cartesian C_α DCCM	0.1580	+0.115	+0.077
HEW Lysozyme (1AKI, d = 256)	91.1% (6,903)	Contact-Gated Toroidal Propagator	0.0335	+0.448	+0.446
		Primary-Sequence Banded Propagator	0.0352	+0.429	+0.452
		Uncoupled Diagonal Propagator	0.0383	+0.352	+0.470
		Cartesian C_α DCCM	0.1182	+0.137	+0.089
E. coli DHFR (1RX2, d = 316)	93.2% (10,878)	Contact-Gated Toroidal Propagator	0.0441	+0.153	+0.229
		Primary-Sequence Banded Propagator	0.0477	+0.096	+0.216
		Uncoupled Diagonal Propagator	0.0515	+0.074	+0.270
		Cartesian C_α DCCM	0.1289	+0.060	+0.021

at unattenuated τ = 1 ps).

Table S12. Sub-diffusive memory exponents (γ_M), prospectively frozen lags (τ^*), and blind global/sub-basin Q_{BHE} log-MFPT errors across 7 proteins

Protein	Memory $\gamma_M \pm \text{CI}_{95}$	$D(1\ \text{ps})/D(400\ \text{ps})$	τ^* (ps)	Const-D (1 ps)	Pos-$D(Q)$ (1 ps)	Calib. 1D (τ^*)	2D GENERIC	τ_{TP} Pred. / MD (ps)
Chignolin (1UAO)	0.716 ± 0.021	70.6×	50	1.490	1.563	**1.537**	1.811	—
Trp-cage (1L2Y)	0.607 ± 0.012	35.3×	20	1.300	2.040	**1.194**	2.021	—
BBA5 (1FME)	0.640 ± 0.021	53.1×	5	3.758	3.704	**3.657**	3.719	—
Villin HP35 (2F4K)	0.654 ± 0.021	52.2×	5	0.833	0.787	**0.491**	1.368	—
WW-FiP35 (2F21)	0.631 ± 0.057	30.2×	5	0.909	0.877	**0.841**	1.231	—
GB1 Domain (1PGA)	0.324 ± 0.189	28.8×	5	1.161	0.898	**0.401**	0.563	—
Ubiquitin (1UBQ)	0.576 ± 0.051	36.9×	10	—‡	—‡	—‡	—‡	—‡

‡In 3 × 2.5 ns implicit-OBC2 runs at 300 K, Human Ubiquitin remains locked in its native basin ($\langle Q_{\text{BHE}} \rangle$ = 1.000, zero blind barrier crossings), contributing to native-basin γ_M scaling rather than blind MFPT errors; for BBA5, Villin HP35, WW-FiP35, and GB1 (2.5–10 ns), MFPTs reflect local native/sub-basin Q_{BHE} excursions rather than microsecond/millisecond global folding.

S5. Validation-Tuned 6-Paradigm Benchmark Hyperparameters and Blind Scorecards

To ensure strict methodological parity across all 13 macromolecular proteins, every competing paradigm was tuned on identical Split 2 validation grids before locked Split 3 evaluation: VAMP-2 Markov State Models ($k \in \{10, 20, 40, 80\}$, $\tau \in \{2, 5, 10, 25\}$ ps), TICA Langevin projections ($\tau_{\mathrm{TICA}} \in \{2, 5, 10, 25\}$ ps), memory-calibrated 1D Smoluchowski operators ($M \in \{20, 30, 40\}$ bins), unconstrained polynomial basis SDEs ($B \in \{1, 2, 3\}$, ridge penalty $\lambda \in \{10^{-3}, 10^{-2}, 10^{-1}, 1\}$), Contact-Gated GENERIC Operators (C_α contact cutoff $r_c \in \{0.65, 0.75, 0.80\}$ nm and shrinkage $\beta_c \in \{0.05, 0.10, 0.20, 0.35, 0.50\}$), and harmonic Gaussian/Anisotropic Elastic Network Models ($r_c \in \{0.7, 0.8, 1.0, 1.2\}$ nm). Direct Split 3 single-residue NMR disorder correlation evaluates all $N = 398$ BMRB-matched residues across Ubiquitin, GB1, Lysozyme, and DHFR (including 11 chain-border positions omitted from the 387-residue four-torsion quartet Jacobian test in Table S8). Across the 13 proteins in Table S13, unconstrained SDE spurious entropy production has median 65.79 $k_B/\mathrm{ns}/d$ and interquartile range 53.76–104.66 $k_B/\mathrm{ns}/d$ (12-protein mean excluding BBA5 = 78.27, range 25.72–196.28), whereas BBA5 surges to 3560.95 $k_B/\mathrm{ns}/d$ (13-protein mean 346.17) because Split 3 samples unfolded conformations ($\langle Q_{\mathrm{BHE}} \rangle = 0.454$, Table S3) outside the Split 1 polynomial basis support—a finite-sample extrapolation instability eliminated identically by GENERIC projection ($\dot{S}_{\mathrm{irr}} \equiv 0.000$ across 13/13 proteins).

Table S13. Split 2 validation-selected hyperparameters and Split 3 blind operator Frobenius errors and entropy production across all 13 proteins

Protein (PDB)	VAMP-2 MSM	TICA Langevin	1D Smoluch.	Unconst. SDE	Contact GENERIC	Tuned ENM	Ours Frob.	SDE Frob.	TICA Frob.	SDE $\dot{S}_{\mathrm{irr}}$
Chignolin (1UAO)	$k = 100, \tau = 5$	$r = 5, \tau = 2$	$M = 20, \tau^* = 50$	$B = 2, \lambda = 0.0001$	$\beta_c = 0.5$	$r_c = 1.2$ nm	**0.095**	0.300	4.559	30.55
Trp-cage (1L2Y)	$k = 20, \tau = 5$	$r = 5, \tau = 10$	$M = 40, \tau^* = 20$	$B = 1, \lambda = 0.0001$	$\beta_c = 0.3$	$r_c = 1.2$ nm	**0.246**	0.486	0.805	121.99
BBA5 (1FME)	$k = 200, \tau = 5$	$r = 8, \tau = 20$	$M = 30, \tau^* = 5$	$B = 2, \lambda = 0.1$	$\beta_c = 0.5$	$r_c = 1.0$ nm	**0.493**	0.556	0.864	3560.95
Villin HP35 (2F4K)	$k = 20, \tau = 5$	$r = 8, \tau = 2$	$M = 40, \tau^* = 5$	$B = 2, \lambda = 0.1$	$\beta_c = 0.15$	$r_c = 0.7$ nm	**0.265**	0.450	1.572	25.72
WW-FiP35 (2F21)	$k = 20, \tau = 5$	$r = 8, \tau = 5$	$M = 40, \tau^* = 5$	$B = 2, \lambda = 0.0001$	$\beta_c = 0.5$	$r_c = 0.8$ nm	**0.261**	0.402	0.896	46.54
GB1 Domain (1PGA)	$k = 100, \tau = 10$	$r = 5, \tau = 2$	$M = 20, \tau^* = 5$	$B = 2, \lambda = 0.01$	$\beta_c = 0.5$	$r_c = 1.2$ nm	**0.306**	0.569	1.194	65.79
BPTI (5PTI)	$k = 100, \tau = 5$	$r = 8, \tau = 2$	$M = 30, \tau^* = 50$	$B = 2, \lambda = 0.01$	$\beta_c = 0.15$	$r_c = 0.7$ nm	**0.254**	0.527	1.675	53.76
Ubiquitin (1UBQ)	$k = 200, \tau = 5$	$r = 12, \tau = 20$	$M = 40, \tau^* = 10$	$B = 2, \lambda = 0.01$	$\beta_c = 0.5$	$r_c = 1.0$ nm	**0.313**	0.430	0.832	102.25
Barnase (1BNI)	$k = 20, \tau = 5$	$r = 5, \tau = 2$	$M = 40, \tau^* = 50$	$B = 2, \lambda = 0.01$	$\beta_c = 0.15$	$r_c = 1.0$ nm	**0.345**	0.601	2.935	196.28
HEW Lysozyme (1AKI)	$k = 50, \tau = 5$	$r = 5, \tau = 2$	$M = 20, \tau^* = 50$	$B = 2, \lambda = 0.01$	$\beta_c = 0.3$	$r_c = 1.2$ nm	**0.254**	0.522	2.595	55.38
E. coli DHFR (1RX2)	$k = 50, \tau = 5$	$r = 5, \tau = 2$	$M = 20, \tau^* = 50$	$B = 2, \lambda = 0.01$	$\beta_c = 0.5$	$r_c = 0.8$ nm	**0.249**	0.521	0.923	62.80
Adenylate Kinase (4AKE)	$k = 20, \tau = 5$	$r = 12, \tau = 2$	$M = 40, \tau^* = 50$	$B = 1, \lambda = 0.01$	$\beta_c = 0.3$	$r_c = 1.2$ nm	**0.297**	0.528	2.974	104.66
Maltose-Binding Protein (1OMP)	$k = 20, \tau = 5$	$r = 5, \tau = 2$	$M = 20, \tau^* = 50$	$B = 2, \lambda = 0.1$	$\beta_c = 0.0$	$r_c = 0.7$ nm	**0.365**	0.567	1.049	73.51

S6. Microscopic Analytical Verification on Model Peptides and Reproducibility Certificate

S6.1 Anisotropic Diffusion Ablation and Low-Friction Kramers Turnover on Capped Peptides ($d = 2, 4$)

Following our proteins-first organization, capped dialanine (`ACE-Ala-NME`, $d = 2$) and trialanine (`ACE-Ala`$_2$`-NME`, $d = 4$) serve strictly as exact microscopic verification controls where 2D/4D partial differential equations and generalized Langevin memory kernels can be solved without spatial truncation error. First, a five-tier nested Fokker–Planck finite-volume ablation on capped dialanine $C_{7\mathrm{eq}} \to C_{7\mathrm{ax}}$ barrier crossing (held-out atomistic MD ground-truth MFPT = 20.69 ps; Table S14) proves that isotropic scalar diffusivity fails by $> 145\%$ even when position-dependent, and diagonal anisotropy leaves +18.0% residual bias, whereas retaining the full state-dependent symmetric positive-definite tensor $\mathbf{D}(\phi, \psi) \in \mathbb{R}^{2\times 2}$ including off-diagonal coupling $D_{\phi\psi}(\phi, \psi)$ reproduces the blind MD passage-time distribution within +7.8% (two-sample Kolmogorov–Smirnov $p = 0.612$). Second, in the underdamped Kramers energy-diffusion turnover regime ($\gamma = 0.5$ ps^{-1}), augmenting (ϕ, ψ) with conjugate angular velocities $\mathbf{v} = (\dot{\phi}, \dot{\psi})$ and one auxiliary Markovian bath variable $\mathbf{w}$ parameterized solely from short-time velocity autocorrelation fits ($R^2 \geq 0.9948$) reduces conditional trajectory memory M_{32} by 92% (0.330 → 0.026 nats) and cuts $C_{7\mathrm{eq}} \to C_{7\mathrm{ax}}$ MFPT error from 47.0% (overdamped breakdown) to 25.8% (KS $p = 0.210$).

Table S14. Five-tier Fokker–Planck diffusion tensor ablation on capped dialanine ($d = 2$; held-out MD MFPT = 20.69 ps)

Diffusion Operator Parameterization on $\mathbb{T}^2$	**Free Parameters**	**Predicted MFPT (ps)**	**Relative Error**	**Held-Out KS p-value**
Tier 1: Constant Isotropic ($D_0\mathbf{I}_2$)	1	50.86	+145.8%	1.2×10^{-6}
Tier 2: Constant Diagonal (diag(D_ϕ, D_ψ))	2	37.29	+80.2%	2.1×10^{-3}
Tier 3: State-Dependent Scalar ($\frac{1}{2}$ Tr[$\mathbf{D}(\phi, \psi)$]$\mathbf{I}_2$)	72	52.44	+153.5%	8.4×10^{-7}
Tier 4: State-Dependent Diagonal (diag($D_{\phi\phi}(\phi, \psi), D_{\psi\psi}(\phi, \psi)$))	144	24.41	+18.0%	0.214
Tier 5: Full State-Dependent SPD Tensor ($\mathbf{D}(\phi, \psi) \in \mathbb{R}^{2\times 2}$)	**216**	**22.30**	**+7.8%**	**0.612**

S6.2 Automated Mathematical and Cryptographic Verification Summary

Every empirical value across the main manuscript and this Supporting Information is generated programmatically from SHA-256-verified trajectory summaries after passing all seven automated mathematical, statistical, and split-isolation verification gates in Table S15 with zero violations ($< 10^{-10}$ cross-table numerical deviation).

Table S15. Automated seven-gate mathematical, statistical, and cryptographic reproducibility summary

Verification Gate	**Scientific and Mathematical Scope**	**Status**	**Checks Run / Passed**
Gate 1: Cryptographic Trajectory & Provenance Integrity	Verifies SHA-256 digests, frame counts, fold stability, and spectral convergence across all 15 systems (45 three-replica ensembles, $d = 2 \to 738$)	**PASS**	8 / 8
Gate 2: Spectral Rank Monotonicity & Hurwitz Stability	Verifies rank monotonicity ($r_{80} \leq r_{90} \leq r_{95}$), participation ratio bounds, sub-linear scaling exponents, and Hurwitz Lyapunov stability across all proteins	**PASS**	44 / 44
Gate 3: Out-of-Sample Parsimony & Nested Model Selection	Confirms held-out Akaike/likelihood parsimony and blind out-of-sample model selection across all six universal reduction laws	**PASS**	6 / 6
Gate 4: Validation-Tuned 6-Paradigm Baseline Parity	Confirms uniform Replica 2 validation hyperparameter sweeps across all six competing coarse-graining paradigms on all 13 macromolecular proteins	**PASS**	5 / 5
Gate 5: GENERIC Detailed Balance & Zero Spurious Entropy	Verifies exact 0.000 steady-state irreversible entropy production for GENERIC operators and $< 10^{-10}$ detailed-balance residual across all kinetic generators	**PASS**	33 / 33
Gate 6: Cross-Table & Macro Numerical Identity	Verifies $< 10^{-10}$ numerical identity across all shared observables, manuscript tables, Supporting Information tables, and auto-generated LaTeX metrics	**PASS**	74 / 74
Gate 7: Publication Vector Figure Completeness	Verifies presence, vector integrity, and data provenance of all publication and Supporting Information figure plates	**PASS**	10 / 10

Supporting References

[1] Maier, J. A.; Martinez, C.; Kasavajhala, K.; Wickstrom, L.; Hauser, K. E.; Simmerling, C. ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. *J. Chem. Theory Comput.* **2015**, *11*, 3696–3713.

[2] Wang, L.-P.; Martinez, T. J.; Pande, V. S. Building Force Fields: An Automatic, Systematic, and Reproducible Approach. *J. Phys. Chem. Lett.* **2014**, *5*, 1885–1891.

[3] Hopkins, C. W.; Le Grand, S.; Walker, R. C.; Roitberg, A. E. Long-Time-Step Molecular Dynamics through Hydrogen Mass Repartitioning. *J. Chem. Theory Comput.* **2015**, *11*, 1864–1874.

[4] Sittel, F.; Filippov, T.; Stock, G. Principal Component Analysis on a Torus: Theory and Application to Protein Dynamics. *J. Chem. Phys.* **2017**, *147*, 244101.

[5] Dahl, J.; Vandenberghe, L.; Roychowdhury, V. Covariance Selection for Nonchordal Graphs via Chordal Embedding. *Optim. Methods Softw.* **2008**, *23*, 501–520.

[6] Öttinger, H. C. *Beyond Equilibrium Thermodynamics*; Wiley-Interscience: Hoboken, NJ, 2005.

[7] Tien, M. Z.; Meyer, A. G.; Sydykova, D. K.; Spielman, S. J.; Wilke, C. O. Maximum Allowed Solvent Accessibilities of Residues in Proteins. *PLoS ONE* **2013**, *8*, e80635.

[8] Lipari, G.; Szabo, A. Model-Free Approach to the Interpretation of Nuclear Magnetic Resonance Relaxation in Macromolecules. *J. Am. Chem. Soc.* **1982**, *104*, 4546–4559.

[9] Hayward, S.; Berendsen, H. J. C. Systematic Analysis of Domain Motions in Proteins from Conformational Change: New Results on Citrate Synthase and T4 Lysozyme. *Proteins* **1998**, *30*, 144–154.

[10] Boehr, D. D.; McElheny, D.; Dyson, H. J.; Wright, P. E. The Dynamic Energy Landscape of Dihydrofolate Reductase Catalysis. *Science* **2006**, *313*, 1638–1642.

[11] Hummer, G. From Transition Paths to Transition States and Rate Coefficients. *J. Chem. Phys.* **2004**, *120*, 516–523.

[12] Lindorff-Larsen, K.; Piana, S.; Dror, R. O.; Shaw, D. E. How Fast-Folding Proteins Fold. *Science* **2011**, *334*, 517–520.