

Chapter 3:

Estimation of the Two-Variable Regression Model (OLS)

Course Supervisor: Dr. Salem

Teaching Assistant: Zoleikani

October 11, 2025

Conceptual Summary (Key Points)

1. Objective of the Ordinary Least Squares (OLS) Method

To find estimates of β_1 and β_2 that *minimize the sum of squared residuals*:

Minimize $\sum e_i^2$, where:

$$e_i = Y_i - \hat{Y}_i$$

2. Normal Equations

Derived by taking derivatives of the SSE with respect to β_1 and β_2 :

$$\sum e_i = 0, \quad \sum X_i e_i = 0$$

3. OLS Estimators

$$\hat{\beta}_2 = \frac{\sum (X_i - \bar{X})(Y_i - \bar{Y})}{\sum (X_i - \bar{X})^2}, \quad \hat{\beta}_1 = \bar{Y} - \hat{\beta}_2 \bar{X}$$

4. Properties of the OLS Regression Line

- Passes through the means $(\bar{X}, \bar{Y})$.
- The sum of residuals equals zero.
- Residuals are uncorrelated with the explanatory variable X .

5. Classical Linear Regression Model (CLRM) Assumptions

For OLS estimators to be valid and BLUE, these must hold:

- $E(U_i | X_i) = 0$ (zero mean of errors)
- $Cov(U_i, U_j) = 0$ (no autocorrelation)
- $Var(U_i) = \sigma^2$ (homoskedasticity)
- $Cov(U_i, X_i) = 0$ (exogeneity)
- Model is correctly specified (no omitted variables or wrong functional form)

6. Properties of OLS Estimators

- **Linearity:** estimates are linear functions of observed data.
- **Unbiasedness:** $E(\hat{\beta}_i) = \beta_i$.
- **Minimum variance among all linear unbiased estimators** (Gauss–Markov theorem $\rightarrow$ OLS is BLUE).
- **Consistency and asymptotic efficiency** for large samples.

7. Estimating the Error Variance σ^2

Since true errors are unobservable, we use sample residuals:

$$\hat{\sigma}^2 = \frac{\sum e_i^2}{n - 2}$$

8. Coefficient of Determination R^2

Measures the proportion of the variation in Y explained by X :

$$R^2 = 1 - \frac{\sum e_i^2}{\sum (Y_i - \bar{Y})^2}$$

Questions

1. Why is the sum of residuals zero in OLS?
2. What's the economic interpretation of β_2 ?
3. Why is OLS considered the *best linear unbiased estimator*?
4. What happens if the homoskedasticity assumption is violated?
5. What's the difference between *unbiasedness* and *consistency*?
6. Why must the regression line pass through the sample means $(\bar{X}, \bar{Y})$?
7. What happens if $Cov(U_i, X_i) \neq 0$?

Problem Set

Problem 3.1 - Computing OLS Estimates

The data below represent weekly income (Y, in \$100) and years of schooling (X):

X	Y
10	30
12	38
14	45
16	50
18	59

Tasks:

1. Compute $\bar{X}$ and $\bar{Y}$.
2. Calculate $\hat{\beta}_2 = \frac{\sum(X_i - \bar{X})(Y_i - \bar{Y})}{\sum(X_i - \bar{X})^2}$.
3. Find $\hat{\beta}_1 = \bar{Y} - \hat{\beta}_2\bar{X}$.
4. Write the estimated regression line.
5. Predict Y when $X = 20$.

Problem 3.2 – Residuals and Goodness of Fit

Using the estimated line from Problem 3.1, compute residuals $e_i = Y_i - \hat{Y}_i$ for each observation. Then calculate:

1. $\sum e_i$ (should equal approximately 0).
2. $R^2 = 1 - \frac{\sum e_i^2}{\sum (Y_i - \bar{Y})^2}$
3. Interpret the R^2 value.

Problem 3.3 – Testing CLRM Assumptions (Conceptual–Quantitative Mix)

For the model $Y_i = \beta_1 + \beta_2 X_i + U_i$, consider:

- $E(U_i | X_i) = 0$
- $Var(U_i | X_i) = \sigma^2$
- $Cov(U_i, X_i) = 0$

Tasks:

1. Explain what happens if $Var(U_i)$ increases with X_i .
2. Suggest how this can be detected using a residual plot.
3. Explain the impact on OLS efficiency.

Problem 3.4 – Variance of the OLS Slope

Suppose the estimated variance of residuals is $\hat{\sigma}^2 = 4$, and the data yield $\sum (X_i - \bar{X})^2 = 50$.

Find $Var(\hat{\beta}_2)$ and the standard error of $\hat{\beta}_2$.

$$Var(\hat{\beta}_2) = \frac{\hat{\sigma}^2}{\sum (X_i - \bar{X})^2}$$