

612303050 Deshmukh Mehmood Rehan's Assignment 1

Q1.Install packages namely 'plyr', 'MASS', 'ggplot2', 'dplyr' etc.

```
#we can use vector to install multiple packages simultaneously
install.packages(c('plyr', 'MASS', 'ggplot2', 'dplyr'), repos =
"http://cran.us.r-project.org")

## Installing packages into 'C:/Users/deshm/AppData/Local/R/win-library/4.3'
## (as 'lib' is unspecified)

## package 'plyr' successfully unpacked and MD5 sums checked
## package 'MASS' successfully unpacked and MD5 sums checked
## package 'ggplot2' successfully unpacked and MD5 sums checked
## package 'dplyr' successfully unpacked and MD5 sums checked
##
## The downloaded binary packages are in
## C:\Users\deshm\AppData\Local\Temp\RtmpUHftca\downloaded_packages
```

Q2.Find answers to $\log_2(2^5)$ and $\log(\exp(1)*\exp(1))$

```
answer1 <- log2(2^5)
answer1

## [1] 5

answer2 <- log(exp(1)*exp(1))
answer2

## [1] 2
```

Q3.Using built-in dataset iris, implement the functions like: Summary, class, type of, head,tail, str, Merge.

```
#loading dataset
data(iris)

#summary
iris_summary <- summary(iris)
iris_summary
```

##	Sepal.Length	Sepal.Width	Petal.Length	Petal.Width
##	Min. :4.300	Min. :2.000	Min. :1.000	Min. :0.100
##	1st Qu.:5.100	1st Qu.:2.800	1st Qu.:1.600	1st Qu.:0.300
##	Median :5.800	Median :3.000	Median :4.350	Median :1.300
##	Mean :5.843	Mean :3.057	Mean :3.758	Mean :1.199
##	3rd Qu.:6.400	3rd Qu.:3.300	3rd Qu.:5.100	3rd Qu.:1.800
##	Max. :7.900	Max. :4.400	Max. :6.900	Max. :2.500
##	Species			
##	setosa :50			

```

## versicolor:50
## virginica :50
##
##
##

#class
iris_class <- class(iris)
iris_class

## [1] "data.frame"

#typeof
iris_typeof <- typeof(iris)
iris_typeof

## [1] "list"

#head
iris_head <- head(iris)
iris_head

##   Sepal.Length Sepal.Width Petal.Length Petal.Width Species
## 1          5.1          3.5          1.4          0.2  setosa
## 2          4.9          3.0          1.4          0.2  setosa
## 3          4.7          3.2          1.3          0.2  setosa
## 4          4.6          3.1          1.5          0.2  setosa
## 5          5.0          3.6          1.4          0.2  setosa
## 6          5.4          3.9          1.7          0.4  setosa

#tail
iris_tail <- tail(iris)
iris_tail

##   Sepal.Length Sepal.Width Petal.Length Petal.Width  Species
## 145          6.7          3.3          5.7          2.5 virginica
## 146          6.7          3.0          5.2          2.3 virginica
## 147          6.3          2.5          5.0          1.9 virginica
## 148          6.5          3.0          5.2          2.0 virginica
## 149          6.2          3.4          5.4          2.3 virginica
## 150          5.9          3.0          5.1          1.8 virginica

#str
iris_str <- str(iris)

## 'data.frame':   150 obs. of  5 variables:
##  $ Sepal.Length: num  5.1 4.9 4.7 4.6 5 5.4 4.6 5 4.4 4.9 ...
##  $ Sepal.Width : num  3.5 3 3.2 3.1 3.6 3.9 3.4 3.4 2.9 3.1 ...
##  $ Petal.Length: num  1.4 1.4 1.3 1.5 1.4 1.7 1.4 1.5 1.4 1.5 ...
##  $ Petal.Width : num  0.2 0.2 0.2 0.2 0.2 0.4 0.3 0.2 0.2 0.1 ...
##  $ Species      : Factor w/ 3 levels "setosa","versicolor",...: 1 1 1 1 1 1
1 1 1 1 ...

```

```
#merge
iris_merge <- merge(iris, iris, by = "Petal.Length")
head(iris_merge)

##   Petal.Length Sepal.Length.x Sepal.Width.x Petal.Width.x Species.x
## 1          1.0           4.6           3.6           0.2    setosa
## 2          1.1           4.3           3.0           0.1    setosa
## 3          1.2           5.8           4.0           0.2    setosa
## 4          1.2           5.8           4.0           0.2    setosa
## 5          1.2           5.0           3.2           0.2    setosa
## 6          1.2           5.0           3.2           0.2    setosa
##   Sepal.Length.y Sepal.Width.y Petal.Width.y Species.y
## 1           4.6           3.6           0.2    setosa
## 2           4.3           3.0           0.1    setosa
## 3           5.8           4.0           0.2    setosa
## 4           5.0           3.2           0.2    setosa
## 5           5.8           4.0           0.2    setosa
## 6           5.0           3.2           0.2    setosa
```

Q4. Write a R program to create a two-dimensional 5×3 array of sequence of even integers greater than

```
answer4 <- array(seq(from = 52, by = 2, length.out = 5*3), dim = c(5, 3))
answer4

##      [,1] [,2] [,3]
## [1,]   52   62   72
## [2,]   54   64   74
## [3,]   56   66   76
## [4,]   58   68   78
## [5,]   60   70   80
```

Q5. Write a R program to create a vector which contains 10 integer values between -50 and +50

```
answer5 <- seq(-50, 50, 10)
answer5

## [1] -50 -40 -30 -20 -10  0  10  20  30  40  50
```

Q6. Suppose the age is a vector containing ages of 10 persons as 22,27,31,41,30,25,19,20,23,35

```
age <- c(22, 27, 31, 41, 30, 25, 19, 20, 23, 35)

#a). Access the age of fourth person
answer6a <- age[4]
answer6a

## [1] 41
```

#b).Create a vector of 'age 30' with a person >30

```
answer6b <- age[age > 30]
```

```
answer6b
```

```
## [1] 31 41 35
```

#c).Access the age of last 3 person

```
answer6c <- tail(age, 3)
```

```
answer6c
```

```
## [1] 20 23 35
```

#d).Find the number of elements in vector age

```
answer6d <- length(age)
```

```
answer6d
```

```
## [1] 10
```

#e).Access the age of person except 5th and 7th

```
answer6e <- age[-c(5, 7)]
```

```
answer6e
```

```
## [1] 22 27 31 41 25 20 23 35
```

#f).Create a vector 'age 2' with a persons between 20 and 25.

```
answer6f <- age[(age >= 20) & (age <= 25)]
```

```
answer6f
```

```
## [1] 22 25 20 23
```

Q7.Create a factor from the following vector data:(1,2,3,2,3,1,4,2,3,NA,5,3,2) and also find levels

```
answer7_factors <- factor(c(1, 2, 3, 2, 3, 1, 4, 2, 3, NA, 5, 3, 2))
```

```
answer7 <- levels(answer7_factors)
```

```
answer7
```

```
## [1] "1" "2" "3" "4" "5"
```

Q8.Write a R program to create a list containing strings, numbers, vectors and a logical values.

```
answer8 <- list("abcd", 123, c(1, 2, 3), TRUE)
```

```
answer8
```

```
## [[1]]
```

```
## [1] "abcd"
```

```
##
```

```
## [[2]]
```

```
## [1] 123
```

```
##
```

```
## [[3]]
```

```
## [1] 1 2 3
```

```
## [[4]]  
## [1] TRUE
```

Q9.Using built-in dataset iris, find out the categorical variables.

```
answer9 <- sapply(iris, is.factor)  
answer9  
  
## Sepal.Length Sepal.Width Petal.Length Petal.Width Species  
## FALSE FALSE FALSE FALSE TRUE
```

Q10.Create a numeric vector c(1:5) and a 5 by 3 matrix with elements from 1 to 15.

```
answer10_vector <- c(1:5)  
answer10_matrix <- matrix(1:15, nrow = 5, ncol = 3)  
answer10_vector  
  
## [1] 1 2 3 4 5  
  
answer10_matrix  
  
##      [,1] [,2] [,3]  
## [1,]    1    6   11  
## [2,]    2    7   12  
## [3,]    3    8   13  
## [4,]    4    9   14  
## [5,]    5   10   15
```

Q11. Create a dataframe of the following dataset

height:140,137,150,147,139,140,150,132,138,140 Weight:55,57,59,62,61,60,60,58,59,57

```
answer11_dataframe <- data.frame(  
  height = c(140, 137, 150, 147, 139, 140, 150, 132, 138, 140),  
  weight = c(55, 57, 59, 62, 61, 60, 60, 58, 59, 57)  
)  
  
#a).Create a vector h1 with height>145cms  
h1 <- answer11_dataframe$height[answer11_dataframe$height > 145]  
  
## [1] 150 147 150  
  
#b).Create a vector h2 with weight>55kgs  
h2 <- answer11_dataframe$weight[answer11_dataframe$weight > 55]  
h2  
  
## [1] 57 59 62 61 60 60 58 59 57  
  
#c).Create a vector h3 with height>140 and weight >60  
h3 <- answer11_dataframe$height[answer11_dataframe$height > 140 &  
  answer11_dataframe$weight > 60]  
h3  
  
## [1] 147
```