

### Choose Parameter Values

$r = 1$  for bias, variance  
 $r = 2$  for distribution, quantiles  
 $c_2 = \begin{cases} 1; & \text{for } r = 1 \\ 0.1; & \text{otherwise} \end{cases}$

### Calculate Bootstrap Estimates

Initial block size,  $\ell = n^{\frac{1}{r+4}}$ ,  
where  $n$  is th sample size  
(1) Calculate  $\hat{\varphi}_n(\ell)$   
(2) Calculate  $\hat{\varphi}_n(2\ell)$

### Estimate $\widehat{\text{VAR}}$

- (1) Set  $m = c_2 n^{1/3} \ell^{2/3}$  (no. of blocks to be deleted)
- (2) Calculate the Jackknife-after-Bootstrap (JAB) Variance estimator  $\widehat{\text{VAR}}$  using  $\hat{\varphi}_n(\ell)$ 
  - (i) Collect the indices set:  
 $I_i^* = \{k: 1 \leq k \leq K, \{ {}_k B_1^*, \dots, {}_k B_b^* \} \cap \{B_i, \dots, B_{i+m-1}\} = \emptyset\}$
  - (ii)  $\hat{\varphi}_n^{(i)}$ :  $i$ -th block-deleted jackknife point value, calculated using  $k \in I_i^*$
  - (iii)  $\tilde{\varphi}_n^{(i)} = m^{-1} [N\hat{\varphi}_n - (N - m)\hat{\varphi}_n^{(i)}]$
  - (iv)  $\widehat{\text{VAR}}_{\text{JAB}}(\hat{\varphi}_n) = \frac{m}{N-m} \frac{1}{M} \sum_{i=1}^M (\tilde{\varphi}_n^{(i)} - \hat{\varphi}_n)^2$ ,  
where,  $N$  = No. of all possible blocks, and  
 $M = N - m + 1$

### Calculate $\hat{C}_1$ and $\hat{C}_2$

$$\hat{C}_1 = n\ell^{-r}\widehat{\text{VAR}}$$
$$\hat{C}_2 = 2\ell\{\hat{\varphi}_n(\ell) - \hat{\varphi}_n(2\ell)\}$$

### Calculate Optimal Block Length, $\hat{\ell}^0$

$$\hat{\ell}^0 = \left[ \frac{2\hat{C}_2^2}{r\hat{C}_1} \right]^{\frac{1}{r+2}} n^{\frac{1}{r+2}}$$