

and (ii)  $a : a - b = c : c - d$

**(5) Theorem of Componendo dividendo**

If  $a : b = c : d$ , then

(i)  $a + b : a - b = c + d : c - d$

and (ii)  $a - b : a + b = c - d : c + d$

**SOLVED EXERCISE 3.4**

1. Prove that  $a : b = c : d$ , if

(i)  $\frac{4a + 5b}{4a - 5b} = \frac{4c + 5d}{4c - 5d}$

*Solution:*

Given  $\frac{4a + 5b}{4a - 5b} = \frac{4c + 5d}{4c - 5d}$

By componendo-dividendo theorem, we have

$$\frac{(4a + 5b) + (4c + 5d)}{(4a - 5b) - (4c - 5d)} = \frac{(4c + 5d) + (4c - 5d)}{(4c + 5d) - (4c - 5d)}$$

$$\frac{4a + 5b + 4c + 5d}{4a + 5b - 4c + 5d} = \frac{4c + 5d + 4c - 5d}{4c + 5d - 4c + 5d}$$

$$\frac{8a}{10b} = \frac{8c}{10d}$$

Multiplying both sides by  $\frac{18}{4}$ , we get

$$\frac{a}{b} = \frac{c}{d}$$

$\Rightarrow a : b = c : d$   
Hence proved

(ii)  $\frac{2a + 9b}{2a - 9b} = \frac{2c + 9d}{2c - 9d}$

*Solution:*

Given

$$\frac{2a + 9b}{2a - 9b} = \frac{2c + 9d}{2c - 9d}$$

By componendo-dividendo theorem, we have

$$\frac{(2a+9b)+(2a+9b)}{(2a+9b)-(2a-9b)} = \frac{(2c+9d)+(2c-9d)}{(2c+9d)-(2c-9d)}$$

$$\frac{2a+9b+2a-9b}{2a+9b-2a+9b} = \frac{2c+9d+2c-9d}{2c+9d-2c+9d}$$

$$\frac{4a}{18b} = \frac{4c}{18d}$$

Multiplying both sides by  $\frac{18}{4}$ , we get

$$\frac{a}{b} = \frac{c}{d}$$

$$\Rightarrow a : b = c : d$$

Hence proved

$$(iii) \frac{ac^2 + bd^2}{ac^2 - bd^2} = \frac{c^3 + d^3}{c^3 - d^3}$$

**Solution:**

Given

$$\frac{ac^2 + bd^2}{ac^2 - bd^2} = \frac{c^3 + d^3}{c^3 - d^3}$$

By componendo-dividendo theorem, we have

$$\frac{(ac^2 + bd^2) + (ac^2 - bd^2)}{(ac^2 + bd^2) - (ac^2 - bd^2)} = \frac{(c^3 + d^3) + (c^3 - d^3)}{(c^3 + d^3) - (c^3 - d^3)}$$

$$\frac{ac^2 + bd^2 + ac^2 - bd^2}{ac^2 + bd^2 - ac^2 + bd^2} = \frac{c^3 + d^3 + c^3 - d^3}{c^3 + d^3 - c^3 + d^3}$$

$$\frac{2ac^2}{2bd^2} = \frac{2c^3}{2d^3}$$

$$\Rightarrow \frac{ac^2}{bd^2} = \frac{c^3}{d^3}$$

Multiplying both sides by  $\frac{d^2}{c^2}$ , we get

$$\frac{a}{b} = \frac{c}{d}$$

$$\Rightarrow a : b = c : d$$

Hence proved

$$(iv) \frac{a^2c + b^2d}{a^2c - b^2d} = \frac{ac^2 + bd^2}{ac^2 - bd^2}$$

**Solution:**

Given

$$\frac{a^2c + b^2d}{a^2c - b^2d} = \frac{ac^2 + bd^2}{ac^2 - bd^2}$$

By componendo-dividendo theorem, we have

$$\frac{(a^2c + b^2d) + (a^2c - b^2d)}{(a^2c + b^2d) - (a^2c - b^2d)} = \frac{(ac^2 + bd^2) + (ac^2 - bd^2)}{(ac^2 + bd^2) - (ac^2 - bd^2)}$$

$$\frac{a^2c + b^2d + a^2c - b^2d}{a^2c + b^2d - a^2c + b^2d} = \frac{ac^2 + bd^2 + ac^2 - bd^2}{ac^2 + bd^2 - ac^2 + bd^2}$$

$$\frac{2a^2c}{2b^2d} = \frac{2ac^2}{2bd^2}$$

$$\Rightarrow \frac{a^3c}{b^2d} = \frac{ac^2}{bd^2}$$

Multiplying both sides by  $\frac{bd}{ac}$ , we get

$$\frac{a}{b} = \frac{c}{d}$$

$$\Rightarrow a : b = c : d$$

Hence proved

$$(v) \quad pa + qb : pa - qb = pc + qd : pc - qd$$

**Solution:**

Given

$$pa + qb : pa - qb = pc + qd : pc - qd$$

$$\frac{pa + qb}{pa - qb} = \frac{pc + qd}{pc - qd}$$

By componendo-dividendo theorem, we have

$$\frac{(pa + qb) + (pa - qb)}{(pa + qb) - (pa - qb)} = \frac{(pc + qd) + (pc - qd)}{(pc + qd) - (pc - qd)}$$

$$\frac{pa + qb + pa - qb}{pa + qb - pa + qb} = \frac{pc + qd + pc - qd}{pc + qd - pc + qd}$$

$$\frac{2pa}{2qb} = \frac{2pc}{2qd}$$

Multiplying both sides by  $\frac{2q}{2p}$ , we get

$$\frac{a}{b} = \frac{c}{d}$$

$$\Rightarrow a : b = c : d$$

Hence proved

$$(vi) \frac{a+b+c+d}{a+b-c-d} = \frac{a-b+c-d}{a-b-c+d}$$

*Solution:*

Given

$$\begin{aligned} \frac{a+b+c+d}{a+b-c-d} &= \frac{a-b+c-d}{a-b-c+d} \\ \frac{(a+b+c+d)+(a+b-c-d)}{(a+b+c+d)-(a+b-c-d)} &= \frac{(a-b+c-d)+(a-b-c+d)}{(a-b+c-d)-(a-b-c+d)} \\ \frac{a+b+c+d+a+b-c-d}{a+b+c+d-a-b+c+d} &= \frac{a-b+c-d+a-b-c+d}{a-b+c-d-a+b+c-d} \\ \frac{2a+2b}{2c+2d} &= \frac{2a-2b}{2c-2d} \\ \frac{2(a+b)}{2(c+d)} &= \frac{2(a-b)}{2(c-d)} \\ \frac{a+b}{c+d} &= \frac{a-b}{c-d} \end{aligned}$$

or  $\frac{a+b}{c-d} = \frac{c+d}{c-d}$

By componendo-dividendo theorem, we have

$$\begin{aligned} \frac{(a+b)+(a-b)}{(a+b)-(a-b)} &= \frac{(c+d)+(c-d)}{(c+d)-(c-d)} \\ \frac{a+b+a-b}{a+b-a+b} &= \frac{c+d+c-d}{c+d-c+d} \\ \frac{2a}{2b} &= \frac{2c}{2d} \end{aligned}$$

$$\Rightarrow \frac{a}{b} = \frac{c}{d}$$

$$\Rightarrow a:b = c:d$$

Hence proved

$$(vii) \frac{2a+3b+2c+3d}{2a+3b-2c-3d} = \frac{2a-3b+2c-3d}{2a-3b-2c+3d}$$

*Solution:*

Given

$$\frac{(2a + 3b + 2c + 3d) + (2a + 3b - 2c - 3d)}{(2a + 3b + 2c + 3d) - (2a + 3b - 2c - 3d)} = \frac{(2a - 3b + 2c - 3d) + (2a - 3b - 2c + 3d)}{(2a - 3b + 2c - 3d) - (2a - 3b - 2c + 3d)}$$

$$\frac{2a + 3b + 2c + 3d + 2a + 3b - 2c - 3d}{2a + 3b + 2c + 3d - 2a - 3b + 2c + 3d} = \frac{2a - 3b + 2c - 3d + 2a - 3b - 2c + 3d}{2a - 3b + 2c - 3d - 2a + 3b - 2c + 3d}$$

$$\frac{4a + 6b}{4c + 6d} = \frac{4a - 6b}{4c - 6d}$$

or  $\frac{4a + 6b}{4c - 6d} = \frac{4c + 6d}{4c - 6d}$

By componendo-dividendo theorem, we have

$$\frac{(4a + 6b) + (4a - 6b)}{(4a + 6b) - (4a - 6b)} = \frac{(4c + 6d) + (4c - 6d)}{(4c + 6d) - (4c - 6d)}$$

$$\frac{4a + 6b + 4a - 6b}{4a + 6b - 4a + 6b} = \frac{4c + 6d + 4c - 6d}{4c + 6d - 4c + 6d}$$

$$\frac{8a}{12b} = \frac{8c}{12d}$$

Multiplying both sides by  $\frac{12}{8}$ , we get

$$\frac{a}{b} = \frac{c}{d}$$

$\Rightarrow a : b = c : d$   
Hence proved

$$(viii) \frac{a^2 + b^2}{a^2 - b^2} = \frac{ac + bd}{ac - bd}$$

**Solution:**

Given

$$\frac{a^2 + b^2}{a^2 - b^2} = \frac{ac + bd}{ac - bd}$$

By componendo-dividendo theorem, we have

$$\frac{(a^2 + b^2) + (a^2 - b^2)}{(a^2 + b^2) - (a^2 - b^2)} = \frac{(ac + bd) + (ac - bd)}{(ac + bd) - (ac - bd)}$$

$$\frac{a^2 + b^2 + a^2 - b^2}{a^2 + b^2 - a^2 + b^2} = \frac{ac + bd + ac - bd}{ac + bd - ac + bd}$$

$$\frac{2a^2}{2b^2} = \frac{2ac}{2bd}$$

Multiplying both sides by  $\frac{2b}{2a}$ , we get

$$\frac{a}{b} = \frac{c}{d}$$

$\Rightarrow a : b = c : d$   
Hence proved

## 2. Using theorem of componendo-dividendo

(i) Find the value of  $\frac{x+2y}{x-2y} + \frac{x+2z}{x-2z}$ , if  $x = \frac{4yz}{y+z}$

*Solution:*

$$\text{Given } x = \frac{4yz}{y+z} \quad \text{or} \quad x = \frac{(2y)(2z)}{y+z}$$

$$\text{or} \quad \frac{x}{2y} = \frac{2z}{y+z}$$

Applying componendo-dividendo theorem, we get.

$$\frac{x+2y}{x-2y} = \frac{2z+(y+z)}{2z-(y+z)}$$

$$\frac{x+2y}{x-2y} = \frac{2z+y+z}{2z-y-z}$$

$$\frac{x+2y}{x-2y} = \frac{y+3z}{z-y} \quad \text{--- (i)}$$

$$\text{Now } x = \frac{4yz}{y+z} \quad \text{or} \quad x = \frac{(2y)(2z)}{y+z}$$

$$\text{or} \quad \frac{x}{2z} = \frac{2y}{y+z}$$

Applying componendo-dividendo theorem, we get.

$$\frac{x+2z}{x-2z} = \frac{2y+(y+z)}{2y-(y+z)}$$

$$\frac{x+2z}{x-2z} = \frac{2y+y+z}{2y-y-z}$$

$$\frac{x+2z}{x-2z} = \frac{3y+z}{y-z}$$

Adding eq. (i) and eq. (ii), we get

$$\begin{aligned}
\frac{x+2y}{x-2y} + \frac{x+2z}{x-2z} &= \frac{y+3z}{z-y} + \frac{3y+z}{y-z} \\
&= \frac{y+3z}{z-y} + \frac{3y+z}{-(z-y)} \\
&= \frac{y+3z}{z-y} - \frac{3y+z}{z-y} \\
&= \frac{(y+3z)-(3y+z)}{z-y} \\
&= \frac{y+3z-3y-z}{z-y} \\
&= \frac{2z-2y}{z-y} \\
&= \frac{2(z-y)}{z-y} = 2
\end{aligned}$$

(ii) Find the value of  $\frac{m+5n}{m-5n} + \frac{m+5p}{m-5p}$ , if  $m = \frac{10np}{n+p}$

**Solution:**

$$\text{Given } m = \frac{10np}{n+p} \quad \text{or} \quad x = \frac{(2p)(5n)}{n+p}$$

$$\text{or} \quad \frac{xm}{5n} = \frac{2p}{n+p}$$

Applying componendo-dividendo theorem, we get.

$$\frac{m+5n}{m-5n} = \frac{2p+(n+p)}{2p-(n+p)}$$

$$\frac{m+5n}{m-5n} = \frac{2p+n+p}{2p-n-p}$$

$$\frac{m+5n}{m-5n} = \frac{3p+n}{p-n} \quad \text{--- (i)}$$

$$\text{Now } m = \frac{10np}{n+p} \quad \text{or} \quad m = \frac{(5p)(2n)}{n+p}$$

$$\text{or} \quad \frac{m}{5p} = \frac{2n}{n+p}$$

Applying componendo - dividendo theorem, we get.

$$\frac{m+5n}{m-5n} = \frac{2n+(n+p)}{2n-(n+p)}$$

$$\frac{m+5n}{m-5n} = \frac{2n+n+p}{2n-n-p}$$

$$\frac{m+5n}{m-5n} + \frac{3n+p}{n-p} \text{ ————— (ii)}$$

Adding eq. (i) and eq. (ii), we get.

$$\begin{aligned} \frac{m+5n}{m-5n} &= \frac{m+5p}{m-5p} = \frac{3p+n}{p-n} + \frac{2n+p}{n-p} \\ &= \frac{3p+n}{p-n} + \frac{2n+p}{-(p-n)} \\ &= \frac{3p+n}{p-n} - \frac{2n+p}{p-n} \\ &= \frac{(3p+n)-(2n+p)}{p-n} \\ &= \frac{3p+n-2n-p}{p-n} \\ &= \frac{2p-2n}{p-n} \\ &= \frac{2(p-n)}{p-n} = 2 \end{aligned}$$

(iii) Find the value of  $\frac{x-6a}{x+6a} - \frac{x-6b}{x+6b}$ , if  $x = \frac{12ab}{a-b}$

**Solution**

$$\text{Given } x = \frac{12ab}{a-b} \quad \text{or } x = \frac{(6a)(2b)}{a-b}$$

$$\text{or } \frac{x}{6a} = \frac{2b}{a-b}$$

Applying componendo-dividendo theorem, we get.

$$\frac{x+6a}{x-6a} = \frac{2b+(a-b)}{2b-(a-b)}$$

$$\frac{x+6a}{x-6a} = \frac{2b+a-b}{2b-a+b}$$

$$\frac{x+6a}{x-6a} = \frac{a+b}{3b-a}$$



$$\text{or } \frac{x-6a}{x+6a} = \frac{3b-a}{a+b} \text{ ————— (i)}$$

$$\text{Now } x = \frac{12ab}{a-b} \quad \text{or} \quad x = \frac{(2a)(6b)}{a-b}$$

$$\text{or } \frac{x}{6b} = \frac{2a}{a-b}$$

Applying componendo- dividend theorem, we get.

$$\frac{x+6b}{x-6b} = \frac{2a+(a-b)}{2a-(a-b)}$$

$$\frac{x+6b}{x-6b} = \frac{2a+a-b}{2a-a+b}$$

$$\frac{x+6b}{x-6b} = \frac{3a+b}{a+b} \text{ ————— (ii)}$$

Subtract eq. (ii) from eq. (i), we have.

$$\begin{aligned} \frac{x-6a}{x+6a} - \frac{x+6b}{x-6b} &= \frac{3b-a}{a+b} - \frac{3a-b}{a+b} \\ &= \frac{(3b-a)-(3a-b)}{a+b} \\ &= \frac{3b-a-3a+b}{a+b} \\ &= \frac{4b-4a}{a+b} \\ &= \frac{4(b-a)}{a+b} \end{aligned}$$

(iv) Find the value of  $\frac{x-3y}{x+3y} - \frac{x-3z}{x+3z}$ , if  $x = \frac{3yz}{y-z}$

*Solution:*

$$\text{Given } x = \frac{3yz}{y-z}$$

$$\text{or } \frac{x}{3y} = \frac{z}{y-z}$$

Applying componendo- dividend theorem, we get.

$$\frac{x+3y}{x-3y} = \frac{z+(y-z)}{z-(y-z)}$$

$$\frac{x+3y}{x-3y} = \frac{z+y-z}{z-y+z}$$

$$\frac{x+3y}{x-3y} = \frac{y}{2z-y}$$

$$\text{or } \frac{x+3y}{x-3y} = \frac{2z-y}{y} \quad \text{--- (i)}$$

Now  $x = \frac{3yz}{y-z}$

$$\frac{x}{3z} = \frac{y}{y-z}$$

Applying componendo-dividendo theorem, we get.

$$\frac{x+3z}{x-3z} = \frac{y+(y-z)}{y-(y-z)}$$

$$\frac{x+3z}{x-3z} = \frac{y+y-z}{y-y+z}$$

$$\frac{x+3z}{x-3z} = \frac{2y-z}{z} \quad \text{--- (ii)}$$

Subtracting eq. (ii) from eq. (i), we get.

$$\begin{aligned} \frac{x+3y}{x+3y} - \frac{x+3z}{x-3z} &= \frac{2z-y}{y} - \frac{2y-z}{z} \\ &= \frac{z(2z-y) - y(2y-z)}{yz} \end{aligned}$$

(v) Find the value of  $\frac{s-3p}{s+3p} + \frac{s-3q}{s+3q}$ , if  $s = \frac{6pq}{s-3q}$

**Solution:**

Given  $s = \frac{6pq}{s-3q}$  or  $s = \frac{(3p)(2q)}{p-q}$  or  $\frac{s}{3p} = \frac{2q}{p-q}$

Applying componendo-dividendo theorem, we get.

$$\frac{s+3p}{s-3p} = \frac{2q+(p-q)}{2q-(p-q)}$$

$$\frac{s+3p}{s-3p} = \frac{2q+p-q}{2q-p+q}$$

$$\frac{s+3p}{s-3p} = \frac{p+q}{3q-p}$$

$$\frac{s+3p}{s-3p} = \frac{3q-p}{p+q} \quad \text{————— (i)}$$

Now  $S = \frac{6pq}{p-q}$  or  $S = \frac{(3q)(2p)}{p-q}$

or  $\frac{S}{3q} = \frac{2p}{p-q}$

Applying componendo-dividendo theorem, we get.

$$\frac{s+3p}{s-3p} = \frac{2p+(p-q)}{2p-(p-q)}$$

$$\frac{s+3p}{s-3p} = \frac{2p+p-q}{2p-p+q}$$

$$\frac{s+3p}{s-3p} = \frac{3p-p}{p+q} \quad \text{————— (ii)}$$

Adding eq, (i) and eq, (ii), we get.

$$\begin{aligned} \frac{s-3p}{s+3p} + \frac{s+3p}{s-3p} &= \frac{3q-p}{p+q} + \frac{3p-p}{p+q} \\ &= \frac{3q-p+3p-q}{p+q} \\ &= \frac{2p=2q}{p+q} \\ &= \frac{2(p+q)}{p+q} = 2 \end{aligned}$$

(vi) Solve  $\frac{(x-2)^2 - (x-4)^2}{(x-2)^2 + (x-4)^2} = \frac{12}{13}$

**Solution:**

$$\frac{(x-2)^2 - (x-4)^2}{(x-2)^2 + (x-4)^2} = \frac{12}{13}$$

Applying componendo-dividendo theorem, we get.

$$\frac{\left[ (x-2)^2 - (x-4)^2 \right] + \left[ (x-2)^2 + (x-4)^2 \right]}{\left[ (x-2)^2 - (x-4)^2 \right] - \left[ (x-2)^2 + (x-4)^2 \right]} = \frac{12+13}{12-13}$$

$$\frac{(x-2)^2 - (x-4)^2 + (x-2)^2 + (x-4)^2}{(x-2)^2 + (x-4)^2 - (x-2)^2 - (x-4)^2} = \frac{25}{-1}$$

$$\frac{2(x-2)^2}{-2(x-4)^2} = -25$$

$$\Rightarrow \frac{(x-2)^2}{(x-4)^2} = 25$$

Taking square root on both sides, we get.

$$\sqrt{\frac{(x-2)^2}{(x-4)^2}} = \pm\sqrt{25}$$

$$\frac{x-2}{x-4} = \pm 5$$

$$\Rightarrow \frac{x-2}{x-4} = -5 \quad \text{or} \quad \frac{x-2}{x-4} = 5$$

$$x-2 = -5(x-4) \quad 5(x-4) = x-2$$

$$x-2 = -5x+20 \quad 5x-20 = x-2$$

$$x+5x = 2+20 \quad 5x-x = 20-2$$

$$6x = 22 \quad 4x = 18$$

$$x = \frac{22}{6} \quad x = \frac{18}{4}$$

$$x = \frac{11}{3} \quad x = \frac{9}{2}$$

$$\text{Thus, solution set} = \left\{ \frac{9}{2}, \frac{11}{3} \right\}$$

$$\text{(vii) Solve } \frac{\sqrt{x^2+2} + \sqrt{x^2-2}}{\sqrt{x^2+2} - \sqrt{x^2-2}} = 2$$

**Solution:**

$$\frac{\sqrt{x^2+2} + \sqrt{x^2-2}}{\sqrt{x^2+2} - \sqrt{x^2-2}} = \frac{2}{1}$$

Applying componendo-dividendo theorem, we get.

$$\frac{\left[\sqrt{x^2+2}+\sqrt{x^2-2}\right]+\left[\sqrt{x^2+2}-\sqrt{x^2-2}\right]}{\left[\sqrt{x^2+2}-\sqrt{x^2-2}\right]-\left[\sqrt{x^2+2}-\sqrt{x^2-2}\right]}=\frac{2+1}{2-1}$$

$$\frac{\sqrt{x^2+2}+\sqrt{x^2-2}+\sqrt{x^2+2}-\sqrt{x^2-2}}{\sqrt{x^2+2}-\sqrt{x^2-2}-\sqrt{x^2+2}+\sqrt{x^2-2}}=\frac{3}{1}$$

$$\frac{2\sqrt{x^2+2}}{2\sqrt{x^2-2}}=3$$

Squaring both sides, we get.

$$\frac{x^2+2}{x^2-2}=9$$

$$9(x^2-2)=x^2+2$$

$$9x^2-18=x^2+2$$

$$9x^2-x^2=18+2$$

$$8x^2=20$$

$$x^2=\frac{20}{8}$$

$$x^2=\frac{5}{2}$$

$$x=\pm\sqrt{\frac{5}{2}}$$

Thus, solution set =  $\left\{\pm\sqrt{\frac{5}{2}}\right\}$

(viii) Solve  $\frac{\sqrt{x^2+8p^2}+\sqrt{x^2-p^2}}{\sqrt{x^2+8p^2}-\sqrt{x^2-p^2}}=\frac{1}{3}$

*Solution:*

$$\frac{\sqrt{x^2+8p^2}+\sqrt{x^2-p^2}}{\sqrt{x^2+8p^2}-\sqrt{x^2-p^2}}=\frac{1}{3}$$

Applying componendo-dividendo theorem, we get.

$$\frac{\left[\sqrt{x^2+8p^2}+\sqrt{x^2-p^2}\right]+\left[\sqrt{x^2+8p^2}+\sqrt{x^2-p^2}\right]}{\left[\sqrt{x^2+8p^2}-\sqrt{x^2-p^2}\right]-\left[\sqrt{x^2+8p^2}+\sqrt{x^2-p^2}\right]}=\frac{1+3}{1-3}$$

$$\frac{\sqrt{x^2 + 8p^2} - \sqrt{x^2 - p^2} + \sqrt{x^2 + 8p^2} + \sqrt{x^2 - p^2}}{\sqrt{x^2 + 8p^2} - \sqrt{x^2 - p^2} - \sqrt{x^2 + 8p^2} - \sqrt{x^2 - p^2}} = \frac{4}{-2}$$

$$\frac{2\sqrt{x^2 + 8p^2}}{-2\sqrt{x^2 - p^2}} = -2$$

$$\Rightarrow \frac{\sqrt{x^2 + 8p^2}}{\sqrt{x^2 - p^2}} = 2$$

Squaring both sides, we get.

$$\frac{x^2 + 8p^2}{x^2 - p^2} = 4$$

$$4(x^2 - p^2) = x^2 + 8p^2$$

$$4x^2 - 4p^2 = x^2 + 8p^2$$

$$4x^2 - x^2 = 8p^2 + 4p^2$$

$$3x^2 = 12p^2$$

$$x^2 = 4p^2$$

$$\Rightarrow x = \pm 2p$$

Thus, solution set =  $\{+2p, -2p\}$

(ix) Solve  $\frac{(x+5)^3 - (x-3)^3}{(x+5)^3 + (x-3)^3} = \frac{13}{14}$

**Solution:**

$$\frac{(x+5)^3 - (x-3)^3}{(x+5)^3 + (x-3)^3} = \frac{13}{14}$$

Applying componendo-dividendo theorem, we get.

$$\frac{[(x+5)^3 - (x-3)^3] + [(x+5)^3 + (x-3)^3]}{[(x+5)^3 - (x-3)^3] - [(x+5)^3 + (x-3)^3]} = \frac{13+14}{13-14}$$

$$\frac{(x+5)^3 - (x-3)^3 + (x+5)^3 + (x-3)^3}{(x+5)^3 - (x-3)^3 - (x+5)^3 - (x-3)^3} = \frac{27}{-1}$$

$$\frac{2(x+5)^3}{-2(x-3)^3} = -27$$

$$\Rightarrow \frac{(x+5)^3}{(x-3)^3} = 27$$

$$\frac{(x+5)^3}{(x-3)^3} = (3)^3$$

Taking power ' $\frac{1}{3}$ ' on both sides, we get.

$$\frac{[(x+5)^3]^{\frac{1}{3}}}{[(x-3)^3]^{\frac{1}{3}}} = [(3)^3]^{\frac{1}{3}}$$

$$\frac{x+5}{x-3} = 3$$

$$3(x-3) = x+5$$

$$3x - 9 = x + 5$$

$$3x - x = 9 + 5$$

$$2x = 14$$

$$\Rightarrow x = 7$$

Thus, solution set = {7}

### (i) Joint variation

A combination of direct and inverse variations of one or more than one variables forms joint variation,

If a variable  $y$  varies directly as  $x$  and varies inversely as  $z$ .

$$\text{Then } y \propto x \text{ and } y \propto \frac{1}{z}$$

In joint variation, we write it as

$$y \propto \frac{x}{z}$$

$$\text{i.e., } y = k \frac{x}{z}$$

Where  $k \neq 0$  is the constant of variation.

## SOLVED EXERCISE 3.5

1. If  $s$  varies directly as  $u^2$  and inversely as  $v$  and  $s = 7$  when  $u = 3$ ,  $v = 2$ . Find the value of  $s$  when  $u = 6$  and  $v = 10$ .

*Solution:*

Given that  $s$  varies directly as  $u^2$ , so

$$S \propto u^2$$

Also given that  $S$  varies inversely as  $V$ , so

$$S \propto \frac{1}{V}$$