

EXERCISE 2.6

2.10 Derivative of General Exponential Function (Page 80)

A function define by

$$f(x)=a^x \text{ where } a > 0, a \neq 1$$

is called general exponential function.

Suppose

$$y=a^x$$

$$\Rightarrow y + \delta y = a^{x+\delta x} \Rightarrow \delta y = a^{x+\delta x} - y$$

$$\Rightarrow \delta y = a^{x+\delta x} - a^x \quad \text{Since } y = a^x$$

$$\Rightarrow \delta y = a^x (a^{\delta x} - 1)$$

Dividing by δx

$$\frac{\delta y}{\delta x} = \frac{a^x (a^{\delta x} - 1)}{\delta x}$$

Taking limit as $\delta x \rightarrow 0$

$$\begin{aligned} \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x} &= \lim_{\delta x \rightarrow 0} \frac{a^x (a^{\delta x} - 1)}{\delta x} \\ \Rightarrow \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} a^x \left(\frac{a^{\delta x} - 1}{\delta x} \right) \Rightarrow \frac{dy}{dx} = a^x \lim_{\delta x \rightarrow 0} \left(\frac{a^{\delta x} - 1}{\delta x} \right) \\ \Rightarrow \boxed{\frac{d}{dx}(a^x) = a^x \cdot \ln a} &\quad \text{Since } \lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \ln a \end{aligned}$$

Derivative of Natural Exponential Function

The exponential function $f(x)=e^x$ where $e = 2.71828...$ is called Natural Exponential Function.

Suppose

$$y = e^x$$

Do yourself ... Just Change a by e in above article. You'll get

$$\boxed{\frac{d}{dx}e^x = e^x}$$

2.11 Derivative of General Logarithmic Function (page 81)

If $a > 0, a \neq 1$ and $x = a^y$, then the function defined by $y = \log_a x$ ($x > 0$) is called General Logarithmic Function.

Suppose

$$y = \log_a x$$

$$\Rightarrow y + \delta y = \log_a (x + \delta x) \Rightarrow \delta y = \log_a (x + \delta x) - y$$

$$\Rightarrow \delta y = \log_a (x + \delta x) - \log_a x$$

$$= \log_a \left(\frac{x + \delta x}{x} \right) \quad \text{Since } \log_a m - \log_a n = \log_a \frac{m}{n}$$

Dividing both sides by δx

$$\frac{\delta y}{\delta x} = \frac{1}{\delta x} \log_a \left(\frac{x + \delta x}{x} \right)$$

Taking limit as $\delta x \rightarrow 0$

$$\begin{aligned} \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x} &= \lim_{\delta x \rightarrow 0} \frac{1}{\delta x} \log_a \left(\frac{x + \delta x}{x} \right) \\ \Rightarrow \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} \frac{1}{\delta x} \log_a \left(1 + \frac{\delta x}{x} \right) \\ &= \lim_{\delta x \rightarrow 0} \frac{x}{x} \cdot \frac{1}{\delta x} \log_a \left(1 + \frac{\delta x}{x} \right) \quad \div \text{ing and } \times \text{ing by } x \end{aligned}$$

$$\begin{aligned}
\Rightarrow \quad \frac{dy}{dx} &= \frac{1}{x} \lim_{\delta x \rightarrow 0} \frac{x}{\delta x} \log_a \left(1 + \frac{\delta x}{x} \right) \\
\Rightarrow \quad \frac{dy}{dx} &= \frac{1}{x} \lim_{\delta x \rightarrow 0} \log_a \left(1 + \frac{\delta x}{x} \right)^{\frac{x}{\delta x}} && \text{Since } m \log_a x = \log_a x^m \\
\Rightarrow \quad \frac{dy}{dx} &= \frac{1}{x} \log_a \left[\lim_{\delta x \rightarrow 0} \left(1 + \frac{\delta x}{x} \right)^{\frac{x}{\delta x}} \right] \\
\Rightarrow \quad \frac{dy}{dx} &= \frac{1}{x} \log_a e && \text{Since } \lim_{x \rightarrow 0} (1+x)^{\frac{1}{x}} = e \\
\Rightarrow \quad \frac{d}{dx} (\log_a x) &= \frac{1}{x} \frac{1}{\log_e a} && \text{Since } \log_a b = \frac{1}{\log_b a} \\
\Rightarrow \quad \frac{d}{dx} (\log_a x) &= \frac{1}{x \ln a} && \text{Since } \log_e a = \ln a
\end{aligned}$$

Derivative of Natural Logarithmic Function

The logarithmic function $f(x) = \log_e x$ where $e = 2.71828\dots$ is called Natural Logarithmic Function. And we write $\ln x$ instead of $\log_e x$ for our ease.

Suppose $y = \ln x$

$$\begin{aligned}
\Rightarrow y + \delta y &= \ln(x + \delta x) \Rightarrow \delta y = \ln(x + \delta x) - y \\
\Rightarrow \delta y &= \ln(x + \delta x) - \ln x \\
\Rightarrow \delta y &= \ln \left(\frac{x + \delta x}{x} \right) && \text{Since } \Rightarrow \ln m - \ln n = \ln \frac{m}{n} \\
&= \ln \left(1 + \frac{\delta x}{x} \right)
\end{aligned}$$

Dividing both sides by δx

$$\frac{\delta y}{\delta x} = \frac{1}{\delta x} \ln \left(1 + \frac{\delta x}{x} \right)$$

Taking limit as $\delta x \rightarrow 0$

$$\begin{aligned}
\lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x} &= \lim_{\delta x \rightarrow 0} \frac{1}{\delta x} \ln \left(1 + \frac{\delta x}{x} \right) \\
\Rightarrow \quad \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} \frac{x}{x} \cdot \frac{1}{\delta x} \ln \left(1 + \frac{\delta x}{x} \right) && \div \text{ing and } \times \text{ing by } x \\
\Rightarrow \quad \frac{dy}{dx} &= \frac{1}{x} \lim_{\delta x \rightarrow 0} \frac{x}{\delta x} \ln \left(1 + \frac{\delta x}{x} \right) \\
\Rightarrow \quad \frac{dy}{dx} &= \frac{1}{x} \lim_{\delta x \rightarrow 0} \ln \left(1 + \frac{\delta x}{x} \right)^{\frac{x}{\delta x}} && \text{Since } m \ln x = \ln x^m \\
\Rightarrow \quad \frac{dy}{dx} &= \frac{1}{x} \ln \left[\lim_{\delta x \rightarrow 0} \left(1 + \frac{\delta x}{x} \right)^{\frac{x}{\delta x}} \right] \\
\Rightarrow \quad \frac{dy}{dx} &= \frac{1}{x} \ln e && \text{Since } \lim_{x \rightarrow 0} (1+x)^{\frac{1}{x}} = e \\
\Rightarrow \quad \frac{d}{dx} (\ln x) &= \frac{1}{x} \cdot 1 && \text{Since } \ln e = \log_e e = 1 \\
\Rightarrow \quad \frac{d}{dx} (\ln x) &= \frac{1}{x}
\end{aligned}$$

Question # 1(i)

$$f(x) = e^{\sqrt{x}-1}$$

Diff. w.r.t x

$$\frac{d}{dx} f(x) = \frac{d}{dx} e^{\sqrt{x}-1}$$

$$\begin{aligned}\Rightarrow f'(x) &= e^{\sqrt{x}-1} \frac{d}{dx} (\sqrt{x}-1) \\ &= e^{\sqrt{x}-1} \left(\frac{1}{2} x^{-\frac{1}{2}} - 0 \right) = \frac{e^{\sqrt{x}-1}}{2\sqrt{x}} \quad \text{Ans.}\end{aligned}$$

Question # 1(ii)

$$f(x) = x^3 e^{\frac{1}{x}}$$

Diff. w.r.t x

$$\frac{d}{dx} f(x) = \frac{d}{dx} x^3 e^{\frac{1}{x}}$$

$$\begin{aligned}\Rightarrow f'(x) &= x^3 \frac{d}{dx} e^{\frac{1}{x}} + e^{\frac{1}{x}} \frac{d}{dx} x^3 \\ &= x^3 e^{\frac{1}{x}} \frac{d}{dx} \left(\frac{1}{x} \right) + e^{\frac{1}{x}} (3x^2) \\ &= x^3 e^{\frac{1}{x}} \left(-\frac{1}{x^2} \right) + e^{\frac{1}{x}} (3x^2) \qquad \because \frac{d}{dx} \left(\frac{1}{x} \right) = \frac{d}{dx} x^{-1} = -x^{-2} = -\frac{1}{x^2} \\ &= -x e^{\frac{1}{x}} + 3x^2 e^{\frac{1}{x}} = x e^{\frac{1}{x}} (3x-1) \quad \text{Ans.}\end{aligned}$$

Question # 1(iii)

$$f(x) = e^x (1 + \ln x)$$

Diff. w.r.t x

$$\frac{d}{dx} f(x) = \frac{d}{dx} e^x (1 + \ln x)$$

$$\begin{aligned}\Rightarrow f'(x) &= e^x \frac{d}{dx} (1 + \ln x) + (1 + \ln x) \frac{d}{dx} e^x \\ &= e^x \left(0 + \frac{1}{x} \right) + (1 + \ln x) e^x \\ \Rightarrow f'(x) &= e^x \left(\frac{1}{x} + 1 + \ln x \right) \quad \text{or} \quad f'(x) = e^x \left(\frac{1 + x(1 + \ln x)}{x} \right)\end{aligned}$$

Question # 1(iv)

$$f(x) = \frac{e^x}{e^{-x} + 1}$$

Diff. w.r.t x

$$\frac{d}{dx} f(x) = \frac{d}{dx} \left(\frac{e^x}{e^{-x} + 1} \right)$$

$$\Rightarrow f'(x) = \frac{(e^{-x} + 1) \frac{d}{dx} e^x - e^x \frac{d}{dx} (e^{-x} + 1)}{(e^{-x} + 1)^2}$$

$$= \frac{(e^{-x} + 1)e^x - e^x(e^{-x}(-1) + 0)}{(e^{-x} + 1)^2} = \frac{e^x(e^{-x} + 1 + e^{-x})}{(e^{-x} + 1)^2}$$

$$\Rightarrow f'(x) = \frac{e^x(2e^{-x} + 1)}{(e^{-x} + 1)^2} \text{ Ans.}$$

Question # 1(v)

$$f(x) = \ln(e^x + e^{-x})$$

Diff. w.r.t x

$$\frac{d}{dx} f(x) = \frac{d}{dx} \ln(e^x + e^{-x})$$

$$\Rightarrow f'(x) = \frac{1}{(e^x + e^{-x})} \frac{d}{dx} (e^x + e^{-x})$$

$$= \frac{1}{(e^x + e^{-x})} (e^x + e^{-x}(-1))$$

$$\Rightarrow f'(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}} \quad \text{or} \quad f'(x) = \tanh x \quad \because \tanh x = \frac{e^x - e^{-x}}{e^x + e^{-x}}$$

Question # 1(vi)

$$f(x) = \frac{e^{ax} - e^{-ax}}{e^{ax} + e^{-ax}}$$

Diff. w.r.t x

$$\frac{d}{dx} f(x) = \frac{d}{dx} \left(\frac{e^{ax} - e^{-ax}}{e^{ax} + e^{-ax}} \right)$$

$$= \frac{(e^{ax} + e^{-ax}) \frac{d}{dx} (e^{ax} - e^{-ax}) - (e^{ax} - e^{-ax}) \frac{d}{dx} (e^{ax} + e^{-ax})}{(e^{ax} + e^{-ax})^2}$$

$$= \frac{(e^{ax} + e^{-ax})(e^{ax}(a) - e^{-ax}(-a)) - (e^{ax} - e^{-ax})(e^{ax} + e^{-ax}(-a))}{(e^{ax} + e^{-ax})^2}$$

$$= \frac{a(e^{ax} + e^{-ax})(e^{ax} + e^{-ax}) - a(e^{ax} - e^{-ax})(e^{ax} - e^{-ax})}{(e^{ax} + e^{-ax})^2}$$

$$= \frac{a[(e^{ax} + e^{-ax})^2 - (e^{ax} - e^{-ax})^2]}{(e^{ax} + e^{-ax})^2}$$

$$= \frac{a[(e^{2ax} + e^{-2ax} + 2e^{ax}e^{-ax}) - (e^{2ax} + e^{-2ax} - 2e^{ax}e^{-ax})]}{(e^{ax} + e^{-ax})^2}$$

$$= \frac{a[e^{2ax} + e^{-2ax} + 2 - e^{2ax} - e^{-2ax} + 2]}{(e^{ax} + e^{-ax})^2} \quad \because e^{ax}e^{-ax} = e^0 = 1$$

$$\Rightarrow f'(x) = \frac{4a}{(e^{ax} + e^{-ax})^2} \text{ Ans.}$$

Question # 1(viii)

$$\begin{aligned}
f(x) &= \sqrt{\ln(e^{2x} + e^{-2x})} \\
\Rightarrow \frac{d}{dx} f(x) &= \frac{d}{dx} \left[\ln(e^{2x} + e^{-2x}) \right]^{\frac{1}{2}} \\
\Rightarrow f'(x) &= \frac{1}{2} \left[\ln(e^{2x} + e^{-2x}) \right]^{-\frac{1}{2}} \frac{d}{dx} \ln(e^{2x} + e^{-2x}) \\
&= \frac{1}{2 \left[\ln(e^{2x} + e^{-2x}) \right]^{\frac{1}{2}}} \cdot \frac{1}{(e^{2x} + e^{-2x})} \frac{d}{dx} (e^{2x} + e^{-2x}) \\
&= \frac{1}{2 \sqrt{\ln(e^{2x} + e^{-2x})}} \cdot \frac{1}{(e^{2x} + e^{-2x})} (e^{2x}(2) + e^{-2x}(-2)) \\
&= \frac{1}{2 \sqrt{\ln(e^{2x} + e^{-2x})}} \cdot \frac{2(e^{2x} - e^{-2x})}{(e^{2x} + e^{-2x})} = \frac{(e^{2x} - e^{-2x})}{(e^{2x} + e^{-2x}) \sqrt{\ln(e^{2x} + e^{-2x})}} \quad \text{Ans.}
\end{aligned}$$

Question # 1(ix)

$$\begin{aligned}
f(x) &= \ln \sqrt{(e^{2x} + e^{-2x})} \\
&= \ln(e^{2x} + e^{-2x})^{\frac{1}{2}} \Rightarrow f(x) = \frac{1}{2} \ln(e^{2x} + e^{-2x}) \quad \because \ln x^m = m \ln x
\end{aligned}$$

Now diff. w.r.t x

$$\frac{d}{dx} f(x) = \frac{1}{2} \frac{d}{dx} \ln(e^{2x} + e^{-2x})$$

Now do yourself

Question # 2(i)

$$\begin{aligned}
y &= x^2 \ln \sqrt{x} \\
\Rightarrow y &= x^2 \ln(x)^{\frac{1}{2}} \Rightarrow y = \frac{1}{2} x^2 \ln x \quad \because \ln x^m = m \ln x
\end{aligned}$$

Now diff. w.r.t x

$$\begin{aligned}
\frac{dy}{dx} &= \frac{1}{2} \frac{d}{dx} x^2 \ln x \\
&= \frac{1}{2} \left(x^2 \frac{d}{dx} \ln x + \ln x \frac{d}{dx} x^2 \right) \\
&= \frac{1}{2} \left(x^2 \cdot \frac{1}{x} + \ln x (2x) \right) = \frac{1}{2} x + x \ln x \quad \text{or} \quad \frac{1}{2} x + 2x \ln \sqrt{x} \quad \text{Ans.}
\end{aligned}$$

Question # 2(ii)

$$y = x \sqrt{\ln x}$$

Diff. w.r.t x

$$\begin{aligned}
\frac{dy}{dx} &= \frac{d}{dx} x (\ln x)^{\frac{1}{2}} \\
&= x \frac{d}{dx} (\ln x)^{\frac{1}{2}} + (\ln x)^{\frac{1}{2}} \frac{d}{dx} (x) \\
&= x \cdot \frac{1}{2} (\ln x)^{-\frac{1}{2}} \frac{d}{dx} (\ln x) + (\ln x)^{\frac{1}{2}} (1) = \frac{x}{2 (\ln x)^{\frac{1}{2}}} \left(\frac{1}{x} \right) + (\ln x)^{\frac{1}{2}} \\
&= \frac{1}{2 \sqrt{\ln x}} + \sqrt{\ln x} = \frac{1 + 2 \ln x}{2 \sqrt{\ln x}} \quad \text{Answer}
\end{aligned}$$

Question # 2(iii)

$$\begin{aligned}
 y &= \frac{x}{\ln x} \\
 \Rightarrow \frac{dy}{dx} &= \frac{d}{dx} \left(\frac{x}{\ln x} \right) \\
 &= \frac{\ln x \frac{dx}{dx} - x \frac{d}{dx} \ln x}{(\ln x)^2} = \frac{\ln x(1) - x \cdot \frac{1}{x}}{(\ln x)^2} = \frac{\ln x - 1}{(\ln x)^2} \quad \text{Answer}
 \end{aligned}$$

Question # 2(iv)

$$y = x^2 \ln \frac{1}{x} \quad \Rightarrow \quad y = x^2 \ln x^{-1} \quad \Rightarrow \quad y = -x^2 \ln x$$

Now do yourself.

Question # 2(v)

$$y = \ln \sqrt{\frac{x^2 - 1}{x^2 + 1}} \quad \Rightarrow \quad y = \ln \left(\frac{x^2 - 1}{x^2 + 1} \right)^{\frac{1}{2}} \quad \Rightarrow \quad y = \frac{1}{2} \ln \left(\frac{x^2 - 1}{x^2 + 1} \right)$$

Now diff. w.r.t x

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{1}{2} \frac{d}{dx} \ln \left(\frac{x^2 - 1}{x^2 + 1} \right) \\
 &= \frac{1}{2} \cdot \frac{1}{\left(\frac{x^2 - 1}{x^2 + 1} \right)} \cdot \frac{d}{dx} \left(\frac{x^2 - 1}{x^2 + 1} \right) \\
 &= \frac{x^2 + 1}{2(x^2 - 1)} \cdot \left(\frac{(x^2 + 1) \frac{d}{dx} (x^2 - 1) - (x^2 - 1) \frac{d}{dx} (x^2 + 1)}{(x^2 + 1)^2} \right) \\
 &= \frac{1}{2(x^2 - 1)} \cdot \left(\frac{(x^2 + 1)(2x) - (x^2 - 1)(2x)}{(x^2 + 1)} \right) \\
 &= \frac{1}{2(x^2 - 1)} \cdot \left(\frac{2x(x^2 + 1 - x^2 + 1)}{(x^2 + 1)} \right) = \frac{1}{(x^2 - 1)} \cdot \left(\frac{x(2)}{(x^2 + 1)} \right) = \frac{2x}{(x^4 - 1)} \quad \text{Ans.}
 \end{aligned}$$

Question # 2(vi)

$$y = \ln \left(x + \sqrt{x^2 + 1} \right)$$

Diff. w.r.t x

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{d}{dx} \ln \left(x + \sqrt{x^2 + 1} \right) \\
 &= \frac{1}{x + \sqrt{x^2 + 1}} \frac{d}{dx} \left(x + \sqrt{x^2 + 1} \right) = \frac{1}{x + \sqrt{x^2 + 1}} \left(1 + \frac{1}{2} (x^2 + 1)^{-\frac{1}{2}} \frac{d}{dx} (x^2 + 1) \right) \\
 &= \frac{1}{x + \sqrt{x^2 + 1}} \left(1 + \frac{1}{2(x^2 + 1)^{\frac{1}{2}}} \cdot (2x) \right) = \frac{1}{x + \sqrt{x^2 + 1}} \left(1 + \frac{x}{\sqrt{x^2 + 1}} \right) \\
 &= \frac{1}{x + \sqrt{x^2 + 1}} \left(\frac{\sqrt{x^2 + 1} + x}{\sqrt{x^2 + 1}} \right) = \frac{1}{\sqrt{x^2 + 1}} \quad \text{Answer}
 \end{aligned}$$

Question # 2(vii)

$$y = \ln(9 - x^2)$$

Diff. w.r.t x

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} \ln(9 - x^2) \\ &= \frac{1}{9 - x^2} \cdot \frac{d}{dx}(9 - x^2) = \frac{1}{9 - x^2} \cdot (-2x) \\ \Rightarrow \frac{dy}{dx} &= \frac{-2x}{9 - x^2}\end{aligned}$$

Question # 2(viii)

$$\begin{aligned}y &= e^{-2x} \sin 2x \\ \Rightarrow \frac{dy}{dx} &= \frac{d}{dx} e^{-2x} \sin 2x \\ &= e^{-2x} \frac{d}{dx} \sin 2x + \sin 2x \frac{d}{dx} e^{-2x} \\ &= e^{-2x} \cos 2x (2) + \sin 2x e^{-2x} (-2) = 2e^{-2x} (\cos 2x - \sin 2x) \quad \text{Answer}\end{aligned}$$

Question # 2(ix)

$$y = e^{-x} (x^3 + 2x^2 + 1)$$

Diff. w.r.t x

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} e^{-x} (x^3 + 2x^2 + 1) \\ &= e^{-x} \frac{d}{dx} (x^3 + 2x^2 + 1) + (x^3 + 2x^2 + 1) \frac{d}{dx} e^{-x} \\ &= e^{-x} (3x^2 + 4x + 0) + (x^3 + 2x^2 + 1) \cdot e^{-x} (-1) \\ &= e^{-x} (3x^2 + 4x) - (x^3 + 2x^2 + 1) \cdot e^{-x} = e^{-x} (3x^2 + 4x - x^3 - 2x^2 - 1) \\ &= e^{-x} (-x^3 + x^2 + 4x - 1) \quad \text{Answer}\end{aligned}$$

Question # 2(x)

$$y = xe^{\sin x}$$

Diff w.r.t x

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} xe^{\sin x} \\ &= x \frac{d}{dx} e^{\sin x} + e^{\sin x} \frac{d}{dx} x \\ &= x \cdot e^{\sin x} \frac{d}{dx} \sin x + e^{\sin x} (1) = x \cdot e^{\sin x} \cos x + e^{\sin x} \\ &= e^{\sin x} (x \cos x + 1) \quad \text{Answer}\end{aligned}$$

Question # 2(xi)*Do yourself***Question # 2(xii)**

$$y = (x+1)^x$$

Taking log on both sides

$$\ln y = \ln(x+1)^x \quad \Rightarrow \quad \ln y = x \ln(x+1)$$

Diff w.r.t x

$$\begin{aligned}
\frac{d}{dx} \ln y &= \frac{d}{dx} x \ln(x+1) \\
\Rightarrow \frac{1}{y} \frac{dy}{dx} &= x \frac{d}{dx} \ln(x+1) + \ln(x+1) \frac{dx}{dx} \\
&= x \cdot \frac{1}{x+1} \frac{d}{dx} (x+1) + \ln(x+1)(1) \\
\Rightarrow \frac{dy}{dx} &= y \left(\frac{x}{x+1} (1) + \ln(x+1) \right) \\
&= (x+1)^x \left(\frac{x}{x+1} + \ln(x+1) \right) \quad \text{Answer}
\end{aligned}$$

Question # 2(xiii)

$$y = (\ln x)^{\ln x}$$

Taking log on both sides

$$\ln y = \ln(\ln x)^{\ln x} \Rightarrow \ln y = (\ln x) \cdot \ln(\ln x)$$

Diff w.r.t x

$$\begin{aligned}
\frac{d}{dx} \ln y &= \frac{d}{dx} (\ln x) \cdot \ln(\ln x) \\
\Rightarrow \frac{1}{y} \frac{dy}{dx} &= (\ln x) \frac{d}{dx} \ln(\ln x) + \ln(\ln x) \frac{d}{dx} (\ln x) \\
&= (\ln x) \cdot \frac{1}{\ln x} \frac{d}{dx} (\ln x) + \ln(\ln x) \cdot \frac{1}{x} \\
&= \frac{1}{x} + \frac{\ln(\ln x)}{x} = \frac{1 + \ln(\ln x)}{x} \\
\Rightarrow \frac{dy}{dx} &= y \left(\frac{1 + \ln(\ln x)}{x} \right) \Rightarrow \frac{dy}{dx} = (\ln x)^{\ln x} \left(\frac{1 + \ln(\ln x)}{x} \right)
\end{aligned}$$

Question # 2(xiv)

$$\begin{aligned}
y &= \frac{\sqrt{x^2-1} (x+1)}{(x^3+1)^{3/2}} \Rightarrow y = \frac{((x+1)(x-1))^{\frac{1}{2}} (x+1)}{\left[(x+1)(x^2-x+1) \right]^{\frac{3}{2}}} \\
\Rightarrow y &= \frac{(x+1)^{\frac{1}{2}} (x-1)^{\frac{1}{2}} (x+1)}{(x+1)^{\frac{3}{2}} (x^2-x+1)^{\frac{3}{2}}} \Rightarrow y = \frac{(x+1)^{\frac{3}{2}} (x-1)^{\frac{1}{2}}}{(x+1)^{\frac{3}{2}} (x^2-x+1)^{\frac{3}{2}}} \\
\Rightarrow y &= \frac{(x-1)^{\frac{1}{2}}}{(x^2-x+1)^{\frac{3}{2}}}
\end{aligned}$$

Taking log on both sides

$$\begin{aligned}
\ln y &= \ln \frac{(x-1)^{\frac{1}{2}}}{(x^2-x+1)^{\frac{3}{2}}} \\
&= \ln(x-1)^{\frac{1}{2}} - \ln(x^2-x+1)^{\frac{3}{2}} \\
\Rightarrow \ln y &= \frac{1}{2} \ln(x-1) - \frac{3}{2} \ln(x^2-x+1)
\end{aligned}$$

Now diff. w.r.t x

$$\frac{d}{dx} \ln y = \frac{1}{2} \frac{d}{dx} \ln(x-1) - \frac{3}{2} \frac{d}{dx} \ln(x^2-x+1)$$

$$\begin{aligned}
\Rightarrow \frac{1}{y} \frac{dy}{dx} &= \frac{1}{2} \frac{1}{x-1} \frac{d}{dx}(x-1) - \frac{3}{2} \frac{1}{(x^2-x+1)} \frac{d}{dx}(x^2-x+1) \\
&= \frac{1}{2(x-1)}(1) - \frac{3}{2(x^2-x+1)}(2x-1) = \frac{1}{2(x-1)} - \frac{3(2x-1)}{2(x^2-x+1)} \\
\Rightarrow \frac{dy}{dx} &= y \left[\frac{x^2-x+1-3(2x-1)(x-1)}{2(x-1)(x^2-x+1)} \right] \\
&= \frac{(x-1)^{\frac{1}{2}}}{(x^2-x+1)^{\frac{3}{2}}} \cdot \left[\frac{x^2-x+1-3(2x^2-x-2x+1)}{2(x-1)(x^2-x+1)} \right] \\
&= \left[\frac{x^2-x+1-6x^2+3x+6x-3}{2(x-1)^{1-\frac{1}{2}}(x^2-x+1)^{\frac{3}{2}+1}} \right] = \frac{-5x^2+8x-2}{2(x-1)^{\frac{1}{2}}(x^2-x+1)^{\frac{5}{2}}} \\
\Rightarrow \frac{dy}{dx} &= \frac{5x^2-8x+2}{2\sqrt{x-1}(x^2-x+1)^{\frac{5}{2}}} \quad \text{Ans.}
\end{aligned}$$

Question # 2(xv)

$$\begin{aligned}
y &= \frac{(x+2)^2 \cdot \sqrt{x-1}}{\sqrt{x^2+x-2}} \\
\Rightarrow y &= \frac{(x+2)^2 \cdot \sqrt{x-1}}{\sqrt{x^2+2x-x-2}} \Rightarrow y = \frac{(x+2)^2 \cdot \sqrt{x-1}}{\sqrt{x(x+2)-1(x+2)}} \\
\Rightarrow y &= \frac{(x+2)^2 \cdot \sqrt{x-1}}{\sqrt{(x+2)(x-1)}} \Rightarrow y = (x+2)^{2-\frac{1}{2}} \Rightarrow y = (x+2)^{\frac{3}{2}}
\end{aligned}$$

Now diff. w.r.t x

$$\frac{dy}{dx} = \frac{d}{dx}(x+2)^{\frac{3}{2}}$$

Do yourself

2.1.3 Derivative of Hyperbolic Function (page 85)

The hyperbolic functions are define by

$$\sinh x = \frac{e^x - e^{-x}}{2}, \quad x \in R; \quad \cosh x = \frac{e^x + e^{-x}}{2}, \quad x \in R$$

$$\text{and} \quad \tanh x = \frac{\sinh x}{\cosh x} = \frac{e^x - e^{-x}}{e^x + e^{-x}}, \quad x \in R$$

The reciprocal of these functions are defined as;

$$\operatorname{csch} x = \frac{1}{\sinh x} = \frac{2}{e^x - e^{-x}}, \quad x \in R - \{0\}; \quad \operatorname{sech} x = \frac{1}{\cosh x} = \frac{2}{e^x + e^{-x}}, \quad x \in R$$

$$\text{and} \quad \operatorname{coth} x = \frac{1}{\tanh x} = \frac{e^x + e^{-x}}{e^x - e^{-x}}, \quad x \in R - \{0\}$$

and there derivatives are

$$(i) \quad \frac{d}{dx}(\sinh x) = \cosh x$$

$$(ii) \quad \frac{d}{dx}(\cosh x) = \sinh x$$

$$(iii) \quad \frac{d}{dx}(\tanh x) = \operatorname{sech}^2 x$$

$$(iv) \quad \frac{d}{dx}(\operatorname{coth} x) = -\operatorname{csch}^2 x$$

$$(v) \quad \frac{d}{dx}(\operatorname{sech} x) = -\operatorname{sech} x \tanh x$$

$$(vi) \quad \frac{d}{dx}(\operatorname{csch} x) = -\operatorname{csch} x \operatorname{coth} x$$

Proof:

$$\begin{aligned} \text{(i)} \quad \frac{d}{dx}(\sinh x) &= \frac{d}{dx} \left(\frac{e^x - e^{-x}}{2} \right) = \frac{d}{dx} \left(\frac{1}{2} (e^x - e^{-x}) \right) = \frac{1}{2} \frac{d}{dx} (e^x - e^{-x}) \\ &= \frac{1}{2} \left(\frac{d}{dx} e^x - \frac{d}{dx} e^{-x} \right) = \frac{1}{2} (e^x - e^{-x}(-1)) = \frac{1}{2} (e^x + e^{-x}) \\ &= \left(\frac{e^x + e^{-x}}{2} \right) = \cosh x \end{aligned}$$

(ii) *Similar as above.*

(iii) *See the below (iv) proof.*

$$\begin{aligned} \text{(iv)} \quad \frac{d}{dx} \coth x &= \frac{d}{dx} \left(\frac{e^x + e^{-x}}{e^x - e^{-x}} \right) \\ &= \frac{(e^x - e^{-x}) \frac{d}{dx} (e^x + e^{-x}) - (e^x + e^{-x}) \frac{d}{dx} (e^x - e^{-x})}{(e^x - e^{-x})^2} \\ &= \frac{(e^x - e^{-x})(e^x + e^{-x}(-1)) - (e^x + e^{-x})(e^x - e^{-x}(-1))}{(e^x - e^{-x})^2} \\ &= \frac{(e^x - e^{-x})(e^x - e^{-x}) - (e^x + e^{-x})(e^x + e^{-x})}{(e^x - e^{-x})^2} \\ &= \frac{(e^x - e^{-x})^2 - (e^x + e^{-x})^2}{(e^x - e^{-x})^2} \\ &= \frac{(e^{2x} + e^{-2x} - 2e^x e^{-x}) - (e^{2x} + e^{-2x} + 2e^x e^{-x})}{(e^x - e^{-x})^2} \\ &= \frac{e^{2x} + e^{-2x} - 2 - e^{2x} - e^{-2x} - 2}{(e^x - e^{-x})^2} \quad \because e^x e^{-x} = e^0 = 1 \\ &= \frac{-4}{(e^x - e^{-x})^2} = - \left(\frac{2}{e^x - e^{-x}} \right)^2 = -\operatorname{csch}^2 x \end{aligned}$$

$$\begin{aligned} \text{(v)} \quad \frac{d}{dx}(\operatorname{sech} x) &= \frac{d}{dx} \left(\frac{2}{e^x + e^{-x}} \right) = \frac{d}{dx} 2(e^x + e^{-x})^{-1} = 2 \frac{d}{dx} (e^x + e^{-x})^{-1} \\ &= 2 \left[(-1)(e^x + e^{-x})^{-1-1} \frac{d}{dx} (e^x + e^{-x}) \right] \\ &= -2(e^x + e^{-x})^{-2} (e^x + e^{-x}(-1)) = \frac{-2}{(e^x + e^{-x})^2} (e^x - e^{-x}) \\ &= \frac{-2(e^x - e^{-x})}{(e^x + e^{-x})(e^x + e^{-x})} = - \frac{2}{(e^x + e^{-x})} \frac{(e^x - e^{-x})}{(e^x + e^{-x})} \\ &= -\operatorname{sech} x \tanh x \end{aligned}$$

(vi) *Do yourself as above (v).*

2.14 Derivative of Inverse Hyperbolic Function (page 86)

$$\begin{array}{ll} \text{(i)} \quad \frac{d}{dx} \sinh^{-1} x = \frac{1}{\sqrt{1+x^2}} & \text{(ii)} \quad \frac{d}{dx} \cosh^{-1} x = \frac{1}{\sqrt{x^2-1}} \\ \text{(iii)} \quad \frac{d}{dx} \tanh^{-1} x = \frac{1}{1-x^2} & \text{(iv)} \quad \frac{d}{dx} \coth^{-1} x = \frac{1}{1-x^2} \\ \text{(v)} \quad \frac{d}{dx} \operatorname{sech}^{-1} x = \frac{-1}{x\sqrt{1-x^2}} & \text{(vi)} \quad \frac{d}{dx} \operatorname{csch}^{-1} x = \frac{-1}{x\sqrt{1+x^2}} \end{array}$$

Proof:

(i) Let $y = \sinh^{-1} x \Rightarrow \sinh y = x$

differentiate w.r.t. x .

$$\frac{d}{dx} \sinh y = \frac{d}{dx} x \Rightarrow \cosh y \frac{dy}{dx} = 1 \Rightarrow \frac{dy}{dx} = \frac{1}{\cosh y}$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{\sqrt{1+\sinh^2 y}} \quad \because \cosh^2 x - \sinh^2 x = 1$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{\sqrt{1+x^2}} \quad \because \sinh y = x$$

(ii) Do yourself as above.

(iii) Do yourself as (iv) below *or* see book at page 88.

(iv) Let $y = \coth^{-1} x \Rightarrow \coth y = x$

differentiate w.r.t. x

$$\frac{d}{dx} \coth y = \frac{d}{dx} x \Rightarrow -\operatorname{csch}^2 y \frac{dy}{dx} = 1 \Rightarrow \frac{dy}{dx} = \frac{1}{-\operatorname{csch}^2 y}$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{-(\coth^2 y - 1)} \quad \because \coth^2 y - 1 = \operatorname{csch}^2 y$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{-\coth^2 y + 1} = \frac{1}{1 - \coth^2 y}$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{1-x^2} \quad \because \coth y = x$$

(v) Suppose $y = \operatorname{sech}^{-1} x \Rightarrow \operatorname{sech} y = x$

differentiate w.r.t. x

$$\frac{d}{dx} \operatorname{sech} y = \frac{d}{dx} x \Rightarrow -\operatorname{sech} y \tanh y \frac{dy}{dx} = 1 \Rightarrow \frac{dy}{dx} = \frac{1}{-\operatorname{sech} y \tanh y}$$

$$\Rightarrow \frac{dy}{dx} = \frac{-1}{\operatorname{sech} y \sqrt{1 - \tanh^2 y}} \quad \because 1 - \tanh^2 y = \operatorname{sech}^2 y$$

$$\Rightarrow \frac{dy}{dx} = \frac{-1}{x\sqrt{1-x^2}} \quad \because \operatorname{sech} y = x$$

(vi) Do yourself as above

Question # 3(i)

$$y = \cosh 2x$$

Diff. w.r.t x

$$\frac{dy}{dx} = \frac{d}{dx} \cosh 2x \Rightarrow \frac{dy}{dx} = \sinh 2x \frac{d}{dx} (2x) \Rightarrow \frac{dy}{dx} = 2 \sinh 2x$$

Question # 3(ii)

Do yourself

Question # 3(iii)

$$y = \tanh^{-1}(\sin x) \Rightarrow \tanh y = \sin x$$

Diff. w.r.t x

$$\frac{d}{dx} \tanh y = \frac{d}{dx} (\sin x)$$

$$\Rightarrow \operatorname{sech}^2 y \frac{dy}{dx} = \cos x \Rightarrow \frac{dy}{dx} = \frac{\cos x}{\operatorname{sech}^2 y}$$

$$\Rightarrow \frac{dy}{dx} = \frac{\cos x}{1 - \tanh^2 y}$$

$$= \frac{\cos x}{1 - \sin^2 x}$$

$$= \frac{\cos x}{\cos^2 x} \Rightarrow \frac{dy}{dx} = \sec x$$

$$\because \cosh^2 \theta - \sinh^2 \theta = 1$$

$$\therefore 1 - \tanh^2 \theta = \operatorname{sech}^2 \theta$$

$$\because \sin x = \tanh y$$

Question # 3(iv)

$$y = \sinh^{-1}(x^3) \Rightarrow \sinh y = x^3$$

$$\Rightarrow \frac{d}{dx} \sinh y = \frac{d}{dx} x^3 \Rightarrow \cosh y \frac{dy}{dx} = 3x^2$$

$$\Rightarrow \frac{dy}{dx} = \frac{3x^2}{\cosh y}$$

$$= \frac{3x^2}{\sqrt{1 + \sinh^2 y}}$$

$$\because \cosh^2 y - \sinh^2 y = 1$$

$$= \frac{3x^2}{\sqrt{1 + (x^3)^2}} = \frac{3x^2}{\sqrt{1 + x^6}} \quad \text{Answer}$$

Question # 3(v)

Do yourself

Question # 3(vi)

$$y = \sinh^{-1}\left(\frac{x}{2}\right) \Rightarrow \sinh y = \frac{x}{2}$$

Now diff w.r.t x

$$\frac{d}{dx} \sinh y = \frac{d}{dx} \left(\frac{x}{2}\right) \Rightarrow \cosh y \frac{dy}{dx} = \frac{1}{2}$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{2 \cosh y}$$

$$\because \cosh^2 y - \sinh^2 y = 1$$

$$= \frac{1}{2 \sqrt{1 + \sinh^2 y}}$$

$$\therefore \cosh^2 y = 1 + \sinh^2 y$$

$$= \frac{1}{2 \sqrt{1 + (x/2)^2}} = \frac{1}{2 \sqrt{(4 + x^2)/2}} = \frac{1}{\sqrt{4 + x^2}} \quad \text{Answer.}$$