

Math 221 Notes

VOGEL, Max

UW-Madison, Summer 2020

Contents

1	Week 1	5
1.1	Day 1	5
1.1.1	Factoring:	5
1.1.2	Rules:	5
1.1.3	Functions:	5
1.1.3.1	Finding domain:	6
1.1.3.2	Picewise:	6
1.1.3.3	Types	6
1.1.3.4	Increasing & Decreasing:	7
1.1.4	Limits	7
1.1.4.1	Infinite Limits	7
1.1.5	Lines	8
1.2	Day 2	8
1.2.1	Limit Laws:	8
1.2.2	Piecewise / (ϵ, δ) definition of limit	9
1.3	Day 3	9
1.3.1	Continuous Function:	9
1.4	Day 4	10
1.4.1	Lines:	10
1.4.1.1	Secant Line:	10
1.4.1.2	Tangent Line:	11
1.4.2	Derivatives:	11

2	Week 2	11
2.1	Day 5	11
2.1.1	Derivatives (cont.):	11
2.2	Day 6	12
2.2.1	Derivatives Rules:	12
2.2.2	Lines:	12
2.3	Day 7	13
2.3.1	Trig Review:	13
2.3.2	Trig Identities:	13
2.3.3	Derivative of Trig Functions:	13
2.3.4	Limit of Trig Functions:	14
3	Week 3	14
3.1	Day 9	14
3.1.1	Composite Function:	14
3.1.2	Decomposite Function:	14
3.1.3	Chain Rule	14
3.2	Day 10	14
3.2.1	Explicit and Implicit:	14
3.3	Day 11	15
3.3.1	Related Rates:	15
3.4	Day 12	15
3.4.1	Maximum and Minimum Values:	15
3.4.2	Increasing/Decreasing Functions:	16
3.4.3	Concavity:	16
3.4.4	Extreme Value Theorem:	17
3.4.5	Fermat's Theorem:	17
3.4.6	Critical Number:	17
3.4.7	Absolute Extreme:	17
4	Week 3	17
4.1	Day 13	17
4.1.1	Rolle's Theorem	17
4.1.1.1	Proof	18
4.1.2	Mean Value Theorem	18
4.2	Day 14	19
4.2.1	Determining Increasing/Decreasing:	19
4.2.2	Local max and min:	20

4.2.3	Second Derivative Test	20
4.2.4	Finding Concavity	20
4.2.4.1	Inflection Point:	20
4.2.4.2	Theorem:	20
4.3	Day 15	21
4.3.1	Limit at ∞	21
4.3.2	Horizontal Asymptote	21
4.3.2.1	Theorem	21
5	Week 5	21
5.1	Day 17	21
5.1.1	Optimization	21
5.2	Day 18	22
5.2.1	Area Under Curve:	22
5.2.2	Absolute Extreme Values	22
5.2.3	Anti-derivative	22
5.2.3.1	Theorem	22
5.2.3.2	Initial Condition Steps	22
5.2.4	Differential Equations	23
5.3	Day 19	23
5.3.1	Definite Integral	23
5.3.1.1	Properties	23
5.3.1.2	Comparison Properties	23
5.4	Day 20	24
5.4.1	Fundamental Theorem of Calculus (1)	24
5.4.2	Fundamental Theorem of Calculus (2)	24
5.4.2.1	Notation:	24
5.4.3	Properties of Definite Integral	24
6	Week 6	24
6.1	Day 21	24
6.1.1	Indefinite Integral	24
6.1.1.1	Indefinite vs Definite Integrals	24
6.1.2	Integral Rules	25
6.1.3	Net Change Theorem	25
6.2	Day 22	25
6.2.1	Integration by Substitution	25
6.2.2	Substitution Rule for Definite Integral	26

6.3	Day 23	26
6.3.1	One-to-One Function	26
6.3.1.1	Horizontal Line Test	26
6.3.1.2	Remark	26
6.3.2	Inverse Function	27
6.3.2.1	Remark	27
6.3.2.2	Steps to find inverse	27
6.3.3	Derivative of Inverse Function	27

1 Week 1

1.1 Day 1

1.1.1 Factoring:

$$(a + b)(a^2 - ab + b^2) = a^3 + b^3$$

$$(a - b)(a^2 + ab + b^2) = a^3 - b^3$$

1.1.2 Rules:

$$\begin{aligned} a < b &\rightarrow -a > -b \\ -ax < b &\rightarrow x > -\frac{b}{a} \\ |x| = b &\rightarrow x = b \vee x = -b \\ |x| < b &\rightarrow -b < x < b \\ |x| = b &\rightarrow x > b \vee x < -b \\ a^m a^n &= a^{m+n} \\ \frac{a^m}{a^n} &= a^{m-n} \\ (ab)^m &= a^m b^m \end{aligned}$$

$$\begin{aligned} \left(\frac{a}{b}\right)^n &= \frac{a^n}{b^n} \\ (a^m)^n &= a^{mn} \\ \sqrt{a} &= a^{\frac{1}{2}} \\ \sqrt[n]{a} &= a^{\frac{1}{n}} \\ (\sqrt[n]{a})^m &= \sqrt[n]{a^m} = a^{\frac{m}{n}} \\ \sqrt{a}\sqrt{b} &= \sqrt{ab} \\ \sqrt[n]{a}\sqrt[n]{b} &= \sqrt[n]{ab} \end{aligned}$$

1.1.3 Functions:

Definition: Assigns each element x in a set D exactly one element, called $f(x)$, in set E .

In terms of a graph, a curve can only be a function if no vertical lines intersect the curve more than once (vertical line test).

- The set D is called the domain – possible x values.
- The set E is called the range – possible y values.
- If f is a function with domain D , then its graph is the set of ordered pairs $\{(x, f(x)) | x \text{ is a element of } D\}$

1.1.3.1 Finding domain:

1. $f(x) = \sqrt{x+2} \rightarrow x+2 \geq 0 \rightarrow x \geq -2 \rightarrow [-2, +\infty)$
2. $f(x) = \frac{1}{x^2-x} \rightarrow x^2 - x \neq 0 \rightarrow x(x-1) \neq 0 \rightarrow x \neq 0 \wedge x \neq 1 \rightarrow (-\infty, 0) \cup (0, 1) \cup (1, +\infty)$

1.1.3.2 Picewise:

- Open circles, \circ , and circle brackets, $()$, are non-inclusive.
- Closed circles, \bullet , and square brackets, $[]$, are inclusive.
- Formatted as:

$$f(x) = \begin{cases} y = 5 : x < 0 \\ y = x^2 : x \geq 0 \end{cases}$$

1.1.3.3 Types

Even:

- A function is even if $f(x) = f(-x)$
- The graph is symmetric with respect to the y-axis.
- Examples: $x^4 - 2, x^{20} + x^6, \cos(x), |x|$

Odd:

- A function is even if $f(x) = -f(x)$
- The graph has rotational symmetry about origin.
- Examples: $x^3, x^7 + x, \sin(x), |x|x$

-
- Even times odd function is always odd.
 - Even times even is always even.
 - Odd times odd is always even.

1.1.3.4 Increasing & Decreasing:

Increasing : A function f is called increasing on an interval if $f(x_1) < f(x_2)$ whenever $x_1 < x_2$. In other words, the slope must always be positive.

Decreasing : A function f is called decreasing on an interval if $f(x_1) > f(x_2)$ whenever $x_1 < x_2$. In other words, the slope must always be negative.

1.1.4 Limits

Definition: Supposing $f(x)$ is defined when x is near the number a , we write $\lim_{x \rightarrow a} f(x) = L$ and say “the limit of $f(x)$, as x approaches a , equals L ”.

- $\lim_{x \rightarrow a} f(x) = L$ if and only if (iff, \iff) $\lim_{x \rightarrow a^-} f(x) = L \wedge \lim_{x \rightarrow a^+} f(x) = L$
- That is, the limit does not exist, \nexists , if x approaches different values when from the left and right sides
- Approaching from left (from $-\infty \rightarrow \infty$) is notated as $x \rightarrow a^-$
- Approaching from right (from $\infty \rightarrow -\infty$) is notated as $x \rightarrow a^+$
- iff, \iff :

$$A \iff B \rightarrow$$

A is necessary and sufficient for $B \rightarrow$

B is necessary and sufficient for $A \rightarrow$

A is equivalent to B

- A vertical asymptote exists if the limit from left side is $+\infty$ or $-\infty$, and the limit from the right side is the opposite.

1.1.4.1 Infinite Limits

- If function limit is $\pm\infty$ (if the denominator is 0 at $f(a)$), you can find whether its $+$ or $-$ by solving the limit for each term. If the term is positive, then it's $+\infty$, and vice-versa, e.g.

$$\lim_{x \rightarrow -2^+} \frac{x-1}{x^2(x+2)}$$

$$\frac{\lim_{x \rightarrow -2^+} [x - 1]}{(\lim_{x \rightarrow -2^+} [x])^2 \cdot \lim_{x \rightarrow -2^+} [(x + 2)]}$$

$$\frac{\ominus}{\oplus \cdot \oplus}$$

$$\ominus \rightarrow -\infty$$

1.1.5 Lines

- Slope-Point form: $y - b = m(x + a) - (a, b)$ will be a point of the equation.
- Slope-Intercept form: $y = mx + b - (0, b)$ will be the y -intercept.
- Vertex form: $y = a(x - h) + k - (h, k)$ will be the vertex.
- Point-Point form: $y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1) - (x_1, y_1)$ and (x_2, y_2) will be points of the equation.
- Intercept form: $\frac{x}{a} + \frac{y}{b} = 1 - (a, b)$ will be a point of the equation.

1.2 Day 2

1.2.1 Limit Laws:

Supposing c is a constant and $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ exists, then

$\lim_{x \rightarrow a} [f(x) \pm g(x)] = \lim_{x \rightarrow a} f(x) \pm \lim_{x \rightarrow a} g(x)$ $\lim_{x \rightarrow a} [cf(x)] = c \cdot \lim_{x \rightarrow a} f(x)$ $\lim_{x \rightarrow a} [f(x)g(x)] = \lim_{x \rightarrow a} f(x) \lim_{x \rightarrow a} g(x)$ $\lim_{x \rightarrow a} \left[\frac{f(x)}{g(x)} \right] = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)} \text{ if } \lim_{x \rightarrow a} g(x) \neq 0$ $\lim_{x \rightarrow a} [f(x)]^n = [\lim_{x \rightarrow a} f(x)]^n$	$\lim_{x \rightarrow a} c = c$ $\lim_{x \rightarrow a} x = a$ $\lim_{x \rightarrow a} x^n = a^n \text{ where } n = \mathbb{Z}^+$ $\lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{a} \text{ where } n = \mathbb{Z}^+$ $\lim_{x \rightarrow a} \sqrt[n]{f(x)} = \sqrt[n]{f(a)} \text{ if } f(a) \geq 0 \text{ and } n_{\text{even}}$
--	---

Direct Substitution Property: If f is a polynomial or rational function, and a is in the domain of f , then $\lim_{x \rightarrow a} f(x) = f(a)$

Theorem 1.6.1: If $f(x) = g(x)$ when $x \neq a$, then $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x)$

Theorem 1.5.1: If $f(x) = L \iff \lim_{x \rightarrow a^-} f(x) = L = \lim_{x \rightarrow a^+} f(x)$


Theorem 1.6.2: If $f(x) \leq g(x)$ when x is near a and the limits of f and g both

exist as x approaches a , then $\lim_{x \rightarrow a} f(x) \leq \lim_{x \rightarrow a} g(x)$

Squeeze Theorem: If $f(x) \leq g(x) \leq h(x)$ when x is near a and $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L$, then $\lim_{x \rightarrow a} g(x) = L$

1.2.2 Piecewise / (ϵ, δ) definition of limit

If for every small number $\epsilon > 0$ there is a number $\delta > 0$ such that if $0 < |x - c| < \delta$ then $|f(x) - L| < \epsilon$.



Solving for δ , rewrite the term defining ϵ to be equal to the term defining δ . E.g., solving for δ if $|x - 2| < \delta$, then $|4x - 8| < \epsilon$, where $\epsilon = 0.1$:

$$|4x - 8| = 4|x - 2| < 0.1$$

$$|x - 2| < \frac{0.1}{4}, \text{ therefore } \delta = \frac{0.1}{4}$$

For non-linear equations, find the lesser and greater δ , and choose the one that results in the smaller ϵ .

1.3 Day 3

1.3.1 Continuous Function:

Definition: A function f is continuous at a number if $\lim_{x \rightarrow a} f(x) = f(a)$. Graphically, a function is continuous if you can draw it without having your pen leave

paper. More formally, $f(x)$ is continuous at $x = a \iff$:

1. $f(a)$ is defined ($a \in D : a$ is in the domain of f).
2. $\lim_{x \rightarrow a} f(x)$ exists, and equals $f(x) = f(a)$.


If one of the aforementioned statements is incorrect, then $f(x)$ is discontinuous at $x = a$

Theorem 1: A function is continuous on an interval if it's continuous at every number in the interval. That is, if f and g are continuous at $x = a$, then the following are also continuous at a :

$$f + g, f - g, cf, fg, \frac{f}{g} \text{ for } g(a) \neq 0$$

Theorem 2: The following types of functions are continuous at every number in their domain: Polynomials, Rational functions, Root functions, & Trig functions.

Theorem 3: Intermediate Value Theorem (IVM): Suppose that f is continuous on the closed interval $[a, b]$ and let N be any number between $f(a)$ and $f(b)$, where $f(a) \neq f(b)$. Then there exists at least one number c in (a, b) such that $f(c) = N$.




1.4 Day 4

1.4.1 Lines:

1.4.1.1 Secant Line: A line that locally intersects two points on a curve.

$$\frac{\text{Rise}}{\text{Run}} = \frac{\Delta y}{\Delta x} = \frac{y_1 - y_0}{x_1 - x_0} = \frac{f(a+h) - f(a)}{(a+h) - a} = \frac{f(a+h) - f(a)}{h}$$



1.4.1.2 Tangent Line: The line through a pair of infinitely close points on the curve so that the line is “just touching”. Slope equation (also known as “Difference Quotient”):

$$\lim_{h \rightarrow 0} \left[\frac{f(a+h) - f(a)}{h} \right]$$

1.4.2 Derivatives:

The derivative of a function f at a number a , denoted by $f'(a)$, is

$$f'(a) = \lim_{h \rightarrow 0} \left[\frac{f(a+h) - f(a)}{h} \right]$$

and the equation of the tangent line to the curve $y = f(x)$ at the point $(a, f(a))$ can be written in point-slope form as

$$y - f(a) = f'(a)(x - a)$$

2 Week 2

2.1 Day 5

2.1.1 Derivatives (cont.):

- Other notation is $\frac{dy}{dx}|_{x=a}$ (Leibniz Notation), $\frac{df}{dx}$, $\frac{d}{dx} f(x)$, $f'(x)$, $Df(x)$, & $D_x f(x)$

- Function $f(x)$ is differentiable at $x = a$ if $f'(a)$ exists (same as Theorem 1.5.1).
- Therefore, Not Continuous \implies Not Differentiable.
- If $f'(a)$ exists, then $\lim_{x \rightarrow a} f(x) = f(a)$
- The derivative is a function, not a constant.
- Because f' is also a function, f' may have a derivative of its own, denoted by $(f')' = f''$ and called the **second derivative** of f . This can also be written as $\frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d^2y}{dx^2}$

2.2 Day 6


2.2.1 Derivatives Rules:

- | | |
|--|---|
| <ul style="list-style-type: none"> • Constant: $\frac{d}{dx}(c) = 0$ • Linear: $\frac{d}{dx}(x) = 1$ | <ul style="list-style-type: none"> • Linear + Constant: $\frac{d}{dx}(ax) = a$ • Power Rule: $\frac{d}{dx}(x^n) = nx^{n-1}$ |
|--|---|
-
- Constant Multiple: $\frac{d}{dx}[c \cdot f(x)] = c \cdot \frac{d}{dx}(f(x))$
 - Sum/Difference: $\frac{d}{dx}[f(x) \pm g(x)] = f' \pm g'$
 - Product Rule: $\frac{d}{dx}[f(x) \cdot g(x)] = f' \cdot g + f \cdot g'$
 - Quotient Rule: $\frac{d}{dx} \left[\frac{f(x)}{g(x)} \right] = \frac{g \cdot f' - f \cdot g'}{g^2} = \frac{\text{lo} \cdot \text{d hi} - \text{hi} \cdot \text{d lo}}{\text{lo} \cdot \text{lo}}$

2.2.2 Lines:

Theorem: If the graph of $y = m_1x + b_1$ is perpendicular to the graph of $y = m_2x + b_2$, then $m_1m_2 = -1$.

Normal Line: The normal line to a curve at point M is the line through M that is perpendicular to the tangent line at M .



2.3 Day 7

2.3.1 Trig Review:

$$\csc = \frac{1}{\sin}, \quad \sec = \frac{1}{\cos}, \quad \cot = \frac{1}{\tan} = \frac{\cos}{\sin}$$

2.3.2 Trig Identities:

$$\sin^2 + \cos^2 = 1$$

$$\text{Dividing by } \sin^2 : 1 + \frac{\cos^2}{\sin^2} = \frac{1}{\sin^2} \rightarrow 1 + \cot^2 = \csc^2$$

$$\text{Dividing by } \cos^2 : \frac{\sin^2}{\cos^2} + 1 = \frac{1}{\cos^2} \rightarrow \tan^2 + 1 = \sec^2$$

2.3.3 Derivative of Trig Functions:

$$(\sin)' = \cos$$

$$(\cos)' = -\sin$$

$$(\tan)' = \sec^2$$

$$(\csc)' = -\cot \cdot \csc$$

$$(\sec)' = \sec \cdot \tan$$

$$(\cot)' = -\csc^2$$

2.3.4 Limit of Trig Functions:

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$$

3 Week 3

3.1 Day 9

3.1.1 Composite Function:

A new function can be composed to two old functions, such as $f \circ g$. Formally, given two functions f and g , the composite function $f \circ g$ and $g \circ f$ (also called the composition of f and g) is defined by $(f \circ g)(x) = f(g(x))$ and $(g \circ f)(x) = g(f(x))$

3.1.2 Decomposite Function:

Many functions can be decomposed into small functions, $h(x) = f(g(x))$

3.1.3 Chain Rule

If the composite function $F(x) = f \cdot g$ is defined by $F(x) = f(g(x))$ is differentiable at x and F' is given by the product

$$F'(x) = f'(g(x)) \cdot g'(x) \text{ or } OUT'(in) \cdot IN'$$

The power rule can be combined with the chain rule as

$$\frac{d}{dx}[g(x)]^n = n[g(x)]^{n-1} \cdot g'(x)$$

3.2 Day 10

3.2.1 Explicit and Implicit:

Explicit: A function given in terms of the independent variable, e.g. $y = \sqrt{x^3 + 1}$.

Implicit: A function given in terms of both dependent and independent variables, e.g. $x^3 - y^2 + 1 = 0$, $y \geq 0$.

Implicit Differentiation: If the function is stuck in implicit form, then differentiate both sides of the equation with respect to x and solve the resulting equation

for y' .

Note:

$$\frac{d}{dx}(y^n) \neq ny^{n-1} \quad \frac{d}{dx}(y^n) = ny^{n-1} \cdot y'$$

3.3 Day 11

3.3.1 Related Rates:

A related rates problem is to compute the rate of change (derivative) of one quantity in terms of the rate of change of another quantity, which is more easily measured.

How to approach a related rates problem:

1. Read the problem, and draw a diagram if possible.
2. Assign symbols to all variables that are a function of time.
3. Find an equation that relates the two variables.
4. Using the Chain Rule to differentiate both sides with respect to time.
5. Substitute the given information into resulting equation and solve for the unknown rate.

3.4 Day 12

3.4.1 Maximum and Minimum Values:

Absolute or Global [Max/Min]: The [max/min] y -value in the domain.

Relative or Local [Max/Min]: The [max/min] y -value in a given range around a x -value. Can only occur when the graph goes from [increasing to decreasing / vice-versa]

Note: The endpoint of a graph can never be a relative [max/min] because one side of the point is always undefined.


3.4.2 Increasing/Decreasing Functions:

Increasing and Decreasing Functions


1. If $f'(x) > 0$ for every x on some interval I , then $f(x)$ is increasing on the interval.

2. If $f'(x) < 0$ for every x on some interval I , then $f(x)$ is decreasing on the interval.

3. If $f'(x) = 0$ for every x on some interval I , then $f(x)$ is constant on the interval.



3.4.3 Concavity:



3.4.4 Extreme Value Theorem:

Suppose that $f(x)$ is continuous on the interval $[a, b]$ then there are two numbers, $a \leq c, d \leq b$ so that $f(c)$ is an absolute maximum for the function and $f(d)$ is an absolute minimum for the function.

3.4.5 Fermat's Theorem:

If f has a local max or min at a , and $f'(a)$ exists, then $f'(a) = 0$.

3.4.6 Critical Number:

A critical number of a function f is a number c in the domain of f such that either $f'(c) = 0$ or $f'(c) = \text{DNE}$

3.4.7 Absolute Extreme:

To find the absolute max and min values of a continuous function f on a closed interval $[a, b]$:

1. Find the critical numbers of the function on (a, b)
2. Find the values of f at the critical numbers of f in (a, b)
3. Find the values of f at the end points of the interval, in other words, find $f(a)$ and $f(b)$.
4. The largest of the values from steps 2 and 3 is the absolute max, the smallest is the absolute min.

4 Week 3

4.1 Day 13


4.1.1 Rolle's Theorem

Let f be a function that satisfies the following three hypotheses:

1. f is continuous on the closed interval $[a, b]$
2. f is differentiable on the open interval (a, b)

3. $f(a) = f(b)$

Then there is at least one number c in (a, b) such that $f'(c) = 0$



4.1.1.1 Proof

- Case 1: $f(x) = k$, a constant. Then, $f'(x) = 0$ so the number c can be taken to be any number in (a, b) .
- Case 2: $f(x) > f(a)$, concave down, for some x in (a, b) . By the extreme value theorem, f has a max value in $[a, b]$. Since $f(a) = f(b)$, it must attain its max value at c ; $c \in (a, b)$. Then, f has a local max at c , f is differentiable at c , therefore $f'(c) = 0$ by Fermat's theorem.
- Case 3: $f(x) < f(a)$, concave up, for some x in (a, b) . By the extreme value theorem, f has a min value in $[a, b]$. Since $f(a) = f(b)$, it must attain its min value at c ; $c \in (a, b)$. Then, f has a local min at c , f is differentiable at c , therefore $f'(c) = 0$ by Fermat's theorem.

4.1.2 Mean Value Theorem

Let f be a function that satisfies the following hypotheses:


- f is continuous on the closed interval $[a, b]$
- f is differentiable on the open interval (a, b)

Then there is at least one number c in (a, b) such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

Slope of tangent line at c = Slope of Secant Line

$$f(b) - f(a) = f'(c)(b - a)$$



4.2 Day 14

4.2.1 Determining Increasing/Decreasing:

1. Find the critical numbers
2. Mark the critical numbers on a real line
3. Pick a test number, c , in each interval:
 - If $f'(c) > 0$, f is increasing at that interval
 - If $f'(c) < 0$, f is decreasing at that interval

4.2.2 Local max and min:

1. Find the critical numbers
2. Mark the critical numbers on a real line
3. Pick a test number, c , in each interval:
 - If f' changes from \oplus to \ominus , then f has a local max at $f(c)$
 - If f' changes from \ominus to \oplus , then f has a local min at $f(c)$
 - If f' doesn't change sign, then f has no local min or max at $f(c)$

4.2.3 Second Derivative Test

Suppose f'' is continuous near c .

1. If $f'(c) = 0$ and $f''(c) < 0$, then f has a local max at c .
2. If $f'(c) = 0$ and $f''(c) > 0$, then f has a local min at c .
3. If $f'(c) = 0$ and $f''(c) = 0$, then the second derivative test says nothing about the point c , a possible inflection point

4.2.4 Finding Concavity

1. $f(x)$ increasing implies $f'(x) > 0$
2. $f'(x)$ increasing implies $f''(x) > 0$
3. $f''(x) > 0$ for all x in I , then the graph of f is concave upwards on I
4. $f''(x) < 0$ for all x in I , then the graph of f is concave downwards on I

4.2.4.1 Inflection Point:

A point p on a curve $f(x)$ is called an inflection point if f is continuous there and the curve changes at p from concave up to concave down or vice versa.

4.2.4.2 Theorem:

At a point of inflection, $(c, f(c))$, either $f''(c) = 0$ or $f''(c)$ DNE

4.3 Day 15

4.3.1 Limit at ∞

Let f be a function defined on some interval $(a, +\infty)$. Then $\lim_{x \rightarrow \infty} f(x) = L$ means that the value of $f(x)$ can be made arbitrarily close to L by making x sufficiently large. In other words, for every $\epsilon > 0$, there is a number N such that if $x < N$ then $|f(x) - L| < \epsilon$. "N": Large negative number.

4.3.2 Horizontal Asymptote

The line $y = L$ is called a horizontal asymptote of the curve $y = f(x)$ if either $\lim_{x \rightarrow +\infty} f(x) = L$ or $\lim_{x \rightarrow -\infty} f(x) = L$

4.3.2.1 Theorem

If $r > 0$ is a rational number, then $\lim_{x \rightarrow \infty} \frac{1}{x^r} = 0$

If $r > 0$ is a rational number such that x^r is defined for all x , then $\lim_{x \rightarrow -\infty} \frac{1}{x^r} = 0$

5 Week 5

5.1 Day 17

5.1.1 Optimization

1. Understand the problem. What is the unknown? What are the given quantities? What are the given conditions?
2. Draw a diagram. Show variables.
3. Introduce notation. Assign Q to the quantity that is to be maximized or minimized.
4. Express Q in terms of other variables.
5. If Q is expressed with more than one variable, use the given info to find a relationship among these variables.
6. Find the absolute max or min value of the Q (the function).

5.2 Day 18

5.2.1 Area Under Curve:

The area A of the region S that lies under the graph of the continuous function f is the limit of the sum of the areas of approximating rectangles:

$$A = \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} (f(x_1)\Delta x + f(x_2)\Delta x + \dots + f(x_n)\Delta x) = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i)\Delta x$$

$$A = \lim_{n \rightarrow \infty} r_n = \lim_{n \rightarrow \infty} (f(x_0)\Delta x + f(x_1)\Delta x + \dots + f(x_{n-1})\Delta x) = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_{i-1})\Delta x$$

5.2.2 Absolute Extreme Values

Suppose that c is a critical number of a continuous function f defined on an interval.

- If $f'(x) > 0$ for all $x < c$, and $f'(x) < 0$ for all $x > c$, then $f(c)$ is the absolute max for values of f .
- If $f'(x) < 0$ for all $x < c$, and $f'(x) > 0$ for all $x > c$, then $f(c)$ is the absolute min for values of f .

5.2.3 Anti-derivative

A function F is called an antiderivative of f on an interval I if $F'(x) = f(x)$ for all x in I .

For example, $F(x) = x^3$ and $f(x) = 3x^2$:

- $f(x)$ is derivative of function $F(x)$
- $F(x)$ is antiderivative of function $f(x)$

5.2.3.1 Theorem If F is an antiderivative of f on an interval I , then the most general antiderivative of f on I is $F(x) + C$ where C is an arbitrary constant.

5.2.3.2 Initial Condition Steps

1. Find the anti-derivative of $f(x)$
2. Plug in the initial condition to evaluate constant C
3. Find the solution

5.2.4 Differential Equations

An equation that involves the derivatives of a function is called a differential equation. The solution of a differential equation is a function that satisfies the equation.

5.3 Day 19

5.3.1 Definite Integral

If f is continuous on $[a, b]$, or if f has only a finite number of jump discontinuities, then f is integrable on $[a, b]$; that is, the definite integral exists.

$$\int_a^b f(x)dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*)\Delta x$$

5.3.1.1 Properties

$$\int_a^a f(x)dx = 0$$

$$\int_a^b f(x)dx = - \int_b^a f(x)dx$$

$$\int_a^b c \cdot dx = c(b - a)$$

$$\int_a^b [f(x) \pm g(x)] dx = \int_a^b f(x)dx \pm \int_a^b g(x)dx$$

$$\int_a^c f(x)dx = \int_a^b f(x)dx + \int_b^c f(x)dx \text{ where } a < b < c$$

5.3.1.2 Comparison Properties

$$f(x) > 0 \text{ for } a < x < b, \text{ then } \int_a^b f(x)dx > 0$$

$$f(x) > g(x) \text{ for } a < x < b, \text{ then } \int_a^b f(x)dx > \int_a^b g(x)dx$$

$$m < f(x) < M \text{ for } a < x < b, \text{ then } m(b - a) < \int_a^b f(x)dx < M(b - a)$$

5.4 Day 20

5.4.1 Fundamental Theorem of Calculus (1)

If $f(x)$ is continuous on $[a, b]$, then the function g defined by $g(x) = \int_a^x f(t)dt$ for $a \leq x \leq b$, is continuous on $[a, b]$ and differentiable on (a, b) , and $g'(x) = f(x)$. In other words,

$$\frac{d}{dx} \left[\int_a^x f(t)dt \right] = f(x)$$

5.4.2 Fundamental Theorem of Calculus (2)

If $f(x)$ is continuous on $[a, b]$, then $\int_a^b f(x)dx = F(b) - F(a)$. where $F(x)$ is any antiderivative of f , that is, a function such that $F' = f$.

5.4.2.1 Notation:

$$F(b) - F(a) = F(x) \Big|_a^b = F(x) \Big]_a^b$$

5.4.3 Properties of Definite Integral

$$\int_{-a}^a f(x)dx = 0 \text{ if } f(x) \text{ is odd}$$

$$\int_{-a}^a f(x)dx = 2 \cdot \int_0^a f(x)dx \text{ if } f(x) \text{ is even}$$

6 Week 6

6.1 Day 21

6.1.1 Indefinite Integral

If $F(x)$ is any anti-derivative of $f(x)$, then the most general anti-derivative of $f(x)$ is called an indefinite integral and denoted as

$$\int f(x)dx = F(x) + C$$

6.1.1.1 Indefinite vs Definite Integrals Definite is a number, where as indefinite is a family of functions

6.1.2 Integral Rules

$$\int cf(x) dx = c \int f(x) dx$$

$$\int [f(x) + g(x)] dx = \int f(x) dx + \int g(x) dx$$

$$\int k dx = kx + C$$

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \quad (n \neq -1)$$

$$\int \frac{1}{x} dx = \ln|x| + C$$

$$\int e^x dx = e^x + C$$

$$\int a^x dx = \frac{a^x}{\ln a} + C$$

$$\int \sin x dx = -\cos x + C$$

$$\int \cos x dx = \sin x + C$$

$$\int \sec^2 x dx = \tan x + C$$

$$\int \csc^2 x dx = -\cot x + C$$

$$\int \sec x \tan x dx = \sec x + C$$

$$\int \csc x \cot x dx = -\csc x + C$$

$$\int \frac{1}{x^2 + 1} dx = \tan^{-1}x + C$$

$$\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1}x + C$$

$$\int \sinh x dx = \cosh x + C$$

$$\int \cosh x dx = \sinh x + C$$

6.1.3 Net Change Theorem

The integral of a rate of change is the net change, $\int_a^b F'(x)dx = F(b) - F(a)$

6.2 Day 22

6.2.1 Integration by Substitution

A function F is called an anti-derivative of f on an interval I if $F'(x) = f(x)$ for all x in I .

1. Let $u = g(x)$, where $g(x)$ is part of the integrand.
2. Calculate $\frac{du}{dx}$ (or $g'(x)$)
3. Use the substitution $u = g(x)$ and $du = g'(x)dx$ to convert the entire integral into one involving only u
4. If indefinite, evaluate the resulting integral and plug in u by $g(x)$.

5. If definite, plug in the lower limit of the integral and the upper limit of the integral into the expression $u = g(x)$ and get the new lower and upper limits. Then, evaluate the new definite integral (involving only u) by finding the anti-derivative of f and use the Fundamental Theorem.

6.2.2 Substitution Rule for Definite Integral


If g' is continuous on $[a, b]$ and f is continuous on the range of $u = g(x)$, then

$$\int_a^b f(g(x)) \cdot g'(x) dx = \int_{g(a)}^{g(b)} f(u) du$$

6.3 Day 23

6.3.1 One-to-One Function

A function f that never takes one a same value twice; that is $f(x_1) \neq f(x_2)$ whenever $x_1 \neq x_2$.



6.3.1.1 Horizontal Line Test If you draw a horizontal line through the graph, it should only intersect once if it's a 1-to-1 function.

6.3.1.2 Remark All strictly increasing/decreasing functions are 1-to-1

- $f'(a) > 0$ means strictly increasing
- $f'(a) < 0$ means strictly decreasing

6.3.2 Inverse Function

Let function f be a 1-to-1 function with domain A and range B . Then its inverse function f^{-1} has domain B and range A and is defined by $f^{-1}(y) = x \iff f(x) = y$ for any y in B .

6.3.2.1 Remark

- The domain of f^{-1} is range of f
- The range of f^{-1} is the domain of f
- $f^{-1}(f(x)) = x$
- $f(f^{-1}(x)) = x$

6.3.2.2 Steps to find inverse

1. Write $y = f(x)$
2. Solve this equation for x in terms of y
3. To express f^{-1} as a function of x , interchange x and y .

6.3.3 Derivative of Inverse Function

If f is a 1-to-1 differentiable function with inverse function f^{-1} , then the inverse function is differentiable and the following is true:

$$(f^{-1})'(x) = \frac{1}{f'(f^{-1}(x))}$$