

Math 221 Notes

VOGEL, Max

UW-Madison, Summer 2020

Contents

1	Week 1	2
1.1	Day 1	2
1.1.1	Factoring:	2
1.1.2	Rules:	2
1.1.3	Functions:	2
1.1.3.1	Finding domain:	3
1.1.3.2	Picewise:	3
1.1.3.3	Types	3
1.1.3.4	Increasing & Decreasing:	4
1.1.4	Limits	4
1.1.4.1	Infinite Limits	4
1.1.5	Lines	5
1.2	Day 2	5
1.2.1	Limit Laws:	5
1.2.2	Piecewise / (ϵ, δ) definition of limit	6
1.3	Day 3	6
1.3.1	Continuous Function:	6
1.4	Day 4	7
1.4.1	Line:	7
1.4.1.1	Secant Line:	7
1.4.1.2	Tangent Line:	8
1.4.1.3	Derivatives:	8

1 Week 1

1.1 Day 1

1.1.1 Factoring:

$$(a + b)(a^2 - ab + b^2) = a^3 + b^3$$

$$(a - b)(a^2 + ab + b^2) = a^3 - b^3$$

1.1.2 Rules:

$$\begin{aligned} a < b &\rightarrow -a > -b \\ -ax < b &\rightarrow x > -\frac{b}{a} \\ |x| = b &\rightarrow x = b \vee x = -b \\ |x| < b &\rightarrow -b < x < b \\ |x| = b &\rightarrow x > b \vee x < -b \\ a^m a^n &= a^{m+n} \\ \frac{a^m}{a^n} &= a^{m-n} \\ (ab)^m &= a^m b^m \end{aligned}$$

$$\begin{aligned} \left(\frac{a}{b}\right)^n &= \frac{a^n}{b^n} \\ (a^m)^n &= a^{mn} \\ \sqrt{a} &= a^{\frac{1}{2}} \\ {}^n\sqrt{a} &= a^{\frac{1}{n}} \\ ({}^n\sqrt{a})^m &= {}^n\sqrt{a^m} = a^{\frac{m}{n}} \\ \sqrt{a}\sqrt{b} &= \sqrt{ab} \\ {}^n\sqrt{a} {}^n\sqrt{b} &= {}^n\sqrt{ab} \end{aligned}$$

1.1.3 Functions:

Definition: Assigns each element x in a set D exactly one element, called $f(x)$, in set E .

In terms of a graph, a curve can only be a function if no vertical lines intersect the curve more than once (vertical line test).

- The set D is called the domain – possible x values.
- The set E is called the range – possible y values.
- If f is a function with domain D , then its graph is the set of ordered pairs $\{(x, f(x)) | x \text{ is a element of } D\}$

1.1.3.1 Finding domain:

1. $f(x) = \sqrt{x+2} \rightarrow x+2 \geq 0 \rightarrow x \geq -2 \rightarrow [-2, +\infty)$
2. $f(x) = \frac{1}{x^2-x} \rightarrow x^2 - x \neq 0 \rightarrow x(x-1) \neq 0 \rightarrow x \neq 0 \wedge x \neq 1 \rightarrow (-\infty, 0) \cup (0, 1) \cup (1, +\infty)$

1.1.3.2 Picewise:

- Open circles, \circ , and circle brackets, $()$, are non-inclusive.
- Closed circles, \bullet , and square brackets, $[]$, are inclusive.
- Formatted as:

$$f(x) = \begin{cases} y = 5 & x < 0 \\ y = x^2 & x \geq 0 \end{cases}$$

1.1.3.3 Types

Even:

- A function is even if $f(x) = f(-x)$
- The graph is symmetric with respect to the y-axis.
- Examples: $x^4 - 2, x^{20} + x^6, \cos(x), |x|$

Odd:

- A function is even if $f(x) = -f(x)$
- The graph has rotational symmetry about origin.
- Examples: $x^3, x^7 + x, \sin(x), |x|x$

-
- Even times odd function is always odd.
 - Even times even is always even.
 - Odd times odd is always even.

1.1.3.4 Increasing & Decreasing:

Increasing : A function f is called increasing on an interval if $f(x_1) < f(x_2)$ whenever $x_1 < x_2$. In other words, the slope must always be positive.

Decreasing : A function f is called decreasing on an interval if $f(x_1) > f(x_2)$ whenever $x_1 < x_2$. In other words, the slope must always be negative.

1.1.4 Limits

Definition: Supposing $f(x)$ is defined when x is near the number a , we write $\lim_{x \rightarrow a} f(x) = L$ and say "the limit of $f(x)$, as x approaches a , equals L ".

- $\lim_{x \rightarrow a} f(x) = L$ if and only if (iff, \iff) $\lim_{x \rightarrow a^-} f(x) = L \wedge \lim_{x \rightarrow a^+} f(x) = L$
- That is, the limit does not exist, \nexists , if x approaches different values when from the left and right sides
- Approaching from left (from $-\infty \rightarrow \infty$) is notated as $x \rightarrow a^-$
- Approaching from right (from $\infty \rightarrow -\infty$) is notated as $x \rightarrow a^+$
- iff, \iff :

$$A \iff B \rightarrow$$

A is necessary and sufficient for $B \rightarrow$

B is necessary and sufficient for $A \rightarrow$

A is equivalent to B

- A vertical asymptote exists if the limit from left side is $+\infty$ or $-\infty$, and the limit from the right side is the opposite.

1.1.4.1 Infinite Limits

- If function limit is $\pm\infty$ (if the denominator is 0 at $f(a)$), you can find whether its $+$ or $-$ by solving the limit for each term. If the term is positive, then it's $+\infty$, and vice-versa, e.g.

$$\lim_{x \rightarrow -2^+} \frac{x-1}{x^2(x+2)}$$

$$\frac{\lim_{x \rightarrow -2^+} [x - 1]}{\lim_{x \rightarrow -2^+} [x^2] \cdot \lim_{x \rightarrow -2^+} [(x + 2)]}$$

$$\frac{\ominus}{\oplus \cdot \oplus}$$

$$\ominus \rightarrow -\infty$$

1.1.5 Lines

- Slope-Point form: $y - b = m(x + a) - (a, b)$ will be a point of the equation.
- Slope-Intercept form: $y = mx + b - (0, b)$ will be the y -intercept.
- Vertex form: $y = a(x - h) + k - (h, k)$ will be the vertex.
- Point-Point form: $y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1) - (x_1, y_1)$ and (x_2, y_2) will be points of the equation.
- Intercept form: $\frac{x}{a} + \frac{y}{b} = 1 - (a, b)$ will be a point of the equation.

1.2 Day 2

1.2.1 Limit Laws:

Supposing c is a constant and $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ exists, then

$\lim_{x \rightarrow a} [f(x) \pm g(x)] = \lim_{x \rightarrow a} f(x) \pm \lim_{x \rightarrow a} g(x)$ $\lim_{x \rightarrow a} [cf(x)] = c \cdot \lim_{x \rightarrow a} f(x)$ $\lim_{x \rightarrow a} [f(x)g(x)] = \lim_{x \rightarrow a} f(x) \lim_{x \rightarrow a} g(x)$ $\lim_{x \rightarrow a} \left[\frac{f(x)}{g(x)} \right] = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)} \text{ if } \lim_{x \rightarrow a} g(x) \neq 0$ $\lim_{x \rightarrow a} [f(x)]^n = [\lim_{x \rightarrow a} f(x)]^n$	$\lim_{x \rightarrow a} c = c$ $\lim_{x \rightarrow a} x = a$ $\lim_{x \rightarrow a} x^n = a^n \text{ where } n = \mathbb{Z}^+$ $\lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{a} \text{ where } n = \mathbb{Z}^+$ $\lim_{x \rightarrow a} \sqrt[n]{f(x)} = \sqrt[n]{f(a)} \text{ if } f(a) \geq 0 \text{ and } n_{\text{even}}$
--	---

Direct Substitution Property: If f is a polynomial or rational function, and a is in the domain of f , then $\lim_{x \rightarrow a} f(x) = f(a)$

Theorem 1.6.1: If $f(x) = g(x)$ when $x \neq a$, then $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x)$

Theorem 1.5.1: If $f(x) = L \iff \lim_{x \rightarrow a^-} f(x) = L = \lim_{x \rightarrow a^+} f(x)$


Theorem 1.6.2: If $f(x) \leq g(x)$ when x is near a and the limits of f and g both

exist as x approaches a , then $\lim_{x \rightarrow a} f(x) \leq \lim_{x \rightarrow a} g(x)$

Squeeze Theorem: If $f(x) \leq g(x) \leq h(x)$ when x is near a and $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L$, then $\lim_{x \rightarrow a} g(x) = L$

1.2.2 Piecewise / (ϵ, δ) definition of limit

If for every small number $\epsilon > 0$ there is a number $\delta > 0$ such that if $0 < |x - a| < \delta$ then $|f(x) - L| < \epsilon$.



1.3 Day 3

1.3.1 Continuous Function:

Definition: A function f is continuous at a number if $\lim_{x \rightarrow a} f(x) = f(a)$. Graphically, a function is continuous if you can draw it without having your pen leave paper. More formally, $f(x)$ is continuous at $x = a$ if :

1. $f(a)$ is defined ($a \in D : a$ is in the domain of f).
2. $\lim_{x \rightarrow a} f(x)$ exists, and equals $f(x) = f(a)$.

If one of the aforementioned statements is incorrect, then $f(x)$ is discontinuous at $x = a$

Theorem 1: A function is continuous on an interval if it's continuous at every number in the interval.


If f and g are continuous at $x = a$, then the following are also continuous at a :

$$f + g, f - g, cf, fg, \frac{f}{g} \text{ for } g(a) \neq 0$$

Theorem 2: The following types of functions are continuous at every number in there domain:

1. Polynomials (always $(-\infty, \infty)$)
2. Rational functions
3. Root functions
4. Trig functions

Theorem 3: Intermediate Value Theorem (IVM): Suppose that f is continuous on the close interval $[a, b]$ and let N be any number between $f(a)$ and $f(b)$, where $f(a) \neq f(b)$. Then there exists at least on number c in (a, b) such that $f(c) = N$




1.4 Day 4

1.4.1 Line:

1.4.1.1 Secant Line: A line that locally intersects two points on a curve.

$$\frac{\text{Rise}}{\text{Run}} = \frac{\Delta y}{\Delta x} = \frac{y_1 - y_0}{x_1 - x_0} = \frac{f(a+h) - f(a)}{(a+h) - a} = \frac{f(a+h) - f(a)}{h}$$



1.4.1.2 Tangent Line: The line through a pair of infinitely close points on the curve so that the line is "just touching". Slope equation (also known as "Difference Quotient"):

$$\lim_{h \rightarrow 0} \left[\frac{f(a+h) - f(a)}{h} \right]$$

1.4.1.3 Derivatives: The derivative of a function f at a number a , denoted by $f'(a)$, is

$$f'(a) = \lim_{h \rightarrow 0} \left[\frac{f(a+h) - f(a)}{h} \right]$$

and the equation of the tangent line to the curve $y = f(x)$ at the point $(a, f(a))$ can be written in point-slope form as

$$y - f(a) = f'(a)(x - a)$$