

# Max\_Vogel\_CMSSDataAnalysis.mlx

This code reads dilepton files from CMS and fits bumps in dilepton mass distributions. The data are available at <http://opendata.cern.ch/>.

Author: Max Vogel

Date: 10/14/2020

Collaborators: John Podczerwinski, Robert Wheatley, & Jaden Sengkhamee -- helped troubleshoot during lab.

Contributions: Professor Carlsmith for the code template and the CERN lab for collecting this data and making it available!

## Table of Contents

Z to two electrons.....	1
Initialize .....	1
Import & Prepare Data.....	1
Mass distribution fit.....	2
Z to two muons.....	3
Initialize .....	3
Import & Prepare Data.....	3
Mass distribution fit.....	4

## Z to two electrons

[Dataset URL](#)

### Initialize

```
clear;% clear variables
delete(findall(0,'Type','figure'))% close all open figure windows
```

### Import & Prepare Data

We'll first import the CSV data file

```
filename='Zee.csv';A=csvread(filename,1,1);nbins=100;
filename ; %drop filename to command window for users benefit
```

Then we'll create vectors for the energy, momentum components, transverse momentum, (pseudo)rapidity, azimuthal angle, & charge in units of proton charge.

```
Event=A(:,2);
E1=A(:,3);px1=A(:,4);py1=A(:,5);pz1=A(:,6);pt1=A(:,7);
eta1=A(:,8);phi1=A(:,9);q1=A(:,10);
E2=A(:,11);px2=A(:,12);py2=A(:,13);pz2=A(:,14);pt2=A(:,15);
eta2=A(:,16);phi2=A(:,17);q2=A(:,18);
M=A(:,19); %pair mass supplied by CMS
```

We can then compute the invariant mass of each lepton

```

m1= (abs(E1.^2-px1.^2-py1.^2-pz1.^2)).^(1/2);% mass of lepton 1
m2= (abs(E2.^2-px2.^2-py2.^2-pz2.^2)).^(1/2);% mass of lepton 2
m3=(abs((E1+E2).^2-(px1+px2).^2-(py1+py2).^2-(pz1+pz2).^2)).^(1/2);

```

...and then create the vector Nevents for the number of entries per bin created by the histogram command along with edges which is a vector of bin edges.

```
[Nevents,edges] = histcounts(m3,nbins);
```

With this data, we can create err containing statistical standard error in each bin equal to the square root of the number of events in that bin ([Poisson statistics](#))

```
err=Nevents.^(1/2);
```

Since edges starts at lower edge of first bin and ends at upper edge of last bin, we'll create the mass equal the vector of bin centers by eliminating last edge and adding a half bin size to each element.

```

mass=edges(1:end-1);
mass=mass +(edges(2)-edges(1))/2;

```

## Mass distribution fit

We'll use the [Relativistic Breit–Wigner distribution](#):  $\frac{k}{(m^2 - m_Z^2)^2 + m_Z^2 \cdot \Gamma_Z^2}$  where  $m_Z$  is the Z-boson mass treated as a parameter,  $\Gamma_Z$  is the width of the Z-boson mass, and  $k$  is a normalization constant.

```
createFitVogelZee(mass, Nevents, err)
```

```


ans =
General model:
ans(x) = k * 1/((x^2 - mz^2)^2 + mz^2 * wz^2)
Coefficients (with 95% confidence bounds):
k = 3.302e+07 (1.209e+07, 5.395e+07)
mz = 88.72 (87.94, 89.5)
wz = -6.512 (-9.019, -4.006)

```

```

title("Z to two electrons: Mass versus Number of events")
ylabel("Number of events")
xlabel("Mass (GeV * c^-2)")
ylim([0 130]); % so table doesn't obstruct view

```



For this fit,  $m_z$  was 88.72 while the true value is 91.1876  $\text{GeV} \cdot \text{c}^{-2}$ ,  $\Gamma_z$  was 6.512 while the true is 2.4952  $\text{GeV} \cdot \text{c}^{-2}$ , and  $k$  was  $3.302 \cdot 10^7$  (which was the constant.) All of the true values can be found on the [PDG's site](#). These discrepancies fall outside of two standard deviations of their respective measurements. The  $m_z$  value controls the center of the "spike" in the graph, so when collecting the data perhaps mass was over-estimated when being measured. Moreover, the line of best fit only considers the lines values with a mass greater than 60 and less than 120 so perhaps those bounds were inaccurate and morphed the  $\Gamma_z$  to be much higher than it should have been.

## Z to two muons

[Dataset URL](#)

### Initialize

```
clear;% clear variables
delete(findall(0,'Type','figure'))% close all open figure windows
```

### Import & Prepare Data

We'll first import the CSV data file

```
filename='Zmumu.csv';A=csvread(filename,1,1);nbins=100;
filename ; %drop filename to command window for users benefit
```

Then we'll create vectors for the energy, momentum components, transverse momentum, (pseudo)rapidity, azimuthal angle, & charge in units of proton charge.

```
Event=A(:,2);
E1=A(:,3);px1=A(:,4);py1=A(:,5);pz1=A(:,6);pt1=A(:,7);
eta1=A(:,8);phi1=A(:,9);q1=A(:,10);
E2=A(:,11);px2=A(:,12);py2=A(:,13);pz2=A(:,14);pt2=A(:,15);
eta2=A(:,16);phi2=A(:,17);q2=A(:,18);
M=A(:,19); %pair mass supplied by CMS
```

We can then compute the invariant mass of each lepton

```
m1= (abs(E1.^2-px1.^2-py1.^2-pz1.^2)).^(1/2);% mass of lepton 1
m2= (abs(E2.^2-px2.^2-py2.^2-pz2.^2)).^(1/2);% mass of lepton 2
m3=(abs((E1+E2).^2-(px1+px2).^2-(py1+py2).^2-(pz1+pz2).^2)).^(1/2);
```

...and then create the vector Nevents for the number of entries per bin created by the histogram command along with edges which is a vector of bin edges.

```
[Nevents,edges] = histcounts(m3,nbins);
```

With this data, we can create err containing statistical standard error in each bin equal to the square root of the number of events in that bin ([Poisson statistics](#))

```
err=Nevents.^(1/2);
```

Since edges starts at lower edge of first bin and ends at upper edge of last bin, we'll create the mass equal the vector of bin centers by eliminating last edge and adding a half bin size to each element.

```
mass=edges(1:end-1);
mass=mass +(edges(2)-edges(1))/2;
```


## Mass distribution fit

We'll use the [Relativistic Breit–Wigner distribution](#):  $\frac{k}{(m^2 - m_Z^2)^2 + m_Z^2 \cdot \Gamma_Z^2}$  where  $m_z$  is the Z-boson mass treated as a parameter,  $\Gamma_Z$  is the width of the Z-boson mass, and  $k$  is a normalization constant.

```
createFitVogelZmumu(mass, Nevents, err)
```

```
ans =
General model:
ans(x) = k * 1/((x^2 - mz^2)^2 + mz^2 * wz^2)
Coefficients (with 95% confidence bounds):
k = 1e+08 (fixed at bound)
mz = 90.9 (90.83, 90.96)
wz = 4.982 (4.935, 5.029)
```

```
title("Z to two muons: Mass versus Number of events")
ylabel("Number of events")
xlabel("Mass (GeV * c^-2)")
ylim([0 650]); % so table doesn't obstruct view
```



For this fit,  $m_z$  was 90.9 while the true value is 91.1876 GeV\*c<sup>-2</sup>,  $\Gamma_z$  was 4.982 while the true is 2.4952 GeV\*c<sup>-2</sup>, and  $k$  was  $1 \cdot 10^8$  (which was the constant.) All of the true values can be found on the [PDG's site](#). Again, these discrepancies fall outside of two standard deviations of their respective measurements. Because the  $k$  value (which is responsible for the amplitude) was fixed/limited, it could have caused the  $\Gamma_z$  value (which also affects the amplitude as well as width of the spike) to be incorrectly estimated.