

# Solving Models of Economic Dynamics with Ridgeless Kernel Regressions

---

Mahdi Ebrahimi Kahou<sup>1</sup>   Jesse Perla<sup>2</sup>   Geoff Pleiss<sup>3,4</sup>

ASSA 2026

<sup>1</sup>Bowdoin College, Econ Dept

<sup>2</sup>University of British Columbia, Vancouver School of Economics

<sup>3</sup>University of British Columbia, Stats Dept

<sup>4</sup>Vector Institute

# Motivation

- Numerical solutions to dynamical systems are central to many quantitative fields in economics.
- Dynamical systems in economics are **boundary value** problems:
  1. The boundary is at **infinity**.
  2. The values at the boundary are potentially **unknown**.
- Resulting from **forward looking** behavior of agents.
- Examples include the transversality and the no-bubble condition.
- Without them, the problems are **ill-posed** and have infinitely many solutions:
  - These forward-looking boundary conditions are a key limitation on increasing dimensionality.

Using kernel method to solve a broad class of infinite-horizon, deterministic, continuous-time model

1. **Minimum-norm alignment:**

- The minimum-norm kernel method aligns with asymptotic boundary conditions.

2. **Learning the right set of steady-states:**


- Kernel machines learn the boundary values, thereby extrapolating outside the training data.

3. **Robustness and speed:**

- Competitive in speed and more stable than traditional methods.

4. **Consistency of ML estimates.**

- **Violation of the boundary conditions:**
  - Sub-optimal solutions explode over time.
  - They have large derivatives.
  - This behavior is due to the **saddle-path** nature of the problem.
- **Minimum-norm solution :**
  - Penalizing large derivatives rules out explosive paths.
  - The remaining solution is the optimal solution.



# The Problem

---

# The class of problems

A differential-algebraic system of equations, coming from an optimization problem:

$$\dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{x}(t), \boldsymbol{\mu}(t), \mathbf{y}(t)) \quad (1)$$

$$\dot{\boldsymbol{\mu}}(t) = r\boldsymbol{\mu}(t) - \boldsymbol{\mu}(t) \odot \mathbf{G}(\mathbf{x}(t), \boldsymbol{\mu}(t), \mathbf{y}(t)) \quad (2)$$

$$\mathbf{0} = \mathbf{H}(\mathbf{x}(t), \boldsymbol{\mu}(t), \mathbf{y}(t)), \quad (3)$$

boundary conditions (at infinity)

$$\mathbf{0} = \lim_{t \rightarrow \infty} e^{-rt} \mathbf{x}(t) \odot \boldsymbol{\mu}(t), \quad (4)$$

initial value  $\mathbf{x}(0) = \mathbf{x}_0$ .

- $\mathbf{x} \in \mathbb{R}^M$ : state variables.
- $\boldsymbol{\mu} \in \mathbb{R}^M$ : co-state variables.
- $\mathbf{y} \in \mathbb{R}^P$ : jump variables.

**Goal:** Find an approximation to  $\mathbf{x}(t)$ ,  $\mu(t)$ , and  $\mathbf{y}(t)$ .

**What is the problem?**

- Initial conditions  $\mathbf{y}_0$  and  $\mu_0$  are unknown.
- The optimal solution follows a **saddle path**.
  - If  $T$  is small, solutions are inaccurate due to premature enforcement of the steady state.
  - If  $T$  is large, the algorithms become increasingly numerically unstable.

## Example: Neoclassical Growth Model

$$\dot{x}(t) = f(x(t)) - \delta x(t) - y(t) := F(x(t), \mu(t), y(t))$$

$$\dot{\mu}(t) = r\mu(t) - \underbrace{\mu(t)(f'(x(t)) - \delta)}_{:= G(x(t), \mu(t), y(t))}$$

$$0 = \mu(t)y(t) - 1 := H(x(t), \mu(t), y(t))$$

$$x(0) = x_0, \quad \lim_{t \rightarrow \infty} e^{-rt} \mu(t)x(t) = 0$$

capital  $x(t)$ , consumption  $y(t)$ , utility  $\log(y)$ , present-value co-state variable  $\mu(t)$ , discount rate  $r > 0$ , depreciation rate  $0 < \delta < 1$ , and production function  $f(x)$ .



## Method

---

## Method: approximation

- Pick a set of points  $\mathcal{D} \equiv \{t_1, \dots, t_N\}$  for some fixed interval  $[0, T]$

$$\hat{\mathbf{x}}(t) = \sum_{j=1}^N \alpha_j^x k(t, t_j), \quad \hat{\mu}(t) = \sum_{j=1}^N \alpha_j^\mu k(t, t_j), \quad \hat{\mathbf{y}}(t) = \sum_{j=1}^N \alpha_j^y k(t, t_j),$$

$$\hat{\mathbf{x}}(t) = \mathbf{x}_0 + \int_0^t \hat{\mathbf{x}}(\tau) d\tau, \quad \hat{\mu}(t) = \mu_0 + \int_0^t \hat{\mu}(\tau) d\tau, \quad \hat{\mathbf{y}}(t) = \mathbf{y}_0 + \int_0^t \hat{\mathbf{y}}(\tau) d\tau.$$

- $\alpha_j^x$ ,  $\alpha_j^\mu$ ,  $\alpha_j^y$ ,  $\mu_0$ , and  $\mathbf{y}_0$  are parameters to be found.
- $k(t, t_j)$  is a kernel that measures “similarity” between  $t$  and  $t_j$ .
- We use a Matérn kernel with smoothness  $\nu$  and length  $\ell$ .

►► Matérn kernel

## Method: Ridgeless kernel regression

We solve

$$\begin{aligned} \min_{\hat{\mathbf{x}}, \hat{\boldsymbol{\mu}}, \hat{\mathbf{y}}} & \left( \sum_{m=1}^M \|\hat{\mathbf{x}}^{(m)}\|_{\mathcal{H}}^2 + \sum_{m=1}^M \|\hat{\boldsymbol{\mu}}^{(m)}\|_{\mathcal{H}}^2 \right) \\ \text{s.t. } & \hat{\mathbf{x}}(t_i) = \mathbf{F}(\hat{\mathbf{x}}(t_i), \hat{\boldsymbol{\mu}}(t_i), \hat{\mathbf{y}}(t_i)), \quad \text{for all } t_i \in \mathcal{D} \\ & \hat{\boldsymbol{\mu}}(t_i) = r \hat{\boldsymbol{\mu}}(t_i) - \hat{\boldsymbol{\mu}}(t_i) \odot \mathbf{G}(\hat{\mathbf{x}}(t_i), \hat{\boldsymbol{\mu}}(t_i), \hat{\mathbf{y}}(t_i)), \quad \text{for all } t_i \in \mathcal{D} \\ & \mathbf{0} = \mathbf{H}(\hat{\mathbf{x}}(t_i), \hat{\boldsymbol{\mu}}(t_i), \hat{\mathbf{y}}(t_i)), \quad \text{for all } t_i \in \mathcal{D}. \end{aligned}$$

- $\|\hat{\mathbf{x}}^{(m)}\|_{\mathcal{H}}^2 = \sum_{i=1}^N \sum_{j=1}^N \alpha_i^{x^{(m)}} \alpha_j^{x^{(m)}} k(t_i, t_j)$  and  $\|\hat{\boldsymbol{\mu}}^{(m)}\|_{\mathcal{H}}^2 = \sum_{i=1}^N \sum_{j=1}^N \alpha_i^{\mu^{(m)}} \alpha_j^{\mu^{(m)}} k(t_i, t_j)$
- TVC is **not imposed**.
- The penalization is used to control the smoothness of the approximating functions.
- For Matérn kernels, it also controls the smoothness of derivatives.

# Applications

---

# Neoclassical growth model

$$\begin{aligned} \max_{y(t)} \int_0^{\infty} e^{-rt} \ln(y(t)) dt \\ \text{s.t. } \dot{x}(t) = f(x(t)) - y(t) - \delta x(t) \end{aligned}$$

for a given  $x_0$ .

- $x(t) \in \mathbb{R}$ : capital,  $y(t) \in \mathbb{R}$ : consumption, and a concave production function  $f(x) = x^a$ .

Constructing the Hamiltonian ...

$$\begin{aligned} \dot{x}(t) &= f(x(t)) - \delta x(t) - y(t), \\ \dot{\mu}(t) &= r\mu(t) - \mu(t)(f'(x(t)) - \delta), \\ 0 &= \mu(t)y(t) - 1, \\ x(0) &= x_0, \quad \lim_{t \rightarrow \infty} e^{-rt} \mu(t)x(t) = 0. \end{aligned}$$

- Last Equation : transversality condition (TVC)

# Why do we need the boundary condition?



Ignoring the transversality condition:

$$\dot{x}(t) = f(x(t)) - \delta x(t) - y(t),$$

$$\dot{\mu}(t) = r\mu(t) - \mu(t)(f'(x(t)) - \delta),$$

$$0 = \mu(t)y(t) - 1,$$

$$x(0) = x_0.$$



# Neoclassical Growth Model: algorithm

$$\min_{\hat{x}, \hat{\mu}, \hat{y}} \left( \|\hat{x}\|_{\mathcal{H}}^2 + \|\hat{\mu}\|_{\mathcal{H}}^2 \right)$$

$$\text{s.t. } \hat{x}(t_i) = f(\hat{x}(t_i)) - \delta \hat{x}(t_i) - \hat{y}(t_i), \quad \text{for all } t_i \in \mathcal{D}$$


$$\hat{\mu}(t_i) = r \hat{\mu}(t_i) - \hat{\mu}(t_i) (f'(\hat{x}(t_i)) - \delta), \quad \text{for all } t_i \in \mathcal{D}$$

$$0 = \hat{\mu}(t_i) \hat{y}(t_i) - 1, \quad \text{for all } t_i \in \mathcal{D}.$$

- $\|\hat{x}\|_{\mathcal{H}}^2 = \sum_{i=1}^N \sum_{j=1}^N \alpha_i^x \alpha_j^x k(t_i, t_j)$  and  $\|\hat{\mu}\|_{\mathcal{H}}^2 = \sum_{i=1}^N \sum_{j=1}^N \alpha_i^{\mu} \alpha_j^{\mu} k(t_i, t_j)$
- TVC is **not imposed**.

# Neoclassical growth model: results

- $\mathcal{D} = \{0, 1, \dots, 40\}$ . ► sparse grids
- $f(x) = x^{\frac{1}{3}}$ ,  $\delta = \frac{1}{3}$ , and  $r = 0.11$ .
- The explosive solutions are ruled out without directly imposing the boundary condition.
- Very accurate approximations, both in the short- and medium-run.
- Learns the **correct steady state**. ► Relative errors





# Neoclassical growth model: learning the steady state

Why does it learn the **correct steady state**?

- A straight line is the “smoothest” solution: it has zero derivatives.
- The kernels we use are zero-reverting:

$$\lim_{t \rightarrow \infty} k(t, t_j) = 0.$$

- We approximate the derivatives using kernels:

$$\lim_{t \rightarrow \infty} \hat{x}(t) = \hat{\mu}(t) = \hat{y}(t) = 0.$$



- This behavior is mainly driven by choosing a large  $t_N$  (e.g.,  $t_N = 40$ ) in  $\mathcal{D}$ .
- **Question:** How accurate are the short-run dynamics when  $t_N$  is small?

# Neoclassical growth model: short-run results

- $\mathcal{D} = \{0, 1, \dots, 10\}$ .
- $f(x) = x^{\frac{1}{3}}$ ,  $\delta = \frac{1}{3}$ , and  $r = 0.11$ .
- Very accurate short-run dynamics. ▶▶ Relative errors

## Conclusion

- For short-run accuracy, we do not need to know the steady state (global condition).
- It suffices that sub-optimal paths diverge fast enough from the optimal path (local condition).



## Extensions

---

# Neoclassical Growth Model: Non-Concave Production Function

- So far we have had a **unique** saddle-path converging to a unique **saddle** steady state.
- What if we have **two** saddle steady states, very close to each other (steady-state multiplicity)?
- Neoclassical growth model with a non-concave production function (threshold externalities):


$$f(x) = A \max\{x^a, b_1 x^a - b_2\}$$

# Non-concave production function: vector field



$$\dot{\mathbf{x}}(t) = f(\mathbf{x}(t)) - \mathbf{y}(t) - \delta \mathbf{x}(t)$$

$$\dot{\mathbf{y}}(t) = \mathbf{y}(t)[f'(\mathbf{x}(t)) - \delta - r]$$

$$\mathbf{x}(0) = \mathbf{x}_0 \text{ given.}$$



# Results



- The approximate solutions approach the right steady states.
- The transversality conditions are satisfied without being directly imposed.
- The steady states are learned. [▶ Full DAE](#)

Linear asset pricing model

$$\dot{x}(t) = c + gx(t)$$

$$\dot{\mu}(t) = r\mu(t) - x(t) := r\mu(t) - \mu(t) \frac{x(t)}{\mu(t)}$$

$$0 = \lim_{t \rightarrow \infty} e^{-rt} \mu(t) x(t).$$

- $x(t) \in \mathbb{R}$ : flow payoffs from a claim to an asset.
- $\mu(t) \in \mathbb{R}$  be the price of a claim to that asset.
- $x_0$  given.

# Why do we need the boundary condition?

$$\dot{\mathbf{x}}(t) = c + g\mathbf{x}(t)$$

$$\dot{\mathbf{y}}(t) = r\mathbf{y}(t) - \mathbf{x}(t)$$


- The solutions:

$$\mathbf{y}(t) = \mathbf{y}_f(t) + \zeta e^{rt}$$

- $\mathbf{y}_f(t) = \int_0^\infty e^{-r\tau} \mathbf{x}(t+s) ds$ : price based on the fundamentals.
- $\zeta e^{rt}$ : explosive bubble terms, it has to be **ruled out** by the boundary condition.
- Triangle inequality:  $\|\mathbf{y}_f\| < \|\mathbf{y}\|$ .
- The price based on the fundamentals has the **lowest norm**.



# Results



$$\mathcal{D} = \{0, 1, \dots, 30\}$$

- The explosive solutions are ruled out without directly imposing the boundary condition.
- Very accurate approximations, both in the short- and medium-run.
- Learns the steady-state.

- Long-run (**global**) conditions can be replaced with appropriate regularization (**local**) to achieve the optimal solutions.
- The minimum-norm implicit bias of large ML models aligns with optimality in economic dynamic models.
- Both kernel and neural network approximations accurately learn the right steady state(s).
- Proceeding with **caution**: can regularization be thought of as an equilibrium selection device?

# Appendix


---

$$K(t, t_j) = C_{\frac{1}{2}}(t, t_j) = \sigma^2 \exp\left(-\frac{|t - t_j|}{\ell}\right),$$


$$K(t, t_j) = C_{\frac{3}{2}}(t, t_j) = \sigma^2 \left(1 + \frac{\sqrt{3}|t - t_j|}{\ell}\right) \exp\left(-\frac{\sqrt{3}|t - t_j|}{\ell}\right),$$

$$K(t, t_j) = C_{\frac{5}{2}}(t, t_j) = \sigma^2 \left(1 + \frac{\sqrt{5}|t - t_j|}{\ell} + \frac{5|t - t_j|^2}{3\ell^2}\right) \exp\left(-\frac{\sqrt{5}|t - t_j|}{\ell}\right).$$


# Neoclassical growth: relative errors



# Neoclassical growth: short-run relative errors



# Neoclassical growth: sparse sampling



$$\mathcal{D} = \{0, 1, 3, 5, 10, 15, 20, 25, 30, 35, 38, 40\}$$

$$\dot{\mathbf{x}}_k(t) = \mathbf{y}_k(t) - \delta_k \mathbf{x}_k(t),$$

$$\dot{\mathbf{x}}_h(t) = \mathbf{y}_h(t) - \delta_h \mathbf{x}_h(t)$$

$$\dot{\mathbf{y}}_c(t) = \mathbf{y}_c(t) [f_1(\mathbf{x}_k(t), \mathbf{x}_h(t)) - \delta_k - r],$$

$$0 = f(\mathbf{x}_k(t), \mathbf{x}_h(t)) - \mathbf{y}_c(t) - \mathbf{y}_k(t) - \mathbf{y}_h(t),$$

$$0 = f_2(\mathbf{x}_k(t), \mathbf{x}_h(t)) - f_1(\mathbf{x}_k(t), \mathbf{x}_h(t)) + \delta_k - \delta_h.$$

$$0 = \lim_{t \rightarrow \infty} e^{-rt} \frac{\mathbf{x}_k(t)}{\mathbf{y}_c(t)}, \quad 0 = \lim_{t \rightarrow \infty} e^{-rt} \frac{\mathbf{x}_h(t)}{\mathbf{y}_c(t)}.$$

- $\mathbf{x}_k$ : physical capital,  $\mathbf{x}_h$ : human capital,  $\mathbf{y}_c$ : consumption,  $\mathbf{y}_k$ : investment in physical capital,  $\mathbf{y}_h$ : investment in human capital
- $f(x_k, x_h) = x_k^{a_k} x_h^{a_h}$



# Results

