

Logistic Regression: Cost Function

To train the parameters w and b , we need to define a cost function.

Recap:

$$\hat{y}^{(i)} = \sigma(w^T x^{(i)} + b), \text{ where } \sigma(z^{(i)}) = \frac{1}{1 + e^{-z^{(i)}}}$$

$x^{(i)}$ the i-th training example

Given $\{(x^{(1)}, y^{(1)}), \dots, (x^{(m)}, y^{(m)})\}$, we want $\hat{y}^{(i)} \approx y^{(i)}$

Loss (error) function:

The loss function measures the discrepancy between the prediction ($\hat{y}^{(i)}$) and the desired output ($y^{(i)}$). In other words, the **loss function computes the error for a single training example**.

$$L(\hat{y}^{(i)}, y^{(i)}) = \frac{1}{2} (\hat{y}^{(i)} - y^{(i)})^2$$

$$L(\hat{y}^{(i)}, y^{(i)}) = -(y^{(i)} \log(\hat{y}^{(i)}) + (1 - y^{(i)}) \log(1 - \hat{y}^{(i)}))$$

- If $y^{(i)} = 1$: $L(\hat{y}^{(i)}, y^{(i)}) = -\log(\hat{y}^{(i)})$ where $\log(\hat{y}^{(i)})$ and $\hat{y}^{(i)}$ should be close to 1
- If $y^{(i)} = 0$: $L(\hat{y}^{(i)}, y^{(i)}) = -\log(1 - \hat{y}^{(i)})$ where $\log(1 - \hat{y}^{(i)})$ and $\hat{y}^{(i)}$ should be close to 0

Cost function

The cost function is the average of the loss function of the entire training set. We are going to find the parameters w and b that minimize the overall cost function.

$$J(w, b) = \frac{1}{m} \sum_{i=1}^m L(\hat{y}^{(i)}, y^{(i)}) = -\frac{1}{m} \sum_{i=1}^m [y^{(i)} \log(\hat{y}^{(i)}) + (1 - y^{(i)}) \log(1 - \hat{y}^{(i)})]$$