



# ΔFosB: A transcriptional regulator of stress and antidepressant responses

Eric J. Nestler\*

Fishberg Department of Neuroscience and Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA



## ARTICLE INFO

### Article history:

Accepted 9 October 2014

Available online 7 November 2014

### Keywords:

Fos  
Nucleus accumbens  
Prefrontal cortex  
Epigenetics

## ABSTRACT

ΔFosB is a member of the Fos family of transcription factors. While other family members are induced rapidly but transiently in response to a host of acute stimuli, ΔFosB is unique in that it accumulates in response to repeated stimulation due to its unusual protein stability. Such prolonged induction of ΔFosB, within nucleus accumbens (NAc), a key brain reward region, has been most studied in animal models of drug addiction, with considerable evidence indicating that ΔFosB promotes reward and motivation and serves as a mechanism of drug sensitization and increased drug self-administration. In more recent years, prolonged induction of ΔFosB has also been observed within NAc in response to chronic administration of certain forms of stress. Increasing evidence indicates that this induction represents a positive, homeostatic adaptation to chronic stress, since overexpression of ΔFosB in this brain region promotes resilience to stress, whereas blockade of its activity promotes stress susceptibility. Chronic administration of several antidepressant medications also induces ΔFosB in the NAc, and this induction is required for the therapeutic-like actions of these drugs in mouse models. Validation of these rodent findings is the demonstration that depressed humans, examined at autopsy, display reduced levels of ΔFosB within the NAc. As a transcription factor, ΔFosB produces this behavioral phenotype by regulating the expression of specific target genes, which are under current investigation. These studies of ΔFosB are providing new insight into the molecular basis of depression and antidepressant action, which is defining a host of new targets for possible therapeutic development.

© 2014 Elsevier B.V. All rights reserved.

## 1. Introduction

The study of transcriptional mechanisms of depression is based on the hypothesis that regulation of gene expression is one important mechanism by which chronic exposure to stress causes depression or related abnormalities in vulnerable individuals (Krishnan and Nestler, 2010). A corollary of this hypothesis is that the ability of a host of antidepressant medications, after prolonged administration, to reduce the symptoms of depression in some individuals is likewise mediated in part by altered gene expression in relevant brain regions.

Work over the past 20 years has provided increasing evidence for a role of gene regulation in depression models, as several transcription factors – proteins that bind to specific response elements in the promoter regions of target genes and regulate those genes' expression – have been implicated in these models.

Examples of transcription factors that have been studied prominently in stress models include CREB (cAMP response element binding protein), glucocorticoid receptor, and NFκB (nuclear factor κB), among others (Nestler et al., 2002; Carlezon et al., 2005; Holsboer and Ising, 2010; Christoffel et al., 2011; Licznarski and Duman, 2013).

The focus of this review is on another transcription factor, ΔFosB, which has been mostly studied in drug addiction models (Nestler, 2008, 2012). More recent work has demonstrated that ΔFosB, a member of the Fos family of proteins, is also regulated in stress and depression models and appears to play a unique role in promoting resilience and antidepressant responses. This discussion also illustrates the types of experimental approaches that have been used to investigate transcriptional mechanisms of depression in mouse models.


## 2. Induction of ΔFosB in nucleus accumbens by chronic stress

ΔFosB is encoded by the *FosB* gene (Fig. 1) and shares homology with other Fos family transcription factors, which include c-Fos, FosB, Fra1, and Fra2 (Morgan and Curran, 1995).

\* Correspondence to: Fishberg Department of Neuroscience, One Gustave L. Levy Place, Box 1065, New York, NY 10029, USA.

Tel.: +1 212 659 5656; fax: +1 212 659 8510.

E-mail address: [eric.nestler@mssm.edu](mailto:eric.nestler@mssm.edu)




**Fig. 1.** Biochemical basis of  $\Delta$ FosB's unique stability.  $\Delta$ FosB and FosB are encoded by the *FosB* gene.  $\Delta$ FosB is generated by alternative splicing and lacks the C-terminal 101 amino acids present in FosB. Two mechanisms are known that account for  $\Delta$ FosB's stability. First,  $\Delta$ FosB lacks two degron domains present in the C-terminus of full length FosB (and found in all other Fos family proteins as well). One of these degron domains targets FosB for ubiquitylation and degradation in the proteasome. The other degron domain targets FosB degradation by a ubiquitin- and proteasome-independent mechanism. Second,  $\Delta$ FosB is phosphorylated by several protein kinases at its N-terminus which further stabilizes the protein. From Nestler, 2008 with permission.

These Fos family proteins heterodimerize with Jun family proteins (c-Jun, JunB, or JunD) to form active AP1 (activator protein-1) transcription factors that bind to AP1 sites (consensus sequence: TGAC/GTCA) present in the promoters of certain genes to regulate their transcription. Fos family proteins are induced rapidly and transiently in specific brain regions after acute administration of several forms of stress (Perrotti et al., 2004). These responses are seen most prominently in nucleus accumbens (NAc), which is best characterized as an important mediator of reward and motivation. All of these Fos family proteins, however, are highly unstable and return to basal levels within hours of the stress exposure Fig. 2.

Very different responses are seen after chronic exposure to stress. Biochemically modified isoforms of  $\Delta$ FosB ( $M_r$  35–37 kD) accumulate within the same brain regions after repeated stress exposure, whereas other Fos family members show desensitization (i.e., reduced induction compared with initial drug exposures) (Perrotti et al., 2004; Vialou et al., 2010a; Lehmann and Herkenham, 2011). Such accumulation of  $\Delta$ FosB has been observed for several forms of active stress, such as chronic restraint stress, chronic unpredictable stress, and chronic social defeat stress (Perrotti et al., 2004; Vialou et al., 2010a; Lehmann and Herkenham, 2011), whereas chronic social isolation of adult animals strikingly causes the opposite effect: a decrease in  $\Delta$ FosB levels in NAc (Vialou et al., 2010a).


For chronic restraint stress and chronic social defeat stress, induction of  $\Delta$ FosB in NAc has been shown to occur in both major subtypes of medium spiny neurons (MSNs)—those that express predominantly the  $D_1$  dopamine receptor ( $D_1$ -type MSNs) or the  $D_2$  dopamine receptor ( $D_2$ -type MSNs), which together represent ~95% of all neurons in this brain region (Perrotti et al., 2004; Lobo et al., 2013). In contrast, no induction is seen in any of several types of interneurons or in non-neural cells. One of the advantages of the social defeat paradigm is that a subset of mice do not develop depression-like behavioral abnormalities, that is, they remain resilient unlike the majority which are susceptible (Krishnan et al., 2007). Interestingly, the modest induction of  $\Delta$ FosB seen in the NAc of susceptible mice occurs primarily in  $D_2$ -type MSNs, where the more robust induction that occurs in resilient mice is specific to  $D_1$ -type MSNs (Lobo et al., 2013). This cellular specificity of  $\Delta$ FosB induction has important functional implications as discussed in Section 4.



**Fig. 2.** Scheme showing the gradual accumulation of  $\Delta$ FosB versus the rapid and transient induction of other Fos family proteins in response to stress or another stimulus. The upper graph shows that several waves of Fos family proteins (comprised of c-Fos, FosB,  $\Delta$ FosB [33 kD isoform], Fra1, and Fra2) are induced in NAc and dorsal striatal neurons by acute stress. Also induced are biochemically modified isoforms of  $\Delta$ FosB (35–37 kD); they are induced at low levels by acute stress, but persist in brain for long periods due to their stability. The lower graph shows that with repeated (e.g., twice daily) stress exposure, each acute stimulus induces a low level of the stable  $\Delta$ FosB isoforms. This is indicated by the lower set of overlapping lines, which indicate  $\Delta$ FosB induced by each acute stimulus. The result is a gradual increase in the total levels of  $\Delta$ FosB with repeated stimuli during a course of chronic treatment. This is indicated by the increasing stepped line in the graph. From Nestler, 2008 with permission.

The 35–37 kD isoforms of  $\Delta$ FosB dimerize predominantly with JunD to form an active and long-lasting AP1 complex within these brain regions (Chen et al., 1997; Hiroi et al., 1998). However, recent in vitro evidence has indicated that  $\Delta$ FosB can also form homodimers with distinct physico-chemical properties compared to  $\Delta$ FosB:JunD heterodimers (Jorissen et al., 2007). An important focus of current research is to determine whether such  $\Delta$ FosB homodimers form in vivo and what physiological function they subserve.

The 35–37 kD  $\Delta$ FosB isoforms accumulate after chronic stress or other repeated stimuli due to their extraordinarily long half-lives, which has been demonstrated both in cultured cells in vitro and within the NAc in vivo (Nestler, 2008). As a result of its stability, therefore, the  $\Delta$ FosB protein persists in neurons for at least several weeks after cessation of stress exposure. We now know that this stability is due to two factors (Fig. 1): (1) the absence from  $\Delta$ FosB of two degron domains which are present at the C-terminus of full length FosB and all other Fos family proteins and target those proteins for rapid degradation, and (2) the phosphorylation of  $\Delta$ FosB at its N-terminus (Ser27) by  $Ca^{2+}$ /calmodulin-dependent protein kinase II (CaMKII) and casein kinase 2 and perhaps by other protein kinases (Ulery et al., 2006; Carle et al., 2007; Ulery-Reynolds, 2009; Robison et al., 2013). The mechanism by which



**Fig. 3.**  $\Delta$ FosB induction in NAc by chronic social defeat stress mediates resilience. (A) Inducible bitransgenic mice overexpressing  $\Delta$ FosB in D<sub>1</sub>-type MSNs do not develop the social aversion which is a hallmark of social defeat stress after either 4 or 10 days of defeat with mice tested on days 5 or 11, respectively. (B) Conversely, overexpression of  $\Delta$ cJun increases susceptibility to social defeat with increased social aversion seen after 4 days of defeat with mice examined on day 5. By day 11, both control and  $\Delta$ cJun-expressing mice exhibit comparable levels of social avoidance. \* $p < 0.05$ . From [Vialou et al., 2010a](#) with permission.

phosphorylation of  $\Delta$ FosB at Ser27 increases its stability remains unknown. The stability of the  $\Delta$ FosB isoforms provides a novel molecular mechanism by which stress-induced changes in gene expression can persist long after the stress exposure.  $\Delta$ FosB is also phosphorylated at two threonine residues near its DNA-binding and transactivation domains, and phosphorylation of one of these sites (Thr149) dramatically increases the transcriptional activity of the protein ([Cates et al., 2014](#)). Further work is needed to define the role of these phosphorylation sites in stress responses.

### 3. Induction of $\Delta$ FosB in nucleus accumbens by chronic antidepressant treatment

Recent studies have shown that chronic administration of fluoxetine, a serotonin-selective reuptake inhibitor antidepressant, induces  $\Delta$ FosB in the NAc ([Vialou et al., 2010a](#)).  $\Delta$ FosB is also induced in response to chronic administration of other antidepressants including the tricyclic antidepressant imipramine (unpublished observations) and the rapidly-acting, novel antidepressant ketamine ([Donahue et al., 2014](#)). Interestingly, similar to the pattern of induction seen in resilient mice after chronic social defeat stress,  $\Delta$ FosB induction in NAc in response to chronic fluoxetine treatment is selective for D<sub>1</sub>-type MSNs, with no induction seen in NAc interneurons or glia ([Lobo et al., 2013](#)). It would be interesting moving forward to characterize the induction of  $\Delta$ FosB in NAc by several other classes of antidepressant medications. Recent work, for example, has shown that environmental enrichment during rearing increases basal levels of  $\Delta$ FosB in the NAc ([Zhang et al., 2014](#)). Since such environmental enrichment is thought to enhance resilience ([Lehmann and Herkenham, 2011](#)), these findings raise the interesting possibility that non-medication antidepressant approaches might induce  $\Delta$ FosB in this brain region as well.

Chronic administration of fluoxetine has recently been shown to increase levels of full-length FosB within the NAc 24 h after the last drug exposure ([Vialou et al., 2014b](#)). This is a surprising finding, since full-length FosB is highly unstable and, in chronic stress or chronic drug abuse models, its levels are not elevated beyond 2–4 h after each exposure. These findings suggest that repeated administration of fluoxetine might cause sustained increases in neural activity in this brain region, which might be expected to drive sustained increases in FosB levels. Further work is needed to study the validity of this possibility as well as to determine if this phenomenon occurs with other antidepressants.

### 4. Role of $\Delta$ FosB in nucleus accumbens in regulating behavioral responses to stress and antidepressant treatments

Insight into the role of  $\Delta$ FosB in stress and antidepressant responses has come from a combination of experimental approaches. First, we have developed bitransgenic mice in which  $\Delta$ FosB can be induced selectively within the NAc and dorsal striatum of adult animals ([Chen et al., 1995](#); [Kelz et al., 1999](#)). Importantly, these mice overexpress  $\Delta$ FosB selectively in D<sub>1</sub>-type MSNs. Second, we have developed a series of viral vectors that selectively overexpress  $\Delta$ FosB in the NAc, without expression in dorsal striatum, although this expression occurs in all neurons, not just D<sub>1</sub>-type MSNs ([Zachariou et al., 2006](#)). Conversely, we have developed bitransgenic mice that overexpress a truncated Jun protein, called  $\Delta$ cJun, within NAc, dorsal striatum, and several other brain regions ([Peakman et al., 2003](#)), as well as viral vectors that selectively overexpress a different truncated Jun protein, called  $\Delta$ JunD, in the NAc ([Winstanley et al., 2007](#)). These truncated Jun proteins serve as dominant negative antagonists of  $\Delta$ FosB- or other AP1-mediated transcription.

By use of these complementary approaches, we have established that increased expression of  $\Delta$ FosB within D<sub>1</sub>-type MSNs of

the NAc reduces an animal's sensitivity to the deleterious effects of chronic stress, promotes stress resilience, and mediates antidepressant-like responses (Fig. 3). Thus, bitransgenic mice inducibly overexpressing  $\Delta$ FosB in D<sub>1</sub>-type MSNs, or mice injected intra-NAc with a viral vector encoding  $\Delta$ FosB show increased resilience to subsequent chronic social defeat stress (Vialou et al., 2010a; Ohnishi et al., 2014). They also display reversal of depression-like behavioral abnormalities after chronic social defeat stress. Conversely, bitransgenic mice inducibly overexpressing  $\Delta$ cJun in NAc and several other brain regions, or mice injected intra-NAc with a viral vector encoding  $\Delta$ JunD show increased susceptibility to sub-threshold levels of social defeat stress. They also fail to respond to chronic fluoxetine administration, which reverses depression-like behavioral abnormalities, after chronic social defeat stress (Vialou et al., 2010a).

Further support for the view that  $\Delta$ FosB induction in NAc promotes resilience is the more recent finding that mice that constitutively lack expression of full length FosB, but show increased expression of  $\Delta$ FosB, display reduced sensitivity to stress (Ohnishi et al., 2011). Likewise, further support for the view that  $\Delta$ FosB induction in NAc elevates mood and promotes antidepressant-like responses are the findings that  $\Delta$ FosB overexpression in D<sub>1</sub>-type MSNs opposes the elevation of brain stimulation reward thresholds induced either by chronic social defeat stress or by a  $\kappa$  opioid agonist which is pro-depressant (Muschamp et al., 2012; Donahue et al., 2014). Moreover,  $\Delta$ FosB overexpression in NAc promotes several rewarding behaviors, including wheel-running, sucrose drinking, consumption of high-fat food, and sexual activity, with most of these effects seen upon selective expression in D<sub>1</sub>-type MSNs, while overexpression of  $\Delta$ cJun or  $\Delta$ JunD exerts the opposite effect (Werme et al., 2002; Olausson et al., 2006; Teegarden et al., 2008; Wallace et al., 2008; Hedges et al., 2009; Pitchers et al., 2010, 2013; Been et al., 2013).

In contrast to the behavioral phenotype mediated by  $\Delta$ FosB induction in D<sub>1</sub>-type MSNs, it has been harder to decipher the effect of  $\Delta$ FosB induction in D<sub>2</sub>-type MSNs. As noted earlier, such induction is seen selectively in mice that are susceptible to chronic social defeat stress. However, bitransgenic mice that inducibly overexpress  $\Delta$ FosB in D<sub>2</sub>-type MSNs (Chen et al., 1995; Werme et al., 2002) do not show a prominent pro-depression-like phenotype in chronic stress models (unpublished observations), although they do show reduced wheel-running activity (Werme et al., 2002). Further complicating the situation is the observation that several rewarding behaviors, including sucrose drinking and being reared in an enriched environment, induce  $\Delta$ FosB roughly equally in NAc D<sub>1</sub>- and D<sub>2</sub>-type MSNs (Lobo et al., 2013). Additional studies are thus needed to better understand the behavioral consequences of  $\Delta$ FosB induction in D<sub>2</sub>-type MSNs. Of note, the recent development of viral vectors, which make it possible for the first time to overexpress  $\Delta$ FosB selectively in either D<sub>1</sub>-type MSNs or in D<sub>2</sub>-type MSNs within the NAc (Grueter et al., 2013), should help in these efforts.

## 5. Mechanism of $\Delta$ FosB induction in nucleus accumbens

The upstream signaling pathways through which chronic stress or antidepressant treatment induces  $\Delta$ FosB in NAc remain largely unknown. Recent work, however, has defined the transcription factors that are required for  $\Delta$ FosB induction by chronic stress. The ability of chronic social defeat stress to induce  $\Delta$ FosB in NAc requires SRF (serum response factor): local knockout of this transcription factor completely blocks stress induction of  $\Delta$ FosB, whereas knockout of another transcription factor, CREB, is without effect (Vialou et al., 2010b). Surprisingly, cocaine induction of  $\Delta$ FosB in this same brain region requires both SRF and CREB: local

knockout of both factors completely blocks the ability of cocaine to induce  $\Delta$ FosB, whereas knockout of either factor alone is without effect (Vialou et al., 2012). These observations demonstrate that different stimuli can invoke different molecular mechanisms to induce  $\Delta$ FosB even within the same brain region and presumably cell type (D<sub>1</sub>-type MSNs). Further work is needed to understand the molecular basis of such stimulus-specific actions.

## 6. Target genes for $\Delta$ FosB in nucleus accumbens

Since  $\Delta$ FosB is a transcription factor, it presumably produces its interesting behavioral phenotypes in NAc by enhancing or repressing expression of other genes. As shown in Fig. 1,  $\Delta$ FosB is a truncated product of the FosB gene that lacks most of the C-terminal transactivation domain present in full-length FosB but retains the dimerization and DNA binding domains (Nestler, 2008, 2012). Some in vitro studies suggest that, because  $\Delta$ FosB lacks much of its transactivation domain, it functions as a negative regulator of AP-1 activity, while several others show that  $\Delta$ FosB can activate transcription at AP1 sites (see Nestler, 2008, 2012). An earlier study using our inducible, bitransgenic mice that overexpress  $\Delta$ FosB or its dominant negative  $\Delta$ cJun, and analyzing gene expression on Affymetrix chips, demonstrated that – in the NAc in vivo –  $\Delta$ FosB functions primarily as a transcriptional activator, while it does serve as a repressor for a smaller subset of genes (McClung and Nestler, 2003). Current research, focused on chromatin mechanisms recruited along with  $\Delta$ FosB to its target genes, is exploring the underlying molecular basis for these opposite, gene-specific actions of  $\Delta$ FosB (Nestler, 2014).

Several target genes of  $\Delta$ FosB have been established using a candidate gene approach. One target is GluA2 (GluR2), an AMPA glutamate receptor subunit.  $\Delta$ FosB overexpression in inducible bitransgenic mice selectively increases GluA2 expression in NAc, with no effect seen for several other AMPA glutamate receptor subunits analyzed (Kelz et al., 1999). AP1 complexes comprised of  $\Delta$ FosB bind a consensus AP1 site present in the GluA2 promoter. Furthermore, GluA2 overexpression in NAc via viral-mediated gene transfer promotes resilience to chronic social defeat stress (Vialou et al., 2010a), much like  $\Delta$ FosB overexpression. Since GluA2-containing AMPA channels are impermeable to Ca<sup>2+</sup> and have a lower overall conductance compared to AMPA channels that do not contain this subunit, the chronic stress- and  $\Delta$ FosB-mediated upregulation of GluA2 in NAc could account, at least in part, for the reduced glutamatergic responses seen in these neurons of resilient mice (Vialou et al., 2010a).

Another candidate target gene of  $\Delta$ FosB in NAc is the opioid peptide, dynorphin, which is suppressed by  $\Delta$ FosB (Zachariou et al., 2006). Recall that  $\Delta$ FosB is induced in the NAc of resilient mice after chronic social defeat stress specifically in D<sub>1</sub>-type MSNs, the subtype of MSNs which also express dynorphin. The suppression of dynorphin by  $\Delta$ FosB has not yet been demonstrated directly in the context of chronic stress, nevertheless, such dynorphin downregulation would be expected to produce pro-resilient- and antidepressant-like responses given the aversive actions of dynorphin acting at  $\kappa$  opioid receptors in this NAc reward circuit. Work is now needed to directly test this possibility in stress models.

We recently showed that *Camk2a* is a target gene for  $\Delta$ FosB within D<sub>1</sub>-type MSNs in drug abuse models. Chronic cocaine administration induces CaMKII in these neurons of the NAc and, interestingly, functions as part of a positive feed-forward loop, whereby CaMKII phosphorylates and stabilizes  $\Delta$ FosB (see above), promoting further CaMKII induction (Robison et al., 2013). In the context of chronic stress and chronic fluoxetine administration, however, the two proteins function very differently (Robison et al., 2014). Chronic social defeat stress does not alter  $\Delta$ FosB binding to the *Camk2a* gene



promoter, while chronic administration of fluoxetine decreases  $\Delta$ FosB binding to the promoter, despite the fact that  $\Delta$ FosB levels are increased under both of these conditions. The depletion of  $\Delta$ FosB from the *Camk2a* promoter in response to chronic fluoxetine occurs in tandem with reduced histone H3 acetylation and increased H3 methylation (at Lys9) at the *Camk2a* promoter and with suppression of *Camk2a* expression. Indeed, we know that these chromatin modifications in the NAc have broad effects on depression-related behavioral abnormalities (Covington et al., 2009, 2011). These findings suggest that specific chromatin modifications at subsets of genes are responsible for gene-specific effects of transcriptional regulation, in this case, with  $\Delta$ FosB binding to and activating *Camk2a* in D<sub>1</sub>-type MSNs after chronic cocaine, but being depleted from the *Camk2a* gene in concert with its repression within the same neuronal cell type after chronic fluoxetine. These actions of fluoxetine appear to be functional important. First, the same pattern of chromatin modifications and CaMKII repression is seen in the NAc of depressed patients treated chronically with antidepressant medications but not in medication-free individuals (Robison et al., 2014). Second, viral-mediated overexpression of CaMKII in the NAc prevents the antidepressant-like effects of fluoxetine in the chronic social defeat procedure (Robison et al., 2014). It will be interesting in future studies to identify the proteins whose reduced phosphorylation as a consequence of suppressed CaMKII levels contributes to antidepressant action.


The second approach used to identify target genes of  $\Delta$ FosB in NAc is unbiased and utilizes genome-wide methods. First, as noted earlier, we have identified genes whose expression levels are up- or down-regulated upon the inducible overexpression of  $\Delta$ FosB (or  $\Delta$ cJun) in D<sub>1</sub>-type NAc MSNs using DNA expression arrays (McClung and Nestler, 2003). More recently, we overlaid this list of  $\Delta$ FosB-regulated genes in

NAc with the list of genes that are altered in the NAc of resilient mice, based on the hypothesis, stated earlier, that  $\Delta$ FosB induction in D<sub>1</sub>-type MSNs mediates resilience. The results identify numerous putative resilience genes whose regulation might be mediated by  $\Delta$ FosB (Fig. 4). One of the genes that is most robustly induced in the NAc under resilience and upon  $\Delta$ FosB overexpression is *Sparcl1*, which encodes Sparc-like 1 (also known as hevin). Sparc1 is an anti-adhesive matrix molecule that is highly expressed in adult brain, where it localizes in the postsynaptic density and is implicated in synaptic plasticity (Lively and Brown, 2008). Indeed, based on our unbiased discovery, we went on to show that Sparc1 levels are downregulated in the NAc of depressed humans and that overexpression of Sparc1 in mouse NAc exerts potent antidepressant-like effects (Fig. 4).

These findings illustrate the potential power of open-ended approaches in driving innovative drug discovery efforts. In parallel, we are now combining such studies of gene expression with chromatin immunoprecipitation (ChIP) coupled with deep sequencing (ChIP-seq) to identify genes where the binding of endogenous  $\Delta$ FosB in NAc is regulated by chronic stress or antidepressant exposure. Together, we expect these unbiased approaches to enable the identification of a large number of previously unappreciated target genes for  $\Delta$ FosB in NAc in the context of depression-related behavioral abnormalities, as we have accomplished in drug abuse models (see Renthal et al., 2009).

## 7. Induction of $\Delta$ FosB in other brain regions

The discussion up to now has focused solely on NAc. While this is a key brain reward region and important for depression and



**Fig. 4.** Sparcl1 has pro-resilience, antidepressant-like effects in NAc. (A) Changes in gene expression observed in NAc during resilience overlap with those observed upon overexpression of  $\Delta$ FosB (comparison of data sets in McClung and Nestler, 2003 and Krishnan et al., 2007). Shown are 106 genes that are significantly regulated ( $> 1.5$  fold;  $*p < 0.05$ ) in NAc by social defeat in resilient mice as compared with controls (upper heat map) and how these genes are regulated both in resilient mice versus susceptible mice (lower heat map) and by overexpression of  $\Delta$ FosB in D<sub>1</sub>-type MSNs of NAc (middle heat map). The position of Sparc1 (SC1) on the heat maps is indicated. (B) Viral overexpression of Sparc1 in NAc reverses the social avoidance induced by chronic (10 days) social defeat. (C) Overexpression of Sparc1 has an antidepressant-like effect as measured by a decrease in time spent immobile and an increase in latency to immobility in the forced swim test. Rats were injected with HSV-GFP or HSV-Sparc1 into NAc before the test. (D) Human NAc samples from depressed individuals show a strong trend for lower Sparc1 concentrations as compared with matched controls. From Vialou et al., 2010a with permission.

antidepressant action, many other brain regions are also crucial. A central question, then, is whether  $\Delta$ FosB acting in other brain regions beyond the NAc may also influence depression-related behavioral abnormalities. Increasing evidence suggests that this is the case.

We have mapped the induction of  $\Delta$ FosB throughout the brain in response to chronic restraint stress or chronic social defeat stress and have demonstrated robust  $\Delta$ FosB induction in numerous brain regions in both stress models (Perrotti et al., 2004; Vialou et al., 2014a). We have also demonstrated  $\Delta$ FosB induction in numerous brain regions in addition to the NAc after chronic fluoxetine administration (Vialou et al., 2014b). A major goal for future research is to carry out studies, analogous to those described above for NAc, to delineate the neural and behavioral phenotype mediated by  $\Delta$ FosB for each of the brain regions implicated. This represents an enormous undertaking, yet it is crucial for understanding the global influence of  $\Delta$ FosB in depression.

An early example of this effort is a recent study where we characterized the influence of  $\Delta$ FosB, acting in the prelimbic region of medial prefrontal cortex (mPFC) in chronic social defeat stress. We had shown previously that decreased neuronal activity, as inferred from immediate early gene expression levels, in medial prefrontal cortex (mPFC) is associated with social defeat-induced depression- and anxiety-like behaviors in mice (Covington et al., 2010). More recently, we demonstrated that  $\Delta$ FosB is induced in the pre-limbic mPFC selectively in susceptible mice, and that overexpression of  $\Delta$ FosB in this region, but not in the nearby infra-limbic area, enhances stress susceptibility (Vialou et al., 2014a).  $\Delta$ FosB produces these effects partly through induction of the cholecystokinin (CCK)-B receptor: CCKB blockade in mPFC induces a resilient phenotype, whereas CCK administration into mPFC mimics the anxiogenic- and depressant-like effects of social stress. We previously found that optogenetic stimulation of mPFC neurons in susceptible mice reverses several behavioral abnormalities seen after chronic social defeat stress (Covington et al., 2010). We therefore hypothesized that optogenetic stimulation of prelimbic mPFC projections would rescue the pathological effects of CCK within this brain region. Indeed, following CCK infusion in mPFC, we optogenetically stimulated mPFC projections to NAc or basolateral amygdala. Stimulation of cortico-NAc projections reversed CCK-induced depression-like but not anxiogenic-like effects, whereas stimulation of cortico-amygdala projections blocked the anxiogenic – but not depression – like effects of CCK (Vialou et al., 2014a). Thus, the pro-susceptible role of  $\Delta$ FosB acting in the prelimbic mPFC is opposite to  $\Delta$ FosB's pro-resilience actions in the NAc, emphasizing the highly region-specific effects that should be expected from many regulatory proteins.

## 8. Future directions

Beyond characterizing the role played by  $\Delta$ FosB, acting in several regions of the brain, in mediating depression- and antidepressant responses, an important question is how this information can be mined to improve clinical management of depression and related disorders. We believe that work on  $\Delta$ FosB can contribute to such clinical efforts. First, it would be interesting to develop imaging ligands for  $\Delta$ FosB, which might be used in conjunction with PET or MRI, for example, to determine an individual's susceptibility to depression and to track antidepressant responses based on levels of  $\Delta$ FosB induction throughout the brain. As well, we are interested in developing small molecules that directly inhibit or potentiate the actions of  $\Delta$ FosB. We have made early progress in this regard (Wang et al., 2012). As test compounds of higher affinity and greater brain penetration are

generated, it would be useful to administer such molecules systemically to determine the net effect of increasing or decreasing  $\Delta$ FosB activity throughout the brain. One striking feature of  $\Delta$ FosB is that it is expressed at many fold higher levels in NAc and dorsal striatum than any other brain region or peripheral tissue studied; we would thus expect effects in NAc to predominate. Moreover, studies of  $\Delta$ FosB are revealing numerous target genes whose encoded proteins show greater region-specific regulation by stress and antidepressants than that exhibited by  $\Delta$ FosB itself. Such targets should be used to help drive antidepressant drug discovery efforts in future years. Finally, studies of  $\Delta$ FosB illustrate the ways in which it is possible to elaborate detailed transcriptional mechanisms of depression and antidepressant action.

## Acknowledgments

Preparation of this review was supported by grants from the National Institute of Mental Health (R01MH51399 and P50MH096890).

## References

- Been, L.E., Hedges, V.L., Vialou, V., Nestler, E.J., Meisel, R.L., 2013. Delta JunD expression in the nucleus accumbens prevents sexual reward in female Syrian hamsters. *Genes Brain Behav.* 12, 666–672.
- Carle, T.L., Ohnishi, Y.N., Ohnishi, Y.H., Alibhai, I.N., Wilkinson, M.B., Kumar, A., Nestler, E.J., 2007. Absence of conserved C-terminal degran domain contributes to  $\Delta$ FosB's unique stability. *Eur. J. Neurosci.* 25, 3009–3019.
- Carlezon Jr., W.A., Duman, R.S., Nestler, E.J., 2005. The many faces of CREB. *Trends Neurosci.* 28, 436–445.
- Cates, H., Thibault, M., Pfau, M., Heller, E., Eagle, A., Gajewski, P., Bagot, R., Colangelo, C., Abbott, T., Rudenko, G., Neve, R., Nestler, E.J., Robison, A.J., 2014. Threonine 149 phosphorylation enhances  $\Delta$ FosB transcriptional activity to control psychomotor responses to cocaine. *J. Neurosci.* 34, 11461–11469.
- Chen, J.S., Nye, H.E., Kelz, M.B., Hiroi, N., Nakabeppu, Y., Hope, B.T., Nestler, E.J., 1995. Regulation of  $\Delta$ FosB and FosB-like proteins by electroconvulsive seizure (ECS) and cocaine treatments. *Mol. Pharmacol.* 48, 880–889.
- Chen, J., Kelz, M.B., Hope, B.T., Nakabeppu, Y., Nestler, E.J., 1997. Chronic FRAs: stable variants of  $\Delta$ FosB induced in brain by chronic treatments. *J. Neurosci.* 17, 4933–4941.
- Christoffel, D.J., Golden, S.A., Dumitriu, D., Robison, A.J., Janssen, W.G., Ahn, H.F., Krishnan, V., Reyes, C.M., Han, M.H., Ables, J.L., Eisch, A.J., Dietz, D.M., Ferguson, D., Neve, R.L., Greengard, P., Kim, Y., Morrison, J.H., Russo, S.J., 2011. Ikb kinase regulates social defeat stress-induced synaptic and behavioral plasticity. *J. Neurosci.* 31, 314–321.
- Covington III, H.E., Maze, I., LaPlant, Q.C., Vialou, V.F., Yoshinori, O.N., Berton, O., Fass, D.M., Renthal, W., Rush III, A.J., Wu, E.Y., Ghose, S., Krishnan, V., Russo, S.J., Tamminga, C., Haggarty, S.J., Nestler, E.J., 2009. Antidepressant actions of HDAC inhibitors. *J. Neurosci.* 29, 11451–11460.
- Covington III, H.E., Lobo, M.K., Maze, I., Vialou, V., Hyman, J.M., Zaman, S., LaPlant, Q., Mouzon, E., Ghose, S., Tamminga, C.A., Neve, R.L., Deisseroth, K., Nestler, E.J., 2010. Antidepressant effect of optogenetic stimulation of the medial prefrontal cortex. *J. Neurosci.* 30, 16082–16090.
- Covington III, H.E., Maze, I., Sun, H.S., Wu, E.Y., Dietz, D., Lobo, M.K., Ghose, S., Neve, R., Tamminga, C.A., Nestler, E.J., 2011. A role for repressive histone methylation in cocaine-induced vulnerability to stress. *Neuron* 71, 656–670.
- Donahue, R.J., Muschamp, J.W., Russo, S.R., Nestler, E.J., Carlezon, W.A., 2014. Effects of striatal  $\Delta$ FosB overexpression and ketamine on social defeat stress-induced anhedonia in mice. *Biol. Psychiatry* (In press).
- Grueter, B.A., Robison, A.J., Neve, R.L., Nestler, E.J., Malenka, R.C., 2013.  $\Delta$ FosB differentially modulates nucleus accumbens direct and indirect pathway function. *Proc. Natl. Acad. Sci. USA* 110, 1923–1928.
- Hedges, V.L., Chakravarty, S., Nestler, E.J., Meisel, R.L., 2009.  $\Delta$ FosB overexpression in the nucleus accumbens enhances sexual reward in female Syrian hamsters. *Genes Brain Behav.* 8, 442–449.
- Hiroi, N., Brown, J., Ye, H., Saudou, F., Vaidya, V.A., Duman, R.S., Greenberg, M.E., Nestler, E.J., 1998. Essential role of the fosB gene in molecular, cellular, and behavioral actions of electroconvulsive seizures. *J. Neurosci.* 18, 6952–6962.
- Holsboer, F., Ising, M., 2010. Stress hormone regulation: biological role and translation into therapy. *Annu. Rev. Psychol.* 61, 81–109.
- Jorissen, H., Ulery, P., Henry, L., Gourneni, S., Nestler, E.J., Rudenko, G., 2007. Dimerization and DNA-binding properties of the transcription factor  $\Delta$ FosB. *Biochemistry* 46, 8360–8372.
- Kelz, M.B., Chen, J.S., Carlezon, W.A., Whisler, K., Gilden, L., Beckmann, A.M., Steffen, C., Zhang, Y.-J., Marotti, L., Self, D.W., Tkatch, R., Baranaukas, G., Surmeier, D.J., Neve, R.L., Duman, R.S., Picciotto, M.R., Nestler, E.J., 1999. Expression of the transcription factor  $\Delta$ FosB in the brain controls sensitivity to cocaine. *Nature* 401, 272–276.

- Krishnan, V., Han, M.-H., Graham, D.L., Berton, O., Renthal, W., Russo, S.J., LaPlant, Q., Graham, A., Lutter, M., Lagace, D.C., Ghose, S., Reister, R., Tannous, P., Green, T.A., Neve, R.L., Chakravarty, S., Kumar, A., Eisch, A.J., Self, D.W., Lee, F.S., Tammimga, C.A., Cooper, D.C., Gershenfeld, H.K., Nestler, E.J., 2007. Susceptibility and resistance to social defeat are mediated through molecular adaptations in brain reward regions. *Cell* 131, 391–404.
- Krishnan, V., Nestler, E.J., 2010. Linking molecules to mood: new insight into the biology of depression. *Am. J. Psychiatry* 167, 1305–1320.
- Lehmann, M.L., Herkenham, M., 2011. Environmental enrichment confers stress resiliency to social defeat through an infralimbic cortex-dependent neuroanatomical pathway. *J. Neurosci.* 31, 6159–6173.
- Licznerski, P., Duman, R.S., 2013. Remodeling of axo-spinous synapses in the pathophysiology and treatment of depression. *Neuroscience* 251, 33–50.
- Lively, S., Brown, I.R., 2008. The extracellular matrix protein SC1/hevin localizes to excitatory synapses following status epilepticus in the rat lithium-pilocarpine seizure model. *J. Neurosci. Res.* 86, 2895–2905.
- Lobo, M.K., Zaman, S., Damez-Werno, D.M., Koo, J.W., Bagot, R.C., DiNieri, J.A., Nugent, A., Finkel, E., Chaudhury, D., Chandra, R., Ribeiro, E., Rabkin, J., Mouzon, E., Cachope, R., Cheer, J.F., Han, M.H., Dietz, D.M., Self, D.W., Hurd, Y.L., Vialou, V., Nestler, E.J., 2013.  $\Delta$ FosB induction in striatal medium spiny neuron subtypes in response to chronic pharmacological, emotional, and optogenetic stimuli. *J. Neurosci.* 33, 18381–18395.
- McClung, C.A., Nestler, E.J., 2003. Regulation of gene expression and cocaine reward by CREB and  $\Delta$ FosB. *Nat. Neurosci.* 11, 1208–1215.
- Morgan, J.I., Curran, T., 1995. Immediate-early genes: ten years on. *Trends Neurosci.* 18, 66–67.
- Muschamp, J.W., Nemeth, C.L., Robison, A.J., Nestler, E.J., Carlezon Jr., W.A., 2012.  $\Delta$ FosB enhances the rewarding effects of cocaine while reducing the pro-depressive effects of the kappa-opioid agonist U50488. *Biol. Psychiatry* 71, 44–50.
- Nestler, E.J., 2008. Transcriptional mechanisms of addiction: role of deltaFosB. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 363, 3245–3255.
- Nestler, E.J., 2012.  $\Delta$ FosB: a molecular switch for reward. *J. Drug Alcohol Res.* 2, 235651.
- Nestler, E.J., 2014. Epigenetic mechanisms of drug addiction. *Neuropharmacology* 76, 259–268.
- Nestler, E.J., Barrot, M., DiLeone, R.J., Eisch, A.J., Gold, S.J., Monteggia, L.M., 2002. Neurobiology of depression. *Neuron* 34, 13–25.
- Ohnishi, Y.N., Ohnishi, Y.H., Hokama, M., Nomaru, H., Yamazaki, K., Tominaga, Y., Sakumi, K., Nestler, E.J., Nakabeppu, Y., 2011. FosB is essential for the enhancement of stress tolerance and antagonizes locomotor sensitization by  $\Delta$ FosB. *Biol. Psychiatry* 70, 487–495.
- Ohnishi, Y.N., Ohnishi, Y.H., Vialou, V., Mouzon, E., LaPlant, Q., Nishi, A., Nestler, E.J., 2014. Functional role of the N-terminal domain of  $\Delta$ FosB in response to stress and drugs of abuse. *Neuroscience* (submitted).
- Olausson, P., Jentsch, J.D., Tronson, N., Neve, R., Nestler, E.J., Taylor, J.R., 2006.  $\Delta$ FosB in the nucleus accumbens regulates food-reinforced instrumental behavior and motivation. *J. Neurosci.* 26, 9196–9204.
- Peakman, M.C., Colby, C., Perrotti, L.I., Tekumalla, P., Carle, T., Ulery, P., Chao, J., Duman, C., Steffen, C., Monteggia, L., Allen, M.R., Stock, J.L., Duman, R.S., McNeish, J.D., Barrot, M., Self, D.W., Nestler, E.J., Schaeffer, E., 2003. Inducible, brain region specific expression of a dominant negative mutant of c-Jun in transgenic mice decreases sensitivity to cocaine. *Brain Res.* 970, 73–86.
- Perrotti, L.I., Hadeishi, Y., Ulery, P., Barrot, M., Monteggia, L., Duman, R.S., Nestler, E.J., 2004. Induction of  $\Delta$ FosB in reward-related brain regions after chronic stress. *J. Neurosci.* 24, 10594–10602.
- Pitchers, K.K., Frohmader, K.S., Vialou, V., Mouzon, E., Nestler, E.J., Lehman, M.N., Coolen, L.M., 2010.  $\Delta$ FosB in the nucleus accumbens is critical for reinforcing effects of sexual reward. *Genes Brain Behav.* 9, 831–840.
- Pitchers, K.K., Vialou, V., Nestler, E.J., Lehman, M.N., Coolen, L.M., 2013. Sexual experience increases amphetamine reward and nucleus accumbens spinogenesis via dopamine D1 receptor activity and induction of deltaFosB. *J. Neurosci.* 33, 3434–3442.
- Renthal, W., Kumar, A., Xiao, G.H., Wilkinson, M., Covington III, H.E., Maze, I., Sikder, D., Robison, A.J., LaPlant, Q., Dietz, D.M., Russo, S.J., Vialou, V., Chakravarty, S., Kodadek, E.J., Stack, A., Kabbaj, M., Nestler, E.J., 2009. Genome wide analysis of chromatin regulation by cocaine reveals a novel role for sirtuins. *Neuron* 62, 335–348.
- Robison, A.J., Vialou, V., Mazei-Robison, M., Feng, J., Kourrich, S., Collins, M., Wee, S.M., Koob, G., Turecki, G., Neve, R., Thomas, M., Nestler, E.J., 2013. Behavioral and structural responses to chronic cocaine require a feed-forward loop involving  $\Delta$ FosB and CaMKII in the nucleus accumbens shell. *J. Neurosci.* 33, 4295–4307.
- Robison, A.J., Vialou, V., Sun, H.S., Labonte, B., Golden, S., Dias, C., Turecki, G., Tammimga, C., Russo, S.J., Mazei-Robison, M., Nestler, E.J., 2014. Fluoxetine epigenetically alters the CaMKII $\alpha$  promoter in nucleus accumbens to regulate  $\Delta$ FosB binding and antidepressant effects. *Neuropsychopharmacology* 39, 1178–1186.
- Teegarden, S.L., Nestler, E.J., Bale, T.L., 2008. Delta FosB-mediated alterations in dopamine signaling are normalized by a palatable high-fat diet. *Biol. Psychiatry* 64, 941–950.
- Ulery, P.G., Rudenko, G., Nestler, E.J., 2006. Regulation of  $\Delta$ FosB stability by phosphorylation. *J. Neurosci.* 26, 5131–5142.
- Ulery-Reynolds, P.G., Castillo, M.A., Vialou, V., Russo, S.J., Nestler, E.J., 2009. Phosphorylation of  $\Delta$ FosB mediates its stability in vivo. *Neuroscience* 158, 369–372.
- Vialou, V., Robison, A.J., LaPlant, Q.C., Covington III, H.E., Dietz, D.M., Ohnishi, Y.N., Mouzon, E., Rush III, A.J., Watts, E.L., Wallace, D.L., Iñiguez, S.D., Ohnishi, Y.H., Steiner, M.A., Warren, B., Krishnan, V., Neve, R.L., Ghose, S., Berton, O., Tammimga, C.A., Nestler, E.J., 2010a.  $\Delta$ FosB in brain reward circuits mediates resilience to stress and antidepressant responses. *Nat. Neurosci.* 13, 745–752.
- Vialou, V., Maze, I., Renthal, W., LaPlant, Q.C., Watts, E.L., Mouzon, E., Ghose, S., Tammimga, C.A., Nestler, E.J., 2010b. Serum response factor promotes resilience to chronic social stress through the induction of  $\Delta$ FosB. *J. Neurosci.* 30, 14585–14592.
- Vialou, V., Feng, J., Robison, A.J., Ku, S.M., Ferguson, D., Scobie, K.N., Mazei-Robison, M.A., Mouzon, E., Nestler, E.J., 2012. Serum response factor and cAMP response element binding protein are both required for cocaine induction of  $\Delta$ FosB. *J. Neurosci.* 32, 7577–7584.
- Vialou, V., Bagot, R.C., Cahill, M.E., Ferguson, D., Robison, A.J., Dietz, D.M., Fallon, B., Mazei-Robison, M., Ku, S.M., Harrigan, E., Winstanley, C.A., Joshi, T., Feng, J., Berton, O., Nestler, E.J., 2014a. Prefrontal cortical circuit for depression- and anxiety-related behaviors mediated by cholecystokinin: role of  $\Delta$ FosB. *J. Neurosci.* 34, 3878–3887.
- Vialou, V., Thibault, M., Kaska, S., Gajewski, P., Eagle, A., Mazei-Robison, M., Nestler, E.J., Robison, A.J., 2014b. Differential induction of FosB isoforms throughout the brain by fluoxetine and chronic stress. *Neuropharmacology* (In press).
- Wallace, D.L., Vialou, V., Rios, L., Carle-Florence, T.L., Chakravarty, S., Kumar, A., Graham, D.L., Green, T.A., Kirk, A., Iniguez, S.D., Perrotti, L.I., Barrot, M., DiLeone, R.J., Nestler, E.J., Bolaños, C.A., 2008. The influence of  $\Delta$ FosB in the nucleus accumbens on natural reward-related behavior. *J. Neurosci.* 28, 10272–10277.
- Wang, Y., Cesena, T., Ohnishi, Y., Burger-Caplan, R., Vivian, L., Kirchhoff, P., Scott, L., Larsen, M., Nestler, E.J., Rudenko, G., 2012. Small molecule screening identifies regulators of the transcription factor DeltaFosB. *ACS Chem. Neurosci.* 3, 546–556.
- Werme, M., Messer, C., Olson, L., Gilden, L., Thorén, P., Nestler, E.J., Brené, S., 2002.  $\Delta$ FosB regulates wheel running. *J. Neurosci.* 22, 8133–8138.
- Winstanley, C.A., LaPlant, Q., Theobald, D.E.H., Green, T.A., Bachtell, R.K., Perrotti, L.I., DiLeone, R.J., Russo, S.J., Garth, W.J., Self, D.W., Nestler, E.J., 2007.  $\Delta$ FosB induction in orbitofrontal cortex mediates tolerance to cocaine-induced cognitive dysfunction. *J. Neurosci.* 27, 10497–10507.
- Zachariou, V., Bolanos, C.A., Selley, D.E., Theobald, D., Cassidy, M.P., Kelz, M.B., Shaw-Lutchmann, T., Berton, O., Sim-Selley, L.J., DiLeone, R.J., Kumar, A., Nestler, E.J., 2006.  $\Delta$ FosB: an essential role for  $\Delta$ FosB in the nucleus accumbens in morphine action. *Nat. Neurosci.* 9, 205–211.
- Zhang, Y.F., Crofton, E.J., Li, D., Lobo, M.K., Fan, X.Z., Nestler, E.J., Green, T.A., 2014. Overexpression of DeltaFosB in nucleus accumbens mimics the protective addiction phenotype, but not the protective depression phenotype, of environmental enrichment. *Front. Behav. Neurosci.* 8, 297.