

Self-Interacting Dynamics with Exponential Memory: Markovian Embedding, Contractive Memory Fibres, and Metastability

H. Horn

Peine Stederdorf, Germany

(Dated: 25 June 2026)

Abstract

We formulate a minimal class of self-interacting stochastic processes in which a visible state x_n is coupled to an explicit memory field ρ_n with exponential forgetting. The visible trajectory is generally non-Markovian: the next-step law depends on the retained memory field, not on x_n alone. The augmented state $z_n = (x_n, \rho_n)$, however, is a time-homogeneous Markov process. We derive the exact exponential memory expansion, prove pathwise contraction of the affine memory fibre along a common visible trajectory, and represent the dynamics by a Markov kernel and its positive unital observable operator. We also describe the associated random skew-product structure and a formal continuum approximation. Metastable “knots” are treated operationally as long-lived regions or slow modes diagnosed by residence times, retained memory weight, autocorrelation, local Ornstein–Uhlenbeck linearization, and finite-rank transfer-operator estimates. Claims about space-time emergence, Lorentz kinematics, quantum dynamics, Standard-Model structure, physical masses, or fundamental constants are outside the scope of this anchor paper and are listed only as future-work directions.

I. INTRODUCTION

Self-interacting stochastic processes provide a natural mathematical language for systems whose future evolution is shaped by their own past. Reinforced random walks, self-interacting diffusions, true self-avoiding walks, and self-repelling Brownian polymers all instantiate the general pattern

trajectory $\longrightarrow$ occupation history $\longrightarrow$ self-induced drift.

Classical examples include self-interacting diffusions [1], reinforced random processes [2], Brownian polymers [3], true self-repelling motion [4], and true self-avoiding walks [5, 6].

The present paper isolates a particularly simple variant: the retained history is not the full local time or a Cesaro-averaged empirical measure, but an explicit exponentially relaxing field. This produces a finite effective memory scale and a direct Markovian embedding. The construction is also close in spirit to Markovian embeddings of memory equations, such as auxiliary variable treatments of generalized Langevin dynamics [7–9]; the difference is that the memory variable here is a self-induced spatial field generat-

ing the drift.

The purpose of this paper is deliberately narrow. We do not attempt to derive spacetime, relativistic kinematics, quantum mechanics, particle spectra, or physical masses. Instead, we establish the model’s mathematical anchor:

define the model
 $\implies$ prove the Markov embedding
 $\implies$ make metastability measurable.

The result is a conservative framework in which later emergence claims can be posed as separate conjectures or companion papers rather than as assumptions inside the anchor model.

II. DISCRETE MODEL

Let $\Omega \subseteq \mathbb{R}^d$ be an effective state space with reference measure dx . The Euclidean structure is used only to write kernels, gradients, and diagnostics; it is not identified with physical space. Let G_σ be a smooth normalized deposition kernel of width σ , and let K be an interaction kernel for which the convolu-

tion

$$(K * \rho)(x) = \int_{\Omega} K(x, y) \rho(y) dy$$

is differentiable in the regimes considered below.

Assumption 1 (Baseline regularity). *For the structural statements in Sections III–IV, it is enough that G_{σ} is a normalized nonnegative density and that the update equations are well-defined. For continuum approximations one should assume enough smoothness and growth control on K and ∇K to make the drift locally Lipschitz or otherwise well posed in the chosen function space.*

The discrete process is

$$x_{n+1} = x_n + \varepsilon \xi_n - \eta \nabla \Phi_n(x_n), \quad (1)$$

$$\rho_{n+1}(x) = (1 - \alpha) \rho_n(x) + \alpha G_{\sigma}(x - x_{n+1}), \quad (2)$$

$$\Phi_n(x) = (K * \rho_n)(x), \quad (3)$$

where $0 < \alpha < 1$, $\varepsilon \geq 0$, $\eta \geq 0$, and ξ_n are independent centered noise variables, typically taken isotropic with unit covariance. If ρ_0 and G_{σ} have unit mass, then every ρ_n has unit mass. If they are nonnegative, every ρ_n is nonnegative. This normalization is a modelling convention; one could equiva-

lently work with finite positive measures and absorb an overall memory scale into η .

The project terminology calls long-lived localized feedback regimes “knots.” In this paper the word has only an operational meaning: a knot is a metastable region or slow mode of the augmented dynamics, not a fundamental particle and not an exact fixed point.

III. EXPONENTIAL MEMORY AND CONTRACTIVE FIBRES

The memory equation can be unrolled exactly.

Proposition 1 (Exponential memory representation). *For $n \geq 1$,*

$$\rho_n = (1 - \alpha)^n \rho_0 + \alpha \sum_{j=0}^{n-1} (1 - \alpha)^j G_\sigma(\cdot - x_{n-j}). \quad (4)$$

Proof. The claim follows by induction. For $n = 1$ it is exactly Eq. (2). If it holds at n , then

$$\rho_{n+1} = (1 - \alpha)^{n+1} \rho_0 + \alpha \sum_{j=1}^n (1 - \alpha)^j G_\sigma(\cdot - x_{n+1-j}) + \alpha G_\sigma(\cdot - x_{n+1}).$$

which is Eq. (4) with $n + 1$. □

The weights are

$$w_j = \alpha(1 - \alpha)^j. \quad (5)$$

For a rest-weight threshold δ , the effective memory depth satisfies

$$j_\delta = \frac{\log \delta}{\log(1 - \alpha)} \sim \frac{|\log \delta|}{\alpha} \quad (\alpha \ll 1). \quad (6)$$

Thus α^{-1} is the natural update-count scale for memory persistence. Figure 1 shows the exponential weights for representative values of α .

Proposition 2 (Pathwise memory-fibre contraction). *Fix a common visible trajectory $x_1, \dots, x_n$. Let ρ_k and $\tilde{\rho}_k$ solve Eq. (2) with the same deposited kernels but different initial memory fields. Then*

$$\rho_n - \tilde{\rho}_n = (1 - \alpha)^n(\rho_0 - \tilde{\rho}_0). \quad (7)$$

Consequently, for any homogeneous norm on the memory space,

$$\|\rho_n - \tilde{\rho}_n\| = (1 - \alpha)^n \|\rho_0 - \tilde{\rho}_0\|. \quad (8)$$




FIG. 1. Exponential memory weights $w_j = \alpha(1 - \alpha)^j$. Vertical guide lines mark $j = \alpha^{-1}$ for each curve.

Proof. Subtract the two memory recursions. The deposited kernels cancel because the visible trajectory is common:

$$\rho_{k+1} - \tilde{\rho}_{k+1} = (1 - \alpha)(\rho_k - \tilde{\rho}_k).$$

Iterating gives Eq. (7). □

This contraction is a statement about the memory fibre under a fixed visible path. It is not yet a global contraction theorem for the full feedback system, because different memory fields can alter future positions through Eq. (1). A full contraction analysis

must control the coupled pair (x_n, ρ_n) .

IV. MARKOVIAN EMBEDDING

Define the augmented state

$$z_n = (x_n, \rho_n) \in \Sigma. \quad (9)$$

Given z_n and a fresh noise draw ξ_n , Eqs. (1) and (2) determine z_{n+1} . Thus the augmented dynamics is a time-homogeneous Markov process.

Definition 1 (Transition kernel). *For measurable $A \subseteq \Sigma$ define*

$$P(z, A) = \mathbb{P}\{z_{n+1} \in A \mid z_n = z\}. \quad (10)$$

The observable Markov operator is

$$(Uf)(z) = \int_{\Sigma} f(z') P(z, dz') = \mathbb{E}[f(z_{n+1}) \mid z_n = z]. \quad (11)$$

The operator U is linear, positive, and unital: $U1 = 1$. Its iterates satisfy $U^{m+n} = U^m U^n$. In the stochastic case it is generally not multiplicative: $U(fg)$ need not equal $(Uf)(Ug)$. The correct alge-

braic object is therefore a positive unital Markov operator on observables, with a dual action on probability measures, rather than a deterministic algebra automorphism.

Proposition 3 (Visible non-Markovian projection). *The projected process x_n is generally not Markov.*

Proof. Take the same visible position x with two memory fields ρ and $\tilde{\rho}$ such that

$$\nabla(K * \rho)(x) \neq \nabla(K * \tilde{\rho})(x).$$

Then the conditional next-step laws of x_{n+1} differ because their drift terms in Eq. (1) differ. Hence conditioning on $x_n = x$ alone does not determine the next-step law. $\square$

The memory update should not be described as algebraically noninvertible in isolation. If x_{n+1} and ρ_{n+1} are known and $\alpha \neq 1$, then

$$\rho_n = \frac{\rho_{n+1} - \alpha G_\sigma(\cdot - x_{n+1})}{1 - \alpha}$$

is a formal rearrangement. The precise irreversibility statement is different: the present augmented

state does not generally encode the full ordered history, and the stochastic dynamics defines a forward Markov semigroup rather than a reversible group action.

V. RANDOM SKEW-PRODUCT STRUCTURE

For each noise realization, write the update as

$$x_{n+1} = F_{\rho_n}(x_n, \xi_n), \quad (12)$$

$$\rho_{n+1} = T_{x_{n+1}}(\rho_n), \quad (13)$$

where

$$T_x(\rho) = (1 - \alpha)\rho + \alpha G_\sigma(\cdot - x).$$

Thus the system is a self-coupled random affine skew product with an exponentially contractive memory fibre along fixed visible paths. For a noise sequence $\omega = (\xi_0, \xi_1, \dots)$ one obtains the random cocycle

$$\varphi(n, \omega, z_0) = F_{\xi_{n-1}(\omega)} \circ \dots \circ F_{\xi_0(\omega)}(z_0),$$

in the standard sense of random dynamical systems [10]. This viewpoint is useful for discussing ran-

dom attractors, path dependence, bifurcations, and metastability under noise, but the present paper uses it only as structural orientation.

VI. FORMAL CONTINUUM APPROXIMATION

Let $t = \alpha n$ and consider a joint scaling

$$\alpha \rightarrow 0, \quad \varepsilon \rightarrow 0, \quad \frac{\varepsilon^2}{2\alpha} \rightarrow D, \quad \frac{\eta}{\alpha} \rightarrow \gamma.$$

Formally, the visible coordinate approaches

$$dX_t = -\gamma \nabla (K * \rho_t)(X_t) dt + \sqrt{2D} dW_t, \quad (14)$$

while the memory field satisfies

$$\partial_t \rho_t = G_\sigma(\cdot - X_t) - \rho_t. \quad (15)$$

More generally one may introduce a memory time τ and write

$$\partial_t \rho_t = \tau^{-1} (G_\sigma(\cdot - X_t) - \rho_t),$$

equivalently

$$\rho_t = \tau^{-1} \int_{-\infty}^t e^{-(t-s)/\tau} G_\sigma(\cdot - X_s) ds$$

after transients. The extended process (X_t, ρ_t) is local in the scaled time variable and Markovian; the visible process X_t alone retains memory.

Equations (14)–(15) are a formal effective description, not a convergence theorem in this paper. A rigorous limit would require specifying topology, tightness estimates, regularity of K , boundary conditions, and the scaling of Ω if the domain changes with α .

VII. METASTABILITY AND KNOT DIAGNOSTICS

A candidate knot is a long-lived localized feedback regime in the augmented dynamics. Operationally, for a region $\mathcal{K} \subseteq \Omega$ one can track retained memory mass

$$M_n(\mathcal{K}) = \int_{\mathcal{K}} \rho_n(x) dx. \quad (16)$$

Persistence over many memory times,

$$\Delta n \gg \alpha^{-1},$$

is a minimal diagnostic for metastability, not a proof of a new particle-like object.

Several diagnostics are useful and mutually constraining:

- residence-time statistics of x_n in candidate regions;
- retained memory weight $M_n(\mathcal{K})$;
- autocorrelation functions of positions and reduced memory features;
- local linearization near a metastable center x_* ;
- slow modes of a transfer-operator estimate on augmented features.

For local linearization, suppose a metastable regime has slowly varying memory ρ and a center x_* with $\nabla(K * \rho)(x_*) = 0$. With

$$H = \nabla \nabla (K * \rho)(x_*)$$

the continuum approximation becomes locally

$$dX_t = -\gamma H(X_t - x_*) dt + \sqrt{2D} dW_t. \quad (17)$$

Positive eigenvalues of γH give local relaxation rates in stable directions. A relaxation-based stability scale can be written as

$$\Gamma_{\text{rel}} = \tau_{\text{rel}}^{-1}, \quad \hat{\Gamma} = \frac{\sigma^2}{D} \Gamma_{\text{rel}}.$$

This is a mass-like or stability proxy only. It is not a calibrated physical mass and should not be identified with mc^2 .

A. Transfer-Operator View

Let $\psi(z_n)$ be a reduced feature vector for the augmented state, for example position, memory centroid, current offset from the memory centroid, memory spread, retained memory mass, or learned/PCA/Fourier features of ρ_n . A finite-rank Ulam or Markov-state approximation estimates a transition matrix at lag ℓ from pairs

$$(\psi(z_n), \psi(z_{n+\ell})).$$

Slow eigenvalues λ_i give implied rates

$$\Gamma_i(\ell) = -\frac{1}{\ell} \log |\lambda_i|. \quad (18)$$

Almost-invariant sets and coherent sets are then candidate metastable knots [11]. Related variational and Koopman approaches are standard in molecular kinetics and operator-theoretic model reduction [12–15].

Figure 2 shows one small reproducible transfer spectrum from the companion pipeline. It is an illustration of the diagnostic workflow, not evidence for a universal phase diagram.

VIII. REPRODUCIBLE NUMERICAL PIPELINE

The repository includes a minimal Paper-0 pipeline with the following steps:

1. run the finite-memory simulation over grids of noise, coupling, memory, kernel scales, kernel amplitudes, and dimension;
2. record reduced augmented-state features $\psi(z_n) = (x_n, \text{summary}(\rho_n))$ at fixed sampling




FIG. 2. Example eigenvalues of a finite-state transfer estimate from reduced augmented features. Slow modes near the unit circle are candidates for metastable structure, subject to lag, discretization, and sampling checks.

intervals;

3. compute residence statistics, covariance/occupancy diagnostics, and autocorrelation functions;
4. estimate a finite transition matrix on augmented features at lag ℓ ;
5. report leading eigenvalues and implied rates $-\ell^{-1} \log |\lambda_i|$;

6. validate candidate metastable sets against residence statistics and autocorrelation decay.

The reduced representation of ρ_n is intentionally labelled as a projection. It is a numerical approximation to the augmented state, not an exact replacement for the field. The point is to make metastability testable without storing a full field at every update.

IX. RELATION TO EXISTING LITERATURE

The closest mathematical family is self-interacting diffusion, where the drift depends on occupation-type memory variables [1]. The main difference is the explicit exponential forgetting in Eq. (2). Reinforced random walks [2], true self-avoiding walks [5], self-repelling motion [4], and Brownian polymers [3] share the feedback pattern but often emphasize repulsion or full/local-time memory.

Generalized Langevin theory and Mori–Zwanzig style embeddings [7, 8] provide the broader idea that nonlocal memory can be made local by adding variables. In the present model the added variable is not an auxiliary friction coordinate but the self-induced memory field itself.

Metastability is naturally formulated through transfer operators, Markov-state models, and Koopman approximations [12–15]. Large-deviation and noise-induced transition theory [16, 17] provides a separate asymptotic language for rare transitions once a small-noise regime has been specified.

X. OPEN MATHEMATICAL PROBLEMS

The structural results above do not settle the main hard questions. The following are open tasks:

- well-posedness of the continuum approximation under explicit kernel and state-space assumptions;
- Feller properties and invariant measures for the augmented Markov process;
- Lyapunov/Harris drift conditions of the form $PV \leq \lambda V + b$ outside compact sets [18];
- conditions for uniqueness, geometric ergodicity, and spectral gaps;

- coupled contraction estimates for (x_n, ρ_n) , not only the memory fibre under a fixed path;
- bifurcation and metastability regimes for attractive, repulsive, and two-scale attractive-repulsive kernels;
- convergence and bias analysis for transfer-operator estimates on reduced memory features.

XI. BOUNDARIES AND FUTURE WORK

The following statements are not claims of this paper:

- that dimension $d = 3$ is selected by the model;
- that exponential memory alone implies a hard finite signal speed;
- that Lorentz invariance follows from the present equations;
- that quantum mechanics, $\hbar$, c , spin, gauge symmetry, or Standard-Model structure emerges here;

- that metastable knots are real particles;
- that relaxation rates are calibrated physical masses.

These questions may be posed later as falsifiable companion programs. A hard finite propagation bound, for example, would require additional assumptions such as finite interaction range, local multi-step transmission, and no direct long-range coupling term. Dimension-selection claims require controlled parameter sweeps, normalized kernel scaling, multiple seeds, and negative controls.

XII. CONCLUSION

The model defines a self-interacting stochastic process with exponential forgetting and a field-mediated self-potential. The visible trajectory is generally non-Markovian, but the explicit memory field yields a Markovian augmented dynamics. The memory fibre is affine and pathwise contractive for a common visible trajectory, while feedback through the self-induced potential can support nontrivial metastable regimes. This is the mathematical anchor needed before stronger emergence claims can

be formulated and tested.

- [1] M. Benaïm and O. Raimond, Probability Theory and Related Fields **122**, 1 (2002).
- [2] R. Pemantle, Probability Surveys **4**, 1 (2007).
- [3] R. T. Durrett and L. C. G. Rogers, Probability Theory and Related Fields **92**, 337 (1992).
- [4] B. Tóth and W. Werner, Probability Theory and Related Fields **111**, 375 (1998).
- [5] D. J. Amit, G. Parisi, and L. Peliti, Physical Review B **27**, 1635 (1983).
- [6] L. Peliti and L. Pietronero, La Rivista del Nuovo Cimento **10**, 1 (1987).
- [7] H. Mori, Progress of Theoretical Physics **33**, 423 (1965).
- [8] R. Zwanzig, *Nonequilibrium Statistical Mechanics* (Oxford University Press, 2001).
- [9] D. Jaganathan and R. N. Valani, Communications in Nonlinear Science and Numerical Simulation 10.1016/j.cnsns.2025.109540 (2025), arXiv:2402.00009.

- [10] L. Arnold, *Random Dynamical Systems* (Springer, 1998).
- [11] G. Froyland, S. Lloyd, and N. Santitissadeekorn, *Physica D* **239**, 1527 (2010).
- [12] F. Noé and F. Nüske, *Multiscale Modeling & Simulation* **11**, 635 (2013).
- [13] F. Noé, H. Wu, J.-H. Prinz, and N. Plattner, *The Journal of Chemical Physics* **139**, 184114 (2013).
- [14] M. O. Williams, I. G. Kevrekidis, and C. W. Rowley, *Journal of Nonlinear Science* **25**, 1307 (2015).
- [15] S. Klus, F. Nüske, P. Koltai, H. Wu, I. Kevrekidis, C. Schütte, and F. Noé, *Journal of Nonlinear Science* **28**, 985 (2018).
- [16] M. Freidlin and A. Wentzell, *Random Perturbations of Dynamical Systems*, 3rd ed. (Springer, 2012).
- [17] H. A. Kramers, *Physica* **7**, 284 (1940).
- [18] S. Meyn and R. L. Tweedie, *Markov Chains and Stochastic Stability*, 2nd ed. (Cambridge University Press, 2009).