

Solutions to Algebraic Curves

Meng-Gen Tsai
plover@gmail.com

March 10, 2021

Contents

| | |
|--|----------|
| Chapter 1: Affine Algebraic Sets | 2 |
| 1.1. Algebraic Preliminaries | 2 |
| Problem 1.1.* | 2 |
| Problem 1.5.* | 3 |
| Problem 1.6.* | 3 |
| 1.2. Affine Space and Algebraic Sets | 4 |
| Problem 1.8.* | 4 |
| Problem 1.11. | 5 |
| Chapter TODO | 7 |
| Section TODO | 7 |
| Problem TODO | 7 |

Chapter 1: Affine Algebraic Sets

1.1. Algebraic Preliminaries

Problem 1.1.*

Let R be a domain.

- (a) If f, g are forms of degree r, s respectively in $R[x_1, \dots, x_n]$, show that fg is a form of degree $r + s$.
- (b) Show that any factor of a form in $R[x_1, \dots, x_n]$ is also a form.

Proof of (a).

- (1) Write

$$f = \sum_{(i)} a_{(i)} x^{(i)},$$

$$g = \sum_{(j)} b_{(j)} x^{(j)},$$

where $\sum_{(i)}$ is the summation over $(i) = (i_1, \dots, i_n)$ with $i_1 + \dots + i_n = r$ and $\sum_{(j)}$ is the summation over $(j) = (j_1, \dots, j_n)$ with $j_1 + \dots + j_n = s$.

- (2) Hence,

$$fg = \sum_{(i)} \sum_{(j)} a_{(i)} b_{(j)} x^{(i)} x^{(j)}$$

$$= \sum_{(i),(j)} a_{(i)} b_{(j)} x^{(k)}$$

where $(k) = (i_1 + j_1, \dots, i_n + j_n)$ with $(i_1 + j_1) + \dots + (i_n + j_n) = r + s$. Each $x^{(k)}$ is the form of degree $r + s$ and $a_{(i)} b_{(j)} \in R$. Hence fg is a form of degree $r + s$.

□

Proof of (b).

- (1) Given any form $f \in R[x_1, \dots, x_n]$, and write $f = gh$. It suffices to show that g is a form as well. (So does h .)
- (2) Write

$$g = g_0 + \dots + g_r, \quad h = h_0 + \dots + h_s$$

where $g_r \neq 0$ and $h_s \neq 0$. So

$$f = gh = g_0h_0 + \cdots + g_rh_s.$$

Since R is a domain, $R[x_1, \dots, x_n]$ is a domain and thus $g_rh_s \neq 0$. The maximality of r and s implies that $\deg f = r + s$. Therefore, by the maximality of $r + s$, $f = g_rh_s$, or $g = g_r$, or g is a form.

□

Problem 1.5.*

Let k be any field. Show that there are an infinitely number of irreducible monic polynomials in $k[x]$. (Hint: Suppose f_1, \dots, f_n were all of them, and factor $f_1 \cdots f_n + 1$ into irreducible factors.)

Proof (Due to Euclid).

- (1) If f_1, \dots, f_n were all irreducible monic polynomials, then we consider

$$g = f_1 \cdots f_n + 1 \in k[x].$$

So there is an irreducible monic polynomial $f = f_i$ dividing g for some i since

$$\deg g = \deg f_1 + \cdots + \deg f_n \geq 1.$$

- (2) However, f would divide the difference

$$g - f_1 \cdots f_{i-1}f_{i+1} \cdots f_n = 1,$$

contrary to $\deg f_i \geq 1$.

□

Problem 1.6.*

Show that any algebraically closed field is infinite. (Hint: The irreducible monic polynomials are $x - a$, $a \in k$.)

Proof (Due to Euclid).

- (1) Let k be an algebraically closed field. If a_1, \dots, a_n were all elements in k , then we consider a monic polynomials

$$f(x) = (x - a_1) \cdots (x - a_n) + 1 \in k[x].$$

- (2) Since k is algebraically closed, there is an element $a \in k$ such that $f(a) = 0$. By assumption, $a = a_i$ for some $1 \leq i \leq n$, and thus $f(a) = f(a_i) = 1$, contrary to the fact that a field is a commutative ring where $0 \neq 1$ and all nonzero elements are invertible.

□

1.2. Affine Space and Algebraic Sets

Problem 1.8.*

Show that the algebraic subsets of $\mathbf{A}^1(k)$ are just the finite subsets, together with $\mathbf{A}^1(k)$ itself.

Proof.

- (1) Show that $k[x]$ is a PID if k is a field.
- (a) Let I be an ideal of $k[x]$.
 - (b) If $I = \{0\}$ then $I = (0)$ and I is principal.
 - (c) If $I \neq \{0\}$, then take f to be a polynomial of minimal degree in I . It suffices to show that $I = (f)$. Clearly, $(f) \subseteq I$ since I is an ideal. Conversely, for any $g \in I$,

$$g(x) = f(x)h(x) + r(x)$$

for some $h, r \in k[x]$ with $r = 0$ or $\deg r < \deg f$. Now as

$$r = g - fh \in I,$$

$r = 0$ (otherwise contrary to the minimality of f), we have $g = fh \in (f)$ for all $g \in I$.

- (2) Let Y be an algebraic subset of $\mathbf{A}^1(k)$, say $Y = V(I)$ for some ideal I of $k[x]$. Since $k[x]$ is a PID, $I = (f)$ for some $f \in k[x]$.
- (a) If $f = 0$, then $I = (0)$ and $Y = V(0) = \mathbf{A}^1(k)$.
 - (b) If $f \neq 0$, then $f(x) = 0$ has finitely many roots in k , say $a_1, \dots, a_m \in k$. Hence,

$$Y = V(I) = V(f) = \{f(a) = 0 : a \in k\} = \{a_1, \dots, a_m\}$$

is a finite subsets of $\mathbf{A}^1(k)$.

By (a)(b), the result is established.

□

Notes.

- (1) By the Hilbert basis theorem, $k[x]$ is Noetherian as k is Noetherian. Hence, for any algebraic subset $Y = V(I)$ of $\mathbf{A}^1(k)$, we can write $I = (f_1, \dots, f_m)$. Note that

$$Y = V(I) = V(f_1) \cap \dots \cap V(f_m).$$

Now apply the same argument to get the same conclusion.

- (2) Suppose $k = \bar{k}$. $\mathbf{A}^1(k)$ is irreducible, because its only proper closed subsets are finite, yet it is infinite (because k is algebraically closed, hence infinite).

Problem 1.11.

Show that the following are algebraic sets:

- (a) $\{(t, t^2, t^3) \in \mathbf{A}^3(k) : t \in k\}$;
- (b) $\{(\cos(t), \sin(t)) \in \mathbf{A}^2(\mathbb{R}) : t \in \mathbb{R}\}$;
- (c) *the set of points in $\mathbf{A}^2(\mathbb{R})$ whose polar coordinates (r, θ) satisfy the equation $r = \sin(\theta)$.*

Proof of (a).

- (1) The twisted cubic curve

$$Y = \{(t, t^2, t^3) \in \mathbf{A}^3(k) : t \in k\} = V(x^2 - y) \cap V(x^3 - z)$$

is algebraic. We say that Y is given by the parametric representation $x = t, y = t^2, z = t^3$.

- (2) The generators for the ideal $I(Y)$ are $x^2 - y$ and $x^3 - z$.
- (3) Y is an affine variety of dimension 1.
- (4) The affine coordinate ring $A(Y)$ is isomorphic to a polynomial ring in one variable over k .

□

Proof of (b). The circle

$$\{(\cos(t), \sin(t)) \in \mathbf{A}^2(\mathbb{R}) : t \in \mathbb{R}\} = V(x^2 - y^2 - 1)$$

is algebraic. □

Proof of (c). The circle

$$\{(r, \theta) : r = \sin(\theta)\} = V(x^2 + y^2 - y)$$

is algebraic again. \square

Chapter TODO

Section TODO

Problem TODO

TODO

Proof.

- (1) TODO