

Notes on the book:
*Apostol, Introduction to Analytic
Number Theory*

Meng-Gen Tsai
plover@gmail.com

August 18, 2021

Contents

Chapter 1: The Fundamental Theorem of Arithmetic	2
Exercise 1.15.	2
Exercise 1.30.	2
Chapter 2: Arithmetical functions and Dirichlet multiplication	4
Exercise 2.1.	4
Exercise 2.3.	5
Supplement 2.3.1. (Chinese remainder theorem)	6
Exercise 2.4.	7

Chapter 1: The Fundamental Theorem of Arithmetic

Exercise 1.15.

Prove that every $n \geq 12$ is the sum of two composite numbers.

Proof. Write $n = 2m$ (resp. $n = 2m + 1$) where $m \in \mathbb{Z}$, $m \geq 6$. Then $n = 8 + 2(m - 4)$ (resp. $n = 9 + 2(m - 4)$) is the sum of two composite numbers. \square

Exercise 1.30.

If $n > 1$ prove that the sum

$$\sum_{k=1}^n \frac{1}{k}$$

is not an integer.

Proof.

(1) (Reductio ad absurdum) Suppose

$$H := \sum_{k=1}^n \frac{1}{k}$$

were an integer.

(2) Let s be the largest integer such that $2^s \leq n$. So the integer number

$$\begin{aligned} 2^{s-1}H &= \sum_{k=1}^n \frac{2^{s-1}}{k} \\ &= 2^{s-1} + 2^{s-2} + \frac{2^{s-1}}{3} + 2^{s-3} + \frac{2^{s-1}}{5} + \frac{2^{s-2}}{3} + \cdots + \frac{1}{2} + \cdots. \end{aligned}$$

has only one term of even denominators (as $n > 1$) if we write all terms in irreducible fractions. That is,

$$2^{s-1}H = \frac{1}{2} + \frac{c}{d} \in \mathbb{Z}$$

where $\frac{c}{d}$ is an irreducible fraction with odd d . Hence it suffices to show that $2 \nmid d$ to get a contradiction.

(3) By

$$\frac{1}{2} + \frac{c}{d} = \frac{d+2c}{2d} \in \mathbb{Z}$$

we have $d + 2c = 2dd'$ for some $d' \in \mathbb{Z}$. Note that 2 is a prime. So $2 \mid (d + 2c)$ or $2 \mid d$, which is absurd.

□

Chapter 2: Arithmetical functions and Dirichlet multiplication

Exercise 2.1.

Find all integers n such that

- (a) $\varphi(n) = \frac{n}{2}$,
- (b) $\varphi(n) = \varphi(2n)$,
- (c) $\varphi(n) = 12$.

Proof of (a).

$$\varphi(n) = n \prod_{p|n} \left(1 - \frac{1}{p}\right) = \frac{n}{2}$$

(Theorem 2.4) implies that $n = 2$. \square

Proof of (b).

- (1) $\varphi(n) = \varphi(2n)$ implies that

$$n \prod_{p|n} \left(1 - \frac{1}{p}\right) = 2n \prod_{p|(2n)} \left(1 - \frac{1}{p}\right).$$

- (2) If $2|n$, then $n = 2n$ or $n = 0$, which is absurd.

- (3) If $2 \nmid n$, then

$$n \prod_{p|n} \left(1 - \frac{1}{p}\right) = 2n \prod_{p|(2n)} \left(1 - \frac{1}{p}\right) = \underbrace{2n \left(1 - \frac{1}{2}\right)}_{=n} \prod_{p|n} \left(1 - \frac{1}{p}\right)$$

is always true. Hence n is odd if $\varphi(n) = \varphi(2n)$.

\square

Proof of (c).

- (1) Show that the solutions of $\varphi(n) = 12$ are $n = 13, 26, 21, 28, 42, 36$. Write $n = p_1^{\alpha_1} \cdots p_r^{\alpha_r}$ where $p_1 < p_2 < \dots$. Then

$$12 = \varphi(n) = \prod_{i=1}^r p_i^{\alpha_i-1} (p_i - 1).$$

(Theorem 2.5). It implies that $p_i \in \{2, 3, 5, 7, 13\}$ if $\alpha_i > 0$. Consider all possible cases of the greatest prime divisor p_r of n as follows.

(2) If $p_r = 13$, then $\alpha_r = 1$ since $13 \nmid 12$. So

$$12 = \varphi(n) = \underbrace{\varphi(13)}_{=12} \varphi\left(\frac{n}{13}\right)$$

or $1 = \varphi\left(\frac{n}{13}\right)$. Hence $\frac{n}{13} = 1, 2$. In this case $n = 13, 26$.

(3) If $p_r = 7$, then $\alpha_r = 1$ since $7 \nmid 12$. So

$$12 = \varphi(n) = \underbrace{\varphi(7)}_{=6} \varphi\left(\frac{n}{7}\right)$$

or $2 = \varphi\left(\frac{n}{7}\right)$. Hence $\frac{n}{7} = 3, 4, 6$. In this case $n = 21, 28, 42$.

(5) If $p_r = 5$, then $\alpha_r = 1$ since $5 \nmid 12$. So $12 = \varphi(5)\varphi\left(\frac{n}{5}\right)$ or $3 = \varphi\left(\frac{n}{5}\right)$, which is impossible.

(6) If $p_r = 3$, then $\alpha_r = 1, 2$. $\alpha_r = 1$ is impossible since $3 \mid 12$. So

$$12 = \varphi(n) = \underbrace{\varphi(3^2)}_{=6} \varphi\left(\frac{n}{3^2}\right)$$

or $2 = \varphi\left(\frac{n}{3^2}\right)$. Hence $\frac{n}{3^2} = 4$. (By assumption $\frac{n}{3^2}$ cannot have any prime factor > 3 .) In this case $n = 36$.

□

Exercise 2.3.

Prove that

$$\frac{n}{\varphi(n)} = \sum_{d \mid n} \frac{\mu(d)^2}{\varphi(d)}.$$

Proof.

(1) Note that fg , f/g and $f * g$ are multiplicative if f and g are multiplicative (Example 5 on page 34 and Theorem 2.14). Hence $\frac{n}{\varphi(n)}$ and $\sum_{d \mid n} \frac{\mu^2(d)}{\varphi(d)}$ are multiplicative. Hence it might assume that $n = p^a$ for some prime p and integer $a \geq 1$. (The case $n = 1$ is trivial.)

(2)

$$\frac{p^a}{\varphi(p^a)} = \frac{p^a}{p^a - p^{a-1}} = \frac{p}{p-1}.$$

(3)

$$\begin{aligned}
\sum_{d|p^a} \frac{\mu(d)^2}{\varphi(d)} &= \frac{\mu(1)^2}{\varphi(1)} + \frac{\mu(p)^2}{\varphi(p)} + \overbrace{\frac{\mu(p^2)^2}{\varphi(p^2)}}^{=0} + \cdots + \overbrace{\frac{\mu(p^a)^2}{\varphi(p^a)}}^{=0} \\
&= 1 + \frac{1}{p-1} + 0 + \cdots + 0 \\
&= \frac{p}{p-1}.
\end{aligned}$$

(4) Or apply Theorems 2.4 and 2.18 to get

$$\begin{aligned}
\sum_{d|n} \frac{\mu(d)^2}{\varphi(d)} &= \prod_{p|n} \left(1 - \frac{\mu(p)}{\varphi(p)}\right) \\
&= \prod_{p|n} \left(1 - \frac{-1}{p-1}\right) \\
&= \prod_{p|n} \frac{p}{p-1} \\
&= \frac{n}{\varphi(n)}.
\end{aligned}$$

□

Supplement 2.3.1. (Chinese remainder theorem)

(Exercise I.3.5 in the textbook: *Jürgen Neukirch, Algebraic Number Theory*.)
The quotient ring \mathcal{O}/\mathfrak{a} of a Dedekind domain by an ideal $\mathfrak{a} \neq 0$ is a principal ideal domain. (Hint: For $\mathfrak{a} = \mathfrak{p}^n$ the only proper ideals of \mathcal{O}/\mathfrak{a} are given by $\mathfrak{p}/\mathfrak{p}^n, \dots, \mathfrak{p}^{n-1}/\mathfrak{p}^n$. Choose $\pi \in \mathfrak{p} \setminus \mathfrak{p}^2$ and show that $\mathfrak{p}^\nu = \mathcal{O}\pi^\nu + \mathfrak{p}^n$.)

Proof.

(1) By the Chinese remainder theorem, it suffices to show the case $\mathfrak{a} = \mathfrak{p}^n$ where \mathfrak{p} is prime.

(2) There is a natural correspondence between

$$\{\text{ideals of } \mathcal{O}/\mathfrak{p}^n\} \longleftrightarrow \{\text{ideals of } \mathcal{O} \text{ containing } \mathfrak{p}^n\}.$$

Hence the proper ideals of $\mathcal{O}/\mathfrak{p}^n$ are given by $\mathfrak{p}/\mathfrak{p}^n, \dots, \mathfrak{p}^{n-1}/\mathfrak{p}^n$.

(3) Similar to Exercise I.3.4, choose $\pi \in \mathfrak{p} \setminus \mathfrak{p}^2$ and thus $\mathfrak{p}^\nu = \mathcal{O}\pi^\nu + \mathfrak{p}^n$ ($\nu = 1, \dots, n-1$) since they have the same prime factorization. Hence $\mathfrak{p}^\nu/\mathfrak{p}^n = (\pi^\nu + \mathfrak{p}^n)$ is principal.

□

Exercise 2.4.

Prove that $\varphi(n) > \frac{n}{6}$ for all n with at most 8 distinct prime factors.

Proof.

(1)

$$\begin{aligned}\varphi(n) &= n \prod_{p|n} \left(1 - \frac{1}{p}\right) && \text{(Theorem 2.4)} \\ &\geq n \left(1 - \frac{1}{2}\right) \left(1 - \frac{1}{3}\right) \left(1 - \frac{1}{5}\right) \left(1 - \frac{1}{7}\right) \\ &\quad \left(1 - \frac{1}{11}\right) \left(1 - \frac{1}{13}\right) \left(1 - \frac{1}{17}\right) \left(1 - \frac{1}{19}\right) \\ &= \frac{55296}{323323} n \\ &> \frac{n}{6}.\end{aligned}$$

(2) The conclusion does not hold if n has more than 9 distinct prime factors.

□