

Notes on the book:
*Apostol, Modular Functions and
Dirichlet Series in Number Theory,
2nd edition*

Meng-Gen Tsai
plover@gmail.com

July 29, 2021

Contents

Chapter 1: Elliptic functions	2
Exercise 1.1.	2
Supplement 1.1.1.	3
Exercise 1.2.	3
Exercise 1.3.	4
Exercise 1.4.	5
Supplement 1.4.1. (Divisor class group)	6
Exercise 1.5.	7
Exercise 1.6.	7
Exercise 1.7.	8
Exercise 1.8.	9
Exercise 1.9.	10
Exercise 1.10.	12
Exercise 1.11.	13
Exercise 1.12.	14
Exercise 1.13.	15
Exercise 1.14. (Lambert series)	16
Exercise 1.15.	19

Chapter 1: Elliptic functions

Exercise 1.1.

Given two pairs of complex numbers (ω_1, ω_2) and (ω'_1, ω'_2) with nonreal ratios ω_1/ω_2 and ω'_1/ω'_2 . Prove that they generate the same set of periods if, and only if, there is a 2×2 matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with integer entries and determinant ± 1 such that

$$\begin{pmatrix} \omega'_2 \\ \omega'_1 \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \omega_2 \\ \omega_1 \end{pmatrix}.$$

Proof.

- (1) (\implies) Suppose (ω_1, ω_2) and (ω'_1, ω'_2) generate the same set of periods.

In particular, there is a 2×2 matrix $A := \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathbf{M}_{2 \times 2}(\mathbb{Z})$ (resp.

$A' := \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} \in \mathbf{M}_{2 \times 2}(\mathbb{Z})$) such that

$$\begin{pmatrix} \omega'_2 \\ \omega'_1 \end{pmatrix} = A \begin{pmatrix} \omega_2 \\ \omega_1 \end{pmatrix}, \quad \begin{pmatrix} \omega_2 \\ \omega_1 \end{pmatrix} = A' \begin{pmatrix} \omega'_2 \\ \omega'_1 \end{pmatrix}.$$

Hence it suffices to show $\det(A) = \pm 1$.

- (2) Note that

$$\begin{pmatrix} \omega'_2 \\ \omega'_1 \end{pmatrix} = AA' \begin{pmatrix} \omega'_2 \\ \omega'_1 \end{pmatrix}.$$

Hence

$$AA' = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Take the determinant on the both sides to get

$$\det(A) \det(A') = 1.$$

Since $\det(\mathbf{M}_{2 \times 2}(\mathbb{Z})) \subseteq \mathbb{Z}$, $\det(A) = \pm 1$.

- (3) (\impliedby) $\Omega(\omega'_1, \omega'_2) \subseteq \Omega(\omega_1, \omega_2)$ is obvious. Note that

$$\begin{pmatrix} \omega_2 \\ \omega_1 \end{pmatrix} = \underbrace{\frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}}_{\in \mathbf{M}_{2 \times 2}(\mathbb{Z})} \begin{pmatrix} \omega'_2 \\ \omega'_1 \end{pmatrix}.$$

Thus $\Omega(\omega_1, \omega_2) \subseteq \Omega(\omega'_1, \omega'_2)$. Therefore $\Omega(\omega_1, \omega_2) = \Omega(\omega'_1, \omega'_2)$.

□

Supplement 1.1.1.

(Exercise I.1.1 in the textbook: *Jürgen Neukirch, Algebraic Number Theory*.)
 $\alpha \in \mathbb{Z}[i]$ is a unit if and only if $N(\alpha) = 1$.

Proof.

- (1) (\implies) Since α is a unit, there is $\beta \in \mathbb{Z}[i]$ such that $\alpha\beta = 1$. So $N(\alpha\beta) = N(1)$, or $N(\alpha)N(\beta) = 1$. Since the image of N is nonnegative integers, $N(\alpha) = 1$.
- (2) (\impliedby) $N(\alpha) = \alpha\bar{\alpha}$, or $1 = \alpha\bar{\alpha}$ since $N(\alpha) = 1$. That is, $\bar{\alpha} \in \mathbb{Z}[i]$ is the inverse of $\alpha \in \mathbb{Z}[i]$. (Or we solve the equation $N(\alpha) = a^2 + b^2 = 1$, and show that all four solutions (± 1 and $\pm i$) are units.)
- (3) Conclusion: a unit $\alpha = a + bi$ of $\mathbb{Z}[i]$ is satisfying the equation $N(\alpha) = a^2 + b^2 = 1$ by (1)(2). That is, the only unit of $\mathbb{Z}[i]$ are ± 1 and $\pm i$.

□

Exercise 1.2.

Let $S(0)$ denote the sum of the zeros of an elliptic function f in a period parallelogram, and let $S(\infty)$ denote the sum of the poles in the same parallelogram. Prove that $S(0) - S(\infty)$ is a period of f . (Hint: Integrate $z \frac{f'(z)}{f(z)}$.)

Proof.

- (1) Similar to Theorem 1.8, the integral

$$\frac{1}{2\pi i} \int_C z \frac{f'(z)}{f(z)}$$

taken around the boundary C of a cell (no zeros or poles on its boundary) counts the difference between the sum of the zeros and the sum of the poles inside the cell, that is,

$$S(0) - S(\infty) = \frac{1}{2\pi i} \int_C z \frac{f'(z)}{f(z)}.$$

(The proof is similar to the proof of the argument principle.)

- (2) Let C_1 be the path from 0 to ω_1 , C_2 be the path from ω_1 to $\omega_1 + \omega_2$, C_3

be the path from $\omega_1 + \omega_2$ to ω_2 , and C_4 be the path from ω_2 to 0. Hence

$$\begin{aligned}
& \frac{1}{2\pi i} \int_{C_1} z \frac{f'(z)}{f(z)} + \frac{1}{2\pi i} \int_{C_3} z \frac{f'(z)}{f(z)} \\
&= \frac{1}{2\pi i} \int_{C_1} z \frac{f'(z)}{f(z)} + \frac{1}{2\pi i} \int_{-C_1} (z + \omega_2) \frac{f'(z + \omega_2)}{f(z + \omega_2)} \\
&= \frac{1}{2\pi i} \int_{C_1} z \frac{f'(z)}{f(z)} - \frac{1}{2\pi i} \int_{C_1} (z + \omega_2) \frac{f'(z)}{f(z)} \\
&= -\omega_2 \frac{1}{2\pi i} \int_{C_1} \frac{f'(z)}{f(z)}
\end{aligned}$$

and

$$\begin{aligned}
& \frac{1}{2\pi i} \int_{C_2} z \frac{f'(z)}{f(z)} + \frac{1}{2\pi i} \int_{C_4} z \frac{f'(z)}{f(z)} \\
&= \frac{1}{2\pi i} \int_{-C_4} (z + \omega_1) \frac{f'(z + \omega_1)}{f(z + \omega_1)} + \frac{1}{2\pi i} \int_{C_4} z \frac{f'(z)}{f(z)} \\
&= -\frac{1}{2\pi i} \int_{C_4} (z + \omega_1) \frac{f'(z)}{f(z)} + \frac{1}{2\pi i} \int_{C_4} z \frac{f'(z)}{f(z)} \\
&= -\omega_1 \frac{1}{2\pi i} \int_{C_4} \frac{f'(z)}{f(z)}
\end{aligned}$$

Therefore

$$S(0) - S(\infty) = -\omega_1 \frac{1}{2\pi i} \int_{C_4} \frac{f'(z)}{f(z)} - \omega_2 \frac{1}{2\pi i} \int_{C_1} \frac{f'(z)}{f(z)}.$$

So it suffices to show that $\frac{1}{2\pi i} \int_{C_1} \frac{f'(z)}{f(z)} \in \mathbb{Z}$. (Other cases are similar.)

(3) By choosing one branch of log, we have

$$\begin{aligned}
\frac{1}{2\pi i} \int_{C_1} \frac{f'(z)}{f(z)} &= \frac{1}{2\pi i} \log \frac{f(\omega_1)}{f(0)} \\
&= \frac{1}{2\pi i} \log(1) && (f(\omega_1) = f(0)) \\
&= \frac{1}{2\pi i} (2\pi i m) \text{ for some } m \in \mathbb{Z} \\
&= m \in \mathbb{Z}.
\end{aligned}$$

□

Exercise 1.3.

(a) Prove that $\wp(u) = \wp(v)$ if, and only if, $u - v$ or $u + v$ is a period of \wp .

- (b) Let a_1, \dots, a_n and b_1, \dots, b_m be complex numbers such that none of the numbers $\wp(a_i) - \wp(b_j)$ is zero. Let

$$f(z) = \frac{\prod_{k=1}^n [\wp(z) - \wp(a_k)]}{\prod_{r=1}^m [\wp(z) - \wp(b_r)]}.$$

Prove that f is an even elliptic function with zeros at a_1, \dots, a_n and poles at b_1, \dots, b_m .

Proof of (a).

- (1) Let Ω be the lattice generated by periods of \wp .
- (2) (\implies) It is equivalent to show that the equation $\wp(u) = \wp(v)$ in terms of u has exactly two roots in some period parallelogram. $u \equiv v \pmod{\Omega}$ is a root clearly and $u \equiv -v \pmod{\Omega}$ is also a root since \wp is even. Since \wp is an elliptic function of order 2 (Theorem 1.8), $u \equiv \pm v \pmod{\Omega}$ is the only two roots of $\wp(u) = \wp(v)$.
- (3) (\impliedby) Obvious.

□

Proof of (b).

- (1) Since \wp is an even elliptic function, f is an even elliptic function too.
- (2) f has zeros at a_1, \dots, a_n and poles at b_1, \dots, b_m (by construction and (a)).

□

Exercise 1.4.

Prove that every even elliptic function f is a rational function of \wp , where periods of \wp are a subset of the periods of f .

Proof.

- (1) Nothing to do if f is constant. Let C be one period parallelogram of f and \wp . Let $\Omega(\omega_1, \omega_2)$ be the lattice generated by periods of \wp . Suppose f has zeros at a_1, \dots, a_n and poles at b_1, \dots, b_m .
- (2) Might assume that $\wp(z) - \wp(a_k)$ (resp. $\wp(z) - \wp(b_r)$) has a simple zero in a_k (resp. b_r) for all k, r . So the function

$$g(z) := f(z) \cdot \frac{\prod_{r=1}^m [\wp(z) - \wp(b_r)]^{\beta_r}}{\prod_{k=1}^n [\wp(z) - \wp(a_k)]^{\alpha_k}}$$

is an elliptic function with no zeros or poles in C where α_k (resp. β_r) is the order of the zero a_k (resp. the pole b_r). By Theorems 1.4 and 1.5, $g(z)$ is a constant. Hence

$$f(z) = C \cdot \frac{\prod_{k=1}^n [\wp(z) - \wp(a_k)]^{\alpha_k}}{\prod_{r=1}^m [\wp(z) - \wp(b_r)]^{\beta_r}}$$

for some constant $C \in \mathbb{C}$.

- (3) Now we consider the case a_k (resp. b_r) is a zero of $\wp'(z)$. Since f is an even elliptic function, the order of a_k (resp. b_r) of f is even. Note that the order of a_k (resp. b_r) of $\wp(z) - \wp(a_k)$ (resp. $\wp(z) - \wp(b_r)$) is 2. Hence the function

$$g(z) := f(z) \cdot \frac{\prod_{\wp'(b_r) \neq 0} [\wp(z) - \wp(b_r)]^{\beta_r}}{\prod_{\wp'(a_k) \neq 0} [\wp(z) - \wp(a_k)]^{\alpha_k}} \cdot \frac{\prod_{\wp'(b_r)=0} [\wp(z) - \wp(b_r)]^{\frac{\beta_r}{2}}}{\prod_{\wp'(a_k)=0} [\wp(z) - \wp(a_k)]^{\frac{\alpha_k}{2}}}$$

is a constant too.

□

Supplement 1.4.1. (Divisor class group)

(Problem 8.6 in the textbook: *William Fulton, Algebraic Curves*.) Let $D(X)$ be the group of divisors on X , $D_0(X)$ the subgroup consisting of divisors of degree zero, and $P(X)$ the subgroup of $D_0(X)$ consisting of divisors of rational functions. Let $C_0(X) = D_0(X)/P(X)$ be the quotient group. It is the **divisor class group** on X . Show that if $X = \mathbf{P}^1$, then $C_0(X) = 0$.

Proof.

- (1) Given a divisor

$$D = \sum_{P \in X} n_P P \in D_0(X)$$

where $n_P \in \mathbb{Z}$ and $\sum_P n_P = 0$.

- (2) Note that $\sum_P n_P = 0$. We can define a rational function $z \in k(X)$ by

$$z = \prod_{P=[a_P : b_P] \in X} (b_P x - a_P y)^{n_P}.$$

Hence $\text{div}(z) = D \in P(X)$. Therefore $C_0(X) = D_0(X)/P(X) = 0$.

□

Exercise 1.5.

Prove that every elliptic function f can be expressed in the form

$$f(z) = R_1[\wp(z)] + \wp'(z)R_2[\wp(z)]$$

where R_1 and R_2 are rational functions and \wp has the same set of periods as f .

Proof.

$$\begin{aligned} f(z) &= \underbrace{\frac{f(z) + f(-z)}{2}}_{\text{even}} + \wp'(z) \underbrace{\frac{f(z) - f(-z)}{2\wp'(z)}}_{\text{even}} \\ &= R_1[\wp(z)] + \wp'(z)R_2[\wp(z)] \text{ for some rational functions } R_1, R_2 \end{aligned}$$

(by Exercise 1.4). \square

Exercise 1.6.

Let f and g be two elliptic functions with the same set of periods. Prove that there exists a polynomial $P(x, y)$, not identically zero, such that

$$P[f(z), g(z)] = C$$

where C is a constant (depending on f and g but not on z).

Proof.

(1) By Exercise 1.5, we have

$$f(z) = R_1[\wp(z)] + \wp'(z)R_2[\wp(z)]$$

for some rational functions R_1, R_2 and \wp has the same set of periods as f . By cleaning the denominators of R_1 and R_2 , we might assume

$$S[\wp(z)]f(z) = R_1[\wp(z)] + \wp'(z)R_2[\wp(z)]$$

for some polynomials R_1, R_2, S .

(2) So

$$\begin{aligned} \wp'(z)R_2[\wp(z)] &= S[\wp(z)]f(z) - R_1[\wp(z)] \\ \implies \wp'(z)^2 R_2[\wp(z)]^2 &= (S[\wp(z)]f(z) - R_1[\wp(z)])^2 \\ \implies (4\wp(z)^3 - 60G_4\wp(z) - 140G_6)R_2[\wp(z)]^2 \\ &= (S[\wp(z)]f(z) - R_1[\wp(z)])^2. \quad (\text{Theorem 1.12}) \\ \implies F(\wp(z), f(z)) &= 0 \end{aligned}$$

for some polynomials $F(x, y) \in \mathbb{C}[x, y]$. Note that $F(x, y)$ is of degree 2 if we view $F \in (\mathbb{C}[x])[y]$.

(3) Similarly,

$$G(\wp(z), g(z)) = 0$$

for some polynomials $G(x, y) \in \mathbb{C}[x, y]$.

(4) Let $P = \text{Res}_x(F, G)$ be the resultant of two polynomials F and G with respect to x to eliminate $\wp(z)$. Note that P is a nonzero polynomial (since F and G are nonzero) and $P[f(z), g(z)] = 0$. So P is our desired polynomial.

□

Exercise 1.7.

The discriminant of the polynomial $f(x) = 4(x - x_1)(x - x_2)(x - x_3)$ is the product $16\{(x_2 - x_1)(x_3 - x_2)(x_3 - x_1)\}^2$. Prove that the discriminant of $f(x) = 4x^3 - ax - b$ is $a^3 - 27b^2$.

Proof.

(1) Since

$$f'(x) = 4(x - x_2)(x - x_3) + 4(x - x_1)(x - x_3) + 4(x - x_1)(x - x_2),$$

we have

$$\begin{aligned} f'(x_1) &= 4(x_1 - x_2)(x_1 - x_3), \\ f'(x_2) &= 4(x_2 - x_1)(x_2 - x_3), \\ f'(x_3) &= 4(x_3 - x_1)(x_3 - x_2). \end{aligned}$$

Hence

$$f'(x_1)f'(x_2)f'(x_3) = -4\text{disc}(f)$$

where $\text{disc}(f)$ is the discriminant of $f(x)$.

(2) As $f(x) = 4x^3 - ax - b$, we have $f'(x) = 12x^2 - a$. So

$$f'(x_1)f'(x_2)f'(x_3) = (12x_1^2 - a)(12x_2^2 - a)(12x_3^2 - a).$$

Note that

$$\begin{aligned} x_1x_2x_3 &= \frac{b}{4}, \\ x_1x_2 + x_2x_3 + x_3x_1 &= -\frac{a}{4}, \\ x_1 + x_2 + x_3 &= 0, \end{aligned}$$

we have

$$\begin{aligned}
x_1^2 x_2^2 x_3^2 &= \frac{b^2}{4^2}, \\
x_1^2 x_2^2 + x_2^2 x_3^2 + x_3^2 x_1^2 &= (x_1 x_2 + x_2 x_3 + x_3 x_1)^2 - 2x_1 x_2 x_3 (x_1 + x_2 + x_3) \\
&= \frac{a^2}{4^2}, \\
x_1^2 + x_2^2 + x_3^2 &= (x_1 + x_2 + x_3)^2 - 2(x_1 x_2 + x_2 x_3 + x_3 x_1) \\
&= \frac{a}{2}.
\end{aligned}$$

(3) Hence

$$\begin{aligned}
f'(x_1)f'(x_2)f'(x_3) &= (12x_1^2 - a)(12x_2^2 - a)(12x_3^2 - a) \\
&= 12^3(x_1^2 x_2^2 x_3^2) - 12^2 a(x_1^2 x_2^2 + x_2^2 x_3^2 + x_3^2 x_1^2) \\
&\quad + 12a^2(x_1^2 + x_2^2 + x_3^2) - a^3 \\
&= 12^3 \cdot \frac{b^2}{4^2} - 12^2 a \cdot \frac{a^2}{4^2} + 12a^2 \cdot \frac{a}{2} - a^3 \\
&= -4(a^3 - 27b^2).
\end{aligned}$$

Therefore

$$\text{disc}(4x^3 - ax - b) = a^3 - 27b^2.$$

□

Exercise 1.8.

The differential equation for \wp shows that $\wp'(z) = 0$ if $z = \frac{\omega_1}{2}, \frac{\omega_2}{2}$ or $\frac{\omega_1 + \omega_2}{2}$. Show that

$$\wp''\left(\frac{\omega_1}{2}\right) = 2(e_1 - e_2)(e_1 - e_3)$$

and obtain corresponding formulas for $\wp''\left(\frac{\omega_2}{2}\right)$ and $\wp''\left(\frac{\omega_1 + \omega_2}{2}\right)$.

Proof.

(1) Differentiation of the equation

$$4\wp(z)^3 - g_2\wp(z) - g_3 = 4(\wp(z) - e_1)(\wp(z) - e_2)(\wp(z) - e_3)$$

in Theorem 1.14 to get

$$\begin{aligned}
12\wp(z)^2\wp'(z) - g_2\wp'(z) &= 4\wp'(z)(\wp(z) - e_2)(\wp(z) - e_3) \\
&\quad + 4\wp'(z)(\wp(z) - e_1)(\wp(z) - e_3) \\
&\quad + 4\wp'(z)(\wp(z) - e_1)(\wp(z) - e_2).
\end{aligned}$$

Since $\wp''(z) = 6\wp(z)^2 - \frac{g_2}{2}$, we have

$$\begin{aligned}\wp''(z) &= 2(\wp(z) - e_2)(\wp(z) - e_3) \\ &\quad + 2(\wp(z) - e_1)(\wp(z) - e_3) \\ &\quad + 2(\wp(z) - e_1)(\wp(z) - e_2).\end{aligned}$$

(2) Hence

$$\begin{aligned}\wp''\left(\frac{\omega_1}{2}\right) &= 2(e_1 - e_2)(e_1 - e_3), \\ \wp''\left(\frac{\omega_2}{2}\right) &= 2(e_2 - e_1)(e_2 - e_3), \\ \wp''\left(\frac{\omega_1 + \omega_2}{2}\right) &= 2(e_3 - e_1)(e_3 - e_2).\end{aligned}$$

□

Exercise 1.9.

According to Exercise 1.4, the function $\wp(2z)$ is a rational function of $\wp(z)$. Prove that, in fact,

$$\wp(2z) = \frac{\{\wp(z)^2 + \frac{1}{4}g_2\}^2 + 2g_3\wp(z)}{4\wp(z)^3 - g_2\wp(z) - g_3} = -2\wp(z) + \frac{1}{4} \left(\frac{\wp''(z)}{\wp'(z)} \right)^2.$$

Proof.

(1) By $\wp'(z)^2 = 4\wp(z)^3 - g_2\wp(z) - g_3$ and $\wp''(z) = 6\wp(z)^2 - \frac{1}{2}g_2$, we have

$$\begin{aligned}& -2\wp(z) + \frac{1}{4} \left(\frac{\wp''(z)}{\wp'(z)} \right)^2 \\ &= -2\wp(z) + \frac{1}{4} \cdot \frac{(6\wp(z)^2 - \frac{1}{2}g_2)^2}{4\wp(z)^3 - g_2\wp(z) - g_3} \\ &= \frac{-2\wp(z)[4\wp(z)^3 - g_2\wp(z) - g_3] + \frac{1}{4}(6\wp(z)^2 - \frac{1}{2}g_2)^2}{4\wp(z)^3 - g_2\wp(z) - g_3} \\ &= \frac{\wp(z)^4 + \frac{1}{2}g_2\wp(z)^2 + \frac{1}{16}g_2^2 + 2g_3\wp(z)}{4\wp(z)^3 - g_2\wp(z) - g_3} \\ &= \frac{\{\wp(z)^2 + \frac{1}{4}g_2\}^2 + 2g_3\wp(z)}{4\wp(z)^3 - g_2\wp(z) - g_3}.\end{aligned}$$

So it suffices to show that $\wp(2z) = -2\wp(z) + \frac{1}{4} \left(\frac{\wp''(z)}{\wp'(z)} \right)^2$.

- (2) Let Ω be the lattice generated by periods of \wp . Suppose the addition theorem of \wp holds, that is,

$$\wp(u) + \wp(v) + \wp(u+v) = \frac{1}{4} \left(\frac{\wp'(u) - \wp'(v)}{\wp(u) - \wp(v)} \right)^2$$

with $u, v, w \not\equiv 0 \pmod{\Omega}$. Then letting v to u , we have

$$\begin{aligned} \wp(2u) &= \lim_{v \rightarrow u} \wp(u+v) \\ &= \lim_{v \rightarrow u} \left\{ -\wp(u) - \wp(v) + \frac{1}{4} \left(\frac{\wp'(u) - \wp'(v)}{\wp(u) - \wp(v)} \right)^2 \right\} \\ &= -2\wp(u) + \frac{1}{4} \lim_{v \rightarrow u} \frac{\wp'(v) - \wp'(u)}{\wp(v) - \wp(u)} \\ &= -2\wp(u) + \frac{1}{4} \left(\frac{\wp''(u)}{\wp'(u)} \right)^2. \end{aligned}$$

The last equality is followed by L'Hospital's rule. So it suffices to show the addition theorem of \wp is true.

- (3) Let $u + v + w = 0$, with $u, v, w \not\equiv 0 \pmod{\Omega}$. Show that

$$\begin{vmatrix} \wp(u) & \wp'(u) & 1 \\ \wp(v) & \wp'(v) & 1 \\ \wp(w) & \wp'(w) & 1 \end{vmatrix} = 0.$$

Consider the elliptic function

$$f(z) := \wp'(z) - \underbrace{\frac{\wp'(u) - \wp'(v)}{\wp(u) - \wp(v)}}_{:=A} \wp(z) - \underbrace{\frac{\wp(u)\wp'(v) - \wp(v)\wp'(u)}{\wp(u) - \wp(v)}}_{:=B}.$$

f has exactly 3 zeros in a period parallelogram as f has order 3. Note that f has a pole at 0 of order 3. By Exercise 1.2, the sum of the zeros is equal to the sum of poles in a period parallelogram. Since u and v are zeros of f (by verifying $f(u) = f(v) = 0$ directly), the third zero must be $-u - v = w$. Hence there is a line

$$y = Ax + B$$

passing through 3 points $(\wp(u), \wp'(u))$, $(\wp(v), \wp'(v))$ and $(\wp(w), \wp'(w))$. So the determinant is zero.

- (4) Now we are going to remove the term $\wp'(w)$ to prove the addition theorem of \wp . By Theorem 1.12, we have the system of equations

$$\begin{cases} y = Ax + B, \\ y^2 = 4x^3 - g_2x - g_3, \end{cases}$$

where $(x, y) = (\wp(z), \wp'(z))$. Hence

$$\begin{aligned}
& (Ax + B)^2 = 4x^3 - g_2x - g_3 \\
& \iff 4x^3 - A^2x^2 - (2AB + g_2)x - (B^2 + g_3) = 0 \\
& \implies \text{sum of three roots of } x \text{ is } \frac{A^2}{4} \\
& \implies \wp(u) + \wp(v) + \wp(w) = \frac{1}{4} \left(\frac{\wp'(u) - \wp'(v)}{\wp(u) - \wp(v)} \right)^2 \\
& \implies \wp(u + v) = -\wp(u) - \wp(v) + \frac{1}{4} \left(\frac{\wp'(u) - \wp'(v)}{\wp(u) - \wp(v)} \right)^2. \quad (\wp: \text{ even})
\end{aligned}$$

The addition theorem of \wp is established.

- (5) Part (4) is similar to defining an addition on a nonsingular cubic in $\mathbf{P}^2(k)$. Actually, if $E \in \mathbf{P}^2(\mathbb{C})$ is the elliptic curve corresponding to the lattice Ω , then there is an isomorphism

$$\alpha : \mathbb{C}/\Omega \longrightarrow E : y^2z = 4x^3 - g_2xz^2 - g_3z^3$$

defined by

$$\alpha(z) = \begin{cases} [\wp(z) : \wp'(z) : 1] & \text{if } z \neq 0 \in \Omega, \\ [0 : 1 : 0] & \text{if } z = 0 \in \Omega, \end{cases}$$

such that α is both analytic (as a mapping of complex manifolds) and algebraic (as a homomorphism of groups).

□

Exercise 1.10.

Let ω_1 and ω_2 be complex numbers with nonreal ratio. Let $f(z)$ be an entire function and assume there are constants a and b such that

$$f(z + \omega_1) = af(z), \quad f(z + \omega_2) = bf(z),$$

for all z . Prove that $f(z) = Ae^{Bz}$, where A and B are constant.

Proof.

- (1) Might assume that $a \neq 0$ and $b \neq 0$ (otherwise $f = 0$ on \mathbb{C}).
- (2) Define

$$g(z) := \frac{f(z)}{e^{Bz}}.$$

It suffices to show g is a constant. Note that $g(z)$ is entire (since f and $e^{Bz} \neq 0$ are entire). By Theorem 1.4, it suffices to show g is doubly periodic, that is, to show

$$g(z + \omega_1) = g(z) \text{ and } g(z + \omega_2) = g(z)$$

for suitable B .

(3) Note that

$$\begin{aligned} g(z + \omega_1) &= g(z) \text{ and } g(z + \omega_2) = g(z) \\ \iff \frac{a}{e^{B\omega_1}} \cdot g(z) &= g(z) \text{ and } \frac{b}{e^{B\omega_2}} \cdot g(z) = g(z) \\ \iff e^{B\omega_1} &= a \text{ and } e^{B\omega_2} = b \\ \iff \exists B \text{ such that } e^{B\omega_1} &= a \text{ and } e^{B\omega_2} = b. \end{aligned}$$

Take B such that $e^{B(\omega_1 - \omega_2)} = \frac{a}{b}$ (since $\frac{a}{b}$ is well-defined, $\omega_1 - \omega_2 \neq 0$, and $z \mapsto \exp(z)$ is a onto map from \mathbb{C} to $\mathbb{C} \setminus \{0\}$). Hence g is doubly periodic.

□

Exercise 1.11.

If $k \geq 2$ and $\tau \in H$ prove that the Eisenstein series

$$G_{2k}(\tau) = \sum_{(m,n) \neq (0,0)} (m + n\tau)^{-2k}$$

has the Fourier expansion

$$G_{2k}(\tau) = 2\zeta(2k) + \frac{2(2\pi i)^{2k}}{(2k-1)!} \sum_{n=1}^{\infty} \sigma_{2k-1}(n) e^{2\pi i n \tau}.$$

Proof.

(1) Let $q = e^{2\pi i \tau}$. Similar to Lemma 1.3 on page 19, we have

$$(2k-1)! \sum_{m=-\infty}^{+\infty} \frac{1}{(\tau + m)^{2k}} = (2\pi i)^{2k} \sum_{r=1}^{\infty} r^{2k-1} q^r.$$

(2) Similar to Theorem 1.18, we have

$$\begin{aligned}
G_{2k}(\tau) &= \sum_{(m,n) \neq (0,0)} (m+n\tau)^{-2k} \\
&= \sum_{\substack{m=-\infty \\ m \neq 0 (n=0)}}^{+\infty} m^{-2k} + \sum_{n=1}^{\infty} \sum_{m=-\infty}^{+\infty} ((m+n\tau)^{-2k} + (m-n\tau)^{-2k}) \\
&= 2\zeta(2k) + 2 \sum_{n=1}^{\infty} \sum_{m=-\infty}^{+\infty} (m+n\tau)^{-2k} \\
&= 2\zeta(2k) + 2 \sum_{n=1}^{\infty} \frac{(2\pi i)^{2k}}{(2k-1)!} \sum_{r=1}^{\infty} r^{2k-1} q^{nr} \\
&= 2\zeta(2k) + \frac{2(2\pi i)^{2k}}{(2k-1)!} \sum_{n=1}^{\infty} \underbrace{\sum_{d|n} d^{2k-1}}_{=\sigma_{2k-1}(n)} q^n.
\end{aligned}$$

In the last double sum we collect together those terms for which nr is constant.

□

Exercise 1.12.

Refer to Exercise 1.11. If $\tau \in H$ prove that

$$G_{2k}\left(-\frac{1}{\tau}\right) = \tau^{2k} G_{2k}(\tau)$$

and deduce that

$$\begin{aligned}
G_{2k}\left(\frac{i}{2}\right) &= (-4)^k G_{2k}(2i) && \text{for all } k \geq 2, \\
G_{2k}(i) &= 0 && \text{if } k \text{ is odd,} \\
G_{2k}(e^{\frac{2\pi i}{3}}) &= 0 && \text{if } k \not\equiv 0 \pmod{3}.
\end{aligned}$$

Proof.

(1)

$$\begin{aligned}
G_{2k}\left(-\frac{1}{\tau}\right) &= \sum_{(m,n) \neq (0,0)} \left(m - \frac{n}{\tau}\right)^{-2k} \\
&= \tau^{2k} \sum_{(m,n) \neq (0,0)} (\tau m - n)^{-2k} \\
&= \tau^{2k} G_{2k}(\tau).
\end{aligned}$$

(2) Let $\tau = 2i$. We have $G_{2k}\left(\frac{i}{2}\right) = (-4)^k G_{2k}(2i)$.

(3) Let $\tau = i$. We have $G_{2k}(i) = (-1)^k G_{2k}(i)$. Hence $G_{2k}(i) = 0$ if k is odd.

(4) Let $\tau = e^{\frac{\pi i}{3}}$. We have $G_{2k}(e^{\frac{2\pi i}{3}}) = e^{\frac{2k\pi i}{3}} G_{2k}(e^{\frac{\pi i}{3}})$. Since

$$e^{\frac{2\pi i}{3}} = -1 + e^{\frac{\pi i}{3}}$$

and each Eisenstein series is a periodic function of τ of period 1, we have $G_{2k}(e^{\frac{2\pi i}{3}}) = G_{2k}(e^{\frac{\pi i}{3}})$. So $G_{2k}(e^{\frac{2\pi i}{3}}) = e^{\frac{2k\pi i}{3}} G_{2k}(e^{\frac{2\pi i}{3}})$. Therefore $G_{2k}(e^{\frac{2\pi i}{3}}) = 0$ if $k \not\equiv 0 \pmod{3}$.

□

Exercise 1.13.

Ramanujan's tau function $\tau(n)$ is defined by the Fourier expansion

$$\Delta(\tau) = (2\pi)^{12} \sum_{n=1}^{\infty} \tau(n) e^{2\pi i n \tau},$$

derived in Theorem 1.19. Prove that

$$\tau(n) = 8000\{(\sigma_3 \circ \sigma_3) \circ \sigma_3\}(n) - 147(\sigma_5 \circ \sigma_5)(n),$$

where $f \circ g$ denotes the Cauchy product of two sequences,

$$(f \circ g)(n) = \sum_{k=0}^n f(k)g(n-k),$$

and $\sigma_\alpha(n) = \sum_{d|n} d^\alpha$ for $n \geq 1$, with $\sigma_3(0) = \frac{1}{240}$, $\sigma_5(0) = -\frac{1}{504}$. (Hint: Theorem 1.18.)

Proof.

(1) Let $q = e^{2\pi i\tau}$. Write

$$g_2(\tau) = \frac{4\pi^4}{3} \left\{ 1 + 240 \sum_{k=1}^{\infty} \sigma_3(k) q^k \right\} = \frac{4\pi^4}{3} \left\{ 240 \sum_{k=0}^{\infty} \sigma_3(k) q^k \right\},$$

$$g_3(\tau) = \frac{8\pi^6}{27} \left\{ 1 - 504 \sum_{k=1}^{\infty} \sigma_5(k) q^k \right\} = \frac{8\pi^6}{27} \left\{ -504 \sum_{k=0}^{\infty} \sigma_5(k) q^k \right\}$$

(Theorem 1.18).

(2) Similar to the proof of Theorem 1.19,

$$\begin{aligned} \Delta(\tau) &= g_2(\tau)^3 - 27g_3(\tau)^2 \\ &= \frac{64\pi^{12}}{27} \left\{ \left(240 \sum_{k=0}^{\infty} \sigma_3(k) q^k \right)^3 - \left(-504 \sum_{k=0}^{\infty} \sigma_5(k) q^k \right)^2 \right\} \\ &= (2\pi)^{12} \left\{ 8000 \left(\sum_{k=0}^{\infty} \sigma_3(k) q^k \right)^3 - 147 \left(\sum_{k=0}^{\infty} \sigma_5(k) q^k \right)^2 \right\} \\ &= (2\pi)^{12} \sum_{n=0}^{\infty} \{ 8000 \{ (\sigma_3 \circ \sigma_3) \circ \sigma_3 \}(n) - 147 (\sigma_5 \circ \sigma_5)(n) \} q^n \\ &= (2\pi)^{12} \sum_{n=1}^{\infty} \{ 8000 \{ (\sigma_3 \circ \sigma_3) \circ \sigma_3 \}(n) - 147 (\sigma_5 \circ \sigma_5)(n) \} q^n. \end{aligned}$$

(Here $8000 \{ (\sigma_3 \circ \sigma_3) \circ \sigma_3 \}(0) - 147 (\sigma_5 \circ \sigma_5)(0) = 0$.)

(3) Therefore

$$\tau(n) = 8000 \{ (\sigma_3 \circ \sigma_3) \circ \sigma_3 \}(n) - 147 (\sigma_5 \circ \sigma_5)(n)$$

for $n \geq 1$.

□

Exercise 1.14. (Lambert series)

A series of the form $\sum_{n=1}^{\infty} f(n) \frac{x^n}{1-x^n}$ is called a **Lambert series**. Assuming absolute convergence, prove that

$$\sum_{n=1}^{\infty} f(n) \frac{x^n}{1-x^n} = \sum_{n=1}^{\infty} F(n) x^n,$$

where

$$F(n) = \sum_{d|n} f(d).$$

Apply this result to obtain the following formulas, valid for $|x| < 1$.

(a)

$$\sum_{n=1}^{\infty} \frac{\mu(n)x^n}{1-x^n} = x.$$

(b)

$$\sum_{n=1}^{\infty} \frac{\varphi(n)x^n}{1-x^n} = \frac{x}{(1-x)^2}.$$

(c)

$$\sum_{n=1}^{\infty} \frac{n^{\alpha}x^n}{1-x^n} = \sum_{n=1}^{\infty} \sigma_{\alpha}(n)x^n.$$

(d)

$$\sum_{n=1}^{\infty} \frac{\lambda(n)x^n}{1-x^n} = \sum_{n=1}^{\infty} x^{n^2}.$$

(e) Use the result in (c) to express $g_2(\tau)$ and $g_3(\tau)$ in terms of Lambert series in $x = e^{2\pi i\tau}$.

Note. In (a), $\mu(n)$ is the Möbius function; In (b), $\varphi(n)$ is Euler's totient; and in (d), $\lambda(n)$ is Liouville's function.

Proof. Similar to the proof of Exercise 1.11.

$$\begin{aligned} \sum_{n=1}^{\infty} f(n) \frac{x^n}{1-x^n} &= \sum_{n=1}^{\infty} f(n) \sum_{r=1}^{\infty} x^{rn} \\ &= \sum_{n=1}^{\infty} \sum_{r=1}^{\infty} f(n) x^{rn} \\ &= \sum_{n=1}^{\infty} \underbrace{\left(\sum_{d|n} f(d) \right)}_{=F(n)} x^n. \end{aligned}$$

□

Proof of (a). Theorem 2.1 in the textbook: *T. M. Apostol, Introduction to Analytic Number Theory* shows that

$$F(n) := \sum_{d|n} \mu(d) = \begin{cases} 1 & \text{if } n = 1, \\ 0 & \text{if } n > 1. \end{cases}$$

Hence

$$\sum_{n=1}^{\infty} \mu(n) \frac{x^n}{1-x^n} = \sum_{n=1}^{\infty} F(n) x^n = x.$$

□

Proof of (b). Theorem 2.2 in the textbook: *T. M. Apostol, Introduction to Analytic Number Theory* shows that $F(n) := \sum_{d|n} \varphi(d) = n$. Hence

$$\sum_{n=1}^{\infty} \varphi(n) \frac{x^n}{1-x^n} = \sum_{n=1}^{\infty} F(n) x^n = \sum_{n=1}^{\infty} n x^n = \frac{x}{(1-x)^2}.$$

□

Proof of (c). Since

$$F(n) := \sum_{d|n} d^\alpha = \sigma_\alpha(n),$$

we have

$$\sum_{n=1}^{\infty} n^\alpha \frac{x^n}{1-x^n} = \sum_{n=1}^{\infty} F(n) x^n = \sum_{n=1}^{\infty} \sigma_\alpha(n) x^n.$$

□

Proof of (d). Theorem 2.19 in the textbook: *T. M. Apostol, Introduction to Analytic Number Theory* shows that

$$F(n) := \sum_{d|n} \lambda(d) = \begin{cases} 1 & \text{if } n \text{ is a square,} \\ 0 & \text{otherwise.} \end{cases}$$

Hence

$$\sum_{n=1}^{\infty} \lambda(n) \frac{x^n}{1-x^n} = \sum_{n=1}^{\infty} F(n) x^n = \sum_{n=1}^{\infty} x^{n^2}.$$

□

Proof of (e).

(1) Let $q = x = e^{2\pi i \tau}$.

$$\begin{aligned} g_2(\tau) &= \frac{4\pi^4}{3} \left\{ 1 + 240 \sum_{k=1}^{\infty} \sigma_3(k) q^k \right\} && \text{(Theorem 1.18)} \\ &= \frac{4\pi^4}{3} \left\{ 1 + 240 \sum_{k=1}^{\infty} \frac{k^3 q^k}{1-q^k} \right\} && ((c)). \end{aligned}$$

(2) Similarly,

$$\begin{aligned} g_3(\tau) &= \frac{8\pi^6}{27} \left\{ 1 - 504 \sum_{k=1}^{\infty} \sigma_5(k) q^k \right\} && \text{(Theorem 1.18)} \\ &= \frac{8\pi^6}{27} \left\{ 1 - 504 \sum_{k=1}^{\infty} \frac{k^5 q^k}{1-q^k} \right\} && ((c)). \end{aligned}$$

□

Exercise 1.15.

Let

$$G(x) = \sum_{n=1}^{\infty} \frac{n^5 x^n}{1 - x^n},$$

and let

$$F(x) = \sum_{\substack{n=1 \\ (n \text{ odd})}}^{\infty} \frac{n^5 x^n}{1 + x^n}.$$

(a) Prove that $F(x) = G(x) - 34G(x^2) + 64G(x^4)$.

(b) Prove that

$$\sum_{\substack{n=1 \\ (n \text{ odd})}}^{\infty} \frac{n^5}{1 + e^{n\pi}} = \frac{31}{504}.$$

(c) Use Theorem 12.17 in the textbook: T. M. Apostol, *Introduction to Analytic Number Theory* to prove the more general result

$$\sum_{\substack{n=1 \\ (n \text{ odd})}}^{\infty} \frac{n^{4k+1}}{1 + e^{n\pi}} = \frac{2^{4k+1} - 1}{8k + 4} B_{4k+2}.$$

Proof of (a).

(1) Consider the general case. Let

$$G(x) = \sum_{n=1}^{\infty} \frac{n^{4k+1} x^n}{1 - x^n},$$

and let

$$F(x) = \sum_{\substack{n=1 \\ (n \text{ odd})}}^{\infty} \frac{n^{4k+1} x^n}{1 + x^n}.$$

Show that $F(x) = G(x) - (2^{4k+1} + 2)G(x^2) + 2^{4k+2}G(x^4)$.

(2) The identity

$$\sum_{n=1}^{\infty} \frac{x^n}{1 + x^n} = \sum_{n=1}^{\infty} \frac{x^n}{1 - x^n} - 2 \sum_{n=1}^{\infty} \frac{x^{2n}}{1 - x^{2n}}$$

is always true. Hence $H(x) := \sum_{n=1}^{\infty} \frac{n^{4k+1} x^n}{1 + x^n} = G(x) - 2G(x^2)$.

(3) Note that

$$\begin{aligned}
H(x) &= \sum_{\substack{n=1 \\ (n \text{ odd})}}^{\infty} \frac{n^{4k+1}x^n}{1+x^n} + \sum_{\substack{n=1 \\ (n \text{ even})}}^{\infty} \frac{n^{4k+1}x^n}{1+x^n} \\
&= F(x) + \sum_{n=1}^{\infty} \frac{(2n)^{4k+1}x^{2n}}{1+x^{2n}} \\
&= F(x) + 2^{4k+1} \sum_{n=1}^{\infty} \frac{n^{4k+1}x^{2n}}{1+x^{2n}} \\
&= F(x) + 2^{4k+1}H(x^2).
\end{aligned}$$

Hence

$$\begin{aligned}
F(x) &= H(x) - 2^{4k+1}H(x^2) \\
&= [G(x) - 2G(x^2)] - 2^{4k+1}[G(x^2) - 2G(x^4)] \\
&= G(x) - (2^{4k+1} + 2)G(x^2) + 2^{4k+2}G(x^4).
\end{aligned}$$

□

Proof of (b). Take $k = 1$ in part (c), we have

$$\sum_{\substack{n=1 \\ (n \text{ odd})}}^{\infty} \frac{n^5}{1+e^{n\pi}} = \frac{31}{12} \cdot \frac{1}{42} = \frac{31}{504}.$$

□

Proof of (c).

(1) Let $q = e^{2\pi i\tau}$. So

$$G_{4k+2}(\tau) = 2\zeta(4k+2) + \frac{2(2\pi i)^{4k+2}}{(4k+1)!} \sum_{n=1}^{\infty} \sigma_{4k+1}(n)q^n \quad (\text{Exercise 1.11})$$

$$= 2\zeta(4k+2) + \frac{2(2\pi i)^{4k+2}}{(4k+1)!} G(q) \quad (\text{Exercise 1.14(c)})$$

Hence

$$\begin{aligned}
& G_{4k+2}(\tau) - (2^{4k+1} + 2)G_{4k+2}(2\tau) + 2^{4k+2}G_{4k+2}(4\tau) \\
&= \left[2\zeta(4k+2) + \frac{2(2\pi i)^{4k+2}}{(4k+1)!} G(q) \right] \\
&\quad - (2^{4k+1} + 2) \left[2\zeta(4k+2) + \frac{2(2\pi i)^{4k+2}}{(4k+1)!} G(q^2) \right] \\
&\quad + 2^{4k+2} \left[2\zeta(4k+2) + \frac{2(2\pi i)^{4k+2}}{(4k+1)!} G(q^4) \right] \\
&= (1 - (2^{4k+1} + 2) + 2^{4k+2}) \cdot 2\zeta(4k+2) \\
&\quad + \frac{2(2\pi i)^{4k+2}}{(4k+1)!} [G(q) - (2^{4k+1} + 2)G(q^2) + 2^{4k+2}G(q^4)] \\
&= (2^{4k+2} - 2)\zeta(4k+2) + \frac{2(2\pi i)^{4k+2}}{(4k+1)!} F(q).
\end{aligned}$$

(2) By taking $\tau = \frac{i}{2}$, we have

$$F(q) = F(e^{-\pi}) = \sum_{\substack{n=1 \\ (n \text{ odd})}}^{\infty} \frac{n^{4k+1}}{1 + e^{n\pi}}$$

and

$$\begin{aligned}
& G_{4k+2}(\tau) - (2^{4k+1} + 2)G_{4k+2}(2\tau) + 2^{4k+2}G_{4k+2}(4\tau) \\
&= G_{4k+2}\left(\frac{i}{2}\right) - (2^{4k+1} + 2)G_{4k+2}(i) + 2^{4k+2}G_{4k+2}(2i) \\
&= (-4)^{2k+1}G_{4k+2}(2i) - (2^{4k+1} + 2) \cdot 0 + 2^{4k+2}G_{4k+2}(2i) \\
&= 0.
\end{aligned}$$

(Exercise 1.12). Hence

$$0 = (2^{4k+2} - 2)\zeta(4k+2) + \frac{2(2\pi i)^{4k+2}}{(4k+1)!} \sum_{\substack{n=1 \\ (n \text{ odd})}}^{\infty} \frac{n^{4k+1}}{1 + e^{n\pi}}.$$

(3) Theorem 12.17 in the textbook: *T. M. Apostol, Introduction to Analytic Number Theory* shows that

$$\zeta(4k+2) = (-1)^{2k+1+1} \frac{(2\pi)^{4k+2} B_{4k+2}}{2(4k+2)!} = \frac{(2\pi)^{4k+2} B_{4k+2}}{2(4k+2)!}.$$

Hence

$$\sum_{\substack{n=1 \\ (n \text{ odd})}}^{\infty} \frac{n^{4k+1}}{1 + e^{n\pi}} = \frac{2^{4k+1} - 1}{8k+4} B_{4k+2}.$$

□