

Chapter 4: Determinants

*Author: Meng-Gen Tsai
Email: plover@gmail.com*

Section 4.1: Determinants of Order 2

Exercise 4.1.1. *Label the following statements as being true or false.*

- (a) *The function $\det : M_{2 \times 2}(F) \rightarrow F$ is a linear transformation.*
- (b) *The determinant of a 2×2 matrix is a linear function of each row of the matrix when the other row is held fixed.*
- (c) *If $A \in M_{2 \times 2}(F)$ and $\det(A) = 0$, then A is invertible.*
- (d) *If u and v are vectors in \mathbb{R}^2 emanating from the origin, then the area of the parallelogram having u and v as adjacent side is*

$$\det \begin{pmatrix} u \\ v \end{pmatrix}.$$

- (e) *A coordinate system is right-handed if and only if its orientation equals 1.*

Proof of (a). False. Example 4.1.1, or take

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \in M_{2 \times 2}(F) \text{ and } B = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \in M_{2 \times 2}(F).$$

Then $\det(A + B) = \det(I_2) = 1 \neq 0 = 0 + 0 = \det(A) + \det(B)$. \square

Proof of (b). True. Proposition 4.1. \square

Proof of (c). False. Proposition 4.2. \square

Proof of (d). False. The area should be

$$O \begin{pmatrix} u \\ v \end{pmatrix} \cdot \det \begin{pmatrix} u \\ v \end{pmatrix} = \left| \det \begin{pmatrix} u \\ v \end{pmatrix} \right|.$$

\square

Proof of (e). True. See Exercise 4.1.12. \square

Exercise 4.1.2. *Compute the determinants of the following elements of $M_{2 \times 2}(\mathbb{R})$.*

$$(a) \begin{pmatrix} 6 & -3 \\ 2 & 4 \end{pmatrix}$$

$$(b) \begin{pmatrix} -5 & 2 \\ 6 & 1 \end{pmatrix}$$

$$(c) \begin{pmatrix} 8 & 0 \\ 3 & -1 \end{pmatrix}$$

Proof of (a).

$$\det \begin{pmatrix} 6 & -3 \\ 2 & 4 \end{pmatrix} = 6 \cdot 4 - (-3) \cdot 2 = 24 + 6 = 30.$$

□

Proof of (b).

$$\det \begin{pmatrix} -5 & 2 \\ 6 & 1 \end{pmatrix} = (-5) \cdot 1 - 2 \cdot 6 = -5 - 12 = -17.$$

□

Proof of (c).

$$\det \begin{pmatrix} 8 & 0 \\ 3 & -1 \end{pmatrix} = 8 \cdot (-1) - 0 \cdot 3 = -8.$$

□

Exercise 4.1.3. *Compute the determinants of the following elements of $M_{2 \times 2}(\mathbb{C})$.*

$$(a) \begin{pmatrix} -1 + i & 1 - 4i \\ 3 + 2i & 2 - 3i \end{pmatrix}$$

$$(b) \begin{pmatrix} 5 - 2i & 6 + 4i \\ -3 + i & 7i \end{pmatrix}$$

$$(c) \begin{pmatrix} 2i & 3 \\ 4 & 6i \end{pmatrix}$$

Proof of (a).

$$\begin{aligned} \det \begin{pmatrix} -1 + i & 1 - 4i \\ 3 + 2i & 2 - 3i \end{pmatrix} &= (-1 + i) \cdot (2 - 3i) - (1 - 4i) \cdot (3 + 2i) \\ &= (1 + 5i) - (11 - 10i) \\ &= -10 + 15i. \end{aligned}$$

□

Proof of (b).

$$\begin{aligned}\det \begin{pmatrix} 5-2i & 6+4i \\ -3+i & 7i \end{pmatrix} &= (5-2i) \cdot (7i) - (6+4i) \cdot (-3+i) \\ &= (14+35i) - (-22-6i) \\ &= 36+41i.\end{aligned}$$

□

Proof of (c).

$$\det \begin{pmatrix} 2i & 3 \\ 4 & 6i \end{pmatrix} = (2i) \cdot (6i) - 3 \cdot 4 = -12 - 12 = -24.$$

□

Exercise 4.1.4. For each of the following pairs of vectors u and v in \mathbb{R}^2 , compute the area of the parallelogram determined by u and v .

(a) $u = (3, -2)$ and $v = (2, 5)$

(b) $u = (1, 3)$ and $v = (-3, 1)$

(c) $u = (4, -1)$ and $v = (-6, -2)$

(d) $u = (3, 4)$ and $v = (2, -6)$

Proof of (a).

$$\left| \det \begin{pmatrix} 3 & -2 \\ 2 & 5 \end{pmatrix} \right| = |19| = 19.$$

□

Proof of (b).

$$\left| \det \begin{pmatrix} 1 & 3 \\ -3 & 1 \end{pmatrix} \right| = |10| = 10.$$

□

Proof of (c).

$$\left| \det \begin{pmatrix} 4 & -1 \\ -6 & -2 \end{pmatrix} \right| = |-14| = 14.$$

□

Proof of (d).

$$\left| \det \begin{pmatrix} 3 & 4 \\ 2 & -6 \end{pmatrix} \right| = |-26| = 26.$$

□

Exercise 4.1.5. *Prove that if B is the matrix obtained by interchanging the rows of a 2×2 matrix A , then $\det(B) = -\det(A)$.*

Proof. Write

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_{2 \times 2}(F).$$

Then

$$B = \begin{pmatrix} c & d \\ a & b \end{pmatrix} \in M_{2 \times 2}(F).$$

Then $\det(B) = cb - ad = -(ad - bc) = -\det(A)$. □

Exercise 4.1.6. *Prove that if the two columns of $A \in M_{2 \times 2}(F)$ are identical, then $\det(A) = 0$.*

Proof. By assumption, write

$$A = \begin{pmatrix} a & a \\ c & c \end{pmatrix} \in M_{2 \times 2}(F).$$

Then $\det(A) = ac - ac = 0$. □

Exercise 4.1.7. *Prove that for any $A \in M_{2 \times 2}(F)$, $\det(A^t) = \det(A)$.*

Proof. Write

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_{2 \times 2}(F),$$

then

$$A^t = \begin{pmatrix} a & c \\ b & d \end{pmatrix} \in M_{2 \times 2}(F).$$

So $\det(A) = ad - bc = ad - cb = \det(A^t)$. □

Exercise 4.1.8. *Prove that if $A \in M_{2 \times 2}(F)$ is upper triangular, then $\det(A)$ equals the product of the diagonal entries of A .*

Proof. Write

$$A = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \in \mathbb{M}_{2 \times 2}(F)$$

since A is upper triangular. Then $\det(A) = ad$, which is equal to the product of the diagonal entries, a and d , of A . \square

Exercise 4.1.9. Prove that for any $A, B \in \mathbb{M}_{2 \times 2}(F)$ we have $\det(AB) = \det(A) \cdot \det(B)$.

Proof. Write

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathbb{M}_{2 \times 2}(F),$$

$$B = \begin{pmatrix} e & f \\ g & h \end{pmatrix} \in \mathbb{M}_{2 \times 2}(F).$$

Then

$$AB = \begin{pmatrix} ae + bg & af + bh \\ ce + dg & cf + dh \end{pmatrix} \in \mathbb{M}_{2 \times 2}(F).$$

A direct calculation shows

$$\begin{aligned} \det(AB) &= (ae + bg)(cf + dh) - (af + bh)(ce + dg) \\ &= (acef + adeh + bcfg + bdgh) - (acef + adfg + bceh + bdgh) \\ &= adeh + bcfg - adfg - bceh \\ &= (ad - bc)(eh - fg) \\ &= \det(A) \det(B). \end{aligned}$$

\square

Exercise 4.1.10. The *classical adjoint* of a 2×2 matrix $A \in \mathbb{M}_{2 \times 2}(F)$ is the matrix

$$C = \begin{pmatrix} A_{22} & -A_{12} \\ -A_{21} & A_{11} \end{pmatrix}.$$

Prove

- (a) $CA = AC = [\det(A)]I$.
- (b) $\det(C) = \det(A)$.
- (c) The classical adjoint of A^t is C^t .
- (d) If A is invertible, then $A^{-1} = [\det(A)]^{-1}C$.

Note that

$$A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}.$$

Proof of (a).

$$\begin{aligned} CA &= \begin{pmatrix} A_{22} & -A_{12} \\ -A_{21} & A_{11} \end{pmatrix} \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \\ &= \begin{pmatrix} A_{22}A_{11} - A_{12}A_{21} & A_{22}A_{12} - A_{12}A_{22} \\ -A_{21}A_{11} + A_{11}A_{21} & -A_{21}A_{12} + A_{11}A_{22} \end{pmatrix} \\ &= \begin{pmatrix} \det(A) & 0 \\ 0 & \det(A) \end{pmatrix} \\ &= [\det(A)]I. \end{aligned}$$

$$\begin{aligned} AC &= \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \begin{pmatrix} A_{22} & -A_{12} \\ -A_{21} & A_{11} \end{pmatrix} \\ &= \begin{pmatrix} A_{11}A_{22} - A_{12}A_{21} & -A_{11}A_{12} + A_{12}A_{11} \\ A_{21}A_{22} - A_{22}A_{21} & -A_{21}A_{12} + A_{22}A_{11} \end{pmatrix} \\ &= \begin{pmatrix} \det(A) & 0 \\ 0 & \det(A) \end{pmatrix} \\ &= [\det(A)]I. \end{aligned}$$

□

Proof of (b).

$$\begin{aligned} \det(C) &= A_{22}A_{11} - (-A_{12})(-A_{21}) \\ &= A_{11}A_{22} - A_{12}A_{21} \\ &= \det(A). \end{aligned}$$

□

Proof of (c).

$$A^t = \begin{pmatrix} A_{11} & A_{21} \\ A_{12} & A_{22} \end{pmatrix}.$$

The classical adjoint of A^t is

$$\begin{pmatrix} A_{22} & -A_{21} \\ -A_{12} & A_{11} \end{pmatrix} = C^t.$$

□

Proof of (d). Proposition 4.2. \square

Exercise 4.1.11. Let $\delta : M_{2 \times 2}(F) \rightarrow F$ be a function with the following three properties.

- (i) δ is a linear function of each row of the matrix when the other row is held fixed.
- (ii) If the two rows of $A \in M_{2 \times 2}(F)$ are identical, then $\delta(A) = 0$.
- (iii) If I is the 2×2 identity matrix, then $\delta(I) = 1$.

Prove that $\delta(A) = \det(A)$ for all $A \in M_{2 \times 2}(F)$. (This result is generalized in Section 4.5.)

Proof. Write

$$A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}.$$

- (1) If u, v are elements of F^2 and k is a scalar, then

$$\delta \begin{pmatrix} u \\ v + ku \end{pmatrix} = \delta \begin{pmatrix} u + kv \\ v \end{pmatrix} = \delta \begin{pmatrix} u \\ v \end{pmatrix}.$$

In fact,

$$\begin{aligned} \delta \begin{pmatrix} u \\ v + ku \end{pmatrix} &= \delta \begin{pmatrix} u \\ v \end{pmatrix} + \delta \begin{pmatrix} u \\ ku \end{pmatrix} && \text{(Property (i))} \\ &= \delta \begin{pmatrix} u \\ v \end{pmatrix} + k\delta \begin{pmatrix} u \\ u \end{pmatrix} && \text{(Property (i))} \\ &= \delta \begin{pmatrix} u \\ v \end{pmatrix}. && \text{(Property (ii))} \end{aligned}$$

Similarly, $\delta \begin{pmatrix} u + kv \\ v \end{pmatrix} = \delta \begin{pmatrix} u \\ v \end{pmatrix}.$

- (2) If u, v are elements of F^2 , then

$$\delta \begin{pmatrix} u \\ v \end{pmatrix} = -\delta \begin{pmatrix} v \\ u \end{pmatrix}.$$

In fact,

$$\begin{aligned} 0 &= \delta \begin{pmatrix} u + v \\ u + v \end{pmatrix} && \text{(Property (ii))} \\ &= \delta \begin{pmatrix} u + v \\ u \end{pmatrix} + \delta \begin{pmatrix} u + v \\ v \end{pmatrix} && \text{(Property (i))} \\ &= \delta \begin{pmatrix} v \\ u \end{pmatrix} + \delta \begin{pmatrix} u \\ v \end{pmatrix}. && ((1)) \end{aligned}$$

(3) If v is an element of F^2 , then

$$\delta \begin{pmatrix} 0 \\ v \end{pmatrix} = 0.$$

In fact,

$$\begin{aligned} \delta \begin{pmatrix} 0 \\ v \end{pmatrix} &= \delta \begin{pmatrix} 0+0 \\ v \end{pmatrix} \\ &= \delta \begin{pmatrix} 0 \\ v \end{pmatrix} + \delta \begin{pmatrix} 0 \\ v \end{pmatrix}. \end{aligned} \quad (\text{Property (i)})$$

$$\text{In particular, } \delta \begin{pmatrix} 0 \\ v \end{pmatrix} = 0 = \det \begin{pmatrix} 0 \\ v \end{pmatrix}.$$

(4) To show $\delta(A) = \det(A)$, we consider three possible cases about the first row: $A_{11} \neq 0$, $A_{12} \neq 0$, or $A_{11} = A_{12} = 0$. The case $A_{11} = A_{12} = 0$ is proved in (3). We prove the rest two cases in (5) and (6). Write

$$u = (A_{11}, A_{12}) \text{ and } v = (A_{21}, A_{22}).$$

(5) Show that $\delta(A) = \det(A)$ if $A_{11} \neq 0$. So

$$\begin{aligned} \delta(A) &= \delta \begin{pmatrix} u \\ v \end{pmatrix} \\ &= \delta \begin{pmatrix} u \\ v - \frac{A_{21}}{A_{11}}u \end{pmatrix} && ((1)) \\ &= \delta \begin{pmatrix} A_{11} & A_{12} \\ 0 & A_{22} - \frac{A_{12}A_{21}}{A_{11}} \end{pmatrix} \\ &= \left(A_{22} - \frac{A_{12}A_{21}}{A_{11}} \right) \delta \begin{pmatrix} A_{11} & A_{12} \\ 0 & 1 \end{pmatrix} && (\text{Property (i)}) \\ &= \left(A_{22} - \frac{A_{12}A_{21}}{A_{11}} \right) \delta \begin{pmatrix} A_{11} & 0 \\ 0 & 1 \end{pmatrix} && ((1)) \\ &= A_{11} \left(A_{22} - \frac{A_{12}A_{21}}{A_{11}} \right) \delta \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} && (\text{Property (i)}) \\ &= \det(A) \delta(I) \\ &= \det(A). && (\text{Property (iii)}) \end{aligned}$$

(6) Show that $\delta(A) = \det(A)$ if $A_{12} \neq 0$. So

$$\begin{aligned}
\delta(A) &= \delta \begin{pmatrix} u \\ v \end{pmatrix} \\
&= \delta \begin{pmatrix} u \\ v - \frac{A_{22}}{A_{12}}u \end{pmatrix} && ((1)) \\
&= \delta \begin{pmatrix} A_{11} & A_{12} \\ A_{21} - \frac{A_{22}A_{11}}{A_{12}} & 0 \end{pmatrix} \\
&= \left(A_{21} - \frac{A_{22}A_{11}}{A_{12}} \right) \delta \begin{pmatrix} A_{11} & A_{12} \\ 1 & 0 \end{pmatrix} && (\text{Property (i)}) \\
&= \left(A_{21} - \frac{A_{22}A_{11}}{A_{12}} \right) \delta \begin{pmatrix} 0 & A_{12} \\ 1 & 0 \end{pmatrix} && ((1)) \\
&= A_{12} \left(A_{21} - \frac{A_{22}A_{11}}{A_{12}} \right) \delta \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} && (\text{Property (i)}) \\
&= -A_{12} \left(A_{21} - \frac{A_{22}A_{11}}{A_{12}} \right) \delta \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} && ((2)) \\
&= \det(A) \delta(I) \\
&= \det(A). && (\text{Property (iii)})
\end{aligned}$$

□

Exercise 4.1.12. Let $\{u, v\}$ be an ordered basis for \mathbb{R}^2 . Prove that

$$O \begin{pmatrix} u \\ v \end{pmatrix} = 1$$

if and only if $\{u, v\}$ forms a right-handed coordinate system. (Hint: Recall the definition of a rotation given in Example 2.1.2.)

If $\beta = \{u, v\}$ is an ordered basis for \mathbb{R}^2 , define the orientation of β as

$$O \begin{pmatrix} u \\ v \end{pmatrix} = \frac{\det \begin{pmatrix} u \\ v \end{pmatrix}}{\left| \det \begin{pmatrix} u \\ v \end{pmatrix} \right|}.$$

A coordinate system $\{u, v\}$ is called right-handed if u can be rotated in a counterclockwise direction through an angle θ ($0 < \theta < \pi$) to coincide with v .

Example 2.1.2. For any angle θ , define $T_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$T_\theta(a_1, a_2) = (a_1 \cos \theta - a_2 \sin \theta, a_1 \sin \theta + a_2 \cos \theta).$$

T_θ is called the rotation by θ .

Proof.

- (1) By Example 2.1.2, for any coordinate system $\{u, v\}$, there is $0 < \theta < 2\pi$ and $\alpha > 0$ such that $v = \alpha T_\theta(u)$. Write $u = (u_1, u_2) \in \mathbb{R}^2, v = (v_1, v_2) \in \mathbb{R}^2$.

- (2) Calculate $\det \begin{pmatrix} u \\ v \end{pmatrix}$.

$$\begin{aligned} \det \begin{pmatrix} u \\ v \end{pmatrix} &= \det \begin{pmatrix} u \\ \alpha T_\theta(u) \end{pmatrix} \\ &= \alpha \det \begin{pmatrix} u \\ T_\theta(u) \end{pmatrix} \\ &= \alpha \det \begin{pmatrix} u_1 & u_2 \\ u_1 \cos \theta - u_2 \sin \theta & u_1 \sin \theta + u_2 \cos \theta \end{pmatrix} \\ &= \alpha(u_1^2 + u_2^2) \sin \theta. \end{aligned}$$

- (3)

$$\begin{aligned} O \begin{pmatrix} u \\ v \end{pmatrix} = 1 &\iff \det \begin{pmatrix} u \\ v \end{pmatrix} = \alpha(u_1^2 + u_2^2) \sin \theta > 0 \\ &\iff \sin \theta > 0 \\ &\iff 0 < \theta < \pi \\ &\iff \{u, v\} \text{ is a right-handed coordinate system.} \end{aligned}$$

□

Section 4.2: Determinants of Order n

Exercise 4.2.2. Find the value of k that satisfies the following equation.

$$\det \begin{pmatrix} 3a_1 & 3a_2 & 3a_3 \\ 3b_1 & 3b_2 & 3b_3 \\ 3c_1 & 3c_2 & 3c_3 \end{pmatrix} = k \det \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix}$$

Proof (Exercise 4.2.25). By Exercise 4.2.25, $\det(3A) = 3^3 \det(A)$ for any $A \in M_{3 \times 3}(F)$, or $k = 3^3 = 27$. □

Exercise 4.2.26. Let $A \in M_{n \times n}(F)$. Under what conditions is $\det(-A) = \det(A)$?

Proof (Exercise 4.2.25). By Exercise 4.2.25, $\det(-A) = (-1)^n \det(A)$ for any $A \in M_{n \times n}(F)$. That is, n is even if and only if $\det(-A) = \det(A)$. \square

Section 4.3: Properties of Determinants

Exercise 4.3.9. A matrix $M \in M_{n \times n}(\mathbb{C})$ is called nilpotent if, for some positive integer k , $M^k = O$, where O is the $n \times n$ zero matrix. Prove that if M is nilpotent, then $\det(M) = 0$.

Proof. Given any nilpotent matrix $M \in M_{n \times n}(\mathbb{C})$ such that $M^k = O$ for some $k \in \mathbb{Z}^+$.

$$\begin{aligned} M^k = O &\implies \det(M^k) = \det(O) \\ &\iff \det(M)^k = 0 && \text{(Theorem 4.7)} \\ &\iff \det(M) = 0. \end{aligned}$$

\square

Exercise 4.3.11. A matrix $Q \in M_{n \times n}(\mathbb{R})$ is called orthogonal if $QQ^t = I$. Prove that if Q is orthogonal, then $\det(Q) = \pm 1$.

Proof. By the orthogonality of Q , $QQ^t = I$. So

$$\begin{aligned} QQ^t = I &\implies \det(QQ^t) = \det(I) \\ &\iff \det(Q) \det(Q^t) = \det(I) && \text{(Theorem 4.7)} \\ &\iff \det(Q) \det(Q) = \det(I) && \text{(Theorem 4.8)} \\ &\iff \det(Q)^2 = 1 && \text{(Example 4.2.4)} \\ &\iff \det(Q) = \pm 1. \end{aligned}$$

\square

Exercise 4.3.14. Prove that if $A, B \in M_{n \times n}(F)$ are similar, then $\det(A) = \det(B)$.

Proof. Since A, B are similar, there exists an invertible matrix Q such that

$B = Q^{-1}AQ$. So

$$\begin{aligned}\det(B) &= \det(Q^{-1}AQ) \\ &= \det(Q^{-1}) \det(A) \det(Q) && \text{(Theorem 4.7)} \\ &= \det(Q) \det(Q^{-1}) \det(A) && (F \text{ is field}) \\ &= \det(QQ^{-1}) \det(A) && \text{(Theorem 4.7)} \\ &= \det(I) \det(A) \\ &= 1 \cdot \det(A) && \text{(Example 4.2.4)} \\ &= \det(A).\end{aligned}$$

□