

Notes on the book:
*P.J. Hilton and U. Stammbach, A
Course in Homological Algebra*

Meng-Gen Tsai
plover@gmail.com

September 1, 2021

Contents

Chapter I: Modules	2
§1. Modules	2
Exercise 1.1. (Diagram chasing)	2
Exercise 1.2. (Five lemma)	6

Chapter I: Modules

§1. Modules

Exercise 1.1. (Diagram chasing)

Complete the proof of Lemma 1.1. Show moreover that α is surjective (resp. injective) if α' , α'' are surjective (resp. injective).

Lemma 1.1. Let $0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0$ and $0 \rightarrow B' \rightarrow B \rightarrow B'' \rightarrow 0$ be two short exact sequences. Suppose that in the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & A' & \xrightarrow{\mu} & A & \xrightarrow{\varepsilon} & A'' \longrightarrow 0 \\ & & \downarrow \alpha' & & \downarrow \alpha & & \downarrow \alpha'' \\ 0 & \longrightarrow & B' & \xrightarrow{\mu'} & B & \xrightarrow{\varepsilon'} & B'' \longrightarrow 0 \end{array}$$

any two of the three homomorphisms α' , α , α'' are isomorphisms. Then the third is an isomorphism, too.

Proof (Diagram chasing).

(1) Show that α is surjective if α' , α'' are surjective.

- (a) Take any $b \in B$, it suffices to find $a \in A$ such that $\alpha a = b$.
- (b) Consider the commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\varepsilon} & A'' \\ \downarrow \alpha & & \downarrow \alpha'' \\ B & \xrightarrow{\varepsilon'} & B'' \end{array}$$

$\varepsilon' b \in B''$. By the surjectivity of α'' , $\exists a'' \in A''$ such that $\alpha'' a'' = \varepsilon' b$. By the surjectivity of ε , $\exists \bar{a} \in A$ such that $\varepsilon \bar{a} = a''$. Hence

$$\begin{aligned} \varepsilon'(b - \alpha \bar{a}) &= \varepsilon' b - \varepsilon' \alpha \bar{a} \\ &= \varepsilon' b - \alpha'' \varepsilon \bar{a} && \text{(The diagram commutes)} \\ &= \varepsilon' b - \alpha'' a'' \\ &= \varepsilon' b - \varepsilon' b \\ &= 0. \end{aligned}$$

- (c) Consider the short exact sequence

$$0 \longrightarrow B' \xrightarrow{\mu'} B \xrightarrow{\varepsilon'} B'' \longrightarrow 0$$

As $\varepsilon'(b - \alpha \bar{a}) = 0$, $\exists b' \in B'$ such that $\mu' b' = b - \alpha \bar{a}$.

(d) Consider the commutative diagram

$$\begin{array}{ccc} A' & \xrightarrow{\mu} & A \\ \downarrow \alpha' & & \downarrow \alpha \\ B' & \xrightarrow{\mu'} & B \end{array}$$

By the surjectivity of α' , $\exists a' \in A'$ such that $\alpha'a' = b'$. Hence

$$\begin{aligned} \alpha(\mu a' + \bar{a}) &= \alpha\mu a' + \alpha\bar{a} \\ &= \mu'\alpha'a' + \alpha\bar{a} && \text{(The diagram commutes)} \\ &= \mu'b' + \alpha\bar{a} \\ &= (b - \alpha\bar{a}) + \alpha\bar{a} \\ &= b. \end{aligned}$$

Therefore, there exists $a := \mu a' + \bar{a}$ such that $\alpha a = b$.

(2) Show that α is injective if α' , α'' are injective.

(a) It suffices to show that $\ker \alpha = 0$. Take $a \in \ker \alpha$. ($\alpha(a) = \alpha a = 0$.)

(b) Consider the commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\varepsilon} & A'' \\ \downarrow \alpha & & \downarrow \alpha'' \\ B & \xrightarrow{\varepsilon'} & B'' \end{array}$$

we have $0 = \varepsilon' \alpha a = \alpha'' \varepsilon a$. By the injectivity of α'' , $\varepsilon a = 0$.

(c) Consider the short exact sequence

$$0 \longrightarrow A' \xrightarrow{\mu} A \xrightarrow{\varepsilon} A'' \longrightarrow 0$$

As $\varepsilon a = 0$, $\exists a' \in A'$ such that $\mu a' = a$.

(d) Consider the commutative diagram

$$\begin{array}{ccc} A' & \xrightarrow{\mu} & A \\ \downarrow \alpha' & & \downarrow \alpha \\ B' & \xrightarrow{\mu'} & B \end{array}$$

$0 = \alpha a = \alpha\mu a' = \mu'\alpha'a'$. By the injectivity of $\mu'\alpha'$, $a' = 0$.
Therefore, $a = \mu a' = 0$.

(3) Suppose α is surjective. Show that α'' is surjective.

(a) Take any $b'' \in B''$, it suffices to find $a'' \in A''$ such that $\alpha'' a'' = b''$.

(b) Consider the commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\varepsilon} & A'' \\ \downarrow \alpha & & \downarrow \alpha'' \\ B & \xrightarrow{\varepsilon'} & B'' \end{array}$$

By the surjectivity of ε' , $\exists b \in B$ such that $\varepsilon'b = b''$. By the surjectivity of α , $\exists a \in A$ such that $\alpha a = b$. Take $a'' := \varepsilon a \in A''$. Hence

$$\begin{aligned} \alpha'' a'' &= \alpha'' \varepsilon a \\ &= \varepsilon' \alpha a && \text{(The diagram commutes)} \\ &= \varepsilon' b \\ &= b''. \end{aligned}$$

(4) Suppose α' is surjective and α is injective. Show that α'' is injective.

(a) It suffices to show that $\ker \alpha'' = 0$. Take $a'' \in \ker \alpha''$. ($\alpha''(a'') = \alpha'' a'' = 0$.)

(b) Consider the commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\varepsilon} & A'' \\ \downarrow \alpha & & \downarrow \alpha'' \\ B & \xrightarrow{\varepsilon'} & B'' \end{array}$$

By the surjectivity of ε , $\exists a \in A$ such that $\varepsilon a = a''$. So

$$\begin{aligned} 0 &= \alpha'' a'' \\ &= \alpha'' \varepsilon a \\ &= \varepsilon' \alpha a. && \text{(The diagram commutes)} \end{aligned}$$

(c) Consider the short exact sequence

$$0 \longrightarrow B' \xrightarrow{\mu'} B \xrightarrow{\varepsilon'} B'' \longrightarrow 0$$

As $\varepsilon'(\alpha a) = 0$, $\exists b' \in B'$ such that $\mu' b' = \alpha a$.

(d) Consider the commutative diagram

$$\begin{array}{ccc} A' & \xrightarrow{\mu} & A \\ \downarrow \alpha' & & \downarrow \alpha \\ B' & \xrightarrow{\mu'} & B \end{array}$$

By surjectivity of α' , $\exists a' \in A'$ such that $\alpha' a' = b'$. So

$$\begin{aligned} \alpha a &= \mu' b' \\ &= \mu' \alpha' a' \\ &= \alpha \mu a'. && \text{(The diagram commutes)} \end{aligned}$$

By the injectivity of α , $a = \mu a'$. Hence

$$a'' = \varepsilon a = \varepsilon \mu a' = 0.$$

Therefore $\ker \alpha'' = 0$.

(5) By (3)(4), α'' is an isomorphism if both α' and α are isomorphisms.

(6) Suppose α is surjective and α'' is injective. Show that α' is surjective.

(a) Take any $b' \in B'$, it suffices to find $a' \in A'$ such that $\alpha' a' = b'$. Let $b := \mu' b' \in B$ and note that $\varepsilon' b = 0$ by the exactness of

$$0 \rightarrow B' \rightarrow B \rightarrow B'' \rightarrow 0.$$

(b) Consider the commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{\varepsilon} & A'' \\ \downarrow \alpha & & \downarrow \alpha'' \\ B & \xrightarrow{\varepsilon'} & B'' \end{array}$$

By the surjectivity of α , $\exists a \in A$ such that $\alpha a = b$. So

$$\begin{aligned} 0 &= \varepsilon' b \\ &= \varepsilon' \alpha a \\ &= \alpha'' \varepsilon a. \end{aligned} \quad (\text{The diagram commutes})$$

By the injectivity of α'' , $\varepsilon a = 0$.

(c) Consider the short exact sequence

$$0 \longrightarrow A' \xrightarrow{\mu} A \xrightarrow{\varepsilon} A'' \longrightarrow 0$$

As $\varepsilon a = 0$, $\exists a' \in A'$ such that $\mu a' = a$.

(d) Consider the commutative diagram

$$\begin{array}{ccc} A' & \xrightarrow{\mu} & A \\ \downarrow \alpha' & & \downarrow \alpha \\ B' & \xrightarrow{\mu'} & B \end{array}$$

Note that

$$\begin{aligned} \mu'(\alpha' a') &= \mu' \alpha' a' \\ &= \alpha \mu a' & (\text{The diagram commutes}) &= \alpha a \\ &= b \\ &= \mu' b'. \end{aligned}$$

By the injectivity of μ' , $b' = \alpha' a'$ for some $a' \in A'$.

(7) Suppose α is injective. Show that α' is injective.

(a) It suffices to show that $\ker \alpha' = 0$. Take $a' \in \ker \alpha'$. ($\alpha'(a') = \alpha'a' = 0$.)

(b) Consider the commutative diagram

$$\begin{array}{ccc} A' & \xrightarrow{\mu} & A \\ \downarrow \alpha' & & \downarrow \alpha \\ B' & \xrightarrow{\mu'} & B \end{array}$$

Note that

$$\begin{aligned} 0 &= \mu'0 \\ &= \mu'\alpha'a' \\ &= \alpha\mu a'. \end{aligned} \quad (\text{The diagram commutes})$$

The injectivity of $\alpha\mu$ shows that $a' = 0$.

(8) By (6)(7), α' is an isomorphism if both α and α'' are isomorphisms.

□

Exercise 1.2. (Five lemma)

Show that, given a commutative diagram

$$\begin{array}{ccccccccc} \cdots & \longrightarrow & A_1 & \longrightarrow & A_2 & \longrightarrow & A_3 & \longrightarrow & A_4 & \longrightarrow & A_5 & \longrightarrow & \cdots \\ & & \downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & \downarrow \varphi_4 & & \downarrow \varphi_5 & & \\ \cdots & \longrightarrow & B_1 & \longrightarrow & B_2 & \longrightarrow & B_3 & \longrightarrow & B_4 & \longrightarrow & B_5 & \longrightarrow & \cdots \end{array}$$

with exact rows, in which $\varphi_1, \varphi_2, \varphi_4, \varphi_5$ are isomorphisms, then φ_3 is also an isomorphism. Can we weaken the hypotheses in a reasonable way?

One reasonable hypotheses:

- (a) If φ_1 is surjective and φ_2, φ_4 is injective, then φ_3 is injective.
- (b) If φ_5 is injective and φ_2, φ_4 is surjective, then φ_3 is surjective.

Proof of (a).

(1) Write

$$\begin{array}{ccccccccc} \cdots & \longrightarrow & A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & A_4 & \xrightarrow{\alpha_4} & A_5 & \longrightarrow & \cdots \\ & & \downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & \downarrow \varphi_4 & & \downarrow \varphi_5 & & \\ \cdots & \longrightarrow & B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & B_4 & \xrightarrow{\beta_4} & B_5 & \longrightarrow & \cdots \end{array}$$

Take $a \in \ker(\varphi_3)$ and then we need to show $a = 0$.

(2) The commutative diagram

$$\begin{array}{ccc} A_3 & \xrightarrow{\alpha_3} & A_4 \\ \downarrow \varphi_3 & & \downarrow \varphi_4 \\ B_3 & \xrightarrow{\beta_3} & B_4 \end{array}$$

implies that $0 = \beta_3 0 = \beta_3 \varphi_3 a = \varphi_4 \alpha_3 a$. The injectivity of φ_4 implies that $\alpha_3 a = 0$.

(3) The exact sequence

$$\cdots \longrightarrow A_2 \xrightarrow{\alpha_2} A_3 \xrightarrow{\alpha_3} A_4 \longrightarrow \cdots$$

shows that $a \in \ker(\alpha_3) = \text{im}(\alpha_2)$. So there exists $a_2 \in A_2$ such that $\alpha_2 a_2 = a$.

(4) The commutative diagram

$$\begin{array}{ccc} A_2 & \xrightarrow{\alpha_2} & A_3 \\ \downarrow \varphi_2 & & \downarrow \varphi_3 \\ B_2 & \xrightarrow{\beta_2} & B_3 \end{array}$$

implies that $0 = \varphi_3 a = \varphi_3 \alpha_2 a_2 = \beta_2 \varphi_2 a_2$.

(5) The exact sequence

$$\cdots \longrightarrow B_1 \xrightarrow{\beta_1} B_2 \xrightarrow{\beta_2} B_3 \longrightarrow \cdots$$

shows that $\varphi_2 a_2 \in \ker(\beta_2) = \text{im}(\beta_1)$. So there exists $b_1 \in B_1$ such that $\varphi_2 a_2 = \beta_1 b_1$.

(6) Consider the commutative diagram

$$\begin{array}{ccc} A_1 & \xrightarrow{\alpha_1} & A_2 \\ \downarrow \varphi_1 & & \downarrow \varphi_2 \\ B_1 & \xrightarrow{\beta_1} & B_2 \end{array}$$

The surjectivity of φ_i implies that $\exists a_1 \in A_1$ such that $\varphi_1 a_1 = b_1$. Hence the commutative diagram implies that $\varphi_2(\alpha_1 a_1) = \varphi_2 \alpha_1 a_1 = \beta_1 \varphi_1 a_1 = \beta_1 b_1 = \varphi_2 a_2$. The injectivity of φ_2 implies that $\alpha_1 a_1 = a_2$.

(7) The exact sequence

$$\cdots \longrightarrow A_1 \xrightarrow{\alpha_1} A_2 \xrightarrow{\alpha_2} A_3 \longrightarrow \cdots$$

shows that $a = \alpha_2 a_2 = \alpha_2 \alpha_1 a_1 = 0$. Therefore φ_3 is injective.

□

Proof of (b).

(1)

□