

Chapter 10: Integration of Differential Forms

Author: Meng-Gen Tsai

Email: plover@gmail.com

Exercise 10.1. Let H be a compact convex set in \mathbb{R}^k , with nonempty interior. Let $f \in \mathcal{C}(H)$, put $f(\mathbf{x}) = 0$ in the complement of H , and define $\int_H f$ as in Definition 10.3. Prove that $\int_H f$ is independent of the order in which the k integrations are carried out. (Hint: Approximate f by functions that are continuous on \mathbb{R}^k and whose supports are in H , as was done in Example 10.4.)

Proof.

(1)

(2)

□

Exercise 10.2. For $i = 1, 2, 3, \dots$, let $\varphi_i \in \mathcal{C}(\mathbb{R}^1)$ have support in $(2^{-i}, 2^{1-i})$, such that $\int \varphi_i = 1$. Put

$$f(x, y) = \sum_{i=1}^{\infty} [\varphi_i(x) - \varphi_{i+1}(x)] \varphi_i(y)$$

Then f has compact support in \mathbb{R}^2 , f is continuous except at $(0, 0)$, and

$$\int dy \int f(x, y) dx = 0 \quad \text{but} \quad \int dx \int f(x, y) dy = 1.$$

Observe that f is unbounded in every neighborhood of $(0, 0)$.

Proof.

(1)

(2)

□

Exercise 10.3.

- (a) If \mathbf{F} is as in Theorem 10.7, put $\mathbf{A} = \mathbf{F}'(\mathbf{0})$, $\mathbf{F}_1(\mathbf{x}) = \mathbf{A}^{-1}\mathbf{F}(\mathbf{x})$. Then $\mathbf{F}_1(\mathbf{0}) = \mathbf{I}$. Show that

$$\mathbf{F}_1(\mathbf{x}) = \mathbf{G}_n \circ \mathbf{G}_{n-1} \circ \cdots \circ \mathbf{G}_1(\mathbf{x})$$

in some neighborhood of $\mathbf{0}$, for certain primitive mappings $\mathbf{G}_1, \dots, \mathbf{G}_n$. This gives another version of Theorem 10.7:

$$\mathbf{F}(\mathbf{x}) = \mathbf{F}'(\mathbf{0})\mathbf{G}_n \circ \mathbf{G}_{n-1} \circ \dots \circ \mathbf{G}_1(\mathbf{x}).$$

- (b) Prove that the mapping $(x, y) \mapsto (y, x)$ of \mathbb{R}^2 onto \mathbb{R}^2 is not the composition of any two primitive mappings, in any neighborhood of the origin. (This shows that the flips B_i cannot be omitted from the statement of Theorem 10.7.)

Proof of (a).

- (1) Suppose \mathbf{F} is a \mathcal{C}' -mapping of an open set $E \subseteq \mathbb{R}^n$ into \mathbb{R}^n , $\mathbf{0} \in E$, $\mathbf{F}(\mathbf{0}) = \mathbf{0}$, and $\mathbf{F}'(\mathbf{0})$ is invertible.
- (2) Similar to the proof of Theorem 10.7. Put $\mathbf{F}_1 = \mathbf{F}$.
- (3) As $m = 1$, there is an open neighborhood $V_1 \subseteq E$ of $\mathbf{0}$ such that $\mathbf{F}_1(\mathbf{0}) = (\mathbf{F}'(\mathbf{0}))^{-1}\mathbf{F}(\mathbf{0}) = \mathbf{0}$, $\mathbf{F}'_1(\mathbf{0}) = \mathbf{I}$ is invertible, and

$$\mathbf{F}_1(\mathbf{x}) = \sum_{i=1}^n \alpha_i(\mathbf{x})\mathbf{e}_i,$$

where $\alpha_1, \dots, \alpha_n$ are real \mathcal{C}' -functions in V_1 . Hence

$$\mathbf{F}'_1(\mathbf{0})\mathbf{e}_1 = \sum_{i=1}^n (D_1\alpha_i)(\mathbf{0})\mathbf{e}_i.$$

Note that $(D_1\alpha_1)(\mathbf{0}) = 1 \neq 0$, and we might pick $B_1 = \mathbf{I}$. Thus we can define

$$\mathbf{G}_1(\mathbf{x}) = \mathbf{x} + [\alpha_1(\mathbf{x}) - x_1]\mathbf{e}_1 \quad (\mathbf{x} \in V_1).$$

Then $\mathbf{G}_1 \in \mathcal{C}'(V_1)$, \mathbf{G}_1 is primitive, and $\mathbf{G}'_1(\mathbf{0}) = \mathbf{I}$ is invertible.

- (4) Now we make the induction hypothesis for $1 \leq m \leq n-1$.
- (5) Since $\mathbf{G}'_m(\mathbf{0}) = \mathbf{I}$ is invertible, the inverse function theorem shows that there is an open set U_m , with $\mathbf{0} \in U_m \subseteq V_m$, such that \mathbf{G}_m is an injective mapping of U_m onto a neighborhood V_{m+1} of $\mathbf{0}$, in which $\mathbf{G}_m^{-1} \in \mathcal{C}'(V_{m+1})$. Define \mathbf{F}_{m+1} by

$$\mathbf{F}_{m+1}(\mathbf{y}) = \mathbf{F}_m \circ \mathbf{G}_m^{-1}(\mathbf{y}) \quad (\mathbf{y} \in V_{m+1}).$$

Then $\mathbf{F}_{m+1} \in \mathcal{C}'(V_{m+1})$, $\mathbf{F}_m(\mathbf{0}) = \mathbf{0}$, and $\mathbf{F}'_{m+1}(\mathbf{0}) = \mathbf{I}$ is invertible by the chain rule and the inverse function theorem. So

$$\mathbf{F}_{m+1}(\mathbf{x}) = P_m\mathbf{x} + \sum_{i=m+1}^n \alpha_i(\mathbf{x})\mathbf{e}_i,$$

where $\alpha_1, \dots, \alpha_n$ are real \mathcal{C}' -functions in V_{m+1} . Hence

$$\mathbf{F}'_{m+1}(\mathbf{0})\mathbf{e}_{m+1} = \sum_{i=m+1}^n (D_{m+1}\alpha_i)(\mathbf{0})\mathbf{e}_i.$$

Note that $(D_{m+1}\alpha_{m+1})(\mathbf{0}) = 1 \neq 0$, and we might pick $B_{m+1} = \mathbf{I}$. Thus we can define

$$\mathbf{G}_{m+1}(\mathbf{x}) = \mathbf{x} + [\alpha_{m+1}(\mathbf{x}) - x_{m+1}]\mathbf{e}_{m+1} \quad (\mathbf{x} \in V_{m+1}).$$

Then $\mathbf{G}_{m+1} \in \mathcal{C}'(V_{m+1})$, \mathbf{G}_{m+1} is primitive, and $\mathbf{G}'_{m+1}(\mathbf{0}) = \mathbf{I}$ is invertible. Our induction hypothesis holds therefore with $m+1$ in place of m .

(6) Note that

$$\mathbf{F}_m(\mathbf{x}) = \mathbf{F}_{m+1}(\mathbf{G}_m(\mathbf{x})) \quad (\mathbf{x} \in U_m).$$

If we apply this with $m = 1, \dots, n-1$, we successively obtain

$$\mathbf{F}_1 = \mathbf{F}_n \circ \mathbf{G}_{n-1} \circ \dots \circ \mathbf{G}_1$$

in some open neighborhood of $\mathbf{0}$. Note that \mathbf{F}_n is primitive since

$$\mathbf{F}_n(\mathbf{x}) = P_{n-1}\mathbf{x} + \alpha_n(\mathbf{x})\mathbf{e}_n.$$

This completes the proof.

□

Proof of (b).

(1) For $(x, y) \in \mathbb{R}^2$, define

$$\mathbf{F}(x, y) = (y, x).$$

(2) (Reductio ad absurdum) If $\mathbf{F} = \mathbf{G}_2 \circ \mathbf{G}_1$ for some primitive mappings \mathbf{G}_i ($i = 1, 2$) in some neighborhood V_i of the origin, $\mathbf{G}_i(\mathbf{0}) = \mathbf{0}$ and \mathbf{G}'_i is invertible, then we may assume that

$$\mathbf{G}_1(x, y) = (x, g_1(x, y)) \quad \text{and} \quad \mathbf{G}_2(x, y) = (g_2(x, y), y).$$

Here the case $\mathbf{G}_1(x, y) = (g_1(x, y), y)$ and $\mathbf{G}_2(x, y) = (x, g_2(x, y))$ is similar to the above case. Besides, $\mathbf{G}_1(x, y) = (x, g_1(x, y))$ and $\mathbf{G}_2(x, y) = (x, g_2(x, y))$ implies that

$$\mathbf{G}_2 \circ \mathbf{G}_1(x, y) = (x, g_2(x, g_1(x, y))) \neq (y, x) = \mathbf{F}(x, y).$$

Same reason for $\mathbf{G}_1(x, y) = (g_1(x, y), y)$ and $\mathbf{G}_2(x, y) = (g_2(x, y), y)$.

(3) Note that

$$\mathbf{F}'(\mathbf{0}) = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

Since

$$\mathbf{F}'(\mathbf{0}) = \mathbf{G}'_2(\mathbf{G}_1(\mathbf{0}))\mathbf{G}'_1(\mathbf{0}) = \mathbf{G}'_2(\mathbf{0})\mathbf{G}'_1(\mathbf{0}),$$

we have

$$\begin{aligned} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} &= \begin{bmatrix} D_1g_2(0,0) & D_2g_2(0,0) \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ D_1g_1(0,0) & D_2g_1(0,0) \end{bmatrix} \\ &= \begin{bmatrix} * & * \\ D_1g_1(0,0) & D_2g_1(0,0) \end{bmatrix}. \end{aligned}$$

So $D_1g_1(0,0) = 1$ and $D_2g_1(0,0) = 0$, and thus $\mathbf{G}'_1(\mathbf{0}) = \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$ is not invertible, which is absurd.

□

Exercise 10.4. For $(x, y) \in \mathbb{R}^2$, define

$$\mathbf{F}(x, y) = (e^x \cos y - 1, e^x \sin y)$$

Prove that $\mathbf{F} = \mathbf{G}_2 \circ \mathbf{G}_1$, where

$$\begin{aligned} \mathbf{G}_1(x, y) &= (e^x \cos y - 1, y) \\ \mathbf{G}_2(u, v) &= (u, (1 + u) \tan v) \end{aligned}$$

are primitive in some neighborhood of $(0, 0)$. Compute the Jacobians of \mathbf{G}_1 , \mathbf{G}_2 , \mathbf{F} at $(0, 0)$. Define

$$\mathbf{H}_2(x, y) = (x, e^x \sin y)$$

and find

$$\mathbf{H}_1(u, v) = (h(u, v), v)$$

so that $\mathbf{F} = \mathbf{H}_1 \circ \mathbf{H}_2$ is in some neighborhood of $(0, 0)$.

Proof.

(1) By Definition 10.5,

$$\begin{aligned} \mathbf{G}_1(x, y) &= (e^x \cos y - 1)\mathbf{e}_1 + y\mathbf{e}_2, \\ \mathbf{G}_2(u, v) &= u\mathbf{e}_1 + ((1 + u) \tan v)\mathbf{e}_2 \end{aligned}$$

are primitive in some neighborhood of $(0, 0)$.

(2) Show that $\mathbf{F} = \mathbf{G}_2 \circ \mathbf{G}_1$. Given any $(x, y) \in \mathbb{R}^2$, we have

$$\begin{aligned}
(\mathbf{G}_2 \circ \mathbf{G}_1)(x, y) &= \mathbf{G}_2(\mathbf{G}_1(x, y)) \\
&= \mathbf{G}_2(e^x \cos y - 1, y) \\
&= (e^x \cos y - 1, (1 + (e^x \cos y - 1)) \tan y) \\
&= (e^x \cos y - 1, e^x \sin y) \\
&= \mathbf{F}(x, y).
\end{aligned}$$

(3) Since

$$\begin{aligned}
J_{\mathbf{G}_1}(x, y) &= \det \begin{bmatrix} e^x \cos y & -e^x \sin y \\ 0 & 1 \end{bmatrix} = e^x \cos y \\
J_{\mathbf{G}_2}(x, y) &= \det \begin{bmatrix} 1 & 0 \\ \tan y & (1 + x) \sec^2 y \end{bmatrix} = (1 + x) \sec^2 y \\
J_{\mathbf{F}}(x, y) &= \det \begin{bmatrix} e^x \cos y & -e^x \sin y \\ e^x \sin y & e^x \cos y \end{bmatrix} = e^{2x},
\end{aligned}$$

$$J_{\mathbf{G}_1}(0, 0) = \det \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = 1$$

$$J_{\mathbf{G}_2}(0, 0) = \det \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = 1$$

$$J_{\mathbf{F}}(0, 0) = \det \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = 1.$$

(4) Define $h(u, v) = \sqrt{e^{2u} - v^2} - 1$ on

$$B\left((0, 0); \frac{1}{64}\right) \subseteq \mathbb{R}^2.$$

$h(u, v)$ is well-defined since $e^{2u} - v^2 > 0$ for all $(u, v) \in B((0, 0); \frac{1}{64})$.

(5) Given any $(x, y) \in \mathbb{R}^2$, we have

$$\begin{aligned}
(\mathbf{H}_1 \circ \mathbf{H}_2)(x, y) &= \mathbf{H}_1(\mathbf{H}_2(x, y)) \\
&= \mathbf{H}_1(x, e^x \sin y) \\
&= (\sqrt{e^{2x} - (e^x \sin y)^2} - 1, e^x \sin y) \\
&= (e^x \cos y - 1, e^x \sin y) \\
&= \mathbf{F}(x, y).
\end{aligned}$$

□

Exercise 10.5. Formulate and prove an analogue of Theorem 10.8, in which K is a compact subset of an arbitrary metric space. (Replace the functions φ_i that occur in the proof of Theorem 10.8 by functions of the type constructed in Exercise 4.22.)

Proof (Theorem 10.8).

- (1) (Partitions of unity.) Suppose K is a compact subset of a metric space X , and $\{V_\alpha\}$ is an open cover of K . Then there exist functions $\psi_1, \dots, \psi_s \in \mathcal{C}(X)$ such that

- (a) $0 \leq \psi_i \leq 1$ for $1 \leq i \leq s$.
- (b) each ψ_i has its support in some V_α , and
- (c) $\psi_1(x) + \dots + \psi_s(x) = 1$ for every $x \in K$.

- (2) It is trivial that some $V_\alpha = X$ by taking $s = 1$ and $\psi_1(x) = 1 \in \mathcal{C}(X)$. Now we assume that all $V_\alpha \subsetneq X$.

- (3) Associate with each $x \in K$ an index $\alpha(x)$ so that $x \in V_{\alpha(x)}$. Then there are open balls $B(x)$ and $W(x)$, centered at x , with

$$x \in B(x) \subseteq \overline{B(x)} \subseteq W(x) \subseteq \overline{W(x)} \subseteq V_{\alpha(x)}$$

(Since $V_{\alpha(x)}$ is open, there exists $r > 0$ such that $B(x; r) \subseteq V_{\alpha(x)}$. Take $B(x) = B(x; \frac{r}{89})$ and $W(x) = B(x; \frac{r}{64})$.)

- (4) Since K is compact, there are finitely many points $x_1, \dots, x_s \in K$ such that

$$K \subseteq B(x_1) \cup \dots \cup B(x_s).$$

Note that

- (a) $\overline{B(x_i)}$ is a nonempty closed set since $x_i \in B(x_i) \subseteq \overline{B(x_i)}$.
- (b) $X - W(x_i) \supseteq X - V_{\alpha(x_i)}$ is a nonempty closed set by the assumption in (2).
- (c) $\overline{B(x_i)} \cap (X - W(x_i)) \subseteq W(x_i) \cap (X - W(x_i)) = \emptyset$.

By Exercise 4.22, there is a function

$$\varphi_i(x) = \frac{\rho_{\overline{B(x_i)}}(x)}{\rho_{\overline{B(x_i)}}(x) + \rho_{X - W(x_i)}(x)} \in \mathcal{C}(X)$$

such that $\varphi_i(x) = 1$ on $\overline{B(x_i)}$, $\varphi_i(x) = 0$ outside $W(x_i)$, and $0 \leq \varphi_i(x) \leq 1$ on X for $1 \leq i \leq s$.

- (5) Define $\psi_1 = \varphi_1 \in \mathcal{C}(X)$ and

$$\psi_{i+1} = (1 - \varphi_1) \cdots (1 - \varphi_i) \varphi_{i+1} \in \mathcal{C}(X)$$

for $1 \leq i \leq s-1$. Properties (a) and (b) in (1) are clear. Also,

$$\psi_1(x) + \cdots + \psi_s(x) = 1 - (1 - \varphi_1(x)) \cdots (1 - \varphi_s(x))$$

by the construction of ψ_i . If $x \in K$, then $x \in B(x_i)$ for some i , hence $\varphi_i(x) = 1$, and the product $(1 - \varphi_1(x)) \cdots (1 - \varphi_s(x)) = 0$. This proves property (c) in (1).

□

Exercise 10.6. *Strengthen the conclusion of Theorem 10.8 by showing that the functions ψ_i can be made differentiable, and even infinitely differentiable. (Use Exercise 8.1 in the construction of the auxiliary functions ψ_i .)*

Proof (Theorem 10.8).

- (1) It is trivial that some $V_\alpha = \mathbb{R}^n$ by taking $s = 1$ and $\psi_1(\mathbf{x}) = 1 \in \mathcal{C}^\infty(\mathbb{R}^n)$. Now we assume that all $V_\alpha \subsetneq \mathbb{R}^n$.
- (2) Associate with each $\mathbf{x} \in K$ an index $\alpha(x)$ so that $\mathbf{x} \in V_{\alpha(x)}$. Then there are open n -cells $B(\mathbf{x})$ and $W(\mathbf{x})$ (Definition 10.1), centered at \mathbf{x} , with

$$\mathbf{x} \in B(\mathbf{x}) \subseteq \overline{B(\mathbf{x})} \subseteq W(\mathbf{x}) \subseteq \overline{W(\mathbf{x})} \subseteq V_{\alpha(\mathbf{x})}$$

(Since $V_{\alpha(\mathbf{x})}$ is open, there exists $r > 0$ such that $B(\mathbf{x}; r) \subseteq V_{\alpha(\mathbf{x})}$. Take

$$B(\mathbf{x}) = I\left(\mathbf{x}; \frac{r}{89\sqrt{n}}\right), \quad W(\mathbf{x}) = I\left(\mathbf{x}; \frac{r}{64\sqrt{n}}\right)$$

where $I(\mathbf{p}; r)$ is the open n -cell centered at $\mathbf{p} = (p_1, \dots, p_n)$ defined by

$$I(\mathbf{p}; r) = (p_1 - r, p_1 + r) \times \cdots \times (p_n - r, p_n + r) \subseteq \mathbb{R}^n.$$

- (3) Define

$$f(y) = \begin{cases} e^{-\frac{1}{y^2}} & (y > 0), \\ 0 & (y \leq 0). \end{cases}$$

$f(y) \in \mathcal{C}^\infty(\mathbb{R}^1)$ by applying the similar argument in Exercise 8.1.

- (4) Given any $\mathbf{x} = (x_1, \dots, x_n) \in K$ and construct $B(\mathbf{x})$ and $W(\mathbf{x})$ as in (2). Define

$$g_{x_j}(y_j) = \frac{f(y_j)}{f(y_j) + f\left(\frac{r}{64\sqrt{n}} - \frac{r}{89\sqrt{n}} - y_j\right)}$$

for $1 \leq j \leq n$. g_{x_j} is well-defined and $g_{x_j} \in \mathcal{C}^\infty(\mathbb{R}^1)$. So

$$g_{x_j}(y_j) = \begin{cases} 0 & \text{if } y_j \leq 0, \\ \text{strictly increasing} & \text{if } 0 \leq y_j \leq \frac{r}{64\sqrt{n}} - \frac{r}{89\sqrt{n}}, \\ 1 & \text{if } y_j \geq \frac{r}{64\sqrt{n}} - \frac{r}{89\sqrt{n}}. \end{cases}$$

Next, define

$$h_{x_j}(y_j) = g_{x_j} \left(y_j - x_j + \frac{r}{64\sqrt{n}} \right) g_{x_j} \left(x_j + \frac{r}{64\sqrt{n}} - y_j \right)$$

for $1 \leq j \leq n$. $h_{x_j} \in \mathcal{C}^\infty(\mathbb{R}^1)$. So

$$h_{x_j}(y_j) = \begin{cases} 0 & \text{if } y_j \leq x_j - \frac{r}{64\sqrt{n}}, \\ \text{strictly increasing} & \text{if } x_j - \frac{r}{64\sqrt{n}} \leq y_j \leq x_j - \frac{r}{89\sqrt{n}}, \\ 1 & \text{if } x_j - \frac{r}{89\sqrt{n}} \leq y_j \leq x_j + \frac{r}{89\sqrt{n}}, \\ \text{strictly decreasing} & \text{if } x_j + \frac{r}{89\sqrt{n}} \leq y_j \leq x_j + \frac{r}{64\sqrt{n}}, \\ 0 & \text{if } y_j \geq x_j + \frac{r}{64\sqrt{n}}. \end{cases}$$

Finally we define $\mathbf{h}_{\mathbf{x}} : \mathbb{R}^n \rightarrow \mathbb{R}^1$ by

$$\mathbf{h}_{\mathbf{x}}(\mathbf{y}) = \prod_{j=1}^n h_{x_j}(y_j)$$

where $\mathbf{y} = (y_1, \dots, y_n) \in \mathbb{R}^n$. Hence, $\mathbf{h}_{\mathbf{x}} \in \mathcal{C}^\infty(\mathbb{R}^n)$ (Theorem 9.21). Also, $\mathbf{h}_{\mathbf{x}}(\mathbf{y}) = 1$ on $B(\mathbf{x})$, $\mathbf{h}_{\mathbf{x}}(\mathbf{y}) = 0$ outside $W(\mathbf{x})$, and $0 \leq \mathbf{h}_{\mathbf{x}}(\mathbf{y}) \leq 1$.

- (5) Since K is compact, there are finitely many points $\mathbf{x}_1, \dots, \mathbf{x}_s \in K$ such that

$$K \subseteq B(\mathbf{x}_1) \cup \dots \cup B(\mathbf{x}_s).$$

Take

$$\varphi_i(\mathbf{x}) = \mathbf{h}_{\mathbf{x}_i}(\mathbf{x}) \in \mathcal{C}^\infty(\mathbb{R}^n)$$

for $1 \leq i \leq s$.

- (6) The rest are the same as the proof of Theorem 10.8 or Exercise 10.5.

□

Exercise 10.7.

- (a) Show that the simplex Q^k is the smallest convex subset of \mathbb{R}^k such that contains $\mathbf{0}, \mathbf{e}_1, \dots, \mathbf{e}_k$.
- (b) Show that affine mappings take convex sets to convex sets.

Proof of (a).

- (1) Show that Q^k contains $\mathbf{0}, \mathbf{e}_1, \dots, \mathbf{e}_k$. Recall

$$Q^k = \{(x_1, \dots, x_k) \in \mathbb{R}^k : x_1 + \dots + x_k \leq 1 \text{ and } x_1, \dots, x_k \geq 0\}$$

(Example 10.14). Hence $\mathbf{0} = (0, \dots, 0) \in Q^k$ and

$$\mathbf{e}_i = (0, \dots, \underbrace{1}_{i\text{th coordinate}}, \dots, 0) \in Q^k.$$

- (2) Show that Q^k is a convex subset of \mathbb{R}^k . Given any $\mathbf{x} = (x_1, \dots, x_k) \in Q^k$, $\mathbf{y} = (y_1, \dots, y_k) \in Q^k$ and $0 < \lambda < 1$. Hence

$$\lambda \mathbf{x} + (1 - \lambda) \mathbf{y} = (\lambda x_1 + (1 - \lambda) y_1, \dots, \lambda x_k + (1 - \lambda) y_k) \in Q^k$$

since each $\lambda x_i + (1 - \lambda) y_i \geq 0$ and

$$\sum_{i=1}^k (\lambda x_i + (1 - \lambda) y_i) = \lambda \sum_{i=1}^k x_i + (1 - \lambda) \sum_{i=1}^k y_i \leq \lambda + (1 - \lambda) = 1.$$

- (3) Given any convex set $E \subseteq \mathbb{R}^k$ containing $\mathbf{0}, \mathbf{e}_1, \dots, \mathbf{e}_k$. Show that $E \supseteq Q^k$.

- (a) Induction on k . Base case: $k = 1$. Given any $\mathbf{x} = (x_1) \in Q^1$. We have $0 \leq x_1 \leq 1$ by the definition of Q^1 . So that $\mathbf{x} = x_1 \mathbf{e}_1 + (1 - x_1) \mathbf{0} \in E$ since $\mathbf{0}, \mathbf{e}_1 \in E$ and E is convex.
- (b) Inductive step: suppose the statement holds for $k = n$. Given any $\mathbf{x} = (x_1, \dots, x_n, x_{n+1}) \in Q^{n+1}$. If $x_{n+1} = 1$, then $x_1 = \dots = x_n = 0$ by the definition of Q^{n+1} . So $\mathbf{x} = \mathbf{e}_{n+1} \in E$ by the assumption of E . If $0 \leq x_{n+1} < 1$, then $x_1 + \dots + x_n \leq 1 - x_{n+1}$ or

$$\frac{x_1}{1 - x_{n+1}} + \dots + \frac{x_n}{1 - x_{n+1}} \leq 1.$$

So the point

$$\left(\frac{x_1}{1 - x_{n+1}}, \dots, \frac{x_n}{1 - x_{n+1}} \right) \in Q^n,$$

or

$$\left(\frac{x_1}{1 - x_{n+1}}, \dots, \frac{x_n}{1 - x_{n+1}}, 0 \right), \text{ say } \hat{\mathbf{x}}, \in E$$

by the induction hypothesis. Note that $\mathbf{e}_{n+1} \in E$. Hence

$$\mathbf{x} = x_{n+1} \mathbf{e}_{n+1} + (1 - x_{n+1}) \hat{\mathbf{x}} \in E$$

by the convexity of E .

- (c) Conclusion: Since both the base case and the inductive step have been proved as true, by mathematical induction the statement holds.

□

Proof of (b).

- (1) Let \mathbf{f} be an affine mapping that carries a vector space X into a vector space Y such that

$$\mathbf{f}(\mathbf{x}) = \mathbf{f}(\mathbf{0}) + A\mathbf{x}$$

for some $A \in L(X, Y)$.

- (2) Given any convex subset C of X . To show that $\mathbf{f}(C)$ is convex, it suffices to show that

$$\lambda \mathbf{y}_1 + (1 - \lambda) \mathbf{y}_2 \in \mathbf{f}(C)$$

for any $\mathbf{y}_1, \mathbf{y}_2 \in \mathbf{f}(C)$ and $0 < \lambda < 1$. Write $\mathbf{y}_1 = \mathbf{f}(\mathbf{x}_1)$, $\mathbf{y}_2 = \mathbf{f}(\mathbf{x}_2)$ for some $\mathbf{x}_1, \mathbf{x}_2 \in C$. Note that $\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2 \in C$ by the convexity of C . Hence

$$\begin{aligned} & \mathbf{f}(\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2) \\ &= \mathbf{f}(\mathbf{0}) + A(\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2) \\ &= \mathbf{f}(\mathbf{0}) + \lambda A \mathbf{x}_1 + (1 - \lambda) A \mathbf{x}_2 & (A \in L(X, Y)) \\ &= \lambda(\mathbf{f}(\mathbf{0}) + A \mathbf{x}_1) + (1 - \lambda)(\mathbf{f}(\mathbf{0}) + A \mathbf{x}_2) \\ &= \lambda \mathbf{f}(\mathbf{x}_1) + (1 - \lambda) \mathbf{f}(\mathbf{x}_2) \\ &= \lambda \mathbf{y}_1 + (1 - \lambda) \mathbf{y}_2 \in \mathbf{f}(C). \end{aligned}$$

□

Exercise 10.8. Let H be the parallelogram in \mathbb{R}^2 whose vertices are $(1, 1)$, $(3, 2)$, $(4, 5)$, $(2, 4)$. Find the affine map T which sends $(0, 0)$ to $(1, 1)$, $(1, 0)$ to $(3, 2)$, $(1, 1)$ to $(4, 5)$, $(0, 1)$ to $(2, 4)$. Show that $J_T = 5$. Use T to convert the integral

$$\alpha = \int_H e^{x-y} dx dy$$

to an integral over I^2 and thus compute α .

Proof.

- (1) By Affine simplexes 10.26,

$$T(\mathbf{x}) = T(\mathbf{0}) + A\mathbf{x},$$

where $A \in L(\mathbb{R}^2, \mathbb{R}^2)$, say $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Note that $T : \begin{bmatrix} 0 \\ 0 \end{bmatrix} \mapsto \begin{bmatrix} 1 \\ 1 \end{bmatrix}$. Thus

$$T : \begin{bmatrix} x \\ y \end{bmatrix} \mapsto \begin{bmatrix} 1 \\ 1 \end{bmatrix} + \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 + ax + by \\ 1 + cx + dy \end{bmatrix}.$$

- (2) By $T : (1, 0) \mapsto (3, 2)$ and $T : (0, 1) \mapsto (2, 4)$, we can solve A as

$$A = \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}.$$

It is easy to verify such

$$T : \underbrace{\begin{bmatrix} x \\ y \end{bmatrix}}_{\mathbf{x}} \mapsto \underbrace{\begin{bmatrix} 1 \\ 1 \end{bmatrix}}_{T(\mathbf{0})} + \underbrace{\begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}}_A \underbrace{\begin{bmatrix} x \\ y \end{bmatrix}}_{\mathbf{x}} = \begin{bmatrix} 1 + 2x + y \\ 1 + x + 3y \end{bmatrix}$$

satisfying our requirement.

(3)

$$J_T = \det \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix} = 5.$$

(4)

$$\begin{aligned} \int_H e^{x-y} dx dy &= \int_{[0,1]^2} e^{(1+2u+v)-(1+u+3v)} |J_T| du dv \\ &= 5 \int_{[0,1]^2} e^{u-2v} du dv \\ &= 5 \left\{ \int_0^1 e^u du \right\} \left\{ \int_0^1 e^{-2v} dv \right\} \quad (\text{Theorem 10.2}) \\ &= \frac{5}{2} (e-1)(1-e^{-2}). \end{aligned}$$

□

Exercise 10.9. Define $(x, y) = T(r, \theta)$ one the rectangle

$$0 \leq r \leq a, \quad 0 \leq \theta \leq 2\pi$$

by the equations

$$x = r \cos \theta, \quad y = r \sin \theta.$$

Show that T maps this rectangle onto the closed disc D with center at $(0, 0)$ and radius a , that T is one-to-one in the interior of the rectangle, and that $J_T(r, \theta) = r$. If $f \in \mathcal{C}(D)$, prove the formula for integration in polar coordinates:

$$\int_D f(x, y) dx dy = \int_0^a \int_0^{2\pi} f(T(r, \theta)) r dr d\theta.$$

(Hint: Let D_0 be the interior of D , minus the interval from $(0, 0)$ to $(0, a)$. As it stands, Theorem 10.9 applies to continuous functions f whose support lies in D_0 . To remove this restriction, proceed as in Example 10.4.)

Proof.

(1)

(2)

□

Exercise 10.10. Let $a \rightarrow \infty$ in Exercise 10.9 and prove that

$$\int_{\mathbb{R}^2} f(x, y) dx dy = \int_0^\infty \int_0^{2\pi} f(T(r, \theta)) r dr d\theta,$$

for continuous functions f that decrease sufficiently rapidly as $|x| + |y| \rightarrow \infty$.
 (Find a more precise formulation.) Apply this to

$$f(x, y) = \exp(-x^2 - y^2)$$

to derive formula

$$\int_{-\infty}^{\infty} e^{-s^2} ds = \sqrt{\pi}.$$

Proof.

(1)

(2)

□

Exercise 10.11. ...

Proof.

(1)

(2)

□

Exercise 10.12. ...

Proof.

(1)

(2)

□

Exercise 10.13. ...

Proof.

(1)

(2)

□

Exercise 10.14 (Levi-Civita symbol). Prove $\varepsilon(j_1, \dots, j_k) = s(j_1, \dots, j_k)$, where

$$s(j_1, \dots, j_k) = \prod_{p < q} \operatorname{sgn}(j_q - j_p).$$

It is usually to define the Levi-Civita symbol by

$$\varepsilon(j_1, \dots, j_k) = \begin{cases} 1 & \text{if } (j_1, \dots, j_k) \text{ is an even permutation of } J, \\ -1 & \text{if } (j_1, \dots, j_k) \text{ is an odd permutation of } J, \\ 0 & \text{otherwise} \end{cases}$$

(Basic k -forms 10.14). Thus, it is the sign of the permutation in the case of a permutation, and zero otherwise. So $\varepsilon(j_1, \dots, j_k)$ is equivalent to an explicit expression $s(j_1, \dots, j_k) = \prod_{p < q} \operatorname{sgn}(j_q - j_p)$.

Proof.

(1) Induction on k . Base case: Show that $\varepsilon(j_1, j_2) = s(j_1, j_2)$. Since

$$\varepsilon(j_1, j_2) = \begin{cases} 1 & \text{if } j_1 < j_2 \\ -1 & \text{if } j_1 > j_2, \end{cases}$$

$$\varepsilon(j_1, j_2) = \operatorname{sgn}(j_2 - j_1) = s(j_1, j_2).$$

(2) Inductive step: Show that for any $s \geq 2$, if $\varepsilon(j_1, \dots, j_s) = s(j_1, \dots, j_s)$ holds, then $\varepsilon(j_1, \dots, j_{s+1}) = s(j_1, \dots, j_{s+1})$ also holds.

$$\begin{aligned} \varepsilon(j_1, \dots, j_{s+1}) &= \varepsilon(j_1, \dots, j_s) \prod_{\substack{1 \leq p \leq s \\ q=s+1}} \operatorname{sgn}(j_q - j_p) \\ &= s(j_1, \dots, j_s) \prod_{\substack{1 \leq p \leq s \\ q=s+1}} \operatorname{sgn}(j_q - j_p) \\ &= \prod_{1 \leq p < q \leq s} \operatorname{sgn}(j_q - j_p) \prod_{\substack{1 \leq p \leq s \\ q=s+1}} \operatorname{sgn}(j_q - j_p) \\ &= \prod_{1 \leq p < q \leq s+1} \operatorname{sgn}(j_q - j_p) \\ &= s(j_1, \dots, j_{s+1}). \end{aligned}$$

(3) Conclusion: Since both the base case and the inductive step have been proved as true, by mathematical induction the statement holds for every integer $k \geq 2$.

□

Exercise 10.15. If ω and λ are k - and m -forms, respectively, prove that

$$\omega \wedge \lambda = (-1)^{km} \lambda \wedge \omega.$$

Proof.

(1) Write

$$\omega = \sum_I b_I(\mathbf{x}) dx_I, \quad \lambda = \sum_J c_J(\mathbf{x}) dx_J$$

in the standard presentations, where I and J range over all increasing k -indices and over all increasing m -indices taken from the set $\{1, \dots, n\}$.

(2) Show that $dx_I \wedge dx_J = (-1)^{km} dx_J \wedge dx_I$.

$$\begin{aligned} dx_I \wedge dx_J &= dx_{i_1} \wedge \dots \wedge dx_{i_k} \wedge dx_J \\ &= (-1)^m dx_{i_1} \wedge \dots \wedge dx_{i_{k-1}} \wedge dx_J \wedge dx_{i_k} \\ &= (-1)^{2m} dx_{i_1} \wedge \dots \wedge dx_{i_{k-2}} \wedge dx_J \wedge dx_{i_{k-1}} \wedge dx_{i_k} \\ &\dots \\ &= (-1)^{km} dx_J \wedge dx_{i_1} \wedge \dots \wedge dx_{i_k} \\ &= (-1)^{km} dx_J \wedge dx_I. \end{aligned}$$

(3)

$$\begin{aligned} \omega \wedge \lambda &= \sum_{I,J} b_I(\mathbf{x}) c_J(\mathbf{x}) dx_I \wedge dx_J \\ &= (-1)^{km} \sum_{J,I} c_J(\mathbf{x}) b_I(\mathbf{x}) dx_J \wedge dx_I \\ &= (-1)^{km} \lambda \wedge \omega. \end{aligned}$$

□

Exercise 10.16. If $k \geq 2$ and $\sigma = [\mathbf{p}_0, \mathbf{p}_1, \dots, \mathbf{p}_k]$ is an oriented affine k -simplex, prove that $\partial^2 \sigma = 0$, directly from the definition of the boundary operator ∂ . Deduce from this that $\partial^2 \Psi = 0$ for every chain Ψ . (Hint: For orientation, do it first for $k = 2$, $k = 3$. In general, if $i < j$, let σ_{ij} be the $(k-2)$ -simplex obtained by deleting \mathbf{p}_i and \mathbf{p}_j from σ . Show that each σ_{ij} occurs twice in $\partial^2 \sigma$, with opposite sign.)

Proof (Brute-force).

- (1) Write the boundary of the oriented affine k -simplex $\sigma = [\mathbf{p}_0, \mathbf{p}_1, \dots, \mathbf{p}_k]$ as

$$\partial\sigma = \sum_{i=0}^k (-1)^i [\mathbf{p}_0, \dots, \widehat{\mathbf{p}}_i, \dots, \mathbf{p}_k]$$

where where the oriented $(k-1)$ -simplex $[\mathbf{p}_0, \dots, \widehat{\mathbf{p}}_i, \dots, \mathbf{p}_k]$ is obtained by deleting σ 's i -th vertex (Boundaries 10.29).

- (2)

$$\begin{aligned} \partial^2\sigma &= \partial \left(\sum_i (-1)^i [\mathbf{p}_0, \dots, \widehat{\mathbf{p}}_i, \dots, \mathbf{p}_k] \right) \\ &= \sum_i (-1)^i \partial [\mathbf{p}_0, \dots, \widehat{\mathbf{p}}_i, \dots, \mathbf{p}_k] \\ &= \sum_{j < i} (-1)^i (-1)^j [\mathbf{p}_0, \dots, \widehat{\mathbf{p}}_j, \dots, \widehat{\mathbf{p}}_i, \dots, \mathbf{p}_k] \\ &\quad + \sum_{j > i} (-1)^i (-1)^{j-1} [\mathbf{p}_0, \dots, \widehat{\mathbf{p}}_i, \dots, \widehat{\mathbf{p}}_j, \dots, \mathbf{p}_k] \\ &= \sum_{j < i} (-1)^{i+j} [\mathbf{p}_0, \dots, \widehat{\mathbf{p}}_j, \dots, \widehat{\mathbf{p}}_i, \dots, \mathbf{p}_k] \\ &\quad - \sum_{j > i} (-1)^{i+j} [\mathbf{p}_0, \dots, \widehat{\mathbf{p}}_i, \dots, \widehat{\mathbf{p}}_j, \dots, \mathbf{p}_k]. \end{aligned}$$

The latter two summations cancel since after switching i and j in the second sum. Therefore $\partial^2\sigma = 0$.

- (3) The boundary of a chain is the linear combination of boundaries of the simplices in the chain. Write $\Psi = \sum_{i=1}^r \sigma_i$. where σ_i is an oriented affine simplex. Then

$$\partial^2\Psi = \partial \left(\partial \sum \sigma_i \right) = \partial \left(\sum \partial\sigma_i \right) = \sum \partial^2\sigma_i = \sum 0 = 0$$

for any affine chain Ψ .

□

Exercise 10.17. Put $J^2 = \tau_1 + \tau_2$, where

$$\tau_1 = [\mathbf{0}, \mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_2], \quad \tau_2 = -[\mathbf{0}, \mathbf{e}_2, \mathbf{e}_2 + \mathbf{e}_1].$$

Explain why it is reasonable to call J^2 the positively oriented unit square in \mathbb{R}^2 . Show that ∂J^2 is the sum of 4 oriented affine 1-simplexes. Find these. What is $\partial(\tau_1 - \tau_2)$?

Proof.

- (1) Note that the unit square $I^2 \in \mathbb{R}^2$ is the union of $\tau_1(Q^2)$ and $\tau_2(Q^2)$, where

$$\begin{aligned}\tau_1(\mathbf{u}) &= ([\mathbf{0}, \mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_2])(\mathbf{u}) \\ &= \mathbf{0} + \alpha_1 \mathbf{e}_1 + \alpha_2 (\mathbf{e}_1 + \mathbf{e}_2) \\ &= \mathbf{0} + (\alpha_1 + \alpha_2) \mathbf{e}_1 + \alpha_2 \mathbf{e}_2 \\ &= \mathbf{0} + \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \mathbf{u}\end{aligned}$$

and

$$\begin{aligned}\tau_2(\mathbf{u}) &= (-[\mathbf{0}, \mathbf{e}_2, \mathbf{e}_2 + \mathbf{e}_1])(\mathbf{u}) \\ &= ([\mathbf{0}, \mathbf{e}_2 + \mathbf{e}_1, \mathbf{e}_2])(\mathbf{u}) \\ &= \mathbf{0} + \alpha_1 (\mathbf{e}_1 + \mathbf{e}_2) + \alpha_2 \mathbf{e}_2 \\ &= \mathbf{0} + \alpha_1 \mathbf{e}_1 + (\alpha_1 + \alpha_2) \mathbf{e}_2 \\ &= \mathbf{0} + \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \mathbf{u}\end{aligned}$$

where $\mathbf{u} = \alpha_1 \mathbf{e}_1 + \alpha_2 \mathbf{e}_2 \in \mathbb{R}^2$ (as in Equation (78)). Both τ_1 and τ_2 have Jacobian $1 > 0$, or positively oriented (Affine simplexes 10.26). So it is reasonable to call J^2 the positively oriented unit square in \mathbb{R}^2 .

- (2)

$$\begin{aligned}\partial\tau_1 &= [\mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_2] - [\mathbf{0}, \mathbf{e}_1 + \mathbf{e}_2] + [\mathbf{0}, \mathbf{e}_1], \\ \partial\tau_2 &= [\mathbf{e}_2 + \mathbf{e}_1, \mathbf{e}_2] - [\mathbf{0}, \mathbf{e}_2] + [\mathbf{0}, \mathbf{e}_2 + \mathbf{e}_1] \\ &= [\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_2] + [\mathbf{e}_2, \mathbf{0}] + [\mathbf{0}, \mathbf{e}_1 + \mathbf{e}_2].\end{aligned}$$

- (3) By (2),

$$\partial J^2 = \partial\tau_1 + \partial\tau_2 = [\mathbf{0}, \mathbf{e}_1] + [\mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_2] + [\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_2] + [\mathbf{e}_2, \mathbf{0}],$$

which is the positively oriented boundary of I^2 .

- (4) By (2),

$$\begin{aligned}\partial(\tau_1 - \tau_2) &= \partial\tau_1 - \partial\tau_2 \\ &= [\mathbf{0}, \mathbf{e}_1] + [\mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_2] + [\mathbf{e}_1 + \mathbf{e}_2, \mathbf{0}] \\ &\quad + [\mathbf{0}, \mathbf{e}_2] + [\mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_2] + [\mathbf{e}_1 + \mathbf{e}_2, \mathbf{0}].\end{aligned}$$

□

Exercise 10.18. Consider the oriented affine 3-simplex

$$\sigma_1 = [\mathbf{0}, \mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3]$$

in \mathbb{R}^3 . Show that σ_1 (regarded as a linear transformation) has determinant 1. Thus σ_1 is positively oriented.

Let $\sigma_2, \dots, \sigma_6$ be five other oriented 3-simplexes, obtained as follows: There are five permutations (i_1, i_2, i_3) of $(1, 2, 3)$, distinct from $(1, 2, 3)$. Associate with each (i_1, i_2, i_3) the simplex

$$s(i_1, i_2, i_3)[\mathbf{0}, \mathbf{e}_{i_1}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2} + \mathbf{e}_{i_3}]$$

where s is the sign that occurs in the definition of the determinant. (This is how τ_2 was obtained from τ_1 in Exercise 10.17.) Show that $\sigma_2, \dots, \sigma_6$ are positively oriented.

Put $J^3 = \sigma_1 + \dots + \sigma_6$. Then J^3 may be called the positively oriented unit cube in \mathbb{R}^3 . Show that ∂J^3 is the sum of 12 oriented affine 2-simplexes. (These 12 triangles cover the surface of the unit cube I^3 .)

Show that $\mathbf{x} = (x_1, x_2, x_3)$ is in the range of σ_1 if and only if $0 \leq x_3 \leq x_2 \leq x_1 \leq 1$.

Show that the range of $\sigma_1, \dots, \sigma_6$ have disjoint interiors, and that their union covers I^3 . (Compared with Exercise 10.13; note that $3! = 6$.)

Proof.

- (1) Show that σ_1 (regarded as a linear transformation) has determinant 1. Given any $\mathbf{u} = \alpha_1 \mathbf{e}_1 + \alpha_2 \mathbf{e}_2 + \alpha_3 \mathbf{e}_3 \in \mathbb{R}^3$, we have

$$\begin{aligned} \sigma_1(\mathbf{u}) &= ([\mathbf{0}, \mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3])(\mathbf{u}) \\ &= \mathbf{0} + \alpha_1 \mathbf{e}_1 + \alpha_2 (\mathbf{e}_1 + \mathbf{e}_2) + \alpha_3 (\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3) \\ &= \mathbf{0} + (\alpha_1 + \alpha_2 + \alpha_3) \mathbf{e}_1 + (\alpha_2 + \alpha_3) \mathbf{e}_2 + \alpha_3 \mathbf{e}_3 \\ &= \mathbf{0} + \underbrace{\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}}_{\text{say } A} \mathbf{u}. \end{aligned}$$

So

$$\det(A) = \det \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} = 1.$$

- (2) Show that $\sigma_2, \dots, \sigma_6$ are positively oriented. Define the permutation matrix $P_{(i_1, i_2, i_3)}$ corresponding to a permutation (i_1, i_2, i_3) of $(1, 2, 3)$ by

$$P_{(i_1, i_2, i_3)} = [\mathbf{e}_{i_1} \quad \mathbf{e}_{i_2} \quad \mathbf{e}_{i_3}].$$

For example,

$$P_{(2,3,1)} = [\mathbf{e}_2 \quad \mathbf{e}_3 \quad \mathbf{e}_1] = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}.$$

Note that the sign $s(i_1, i_2, i_3)$ of the permutation (i_1, i_2, i_3) is exactly the same as the determinant of the permutation matrix $P_{(i_1, i_2, i_3)}$. Define a permutation $(j_1, j_2, 3)$ of $(1, 2, 3)$ (for swapping the first and the second coordinates of \mathbf{u}) by

$$(j_1, j_2, 3) = \begin{cases} (1, 2, 3) & \text{if } s(i_1, i_2, i_3) = 1, \\ (2, 1, 3) & \text{if } s(i_1, i_2, i_3) = -1. \end{cases}$$

Write

$$\sigma_{(i_1, i_2, i_3)} = s(i_1, i_2, i_3)[\mathbf{0}, \mathbf{e}_{i_1}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2} + \mathbf{e}_{i_3}].$$

(So that $\sigma_1 = \sigma_{(1,2,3)}$.) Hence,

$$\begin{aligned} & \sigma_{(i_1, i_2, i_3)}(\mathbf{u}) \\ &= \mathbf{0} + \alpha_{j_1} \mathbf{e}_{i_1} + \alpha_{j_2}(\mathbf{e}_{i_1} + \mathbf{e}_{i_2}) + \alpha_3(\mathbf{e}_{i_1} + \mathbf{e}_{i_2} + \mathbf{e}_{i_3}) \\ &= \mathbf{0} + (\alpha_{j_1} + \alpha_{j_2} + \alpha_3)\mathbf{e}_{i_1} + (\alpha_{j_2} + \alpha_3)\mathbf{e}_{i_2} + \alpha_3\mathbf{e}_{i_3} \\ &= \mathbf{0} + P_{(i_1, i_2, i_3)}AP_{(j_1, j_2, 3)}\mathbf{u} \end{aligned}$$

where $\mathbf{u} = \alpha_1\mathbf{e}_1 + \alpha_2\mathbf{e}_2 + \alpha_3\mathbf{e}_3 \in \mathbb{R}^3$. For example,

$$P_{(2,3,1)}AP_{(1,2,3)} = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

So

$$\begin{aligned} \det(P_{(i_1, i_2, i_3)}AP_{(j_1, j_2, 3)}) &= \det(P_{(i_1, i_2, i_3)}) \det(A) \det(P_{(j_1, j_2, 3)}) \\ &= s(i_1, i_2, i_3) \cdot 1 \cdot s(i_1, i_2, i_3) \\ &= 1. \end{aligned}$$

(3) Show that ∂J^3 is the sum of 12 oriented affine 2-simplexes. Note that

$$\begin{aligned} \sum_{(i_1, i_2, i_3)} \sigma_{(i_1, i_2, i_3)} &= \sum_{\substack{(i_1, i_2, i_3) \\ i_1 > i_2}} \sigma_{(i_1, i_2, i_3)} + \sum_{\substack{(i_1, i_2, i_3) \\ i_1 < i_2}} \sigma_{(i_1, i_2, i_3)} \\ &= \sum_{\substack{(i_1, i_2, i_3) \\ i_1 > i_2}} s(i_1, i_2, i_3)[\mathbf{0}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3] \\ &\quad + \sum_{\substack{(i_1, i_2, i_3) \\ i_2 > i_1}} -s(i_2, i_1, i_3)[\mathbf{0}, \mathbf{e}_{i_2} + \mathbf{e}_{i_1}, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3] \\ &= \mathbf{0} \end{aligned}$$

and

$$\begin{aligned}
\sum_{(i_1, i_2, i_3)} \sigma_{(i_1, i_2, i_3)} &= \sum_{\substack{(i_1, i_2, i_3) \\ i_2 > i_3}} \sigma_{(i_1, i_2, i_3)} + \sum_{\substack{(i_1, i_2, i_3) \\ i_2 < i_3}} \sigma_{(i_1, i_2, i_3)} \\
&= \sum_{\substack{(i_1, i_2, i_3) \\ i_2 > i_3}} s(i_1, i_2, i_3) [\mathbf{0}, \mathbf{e}_{i_1}, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3] \\
&\quad + \sum_{\substack{(i_1, i_2, i_3) \\ i_3 > i_2}} -s(i_1, i_3, i_2) [\mathbf{0}, \mathbf{e}_{i_1}, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3] \\
&= \mathbf{0}.
\end{aligned}$$

So

$$\begin{aligned}
\partial J^3 &= \sum_{(i_1, i_2, i_3)} \partial \sigma_{(i_1, i_2, i_3)} \\
&= \sum_{(i_1, i_2, i_3)} s(i_1, i_2, i_3) [\mathbf{e}_{i_1}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2} + \mathbf{e}_{i_3}] \\
&\quad - s(i_1, i_2, i_3) [\mathbf{0}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2} + \mathbf{e}_{i_3}] \\
&\quad + s(i_1, i_2, i_3) [\mathbf{0}, \mathbf{e}_{i_1}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2} + \mathbf{e}_{i_3}] \\
&\quad - s(i_1, i_2, i_3) [\mathbf{0}, \mathbf{e}_{i_1}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}] \\
&= \sum_{(i_1, i_2, i_3)} s(i_1, i_2, i_3) [\mathbf{e}_{i_1}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3] \\
&\quad - \underbrace{\sum_{(i_1, i_2, i_3)} s(i_1, i_2, i_3) [\mathbf{0}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3]}_{=0} \\
&\quad + \underbrace{\sum_{(i_1, i_2, i_3)} s(i_1, i_2, i_3) [\mathbf{0}, \mathbf{e}_{i_1}, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3]}_{=0} \\
&\quad - \sum_{(i_1, i_2, i_3)} s(i_1, i_2, i_3) [\mathbf{0}, \mathbf{e}_{i_1}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}].
\end{aligned}$$

Thus,

$$\begin{aligned}
\partial J^3 &= \sum_{(i_1, i_2, i_3)} s(i_1, i_2, i_3) [\mathbf{e}_{i_1}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3] \\
&\quad - \sum_{(i_1, i_2, i_3)} s(i_1, i_2, i_3) [\mathbf{0}, \mathbf{e}_{i_1}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}]
\end{aligned}$$

is the sum of 12 oriented affine 2-simplexes. (Note that $3! = 6$.)

- (4) Show that $\mathbf{x} = (x_1, x_2, x_3)$ is in the range of σ_1 if and only if $0 \leq x_3 \leq x_2 \leq x_1 \leq 1$.

- (a) By (1), \mathbf{x} is in the range of σ_1 if and only if $\mathbf{x} = A\mathbf{u}$ for $\mathbf{u} = (u_1, u_2, u_3) \in Q^3$, or

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix} = \begin{bmatrix} u_1 + u_2 + u_3 \\ u_2 + u_3 \\ u_3 \end{bmatrix}.$$

- (b) Since $\mathbf{u} = (u_1, u_2, u_3) \in Q^3$, $u_1 + u_2 + u_3 \leq 1$ and $u_1, u_2, u_3 \geq 0$. Hence $0 \leq u_3 \leq u_2 + u_3 \leq u_1 + u_2 + u_3 \leq 1$ or $0 \leq x_3 \leq x_2 \leq x_1 \leq 1$.
(c) Conversely, if $0 \leq x_3 \leq x_2 \leq x_1 \leq 1$, we define

$$\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} x_1 - x_2 \\ x_2 - x_3 \\ x_3 \end{bmatrix}.$$

Clearly, $\mathbf{v} \in Q^3$.

- (5) Show that the range of $\sigma_1, \dots, \sigma_6$ have disjoint interiors, and that their union covers I^3 . Similar to (4). By (2), $\mathbf{x} = P_{(i_1, i_2, i_3)} A P_{(j_1, j_2, 3)} \mathbf{u}$, or $P_{(i_1, i_2, i_3)}^{-1} \mathbf{x} = A P_{(j_1, j_2, 3)} \mathbf{u}$, or

$$\begin{bmatrix} x_{i_1} \\ x_{i_2} \\ x_{i_3} \end{bmatrix} = \begin{bmatrix} u_1 + u_2 + u_3 \\ u_{j_2} + u_3 \\ u_3 \end{bmatrix}.$$

In any case, we always have $0 \leq u_3 \leq u_{j_2} + u_3 \leq u_1 + u_2 + u_3 \leq 1$. Hence $\mathbf{x} = (x_1, x_2, x_3)$ is in the range of $\sigma_{(i_1, i_2, i_3)}$ if and only if

$$0 \leq x_{i_3} \leq x_{i_2} \leq x_{i_1} \leq 1.$$

The interior of $\sigma_{(i_1, i_2, i_3)}$ is

$$\{\mathbf{x} \in \mathbb{R}^3 : 0 < x_{i_3} < x_{i_2} < x_{i_1} < 1\},$$

and thus the range of $\sigma_1, \dots, \sigma_6$ have disjoint interiors. Also, any $\mathbf{x} \in I^3$ has the relation

$$0 \leq x_{i_3} \leq x_{i_2} \leq x_{i_1} \leq 1$$

for some permutation (i_1, i_2, i_3) of $(1, 2, 3)$. Hence

$$I^3 = \bigcup_{(i_1, i_2, i_3)} \sigma_{(i_1, i_2, i_3)}(Q^3) = \bigcup_{i=1}^6 \sigma_i(Q^3).$$

□

Exercise 10.19. Let J^2 and J^3 be as in Exercise 10.17 and Exercise 10.18. Define

$$\begin{aligned} B_{01}(u, v) &= (0, u, v), & B_{11}(u, v) &= (1, u, v), \\ B_{02}(u, v) &= (u, 0, v), & B_{12}(u, v) &= (u, 1, v), \\ B_{03}(u, v) &= (u, v, 0), & B_{13}(u, v) &= (u, v, 1). \end{aligned}$$

These are affine, and map \mathbb{R}^2 into \mathbb{R}^3 . Put $\beta_{ri} = B_{ri}(J^2)$, for $r = 0, 1$, $i = 1, 2, 3$. Each β_{ri} is an affine-oriented 2-chain. (See Section 10.30.) Verify that

$$\partial J^3 = \sum_{i=1}^3 (-1)^i (\beta_{0i} - \beta_{1i}),$$

in agreement with Exercise 10.18.)

Proof.

(1) A direct calculation shows that

$$\begin{aligned} B_{01}(\tau_1) - B_{11}(\tau_1) &= [\mathbf{0}, \mathbf{e}_2, \mathbf{e}_2 + \mathbf{e}_3] - [\mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3] \\ B_{02}(\tau_1) - B_{12}(\tau_1) &= [\mathbf{0}, \mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_3] - [\mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3] \\ B_{03}(\tau_1) - B_{13}(\tau_1) &= [\mathbf{0}, \mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_2] - [\mathbf{e}_3, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3] \\ B_{01}(\tau_2) - B_{11}(\tau_2) &= -[\mathbf{0}, \mathbf{e}_3, \mathbf{e}_2 + \mathbf{e}_3] + [\mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3] \\ B_{02}(\tau_2) - B_{12}(\tau_2) &= -[\mathbf{0}, \mathbf{e}_3, \mathbf{e}_1 + \mathbf{e}_3] + [\mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3] \\ B_{03}(\tau_2) - B_{13}(\tau_2) &= -[\mathbf{0}, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_2] + [\mathbf{e}_3, \mathbf{e}_2 + \mathbf{e}_3, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3]. \end{aligned}$$

(2) To express the formula in (1) clearly, we define

$$\omega_{(i_1, i_2, i_3)} = [\mathbf{e}_{i_1}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3] - [\mathbf{0}, \mathbf{e}_{i_2}, \mathbf{e}_{i_2} + \mathbf{e}_{i_3}],$$

and thus

$$\begin{aligned} -(B_{01}(\tau_1) - B_{11}(\tau_1)) &= s(1, 2, 3)\omega_{(1, 2, 3)} \\ B_{02}(\tau_1) - B_{12}(\tau_1) &= s(2, 1, 3)\omega_{(2, 1, 3)} \\ -(B_{03}(\tau_1) - B_{13}(\tau_1)) &= s(3, 1, 2)\omega_{(3, 1, 2)} \\ -(B_{01}(\tau_2) - B_{11}(\tau_2)) &= s(1, 3, 2)\omega_{(1, 3, 2)} \\ B_{02}(\tau_2) - B_{12}(\tau_2) &= s(2, 3, 1)\omega_{(2, 3, 1)} \\ -(B_{03}(\tau_2) - B_{13}(\tau_2)) &= s(3, 2, 1)\omega_{(3, 2, 1)}. \end{aligned}$$

(3) Note that

$$\begin{aligned} \beta_{0i} - \beta_{1i} &= B_{0i}(J^2) - B_{1i}(J^2) \\ &= B_{0i}(\tau_1 + \tau_2) - B_{1i}(\tau_1 + \tau_2) \\ &= B_{0i}(\tau_1) + B_{0i}(\tau_2) - B_{1i}(\tau_1) - B_{1i}(\tau_2) \\ &= (B_{0i}(\tau_1) - B_{1i}(\tau_1)) + (B_{0i}(\tau_2) - B_{1i}(\tau_2)). \end{aligned}$$

Thus,

$$\begin{aligned}
& \sum_{i=1}^3 (-1)^i (\beta_{0i} - \beta_{1i}) \\
&= \sum_{i=1}^3 (-1)^i (B_{0i}(\tau_1) - B_{1i}(\tau_1)) + \sum_{i=1}^3 (-1)^i (B_{0i}(\tau_2) - B_{1i}(\tau_2)) \\
&= \sum_{(i_1, i_2, i_3)} s(i_1, i_2, i_3) \omega_{(i_1, i_2, i_3)} \\
&= \sum_{(i_1, i_2, i_3)} s(i_1, i_2, i_3) [\mathbf{e}_{i_1}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3] \\
&\quad - \sum_{(i_1, i_2, i_3)} s(i_1, i_2, i_3) [\mathbf{0}, \mathbf{e}_{i_1}, \mathbf{e}_{i_1} + \mathbf{e}_{i_2}] \\
&= \partial J^3.
\end{aligned}$$

□

Exercise 10.20. *State conditions under which the formula*

$$\int_{\Phi} f d\omega = \int_{\partial\Phi} f\omega - \int_{\Phi} (df) \wedge \omega$$

is valid, and show that it generalizes the formula for integration by parts. (Hint: $d(f\omega) = (df) \wedge \omega + f d\omega$.)

Proof.

(1) *If*

- (a) Φ is a k -chain of class \mathcal{C}'' in an open set $V \subseteq \mathbb{R}^m$,
- (b) ω is a $(k-1)$ -form of class \mathcal{C}' in V ,
- (c) f is a 0-form of class \mathcal{C}' in V ,

then

$$\int_{\Phi} f d\omega = \int_{\partial\Phi} f\omega - \int_{\Phi} (df) \wedge \omega$$

(2) Theorem 10.20(a) implies that

$$d(f\omega) = (df) \wedge \omega + f d\omega.$$

(3) The Stokes' theorem (Theorem 10.33) shows that

$$\int_{\Phi} d(f\omega) = \int_{\partial\Phi} f\omega.$$

Hence

$$\int_{\Phi} f d\omega = \int_{\Phi} d(f\omega) - \int_{\Phi} (df) \wedge \omega = \int_{\partial\Phi} f\omega - \int_{\Phi} (df) \wedge \omega.$$

(4) Define $\Phi : Q^1 = [0, 1] \rightarrow [a, b]$ by

$$\Phi(\alpha) = a + \alpha(b - a).$$

Φ is a 1-simplex of class \mathcal{C}'' in an open set $V \supseteq [a, b]$. Also,

$$\partial\Phi = [b] - [a].$$

Let $\omega = g$ be a 0-form of class $\mathcal{C}'(V)$.

(5) Note that

$$\begin{aligned} \int_{\Phi} f d\omega &= \int_{\Phi} f dg = \int_0^1 f(\Phi(t))g'(\Phi(t))\Phi'(t)dt = \int_a^b f(u)g'(u)du, \\ \int_{\partial\Phi} f\omega &= \int_{[b]} fg + \int_{-[a]} fg = f(b)g(b) + (-1)f(a)f(a), \\ \int_{\Phi} (df) \wedge \omega &= \int_{\Phi} (df)g = \int_0^1 f'(\Phi(t))g(\Phi(t))\Phi'(t)dt = \int_a^b f'(u)g(u)du. \end{aligned}$$

Hence

$$\int_a^b f(u)g'(u)du = f(b)g(b) - f(a)f(a) - \int_a^b f'(u)g(u)du,$$

which is the same as the integration by parts (Theorem 6.22).

□

Exercise 10.21. *As in Example 10.36, consider the 1-form*

$$\eta = \frac{xdy - ydx}{x^2 + y^2}$$

in $\mathbb{R}^2 - \{\mathbf{0}\}$.

(a) *Carry out the computation that leads to*

$$\int_{\gamma} \eta = 2\pi \neq 0,$$

and prove that $d\eta = 0$.

(b)

- (c) Take $\Gamma(t) = (a \cos t, b \sin t)$ where $a > 0$, $b > 0$ are fixed. Use part (b) to show that

$$\int_0^{2\pi} \frac{ab}{a^2 \cos^2 t + b^2 \sin^2 t} dt = 2\pi.$$

- (d) Show that

$$\eta = d \left(\arctan \frac{y}{x} \right)$$

in any convex open set in which $x \neq 0$, and that

$$\eta = d \left(-\arctan \frac{x}{y} \right)$$

in any convex open set in which $y \neq 0$. Explain why this justifies the notation $\eta = d\theta$, in spite of the fact that η is not exact in $\mathbb{R}^2 - \{0\}$.

- (e) Show that (b) can be derived from (d).

- (f) If Γ is any closed \mathcal{C}' -curve in $\mathbb{R}^2 - \{0\}$, prove that

$$\frac{1}{2\pi} \int_{\Gamma} \eta = \text{Ind}(\Gamma).$$

(See Exercise 8.23 for the definition of the index of a curve.)

Proof of (a).

(1)

$$\begin{aligned} \int_{\gamma} \eta &= \int_0^{2\pi} \frac{(r \cos t)d(r \sin t) - (r \sin t)d(r \cos t)}{(r \cos t)^2 + (r \sin t)^2} \\ &= \int_0^{2\pi} \frac{(r \cos t)(r \cos t) - (r \sin t)(-r \sin t)}{(r \cos t)^2 + (r \sin t)^2} dt \\ &= \int_0^{2\pi} dt \\ &= 2\pi. \end{aligned}$$

(2)

$$\begin{aligned}
d\eta &= d\left(\frac{xdy - ydx}{x^2 + y^2}\right) \\
&= d\left(\frac{x}{x^2 + y^2}\right) \wedge dy + \frac{x}{x^2 + y^2} \wedge d^2y \\
&\quad - d\left(\frac{y}{x^2 + y^2}\right) \wedge dx - \frac{y}{x^2 + y^2} \wedge d^2x \\
&= d\left(\frac{x}{x^2 + y^2}\right) \wedge dy - d\left(\frac{y}{x^2 + y^2}\right) \wedge dx \quad (d^2 = 0) \\
&= \left\{D_1\left(\frac{x}{x^2 + y^2}\right)dx + D_2\left(\frac{y}{x^2 + y^2}\right)dy\right\} \wedge dy \\
&\quad - \left\{D_1\left(\frac{x}{x^2 + y^2}\right)dx + D_2\left(\frac{y}{x^2 + y^2}\right)dy\right\} \wedge dx \\
&= D_1\left(\frac{x}{x^2 + y^2}\right)dx \wedge dy \quad (dy \wedge dy = 0) \\
&\quad - D_2\left(\frac{y}{x^2 + y^2}\right)dy \wedge dx \quad (dx \wedge dx = 0) \\
&= 0
\end{aligned}$$

□

Note.

- (1) η is closed and locally exact, that is, $\eta = dt$ on $\mathbb{R}^2 - L$ where L is a half-line issuing from $\mathbf{0}$. η is not exact since $\int_\gamma \eta = 2\pi \neq 0$.
- (2) (*Poincaré's Lemma for 1-form.*) Let $\omega = \sum a_i dx_i$ be defined in an open set $U \subseteq \mathbb{R}^n$. Then $d\omega = 0$ if and only if for each $p \in U$ there is a neighborhood $V \subseteq U$ of p and a differentiable function $f : V \rightarrow \mathbb{R}^1$ with $df = \omega$ (i.e., ω is locally exact).

Proof of (b).

(1)

(2)

□

Proof of (c).

- (1) Γ satisfies all conditions described in (b). So

$$\int_\Gamma \eta = 2\pi.$$

(2) A direct calculation shows that

$$\begin{aligned}
2\pi &= \int_{\Gamma} \eta = \int_{\Gamma} \frac{xdy - ydx}{x^2 + y^2} \\
&= \int_0^{2\pi} \frac{a \cos(t) d(b \sin(t)) - b \sin(t) d(a \cos(t))}{(a \cos(t))^2 + (b \sin(t))^2} \\
&= \int_0^{2\pi} \frac{ab(\cos^2 t + \sin^2 t)}{a^2 \cos^2 t + b^2 \sin^2 t} \\
&= \int_0^{2\pi} \frac{ab}{a^2 \cos^2 t + b^2 \sin^2 t}.
\end{aligned}$$

□

Proof of (d).

(1) In any convex open set in which $x \neq 0$, we have

$$\begin{aligned}
d\left(\arctan \frac{y}{x}\right) &= \left(D_1 \arctan \frac{y}{x}\right) dx + \left(D_2 \arctan \frac{y}{x}\right) dy \\
&= -\frac{y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy \\
&= \eta.
\end{aligned}$$

(2) In any convex open set in which $y \neq 0$, we have

$$\begin{aligned}
d\left(-\arctan \frac{x}{y}\right) &= \left(D_1 \left(-\arctan \frac{x}{y}\right)\right) dx + \left(D_2 \left(-\arctan \frac{x}{y}\right)\right) dy \\
&= -\frac{y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy \\
&= \eta.
\end{aligned}$$

(3) By (1)(2), η is locally exact. Note that $\theta_1 = \arctan \frac{y}{x}$ and $\theta_2 = -\arctan \frac{x}{y}$ cannot be patched together to defined a global 0-form θ on $\mathbb{R}^2 - \{\mathbf{0}\}$.

□

Proof of (e).

(1)

(2)

□

Proof of (f).

(1)

(2)

□

Exercise 10.22. *As in Example 10.37, define ζ in $\mathbb{R}^3 - \{\mathbf{0}\}$ by*

$$\zeta = \frac{xdy \wedge dz + ydz \wedge dx + zdx \wedge dy}{r^3}$$

where $r = (x^2 + y^2 + z^2)^{\frac{1}{2}}$,

Proof of (a).

(1)

(2)

□

Proof of (b).

(1)

(2)

□

Proof of (c).

(1)

(2)

□

Proof of (d).

(1)

(2)

□

Proof of (e).

(1)

(2)

□

Proof of (f).

(1)

(2)

□

Proof of (g).

(1)

(2)

□

Exercise 10.23. ...

Proof of (a).

(1)

(2)

□

Proof of (b).

(1)

(2)

□

Proof of (c).

(1)

(2)

□

Proof of (d).

(1)

(2)

□

Exercise 10.24. Let $\omega = \sum a_i(\mathbf{x})dx_i$ be a 1-form of class \mathcal{C}'' in a convex open set $E \subseteq \mathbb{R}^n$. Assume $d\omega = 0$ and prove that ω is exact in E , by completing the following outline:

Fix $\mathbf{p} \in E$. Define

$$f(\mathbf{x}) = \int_{[\mathbf{p}, \mathbf{x}]} \omega \quad (\mathbf{x} \in E).$$

Apply Stokes' theorem to affine-oriented 2-simplexes $[\mathbf{p}, \mathbf{x}, \mathbf{y}]$ in E . Deduce that

$$f(\mathbf{y}) - f(\mathbf{x}) = \sum_{i=1}^n (y_i - x_i) \int_0^1 a_i((1-t)\mathbf{x} + t\mathbf{y}) dt$$

for $\mathbf{x} \in E, \mathbf{y} \in E$. Hence $(D_i f)(\mathbf{x}) = a_i(\mathbf{x})$.

Proof.

(1) Fix $\mathbf{p} \in E$. Define

$$f(\mathbf{x}) = \int_{[\mathbf{p}, \mathbf{x}]} \omega \quad (\mathbf{x} \in E).$$

(2) Given any $\mathbf{x} \in E, \mathbf{y} \in E$, and $\mathbf{x} \neq \mathbf{y}$. The affine-oriented 2-simplex $\Psi = [\mathbf{p}, \mathbf{x}, \mathbf{y}]$ is in E by the convexity of E . (If E is open but not convex, we can show that $\omega = df$ **locally** as the note in Exercise 10.21(a). That is why we say that ω is locally exact. The proof is exactly the same.)

(3) Note that

$$\partial\Psi = \partial[\mathbf{p}, \mathbf{x}, \mathbf{y}] = [\mathbf{x}, \mathbf{y}] - [\mathbf{p}, \mathbf{y}] + [\mathbf{p}, \mathbf{x}].$$

The Stokes' theorem (Theorem 10.33) implies that

$$\begin{aligned} \int_{\Psi} d\omega &= \int_{\partial\Psi} \omega \iff \int_{\Psi} 0 = \int_{[\mathbf{x}, \mathbf{y}]} \omega - \int_{[\mathbf{p}, \mathbf{y}]} \omega + \int_{[\mathbf{p}, \mathbf{x}]} \omega \\ &\iff 0 = \int_{[\mathbf{x}, \mathbf{y}]} \omega - f(\mathbf{y}) + f(\mathbf{x}) \\ &\iff f(\mathbf{y}) - f(\mathbf{x}) = \int_{[\mathbf{x}, \mathbf{y}]} \omega. \end{aligned}$$

(4) Define $\gamma : [0, 1] \rightarrow E$ by

$$\begin{aligned} \gamma(t) &= \mathbf{x} + t(\mathbf{y} - \mathbf{x}) \\ &= \sum_{i=1}^n x_i + t(y_i - x_i) \end{aligned}$$

(where $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_n)$). Hence $[0, 1]$ is the parameter domain of $[\mathbf{x}, \mathbf{y}]$ with respect to γ . So

$$\begin{aligned}\int_{[\mathbf{x}, \mathbf{y}]} \omega &= \int_0^1 \sum_{i=1}^n a_i(\gamma(t)) \frac{\partial(x_i + t(y_i - x_i))}{\partial t} dt \\ &= \int_0^1 \sum_{i=1}^n a_i(\mathbf{x} + t(\mathbf{y} - \mathbf{x}))(y_i - x_i) dt \\ &= \sum_{i=1}^n (y_i - x_i) \int_0^1 a_i(\mathbf{x} + t(\mathbf{y} - \mathbf{x})) dt.\end{aligned}$$

Thus,

$$f(\mathbf{y}) - f(\mathbf{x}) = \sum_{i=1}^n (y_i - x_i) \int_0^1 a_i(\mathbf{x} + t(\mathbf{y} - \mathbf{x})) dt.$$

(5) Note that

$$\begin{aligned}f(\mathbf{x} + h\mathbf{e}_j) - f(\mathbf{x}) &= \sum_{i=1}^n ((x_i + h\delta_{ij}) - x_i) \int_0^1 a_i(\mathbf{x} + t((\mathbf{x} + h\mathbf{e}_j) - \mathbf{x})) dt \\ &= \sum_{i=1}^n h\delta_{ij} \int_0^1 a_i(\mathbf{x} + t h\mathbf{e}_j) dt \\ &= h \int_0^1 a_j(\mathbf{x} + t h\mathbf{e}_j) dt.\end{aligned}$$

(Here δ_{ij} is the Kronecker delta.) So

$$\begin{aligned}(D_j f)(\mathbf{x}) &= \lim_{h \rightarrow 0} \frac{f(\mathbf{x} + h\mathbf{e}_j) - f(\mathbf{x})}{h} \\ &= \lim_{h \rightarrow 0} \int_0^1 a_j(\mathbf{x} + t h\mathbf{e}_j) dt \\ &= \int_0^1 a_j(\mathbf{x}) dt \quad (a_j \in \mathcal{C}'') \\ &= a_j(\mathbf{x}).\end{aligned}$$

Thus,

$$df = \sum_{j=1}^n (D_j f)(\mathbf{x}) dx_j = \sum_{j=1}^n a_j(\mathbf{x}) dx_j = \omega,$$

or ω is exact in E .

□

Exercise 10.25. Assume ω is a 1-form in an open set $E \subseteq \mathbb{R}^n$ such that

$$\int_{\gamma} \omega = 0$$

for every closed curve γ in E , of class \mathcal{C}' . Prove that ω is exact in E , by imitating part of the argument sketched in Exercise 10.24.

Proof.

- (1) Assume that E is a **connected** open subset of \mathbb{R}^n . Show that ω is exact in E if $\int_{\gamma} \omega = 0$ for every closed curve γ in E , of class \mathcal{C}' .

- (2) Fix $\mathbf{p} \in E$. Define

$$f(\mathbf{x}) = \int_{[\mathbf{p}, \mathbf{x}]} \omega \quad (\mathbf{x} \in E).$$

It is well-defined since E is connected and $\int_{\gamma} \omega = 0$ for every closed curve γ in E .

- (3) Given any $\mathbf{x} \in E$, $\mathbf{y} \in E$, and $\mathbf{x} \neq \mathbf{y}$. Let

$$\gamma = [\mathbf{x}, \mathbf{y}] - [\mathbf{p}, \mathbf{y}] + [\mathbf{p}, \mathbf{x}]$$

be a closed curve in E . Hence,

$$\begin{aligned} 0 &= \int_{\gamma} \omega && \text{(Assumption)} \\ &= \int_{[\mathbf{x}, \mathbf{y}]} \omega - \int_{[\mathbf{p}, \mathbf{y}]} \omega + \int_{[\mathbf{p}, \mathbf{x}]} \omega \\ &= \int_{[\mathbf{x}, \mathbf{y}]} \omega - f(\mathbf{y}) + f(\mathbf{x}). \end{aligned}$$

So

$$f(\mathbf{y}) - f(\mathbf{x}) = \int_{[\mathbf{x}, \mathbf{y}]} \omega$$

- (4) Similar to (4)(5) in the proof of Exercise 10.24, we have $df = \omega$. So the statement in (1) is proved. In general, we can define each f_{α} on each connected component E_{α} (which is open) of E such that $df_{\alpha} = \omega$ on E_{α} . Take

$$f|_{E_{\alpha}} = f_{\alpha}$$

on E . Hence, $df = \omega$ on the whole E .

□

Exercise 10.26. Assume ω is a 1-form in $\mathbb{R}^3 - \{\mathbf{0}\}$, of class \mathcal{C}' and $d\omega = 0$. Prove that ω is exact in $\mathbb{R}^3 - \{\mathbf{0}\}$. (Hint: Every closed continuously differentiable curve in $\mathbb{R}^3 - \{\mathbf{0}\}$ is the boundary of a 2-surface in $\mathbb{R}^3 - \{\mathbf{0}\}$. Apply Stokes' theorem and Exercise 10.25.)

Proof.

- (1) Let $E = \mathbb{R}^3 - \{\mathbf{0}\}$. By Exercise 10.25, it suffices to show that

$$\int_{\gamma} \omega = 0$$

for every closed curve γ in E , of class \mathcal{C}' .

- (2) Intuitively, every closed continuously differentiable curve in $\mathbb{R}^3 - \{\mathbf{0}\}$ is the boundary of a 2-surface in $\mathbb{R}^3 - \{\mathbf{0}\}$. So there is some 2-surface Ψ such that $\partial\Psi = \gamma$. The Stokes' theorem (Theorem 10.33) implies that

$$\int_{\gamma} \omega = \int_{\partial\Psi} \omega = \int_{\Psi} d\omega = \int_{\Psi} 0 = 0.$$

□

Exercise 10.27. ...

Proof.

- (1)

- (2)

□

Exercise 10.28. Fix $b > a > 0$, define

$$\Phi(r, \theta) = (r \cos \theta, r \sin \theta)$$

for $a \leq r \leq b$, $0 \leq \theta \leq 2\pi$. (The range of Φ is an annulus in \mathbb{R}^2 .) Put $\omega = x^3 dy$, and compute both

$$\int_{\Phi} d\omega \quad \text{and} \quad \int_{\partial\Phi} \omega$$

to verify that they are equal.

Proof.

(1) Note that

$$\frac{\partial(x, y)}{\partial(r, \theta)} = \det \begin{bmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{bmatrix} = r.$$

So

$$\begin{aligned} \int_{\Phi} d\omega &= \int_{\Phi} 3x^2 dx \wedge dy & (dy \wedge dy = 0) \\ &= \int_{[a, b] \times [0, 2\pi]} 3(r \cos \theta)^2 \frac{\partial(x, y)}{\partial(r, \theta)} dr d\theta \\ &= \int_a^b \int_0^{2\pi} 3r^3 (\cos \theta)^2 dr d\theta \\ &= \frac{3\pi}{4} (b^4 - a^4). \end{aligned}$$

(2) Similar to Exercise 10.21(b), write

$$\partial\Phi = \Gamma - \gamma,$$

where $\Gamma(t) = (b \cos t, b \sin t)$ on $[0, 2\pi]$ and $\gamma(t) = (a \cos t, a \sin t)$ on $[0, 2\pi]$.
Hence

$$\begin{aligned} \int_{\partial\Phi} \omega &= \int_{\Gamma} \omega - \int_{\gamma} \omega \\ &= \int_{\Gamma} x^3 dy - \int_{\gamma} x^3 dy \\ &= \int_{[0, 2\pi]} (b \cos \theta)^3 \frac{\partial y}{\partial \theta} d\theta - \int_{[0, 2\pi]} (a \cos \theta)^3 \frac{\partial y}{\partial \theta} d\theta \\ &= \int_0^{2\pi} b^4 (\cos \theta)^4 d\theta - \int_0^{2\pi} a^4 (\cos \theta)^4 d\theta \\ &= \frac{3\pi}{4} (b^4 - a^4). \end{aligned}$$

(3)

$$\int_{\Phi} d\omega = \int_{\partial\Phi} \omega = \frac{3\pi}{4} (b^4 - a^4).$$

□

Exercise 10.29. ...

Proof.

(1)

(2)

□

Exercise 10.30. If \mathbf{N} is the vector given by

$$\mathbf{N} = (\alpha_2\beta_3 - \alpha_3\beta_2)\mathbf{e}_1 + (\alpha_3\beta_1 - \alpha_1\beta_3)\mathbf{e}_2 + (\alpha_1\beta_2 - \alpha_2\beta_1)\mathbf{e}_3$$

(Equation (135)), prove that

$$\det \begin{bmatrix} \alpha_1 & \beta_1 & \alpha_2\beta_3 - \alpha_3\beta_2 \\ \alpha_2 & \beta_2 & \alpha_3\beta_1 - \alpha_1\beta_3 \\ \alpha_3 & \beta_3 & \alpha_1\beta_2 - \alpha_2\beta_1 \end{bmatrix} = |\mathbf{N}|^2$$

Also, verify

$$\mathbf{N} \cdot (T\mathbf{e}_1) = \mathbf{N} \cdot (T\mathbf{e}_2)$$

(Equation (137)).

Proof.

(1) By Laplace's expansion along the third column,

$$\begin{aligned} & \det \begin{bmatrix} \alpha_1 & \beta_1 & \alpha_2\beta_3 - \alpha_3\beta_2 \\ \alpha_2 & \beta_2 & \alpha_3\beta_1 - \alpha_1\beta_3 \\ \alpha_3 & \beta_3 & \alpha_1\beta_2 - \alpha_2\beta_1 \end{bmatrix} \\ &= (-1)^{1+3}(\alpha_2\beta_3 - \alpha_3\beta_2) \det \begin{bmatrix} \alpha_2 & \beta_2 \\ \alpha_3 & \beta_3 \end{bmatrix} \\ & \quad + (-1)^{2+3}(\alpha_3\beta_1 - \alpha_1\beta_3) \det \begin{bmatrix} \alpha_1 & \beta_1 \\ \alpha_3 & \beta_3 \end{bmatrix} \\ & \quad + (-1)^{3+3}(\alpha_1\beta_2 - \alpha_2\beta_1) \det \begin{bmatrix} \alpha_1 & \beta_1 \\ \alpha_2 & \beta_2 \end{bmatrix} \\ &= (\alpha_2\beta_3 - \alpha_3\beta_2)^2 + (\alpha_3\beta_1 - \alpha_1\beta_3)^2 + (\alpha_1\beta_2 - \alpha_2\beta_1)^2 \\ &= |\mathbf{N}|^2. \end{aligned}$$

(2)

$$\begin{aligned} \mathbf{N} \cdot (T\mathbf{e}_1) &= (\alpha_2\beta_3 - \alpha_3\beta_2, \alpha_3\beta_1 - \alpha_1\beta_3, \alpha_1\beta_2 - \alpha_2\beta_1) \cdot (\alpha_1, \alpha_2, \alpha_3) \\ &= (\alpha_2\beta_3 - \alpha_3\beta_2)\alpha_1 + (\alpha_3\beta_1 - \alpha_1\beta_3)\alpha_2 + (\alpha_1\beta_2 - \alpha_2\beta_1)\alpha_3 \\ &= (\alpha_3\alpha_2 - \alpha_2\alpha_3)\beta_1 + (\alpha_1\alpha_3 - \alpha_3\alpha_1)\beta_2 + (\alpha_2\alpha_1 - \alpha_1\alpha_2)\beta_3 \\ &= 0. \end{aligned}$$

(3)

$$\begin{aligned} \mathbf{N} \cdot (T\mathbf{e}_2) &= (\alpha_2\beta_3 - \alpha_3\beta_2, \alpha_3\beta_1 - \alpha_1\beta_3, \alpha_1\beta_2 - \alpha_2\beta_1) \cdot (\beta_1, \beta_2, \beta_3) \\ &= (\alpha_2\beta_3 - \alpha_3\beta_2)\beta_1 + (\alpha_3\beta_1 - \alpha_1\beta_3)\beta_2 + (\alpha_1\beta_2 - \alpha_2\beta_1)\beta_3 \\ &= (\beta_2\beta_3 - \beta_3\beta_2)\alpha_1 + (\beta_3\beta_1 - \beta_1\beta_3)\alpha_2 + (\beta_1\beta_2 - \beta_2\beta_1)\alpha_3 \\ &= 0. \end{aligned}$$

□

Exercise 10.31. ...

Proof of (a).

(1)

(2)

□

Proof of (b).

(1)

(2)

□

Proof of (c).

(1)

(2)

□

Proof of (d).

(1)

(2)

□

Exercise 10.32. ...

Proof.

(1)

(2)

□