

Solutions to the book: *Fulton, Algebraic Curves*

Meng-Gen Tsai
plover@gmail.com

March 15, 2021

Contents

Chapter 1: Affine Algebraic Sets	5
1.1. Algebraic Preliminaries	5
Problem 1.1.*	5
Problem 1.2.*	6
Problem 1.3.*	7
Problem 1.4.*	8
Problem 1.5.*	9
Problem 1.6.*	9
Problem 1.7.*	10
1.2. Affine Space and Algebraic Sets	12
Problem 1.8.*	12
Problem 1.9.	13
Problem 1.10.	13
Problem 1.11.	13
Problem 1.12.	14
Problem 1.13.	15
Problem 1.14.*	17
Problem 1.15.*	19
1.3. The Ideal of a Set of Points	19
Problem 1.16.*	19
Problem 1.17.*	20
Problem 1.18.*	21
Problem PLACEHOLDER	22
Problem 1.20.*	23
Problem 1.21.*	23
1.4. The Hilbert Basis Theorem	24
Problem PLACEHOLDER	24
1.5. Irreducible Components of an Algebraic Set	24
Problem PLACEHOLDER	24

Problem PLACEHOLDER	25
Problem PLACEHOLDER	25
Problem PLACEHOLDER	25
Problem PLACEHOLDER	25
Problem PLACEHOLDER	26
Problem PLACEHOLDER	26
1.6. Algebraic Subsets of the Plane	26
Problem PLACEHOLDER	26
Problem PLACEHOLDER	26
1.7. Hilbert's Nullstellensatz	27
Problem PLACEHOLDER	27
Problem PLACEHOLDER	27
Problem PLACEHOLDER	27
Problem PLACEHOLDER	27
Problem PLACEHOLDER	28
Problem PLACEHOLDER	28
Problem PLACEHOLDER	28
Problem PLACEHOLDER	28
Problem PLACEHOLDER	29
1.8. Modules; Finiteness Conditions	29
Problem 1.41.*	29
Problem 1.42.	29
Problem 1.43.* (WIP)	30
Problem PLACEHOLDER	30
Problem PLACEHOLDER	30
1.9. Integral Elements	30
Problem PLACEHOLDER	30
Problem PLACEHOLDER	31
Problem PLACEHOLDER	31
Problem PLACEHOLDER	31
Problem PLACEHOLDER	31
1.10. Field Extensions	32
Problem PLACEHOLDER	32
Problem PLACEHOLDER	32
Problem PLACEHOLDER	32
Problem PLACEHOLDER	32
Chapter 2: Affine Varieties	34
2.1. Coordinate Rings	34
Problem 2.1.*	34
Problem PLACEHOLDER	34
2.2. Polynomial Maps	35
2.3. Coordinate Changes	35
2.4. Rational Functions and Local Rings	35
2.5. Discrete Valuation Rings	35
2.6. Forms	35

2.7. Direct Products of Rings	35
2.8. Operations with Ideals	35
Problem 2.39.*	35
Problem 2.41.*	37
2.9. Ideals with a Finite Number of Zeros	39
2.10. Quotient Modules and Exact Sequences	39
Problem 2.51.	39
2.11. Free Modules	40
Chapter 3: Local Properties of Plane Curves	41
3.1. Multiple Points and Tangent Lines	41
Problem PLACEHOLDER	41
3.2. Multiplicities and Local Rings	41
3.3. Intersection Numbers	41
Chapter 4: Projective Varieties	42
4.1. Projective Space	42
Problem PLACEHOLDER	42
4.2. Projective Algebraic Sets	42
4.3. Affine and Projective Varieties	42
4.4. Multiprojective Space	42
Chapter 5: Projective Plane Curves	43
5.1. Definitions	43
Problem PLACEHOLDER	43
5.2. Linear Systems of Curves	43
5.3. Bézout's Theorem	43
5.4. Multiple Points	43
5.5. Max Noether's Fundamental Theorem	43
5.6. Applications of Noether's Theorem	43
Chapter 6: Varieties, Morphisms, and Rational Maps	44
6.1. The Zariski Topology	44
6.2. Varieties	44
6.3. Morphisms of Varieties	44
6.4. Products and Graphs	44
6.5. Algebraic Function Fields and Dimension of Varieties	44
6.6. Rational Maps	44
Chapter 7: Resolution of Singularities	45
7.1. Rational Maps of Curves	45
Problem PLACEHOLDER	45
7.2. Blowing up a Point in \mathbf{A}^2	45
7.3. Blowing up a Point in \mathbf{P}^2	45
7.4. Quadratic Transformations	45
7.5. Nonsingular Models of Curves	45

Chapter 8: Riemann-Roch Theorem	46
8.1. Divisors	46
Problem PLACEHOLDER	46
8.1. The Vector Spaces $L(D)$	46
8.1. Riemann's Theorem	46
8.1. Derivations and Differentials	46
8.1. Canonical Divisors	46
8.6. Riemann-Roch Theorem	46

Chapter 1: Affine Algebraic Sets

1.1. Algebraic Preliminaries

Problem 1.1.*

Let R be a domain.

- (a) If f, g are forms of degree r, s respectively in $R[x_1, \dots, x_n]$, show that fg is a form of degree $r + s$.
- (b) Show that any factor of a form in $R[x_1, \dots, x_n]$ is also a form.

Proof of (a).

- (1) Write

$$f = \sum_{(i)} a_{(i)} x^{(i)},$$
$$g = \sum_{(j)} b_{(j)} x^{(j)},$$

where $\sum_{(i)}$ is the summation over $(i) = (i_1, \dots, i_n)$ with $i_1 + \dots + i_n = r$ and $\sum_{(j)}$ is the summation over $(j) = (j_1, \dots, j_n)$ with $j_1 + \dots + j_n = s$.

- (2) Hence,

$$fg = \sum_{(i)} \sum_{(j)} a_{(i)} b_{(j)} x^{(i)} x^{(j)}$$
$$= \sum_{(i), (j)} a_{(i)} b_{(j)} x^{(k)}$$

where $(k) = (i_1 + j_1, \dots, i_n + j_n)$ with $(i_1 + j_1) + \dots + (i_n + j_n) = r + s$. Each $x^{(k)}$ is the form of degree $r + s$ and $a_{(i)} b_{(j)} \in R$. Hence fg is a form of degree $r + s$.

□

Proof of (b).

- (1) Given any form $f \in R[x_1, \dots, x_n]$, and write $f = gh$. It suffices to show that g is a form as well. (So does h .)
- (2) Write

$$g = g_0 + \dots + g_r, \quad h = h_0 + \dots + h_s$$

where $g_r \neq 0$ and $h_s \neq 0$. So

$$f = gh = g_0h_0 + \cdots + g_rh_s.$$

Since R is a domain, $R[x_1, \dots, x_n]$ is a domain and thus $g_rh_s \neq 0$. The maximality of r and s implies that $\deg f = r + s$. Therefore, by the maximality of $r + s$, $f = g_rh_s$, or $g = g_r$, or g is a form.

□

Problem 1.2.*

Let R be a UFD, K the quotient field of R . Show that every element z of K may be written $z = a/b$, where $a, b \in R$ have no common factors; this representative is unique up to units of R .

Proof.

- (1) Show that every element z of K may be written $z = a/b$, where $a, b \in R$ have no common factors. Given any $z = a/b \in K$ where $a, b \in R$. Write

$$\begin{aligned} a &= p_1 \cdots p_n, \\ b &= q_1 \cdots q_m \end{aligned}$$

where all $p_1, \dots, p_n, q_1, \dots, q_m$ are irreducible in R . (It is possible since R is a UFD.) For each i , suppose $p_i \mid q_j$ for some i, j . Write $q_j = p_i u$ for some $u \in R$. By the irreducibility of p_i and q_j , u is a unit. So

$$z = \frac{a}{b} = \frac{p_1 \cdots \widehat{p_i} \cdots p_n}{q_1 \cdots \widehat{q_j} \cdots q_m} = \frac{p_1 \cdots \widehat{p_i} \cdots p_n}{u q_1 \cdots \widehat{q_j} \cdots q_m}.$$

Continue this method we can write $z = \frac{a'}{b'}$ where a' and b' have no common factors.

- (2) Write $z = a/b = a'/b'$ where

- (a) $a, b, a', b' \in R$,
- (b) a and b have no common factors,
- (c) a' and b' have no common factors.

Write

$$\begin{aligned} a &= p_1 \cdots p_n, \\ b &= q_1 \cdots q_m, \\ a' &= p'_1 \cdots p'_{n'}, \\ b' &= q'_1 \cdots q'_{m'} \end{aligned}$$

where all $p_i, q_j, p'_{i'}, q'_{j'}$ are irreducible in R . As $z = a/b = a'/b'$, $ab' = a'b$ or

$$p_1 \cdots p_n q'_1 \cdots q'_{m'} = p'_1 \cdots p'_{n'} q_1 \cdots q_m.$$

- (3) For $i = 1$, $p_1 = u_1 p'_{i'}$ for some unit $u_1 \in R$ since a and b have no common factors and all $p_1, q_j, p'_{i'}$ are irreducible. Hence

$$u_1 \widehat{p_1} p_2 \cdots p_n q'_1 \cdots q'_{m'} = p'_1 \cdots \widehat{p'_{i'}} \cdots p'_{n'} q_1 \cdots q_m.$$

Continue this method, we have $n \leq n'$ and all p_1, \dots, p_n are canceled.

- (4) Conversely, we can apply the argument in (3) to $i' = 1, \dots, n'$ to conclude that $n' \leq n$. Therefore, $n = n'$ and

$$\underbrace{u_1 \cdots u_n}_{\text{a unit in } R} q'_1 \cdots q'_{m'} = q_1 \cdots q_m.$$

Hence, $b = ub'$ where $u = u_1 \cdots u_n$ is a unit in R . Similarly, $a = va'$ where v is a unit in R . So the representative of $z \in K$ is unique up to units of R .

□

Problem 1.3.*

Let R be a PID. Let \mathfrak{p} be a nonzero, proper, prime ideal in R .

- (a) Show that \mathfrak{p} is generated by an irreducible element.
- (b) Show that \mathfrak{p} is maximal.

Proof of (a).

- (1) Let $\mathfrak{p} = (a)$ be a nonzero, proper, prime ideal in R . It suffices to show that a is irreducible.
- (2) Suppose $a = bc$. By the primality of \mathfrak{p} , $b \in \mathfrak{p}$ or $c \in \mathfrak{p}$. Suppose $b \in \mathfrak{p} = (a)$. (The case $c \in \mathfrak{p}$ is similar.) Then there is a $d \in R$ such that $b = ad$. Hence, $a = bc = adc$ or $(1 - dc)a = 0$.
- (3) Since R is a domain, $1 = dc$ or $a = 0$. $a = 0$ implies that $\mathfrak{p} = (0)$ is a zero ideal, contrary to the assumption. Therefore, $1 = dc$, or c is a unit, or a is irreducible.

□

Proof of (b).

- (1) Given any ideal $I = (b)$ of R containing $\mathfrak{p} = (a)$. As the generator a of \mathfrak{p} is in $\mathfrak{p} \subseteq I$, there is some $c \in R$ such that $a = bc$. By the irreducibility of a (in (a)), b is a unit or c is a unit.
- (2) b is a unit implies that $I = R$. c is a unit implies that $I = \mathfrak{p}$. In any case, we conclude that \mathfrak{p} is maximal.

□

Problem 1.4.*

Let k be an infinite field, $f \in k[x_1, \dots, x_n]$. Suppose $f(a_1, \dots, a_n) = 0$ for all $a_1, \dots, a_n \in k$. Show that $f = 0$. (Hint: Write

$$f = \sum f_i x_n^i, \quad f_i \in k[x_1, \dots, x_{n-1}].$$

Use induction on n , and the fact that $f(a_1, \dots, a_{n-1}, x_n)$ has only a finite number of roots if any $f_i(a_1, \dots, a_{n-1}) \neq 0$.)

Proof.

- (1) Induction on n . The case $n = 1$. (Reductio ad absurdum) If there were a nonzero $f \in k[x_1]$ such that $f(a) = 0$ for all $a \in k$. Note that f has at most $\deg f < \infty$ roots, contrary to the infinity of k .
- (2) Assume that the conclusion holds for $n - 1$, then for any $f \in k[x_1, \dots, x_n]$ we can write

$$f = \sum f_i x_n^i, \quad f_i \in k[x_1, \dots, x_{n-1}]$$

as $f \in (k[x_1, \dots, x_{n-1}])[x_n]$. Suppose $f(a_1, \dots, a_n) = 0$ for all $a_1, \dots, a_n \in k$. For fixed a_1, \dots, a_{n-1} , the polynomial $f(a_1, \dots, a_{n-1}, x_n) \in k[x_n]$ has all distinct roots in an infinite field k . By (1), $f(a_1, \dots, a_{n-1}, x_n) = 0 \in k[x_n]$, or each $f_i(a_1, \dots, a_{n-1}) = 0$. As all a_1, \dots, a_{n-1} run over k , we can apply the induction hypothesis each $f_i(x_1, \dots, x_{n-1}) = 0 \in k[x_1, \dots, x_{n-1}]$. Hence, $f = 0 \in k[x_1, \dots, x_n]$.

□

Note. If k is a finite field of order $q = p^k$, then the polynomial $f(x) = x^q - x$ has q distinct roots in k .

Problem 1.5.*

Let k be any field. Show that there are an infinitely number of irreducible monic polynomials in $k[x]$. (Hint: Suppose f_1, \dots, f_n were all of them, and factor $f_1 \cdots f_n + 1$ into irreducible factors.)

Proof (Due to Euclid).

- (1) If f_1, \dots, f_n were all irreducible monic polynomials, then we consider

$$g = f_1 \cdots f_n + 1 \in k[x].$$

So there is an irreducible monic polynomial $f = f_i$ dividing g for some i since

$$\deg g = \deg f_1 + \cdots + \deg f_n \geq 1$$

and $k[x]$ is a UFD.

- (2) However, f would divide the difference

$$g - f_1 \cdots f_{i-1} f_{i+1} \cdots f_n = 1,$$

contrary to $\deg f_i \geq 1$.

□

Problem 1.6.*

Show that any algebraically closed field is infinite. (Hint: The irreducible monic polynomials are $x - a$, $a \in k$.)

Proof (Due to Euclid).

- (1) Let k be an algebraically closed field. If a_1, \dots, a_n were all elements in k , then we consider a monic polynomials

$$f(x) = (x - a_1) \cdots (x - a_n) + 1 \in k[x].$$

- (2) Since k is algebraically closed, there is an element $a \in k$ such that $f(a) = 0$. By assumption, $a = a_i$ for some $1 \leq i \leq n$, and thus $f(a) = f(a_i) = 1$, contrary to the fact that a field is a commutative ring where $0 \neq 1$ and all nonzero elements are invertible.

□

Problem 1.7.*

Let k be a field, $f \in k[x_1, \dots, x_n]$, $a_1, \dots, a_n \in k$.

(a) Show that

$$f = \sum \lambda_{(i)} (x_1 - a_1)^{i_1} \cdots (x_n - a_n)^{i_n}, \quad \lambda_{(i)} \in k.$$

(b) If $f(a_1, \dots, a_n) = 0$, show that $f = \sum_{i=1}^n (x_i - a_i)g_i$ for some (not unique) g_i in $k[x_1, \dots, x_n]$.

Proof of (a).

(1) Regard $k[x_1, \dots, x_n]$ as $(k[x_1, \dots, x_{n-1}])[x_n]$. Since $(k[x_1, \dots, x_{n-1}])[x_n]$ is a Euclidean domain with a function

$$f \in (k[x_1, \dots, x_{n-1}])[x_n] \mapsto \deg_{x_n} f \in \mathbb{Z}_{\geq 0}$$

satisfying the division-with-remainder property.

(2) Apply the division algorithm for f and nonzero $x_n - a_n$ to produce a quotient q and remainder r with $f = (x_n - a_n)q + r$ and either $r = 0$ or $\deg_{x_n}(r) < \deg_{x_n}(x_n - a_n) = 1$. That is, $r \in k[x_1, \dots, x_{n-1}]$ is a constant in $(k[x_1, \dots, x_{n-1}])[x_n]$. Continue this process to get that f is of the form

$$f = \sum_{i_n} f_{i_n} (x_n - a_n)^{i_n}$$

where $f_{i_n} \in k[x_1, \dots, x_{n-1}]$.

(3) Use the same argument in (2) for each $f_{i_n} \in k[x_1, \dots, x_{n-1}]$, we have

$$\begin{aligned} f_{i_n} &= \sum_{\substack{i_{n-1} \\ \in k[x_1, \dots, x_{n-2}]}} \underbrace{f_{i_n, i_{n-1}}}_{\in k[x_1, \dots, x_{n-2}]} (x_{n-1} - a_{n-1})^{i_{n-1}} \\ f_{i_n, i_{n-1}} &= \sum_{\substack{i_{n-2} \\ \in k[x_1, \dots, x_{n-3}]}} \underbrace{f_{i_n, i_{n-1}, i_{n-2}}}_{\in k[x_1, \dots, x_{n-3}]} (x_{n-2} - a_{n-2})^{i_{n-2}}, \\ &\dots \\ f_{i_n, \dots, i_2} &= \sum_{\substack{i_1 \\ \in k}} \underbrace{f_{i_n, \dots, i_1}}_{\in k} (x_1 - a_1)^{i_1}. \end{aligned}$$

Note that $f_{i_n, \dots, i_1} \in k$, we can write

$$f = \sum \lambda_{(i)} (x_1 - a_1)^{i_1} \cdots (x_n - a_n)^{i_n}, \quad \lambda_{(i)} \in k.$$

by replacing all f_{i_n, \dots, i_k} by $f_{i_n, \dots, i_{k-1}}$ for $k = n, n-1, \dots, 2$.

(4) Or use the induction on n .

□

Proof of (b).

(1) Write

$$f = \sum \lambda_{(i)} (x_1 - a_1)^{i_1} \cdots (x_n - a_n)^{i_n}, \quad \lambda_{(i)} \in k$$

by (a).

(2) As $f(a_1, \dots, a_n) = 0$, $\lambda_{(i)} = 0$ if all i_1, \dots, i_n are zero, that is, there is no nonzero constant term in the representation of f . Hence, for each term

$$f_{(i)} := \lambda_{(i)} (x_1 - a_1)^{i_1} \cdots (x_n - a_n)^{i_n}$$

with $\lambda_{(i)} \neq 0$, there exists one $i_k > 0$ for some $1 \leq k \leq n$. So we can write

$$f_{(i)} = (x_k - a_k) \underbrace{(\lambda_{(i)} (x_1 - a_1)^{i_1} \cdots (x_k - a_k)^{i_k-1} \cdots (x_n - a_n)^{i_n})}_{:= g_{(i)} \in k[x_1, \dots, x_n]}.$$

Note that the expression of $f_{(i)}$ is not unique since there may exist more than one $i_k > 0$ as $1 \leq k \leq n$.

(3) Now we iterate each nonzero term in f , apply the factorization in (2), and then group by each $x_k - a_k$. Therefore, we can write

$$f = \sum_{i=1}^n (x_i - a_i) g_i$$

for some $g_i \in k[x_1, \dots, x_n]$.

(4) The expression of f is not unique. For example, take $f(x, y) = x^2 + 2xy + y^2 \in k[x, y]$. As $f(0, 0) = 0$, we can write

$$\begin{aligned} f(x, y) &= x \cdot \underbrace{(x + 2y)}_{g_1} + y \cdot \underbrace{y}_{g_2}, \text{ or} \\ &= x \cdot \underbrace{(x + y)}_{g_1} + y \cdot \underbrace{(x + y)}_{g_2}, \text{ or} \\ &= x \cdot \underbrace{x}_{g_1} + y \cdot \underbrace{(2x + y)}_{g_2}. \end{aligned}$$

□

1.2. Affine Space and Algebraic Sets

Problem 1.8.*

Show that the algebraic subsets of $\mathbf{A}^1(k)$ are just the finite subsets, together with $\mathbf{A}^1(k)$ itself.

Proof.

(1) Show that $k[x]$ is a PID if k is a field.

- (a) Let I be an ideal of $k[x]$.
- (b) If $I = \{0\}$ then $I = (0)$ and I is principal.
- (c) If $I \neq \{0\}$, then take f to be a polynomial of minimal degree in I . It suffices to show that $I = (f)$. Clearly, $(f) \subseteq I$ since I is an ideal. Conversely, for any $g \in I$,

$$g(x) = f(x)h(x) + r(x)$$

for some $h, r \in k[x]$ with $r = 0$ or $\deg r < \deg f$ (as $k[x]$ is a Euclidean domain). Now as

$$r = g - fh \in I,$$

$r = 0$ (otherwise contrary to the minimality of f), we have $g = fh \in (f)$ for all $g \in I$.

(2) Let Y be an algebraic subset of $\mathbf{A}^1(k)$, say $Y = V(I)$ for some ideal I of $k[x]$. Since $k[x]$ is a PID, $I = (f)$ for some $f \in k[x]$.

- (a) If $f = 0$, then $I = (0)$ and $Y = V(0) = \mathbf{A}^1(k)$.
- (b) If $f \neq 0$, then $f(x) = 0$ has finitely many roots in k , say $a_1, \dots, a_m \in k$. Hence,

$$Y = V(I) = V(f) = \{f(a) = 0 : a \in k\} = \{a_1, \dots, a_m\}$$

is a finite subsets of $\mathbf{A}^1(k)$.

By (a)(b), the result is established.

□

Notes.

- (1) By the Hilbert basis theorem, $k[x]$ is Noetherian as k is Noetherian. Hence, for any algebraic subset $Y = V(I)$ of $\mathbf{A}^1(k)$, we can write $I = (f_1, \dots, f_m)$. Note that

$$Y = V(I) = V(f_1) \cap \dots \cap V(f_m).$$

Now apply the same argument to get the same conclusion.

- (2) Suppose $k = \bar{k}$. $\mathbf{A}^1(k)$ is irreducible, because its only proper closed subsets are finite, yet it is infinite (because k is algebraically closed, hence infinite).

Problem 1.9.

If k is a finite field, show that every subset of $\mathbf{A}^n(k)$ is algebraic.

Proof.

- (1) Every subset of $\mathbf{A}^n(k)$ is finite since $|\mathbf{A}^n(k)| = |k|^n$ is finite.
- (2) Note that $V(x_1 - a_1, \dots, x_n - a_n) = \{(a_1, \dots, a_n)\} \subseteq \mathbf{A}^n(k)$ (property (5) in this section) and any finite union of algebraic sets is algebraic (property (4) in this section). Thus, every subset of $\mathbf{A}^n(k)$ is algebraic (by (1)).

□

Problem 1.10.

Give an example of a countable collection of algebraic sets whose union is not algebraic.

Proof.

- (1) Let $k = \mathbb{Q}$ be an infinite field. $V(x - a) = \{a\}$ is an algebraic sets for all $a \in \mathbb{Q}$. In particular, $V(x - a) = \{a\}$ is algebraic for all $a \in \mathbb{Z}$.
- (2) Note that

$$Y := \bigcup_{a \in \mathbb{Z}} V(x - a) = \mathbb{Z}$$

is a countable union of algebraic sets. Since Y is a proper subset of $k = \mathbb{Q}$, it cannot be algebraic by Problem 1.8.

□

Problem 1.11.

Show that the following are algebraic sets:

- (a) $\{(t, t^2, t^3) \in \mathbf{A}^3(k) : t \in k\}$;
- (b) $\{(\cos(t), \sin(t)) \in \mathbf{A}^2(\mathbb{R}) : t \in \mathbb{R}\}$;
- (c) the set of points in $\mathbf{A}^2(\mathbb{R})$ whose polar coordinates (r, θ) satisfy the equation $r = \sin(\theta)$.

Proof of (a).

- (1) The twisted cubic curve

$$Y = \{(t, t^2, t^3) \in \mathbf{A}^3(k) : t \in k\} = V(x^2 - y) \cap V(x^3 - z)$$

is algebraic. We say that Y is given by the parametric representation $x = t, y = t^2, z = t^3$.

- (2) The generators for the ideal $I(Y)$ are $x^2 - y$ and $x^3 - z$.
 (3) Y is an affine variety of dimension 1.
 (4) The affine coordinate ring $A(Y)$ is isomorphic to a polynomial ring in one variable over k .

□

Proof of (b). The circle

$$\{(\cos(t), \sin(t)) \in \mathbf{A}^2(\mathbb{R}) : t \in \mathbb{R}\} = V(x^2 + y^2 - 1)$$

is algebraic. □

Proof of (c). The circle

$$\{(r, \theta) : r = \sin(\theta)\} = V(x^2 + y^2 - y)$$

is algebraic again. □

Problem 1.12.

Suppose C is an affine plane curve, and L is a line in $\mathbf{A}^2(k)$, $L \not\subseteq C$. Suppose $C = V(f)$, $f \in k[x, y]$ a polynomial of degree n . Show that $L \cap C$ is a finite set of no more than n points. (Hint: Suppose $L = V(y - (ax + b))$, and consider $f(x, ax + b) \in k[x]$.)

Proof.

- (1) Say $L = V(y - (ax + b))$ be a line in $\mathbf{A}^2(k)$. (The case $L = V(x - (ay + b))$ is similar.)
 (2) Note that $L \not\subseteq C$ implies that $(y - (ax + b)) \nmid f$. Hence, the polynomial

$$g : x \mapsto f(x, ax + b) \in k[x]$$

is nonzero and $\deg g \leq n$. Therefore, the number of roots of g in k is no more than n .

(3) Hence,

$$\begin{aligned}
L \cap C &= V(y - (ax + b)) \cap V(f) \\
&= \{(x, y) \in \mathbb{A}^2(k) : y = ax + b \text{ and } f(x, y) = 0\} \\
&= \{(x, y) \in \mathbb{A}^2(k) : f(x, ax + b) = 0\}
\end{aligned}$$

is finite of no more than n points.

□

Problem 1.13.

Show that each of the following sets is not algebraic:

- (a) $\{(x, y) \in \mathbf{A}^2(\mathbb{R}) : y = \sin(x)\}$.
- (b) $\{(z, w) \in \mathbf{A}^2(\mathbb{C}) : |z|^2 + |w|^2 = 1\}$, where $|x + iy|^2 = x^2 + y^2$ for $x, y \in \mathbb{R}$.
- (c) $\{(\cos(t), \sin(t), t) \in \mathbf{A}^3(\mathbb{R}) : t \in \mathbb{R}\}$.

Proof of (a).

- (1) (Reductio ad absurdum) If

$$Y := \{(x, y) \in \mathbf{A}^2(\mathbb{R}) : y = \sin(x)\}$$

were algebraic, then there is a subset S of $\mathbb{R}[x, y]$ such that

$$Y = V(S) = \bigcap_{f \in S} V(f).$$

- (2) $S \neq \emptyset$ since $Y \neq \mathbf{A}^2(\mathbb{R})$. ($(89, 64) \in \mathbf{A}^2(\mathbb{R}) - Y$.)
- (3) Take a fixed line $L = V(y)$ in $\mathbf{A}^2(\mathbb{R})$. For each affine curve $f \in S$, we have

$$V(f) \cap L \supseteq \bigcap_{f \in S} V(f) \cap L = Y \cap L = \{(n\pi, 0) \in \mathbf{A}^2(\mathbb{R}) : n \in \mathbb{Z}\},$$

which is infinite. By problem 1.12, $y \mid f$. As f runs over S , $Y \subseteq V(y) = L$, contradicts that $(0, \frac{\pi}{2}) \in L - Y$.

□

Proof of (b).

- (1) Similar to (a). (Reductio ad absurdum) If

$$Y := \{(x, y) \in \mathbf{A}^2(\mathbb{C}) : |x|^2 + |y|^2 = 1\}$$

were algebraic, then there is a subset S of $\mathbb{C}[x, y]$ such that

$$Y = V(S) = \bigcap_{f \in S} V(f).$$

- (2) $S \neq \emptyset$ since $Y \neq \mathbf{A}^2(\mathbb{C})$. $((89, 64) \in \mathbf{A}^2(\mathbb{C}) - Y)$
 (3) Take a fixed line $L = V(x)$ in $\mathbf{A}^2(\mathbb{C})$. For each affine curve $f \in S$, we have

$$V(f) \cap L \supseteq \bigcap_{f \in S} V(f) \cap L = Y \cap L = \{(0, y) \in \mathbf{A}^2(\mathbb{C}) : |y| = 1\},$$

which is infinite (since Y contains a unit circle in the complex plane). By problem 1.12, $x \mid f$. As f runs over S , $Y \subseteq V(x) = L$, contradicts that the origin $(0, 0) \in L - Y$.

□

Proof of (c).

- (1) Similar to (a) and (b).
 (2) Suppose C is an affine plane curve, and L is a line in $\mathbb{A}^3(k)$, $L \not\subseteq C$. Suppose $C = V(f)$, $f \in k[x, y, z]$ a polynomial of degree n . Show that $L \cap C$ is a finite set of no more than n points. The proof is similar to Problem 1.12.
 (a) Say $L = V(y - (ax + b), z - (cx + d))$ be a line in $\mathbb{A}^3(k)$.
 (b) Note that $L \not\subseteq C$ implies that $(y - (ax + b)) \nmid f$ and $(z - (cx + d)) \nmid f$. Hence, the polynomial

$$g : x \mapsto f(x, ax + b, cx + d) \in k[x]$$

is nonzero and $\deg g \leq n$. Therefore, the number of roots of g in k is no more than n .

- (c) Hence,

$$\begin{aligned} L \cap C &= V(y - (ax + b), z - (cx + d)) \cap V(f) \\ &= \{(x, y) \in \mathbb{A}^2(k) : y = ax + b, z = cx + d \text{ and } f(x, y) = 0\} \\ &= \{(x, y) \in \mathbb{A}^2(k) : f(x, ax + b, cx + d) = 0\} \end{aligned}$$

is finite of no more than n points.

(3) (Reductio ad absurdum) If

$$Y := \{(\cos(t), \sin(t), t) \in \mathbf{A}^3(\mathbb{R}) : t \in \mathbb{R}\}$$

were algebraic, then there is a subset S of $\mathbb{R}[x, y, z]$ such that

$$Y = V(S) = \bigcap_{f \in S} V(f).$$

(4) $S \neq \emptyset$ since $Y \neq \mathbf{A}^3(\mathbb{R})$. ((1989, 6, 4) $\in \mathbf{A}^3(\mathbb{R}) - Y$.)

(5) Take a fixed line $L = V(x - 1, y)$ in $\mathbf{A}^3(\mathbb{R})$. For each affine curve $f \in S$, we have

$$V(f) \cap L \supseteq \bigcap_{f \in S} V(f) \cap L = Y \cap L = \{(1, 0, 2n\pi) \in \mathbf{A}^3(\mathbb{R}) : n \in \mathbb{Z}\},$$

which is infinite. By (2), $(x - 1) \mid f$ and $y \mid f$. As f runs over S , $Y \subseteq V(x - 1, y) = L$, contradicts that $(1, 0, \pi) \in L - Y$.

□

Supplement. A circular disk of radius 1 in the plane xy rolls without slipping along the x axis. The figure described by a point of the circumference of the disk is called a **cycloid**. The parametrized curve $\alpha : \mathbb{R} \rightarrow \mathbb{R}^2$ is

$$\begin{cases} x = t - \sin t \\ y = 1 - \cos t. \end{cases}$$

The cycloid is not algebraic (as (a)).

Problem 1.14.*

Let f be a nonconstant polynomial in $k[x_1, \dots, x_n]$, k algebraically closed. Show that $\mathbf{A}^n(k) - V(f)$ is infinite if $n \geq 1$, and $V(f)$ is infinite if $n \geq 2$. Conclude that the complement of any proper algebraic set is infinite. (Hint: See Problem 1.4.)

Proof.

(1) Show that $\mathbf{A}^n(k) - V(f)$ is infinite if $n \geq 1$. Since f is a nonconstant polynomial in $k[x_1, \dots, x_n]$, we may assume that $\deg_{x_n}(f) > 0$. Hence

$$x_n \mapsto f(1, \dots, 1, x_n)$$

is a nonconstant polynomial of degree $\deg_{x_n}(f) > 0$ in $k[x_n]$. So f has finitely many roots in k , say ξ_1, \dots, ξ_m ($m \geq 0$). Hence,

$$(1, \dots, 1, x_n) \neq 0$$

whenever $x_n \neq \xi_m$. Such subset in $\mathbf{A}^1(k)$ is infinite since $k = \bar{k}$ (Problem 1.6). Therefore,

$$\begin{aligned}\mathbf{A}^n(k) - V(f) &= \{(a_1, \dots, a_n) \in \mathbf{A}^n(k) : f(a_1, \dots, a_n) \neq 0\} \\ &\supseteq \{a_n \in \mathbf{A}^1(k) : f(1, \dots, 1, x_n) \neq 0\}\end{aligned}$$

is infinite.

(2) Show that $V(f)$ is infinite if $n \geq 2$.

(a) Similar to (1). Since f is a nonconstant polynomial in $k[x_1, \dots, x_n]$, we may assume that $m := \deg_{x_n}(f) > 0$. Write

$$f = \sum_{i=0}^m f_i(x_1, \dots, x_{n-1})x_n^i.$$

Note that each f_i is well-defined since $n \geq 2$.

(b) If f_n is constant in $k[x_1, \dots, x_{n-1}]$, then f_n is nonzero (since $m > 0$) or $V(f_n) = \emptyset$. If f_n is nonconstant in $k[x_1, \dots, x_{n-1}]$, then the set $\mathbf{A}^{n-1}(k) - V(f_n)$ is infinite by (1). In any case,

$$\mathbf{A}^{n-1}(k) - V(f_n)$$

is infinite.

(c) For each $P = (a_1, \dots, a_{n-1}) \in \mathbf{A}^{n-1}(k) - V(f_n)$,

$$g_P : x_n \mapsto f(P, x_n) = f(a_1, \dots, a_{n-1}, x_n)$$

defines a polynomial in $k[x_n]$ of degree $m > 0$. Since $k = \bar{k}$, g_P has at least one root $Q \in k$. Hence

$$V(f) \supseteq \{(P, Q) \in \mathbf{A}^n(k) : P \in \mathbf{A}^{n-1}(k) - V(f_n), g_P(Q) = 0\}$$

is infinite since the set $\mathbf{A}^{n-1}(k) - V(f_n)$ is infinite.

Note. It is not true if $k \neq \bar{k}$. For example, $V(x^2 + y^2 + 1) = \emptyset$ in $\mathbf{A}^2(\mathbb{R})$.

(3) Note that

$$\mathbf{A}^n(k) - V(S) = \mathbf{A}^n(k) - \bigcap_{f \in S} V(f) = \bigcup_{f \in S} (\mathbf{A}^n(k) - V(f)).$$

Thus the complement of any proper algebraic set is infinite by (1).

□

Problem 1.15.*

Let $V \subseteq \mathbf{A}^n(k)$, $W \subseteq \mathbf{A}^m(k)$ be algebraic sets. Show that

$$V \times W = \{(a_1, \dots, a_n, b_1, \dots, b_m) : (a_1, \dots, a_n) \in V, (b_1, \dots, b_m) \in W\}$$

is an algebraic set in $\mathbf{A}^{n+m}(k)$. It is called the **product** of V and W .

Proof.

(1) Write

$$\begin{aligned} V &= V(S_V) = \{P \in \mathbf{A}^n(k) : f(P) = 0 \forall f \in S_V\} \\ W &= V(S_W) = \{Q \in \mathbf{A}^m(k) : g(Q) = 0 \forall g \in S_W\}, \end{aligned}$$

where $S_V \subseteq k[x_1, \dots, x_n]$ and $S_W \subseteq k[y_1, \dots, y_m]$. It suffices to show that

$$V \times W = V(S),$$

where $S \subseteq k[x_1, \dots, x_n, y_1, \dots, y_m]$ is the union of S_V and S_W .

(2) Here we can identify S_V with the subset of $k[x_1, \dots, x_n, y_1, \dots, y_m]$ by noting that

$$k[x_1, \dots, x_n] \hookrightarrow (k[y_1, \dots, y_m])[x_1, \dots, x_n] = k[x_1, \dots, x_n, y_1, \dots, y_m].$$

Here we regard k as a subring of $k[y_1, \dots, y_m]$. Similar treatment to S_W .

(3) By construction, $V \times W \subseteq V(S)$. Conversely, given any $(P, Q) \in V(S) \subseteq \mathbf{A}^{n+m}(k)$, we have $h(P, Q) = 0$ for all $h \in S = S_V \cup S_W$ (by (2)). By construction, $f(P) = 0$ for all $f \in S_V$ since f only involve x_1, \dots, x_n . Hence, $P \in V$. Similarly, $Q \in W$. Therefore, $(P, Q) \in V \times W$.

□

1.3. The Ideal of a Set of Points

Problem 1.16.*

Let V, W be algebraic sets in $\mathbf{A}^n(k)$. Show that $V = W$ if and only if $I(V) = I(W)$.

Proof.

(1) (Proof of Equation (6) in this section.) Show that if $X \subseteq Y$, then $I(X) \supseteq I(Y)$. If $f \in I(Y)$ then $f(P) = 0$ for all $P \in Y$. So $f(P) = 0$ for all $P \in X \subseteq Y$ or $f \in I(X)$.

- (2) (Proof of Equation (8) in this section.) $I(V(S)) \supseteq S$ for any set S of polynomials; $V(I(X)) \supseteq X$ for any set X of points.
- (a) If $f \in S$ then f vanishes on $V(S)$, hence $f \in I(V(S))$.
 - (b) If $P \in X$ then every polynomial in $I(X)$ vanishes at P , so P belongs to the zero set of $I(X)$.
- (3) (Proof of Equation (9) in this section.) $V(I(V(S))) = V(S)$ for any set S of polynomials, and $I(V(I(X))) = I(X)$ for any set X of points. So if V is an algebraic set, $V = V(I(V))$, and if I is the ideal of an algebraic set, $I = I(V(I))$.
- (a) In each case, it suffices to show that the left side is a subset of the right side. (by Equations (6)(8) in this section).
 - (b) If $P \in V(S)$ then $f(P) = 0$ for all $f \in I(V(S))$, so $P \in V(I(V(S)))$.
 - (c) If $f \in I(X)$ then $f(P) = 0$ for all $P \in V(I(X))$. Thus f vanishes on $V(I(X))$, so $f \in I(V(I(X)))$.
- (4) Show that $V = W$ if and only if $I(V) = I(W)$.
- (a) By Equation (6) in this section, $I(V) \supseteq I(W)$ if $V \subseteq W$ and $I(V) \subseteq I(W)$ if $V \supseteq W$. Thus, $I(V) = I(W)$ if $V = W$.
 - (b) Conversely, $I(V) = I(W)$ implies that $V(I(V)) = V(I(W))$ by Equation (3) in the previous section and similar argument in (a). By Equation (9) in this section, $V(I(V)) = V$ and $V(I(W)) = W$. Thus, $V = W$.

□

Problem 1.17.*

- (a) Let V be an algebraic set in $\mathbf{A}^n(k)$, $P \in \mathbf{A}^n(k)$ a point not in V . Show that there is a polynomial $f \in k[x_1, \dots, x_n]$ such that $f(Q) = 0$ for all $Q \in V$, but $f(P) = 1$. (Hint: $I(V) \neq I(V \cup \{P\})$.)
- (b) Let P_1, \dots, P_r be distinct points in $\mathbf{A}^n(k)$, not in an algebraic set V . Show that there are polynomials $f_1, \dots, f_r \in I(V)$ such that $f_i(P_j) = 0$ if $i \neq j$, and $f_i(P_i) = 1$. (Hint: Apply (a) to the union of V and all but one point.)
- (c) With P_1, \dots, P_r and V as in (b), and $a_{ij} \in k$ for $1 \leq i, j \leq r$, show that there are $g_i \in I(V)$ with $g_i(P_j) = a_{ij}$ for all i and j . (Hint: Consider $\sum_j a_{ij} f_j$.)

Proof of (a).

- (1) Since $I(V) \supsetneq I(V \cup \{P\})$ (by Problem 1.16), there is a polynomial $f \in k[x_1, \dots, x_n]$ such that $f(Q) = 0$ for all $Q \in V$, but $f(P) \neq 0$.
- (2) Since k is a field, $(f(P))^{-1} \in k$. Consider the polynomial $(f(P))^{-1}f \in k[x_1, \dots, x_n]$. It is well-defined. Also, $((f(P))^{-1}f)(Q) = (f(P))^{-1}f(Q) = 0$ for all $Q \in V$, but $(f(P))^{-1}f(P) = (f(P))^{-1}f(P) = 1$.

□

Proof of (b).

- (1) For $1 \leq i \leq$, define

$$\begin{aligned} W &= V \cup \{P_1, \dots, P_r\} \\ W_i &= V \cup \{P_1, \dots, \widehat{P}_i, \dots, P_r\}. \end{aligned}$$

Here $W = W_i \cup \{P_i\} \neq W_i$.

- (2) By (a), there is a polynomial $f_i \in k[x_1, \dots, x_n]$ such that $f_i(Q) = 0$ for all $Q \in W_i$, but $f_i(P_i) = 1$. Here $f_i \in I(V)$ and $f_i(P_j) = \delta_{ij}$ where δ_{ij} is the Kronecker delta.

□

Proof of (c).

- (1) For each $1 \leq i \leq r$, define

$$g_i = \sum_j a_{ij} f_j \in k[x_1, \dots, x_n].$$

- (2) $g_i \in I(V)$ since g_i is a linear combination of f_j and $I(V)$ is an ideal.

- (3) Also,

$$g_i(P_j) = \sum_{j'} a_{ij'} f_{j'}(P_j) = \sum_{j'} a_{ij'} \delta_{j'j} = a_{ij}.$$

□

Problem 1.18.*

Let I be an ideal in a ring R . If $a^n \in I$, $b^m \in I$, show that $(a+b)^{n+m} \in I$. Show that $\text{rad}(I)$ is an ideal, in fact a radical ideal. Show that any prime ideal is radical.

Proof.

- (1) Show that $(a + b)^{n+m} \in I$ if $a^n \in I$, $b^m \in I$. By the binomial theorem,

$$(a + b)^{n+m} = \sum_{i=0}^{n+m} a^i b^{n+m-i}.$$

For each term $a^i b^{n+m-i}$, either $i \geq n$ holds or $n + m - i \geq m$ holds, and thus $a^i b^{n+m-i} \in I$ (since $a^n \in I$, $b^m \in I$ and I is an ideal). Hence, the result is established.

- (2) Show that $\text{rad}(I)$ is an ideal.

(a) $0 \in \text{rad}(I)$ since $0 = 0^1 \in I$ for any ideal in R .

(b) $(a + b)^{n+m} \in I$ if $a^n \in I$, $b^m \in I$ by (1).

(c) $(-a)^{2n} = (a^n)^2 \in I$ if $a^n \in I$ (since I is an ideal).

(d) $(ra)^n = r^n a^n \in I$ if $a^n \in I$ and $r \in R$ (since I is an ideal and R is commutative).

- (3) Show that $\text{rad}(\text{rad}(I)) = \text{rad}(I)$. It suffices to show $\text{rad}(\text{rad}(I)) \subseteq \text{rad}(I)$. Given any $a \in \text{rad}(\text{rad}(I))$. By definition $a^n \in \text{rad}(I)$ for some positive integer n . Again by definition $(a^n)^m = a^{nm} \in I$ for some positive integer m . As nm is a positive integer, $a \in \text{rad}(I)$.

- (4) Show that every prime ideal \mathfrak{p} is radical. Given any $a \in \text{rad}(\mathfrak{p})$, that is, $a^n \in \mathfrak{p}$ for some positive integer. Write $a^n = aa^{n-1}$ if $n > 1$. By the primality of \mathfrak{p} , $a \in \mathfrak{p}$ or $a^{n-1} \in \mathfrak{p}$. If $a \in \mathfrak{p}$, we are done. If $a^{n-1} \in \mathfrak{p}$, we continue this descending argument (or the mathematical induction) until the power of a is equal to 1. Hence \mathfrak{p} is radical.

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

- (1) PLACEHOLDER

□

Problem 1.20.*

Show that for any ideal I in $k[x_1, \dots, x_n]$, $V(I) = V(\text{rad}(I))$, and $\text{rad}(I) \subseteq I(V(I))$.

Proof.

- (1) Show that $V(I) = V(\text{rad}(I))$. Since $I \subseteq \text{rad}(I)$, it suffices to show that $V(I) \subseteq V(\text{rad}(I))$. Given any $P \in V(I)$. For any $f \in \text{rad}(I)$, $f^n \in I$ for some positive integer $n > 0$. Note that

$$0 = (f^n)(P) = f(P)^n$$

since $f^n \in I$ and $P \in V(I)$. As k is a domain, $f(P)^n = 0$ implies $f(P) = 0$. So $P \in V(\text{rad}(I))$.

- (2) By Equations (6) and (8) in this section,

$$I(V(I)) = I(V(\text{rad}(I))) \supseteq \text{rad}(I).$$

□

Note.

- (1) By the Hilbert's Nullstellensatz, $I(V(I)) = \text{rad}(I)$ if $k = \bar{k}$.
 (2) Take $I = (x^2 + 1)$ as an ideal in $\mathbb{R}[x]$. Note that $I(V(I)) = I(\emptyset) = (1)$ and $\text{rad}(I) = I = (x^2 + 1)$. So the equality in $\text{rad}(I) \subseteq I(V(I))$ might not hold if $k \neq \bar{k}$.

Problem 1.21.*

Show that $I = (x_1 - a_1, \dots, x_n - a_n) \subseteq k[x_1, \dots, x_n]$ is a maximal ideal, and that the natural homomorphism from k to $k[x_1, \dots, x_n]/I$ is an isomorphism.

Proof.

- (1) Show that I is a maximal ideal. Suppose that J is an ideal such that $J \supsetneq I$. Take any $f \in J - I$. By Problem 1.7(a),

$$f = \sum \lambda_{(i)} (x_1 - a_1)^{i_1} \cdots (x_n - a_n)^{i_n}.$$

As $f \notin I$, there is a nonzero constant term in f , say $\lambda \in k - \{0\}$. Note that $f - \lambda \in I \subseteq J$. Hence,

$$\lambda = f - (f - \lambda) \in J$$

since J is an ideal. As $\lambda \neq 0$, $J = k[x_1, \dots, x_n]$ is not a proper ideal containing I .

- (2) Let $\varphi : k \rightarrow k[x_1, \dots, x_n]/I$ be the natural homomorphism. (That is, $\varphi : \lambda \rightarrow \lambda + I \in k[x_1, \dots, x_n]/I$.)
- (3) Show that φ is surjective. Given any $f + I \in k[x_1, \dots, x_n]/I$. By Problem 1.7(a),

$$f = \sum \lambda_{(i)} (x_1 - a_1)^{i_1} \cdots (x_n - a_n)^{i_n}.$$

So

$$\begin{aligned} f + I &= \sum \lambda_{(i)} (x_1 - a_1)^{i_1} \cdots (x_n - a_n)^{i_n} + I \\ &= \left(f(a_1, \dots, a_n) + \sum_{\text{nonconstant}} \lambda_{(i)} (x_1 - a_1)^{i_1} \cdots (x_n - a_n)^{i_n} \right) + I \\ &= f(a_1, \dots, a_n) + I. \end{aligned}$$

(Here the summation over all nonconstant terms is in I .) Hence

$$\varphi : f(a_1, \dots, a_n) \in k \mapsto f + I.$$

- (4) Show that φ is injective. $\ker(\varphi) = \{\lambda \in k : \lambda \in I\} = k \cap I = \{0\}$ since I is a proper ideal.
- (5) By (2)(3)(4), $\varphi : k \rightarrow k[x_1, \dots, x_n]/(x_1 - a_1, \dots, x_n - a_n)$ is an isomorphism.

□

1.4. The Hilbert Basis Theorem

Problem PLACEHOLDER

PLACEHOLDER

Proof.

- (1) PLACEHOLDER

□

1.5. Irreducible Components of an Algebraic Set

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

1.6. Algebraic Subsets of the Plane

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

1.7. Hilbert's Nullstellensatz

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

- (1) PLACEHOLDER

□

1.8. Modules; Finiteness Conditions

Problem 1.41.*

If S is module-finite over R , then S is ring-finite over R .

Proof.

- (1) $S = \sum Rs_i$ for some $s_1, \dots, s_n \in S$ since S is module-finite over R .
(2) Let I be the minimal subset of $\{s_1, \dots, s_n\}$ which also spans S , say $\{t_1, \dots, t_m\}$ with $m \leq n$. Clearly we can write

$$S = R[t_1, \dots, t_m],$$

that is, S is ring-finite over R .

- (3) The converse is not true (Problem 1.42).

□

Problem 1.42.

Show that $S = R[x]$ (the ring of polynomials in one variable) is ring-finite over R , but not module-finite.

Proof.

- (1) $S = R[x]$ is ring-finite over R by definition (as $x \in S$).
(2) (Reductio ad absurdum) If $S = \sum Rs_i$ for some $s_1, \dots, s_n \in S$ were module-finite over R . Any element $s \in \sum Rs_i$ is of degree

$$\deg s \leq \max_{1 \leq i \leq n} \deg s_i := m.$$

So that $x^{m+1} \in S = R[x]$ but not in $\sum Rs_i$, which is absurd.

□

Problem 1.43.* (WIP)

If L is ring-finite over K (K, L fields) then L is a finitely generated field extension of K .

Proof.

- (1) $L = K[v_1, \dots, v_n]$ for some $v_i \in L$. To show $L = K[v_1, \dots, v_n] = K(v_1, \dots, v_n)$, it suffices to show that all v_i are algebraic over L .

- (2)

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

- (1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

- (1) PLACEHOLDER

□

1.9. Integral Elements

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

1.10. Field Extensions

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

Chapter 2: Affine Varieties

2.1. Coordinate Rings

Problem 2.1.*

Show that the map which associates to each $f \in k[x_1, \dots, x_n]$ a polynomial function in $\mathcal{F}(V, k)$ is a ring homomorphism whose kernel is $I(V)$.

Proof.

- (1) Define a map $\alpha : k[x_1, \dots, x_n] \rightarrow \mathcal{F}(V, k)$. Every polynomial $f \in k[x_1, \dots, x_n]$ defines a function from V to k by

$$\alpha(f)(a_1, \dots, a_n) = f(a_1, \dots, a_n)$$

for all $(a_1, \dots, a_n) \in V$.

- (2) α is a ring homomorphism by construction in (1).
(3) Show that $\ker(\alpha) = I(V)$. In fact, given any $f \in k[x_1, \dots, x_n]$, we have $\alpha(f) = 0$ (sending all $a \in V$ to $0 \in k$) if and only if $f(a) = 0$ for all $a \in V$ if and only if $f \in I(V)$.
(4) Hence $k[x_1, \dots, x_n]/I(V) = \Gamma(V) \hookrightarrow \mathcal{F}(V, k)$ is an injective homomorphism.

□

Problem PLACEHOLDER

PLACEHOLDER

Proof.

- (1) PLACEHOLDER

2.2. Polynomial Maps

2.3. Coordinate Changes

2.4. Rational Functions and Local Rings

2.5. Discrete Valuation Rings

2.6. Forms

2.7. Direct Products of Rings

2.8. Operations with Ideals

Problem 2.39.*

Prove the following relations among ideals I_i , J in a ring R :

(a) $(I_1 + I_2)J = I_1J + I_2J$.

(b) $(I_1 \cdots I_N)^n = I_1^n \cdots I_N^n$.

Proof of (a).

(1) Note that $(I_1 + I_2)J$ and $I_1J + I_2J$ are ideals.

(2) Show that $(I_1 + I_2)J \subseteq I_1J + I_2J$. Given any

$$(x_1 + x_2)y \in (I_1 + I_2)J$$

where $x_i \in I_i$ and $y \in J$. It suffices to show that $(x_1 + x_2)y \in I_1J + I_2J$ (by (1)). In fact,

$$(x_1 + x_2)y = x_1y + x_2y \in I_1J + I_2J.$$

(3) Show that $(I_1 + I_2)J \supseteq I_1J + I_2J$. Given any

$$x_1y_1 + x_2y_2 \in I_1J + I_2J$$

where $x_i \in I_i$ and $y_i \in J$. It suffices to show that $x_1y_1 + x_2y_2 \in (I_1 + I_2)J$ (by (1)). In fact,

$$x_1y_1 + x_2y_2 = (x_1 + \underbrace{0}_{\in I_2})y_1 + (\underbrace{0}_{\in I_1} + x_2)y_2 \in (I_1 + I_2)J$$

since $(I_1 + I_2)J$ is an ideal.

□

Proof of (b).

- (1) Note that $(I_1 \cdots I_N)^n$ and $I_1^n \cdots I_N^n$ are ideals.
(2) Show that $(I_1 \cdots I_N)^n \subseteq I_1^n \cdots I_N^n$. Given any

$$x = x_1 \cdots x_n$$

where $x_i \in I_1 \cdots I_N$. It suffices to show that $x \in I_1^n \cdots I_N^n$ (by (1)). For each $x_i \in I_1 \cdots I_N$, write

$$x_i = \sum_{j(i)} x_{j(i),1} \cdots x_{j(i),N}$$

where $x_{j(i),k} \in I_k$ for $1 \leq k \leq N$. Hence

$$\begin{aligned} x &= x_1 \cdots x_n \\ &= \left(\sum_{j(1)} x_{j(1),1} \cdots x_{j(1),N} \right) \cdots \left(\sum_{j(n)} x_{j(n),1} \cdots x_{j(n),N} \right) \\ &= \sum_{j(1), \dots, j(n)} (x_{j(1),1} \cdots x_{j(1),N}) \cdots (x_{j(n),1} \cdots x_{j(n),N}) \\ &= \sum_{j(1), \dots, j(n)} \underbrace{(x_{j(1),1} \cdots x_{j(n),1})}_{\in I_1^n} \cdots \underbrace{(x_{j(1),N} \cdots x_{j(n),N})}_{\in I_N^n} \\ &\in I_1^n \cdots I_N^n. \end{aligned}$$

- (3) Show that $(I_1 \cdots I_N)^n \supseteq I_1^n \cdots I_N^n$. Given any

$$x = x_1 \cdots x_N \in I_1^n \cdots I_N^n$$

where $x_i \in I_i^n$ ($1 \leq i \leq N$). It suffices to show that $x \in (I_1 \cdots I_N)^n$ (by (1)). For each $x_i \in I_i^n$, write

$$x_i = \sum_{j(i)} x_{j(i),1} \cdots x_{j(i),n}$$

where $x_{j(i),k} \in I_i$ for $1 \leq k \leq n$. Hence

$$\begin{aligned}
x &= x_1 \cdots x_N \\
&= \left(\sum_{j(1)} x_{j(1),1} \cdots x_{j(1),n} \right) \cdots \left(\sum_{j(N)} x_{j(N),1} \cdots x_{j(N),n} \right) \\
&= \sum_{j(1), \dots, j(N)} (x_{j(1),1} \cdots x_{j(1),n}) \cdots (x_{j(N),1} \cdots x_{j(N),n}) \\
&= \sum_{j(1), \dots, j(N)} \underbrace{(x_{j(1),1} \cdots x_{j(N),1})}_{\in I_1 \cdots I_N} \cdots \underbrace{(x_{j(1),n} \cdots x_{j(N),n})}_{\in I_1 \cdots I_N} \\
&\in (I_1 \cdots I_N)^n.
\end{aligned}$$

□

Problem 2.41.*

Let I, J be ideals in R . Suppose I is finitely generated and $I \subseteq \text{rad}(J)$. Show that $I^n \subseteq J$ for some n .

Proof.

- (1) Let I be generated by $x_1, \dots, x_m \in I$. As $I \subseteq \text{rad}(J)$, there are integers $n_i > 0$ such that $x_i^{n_i} \in J$.
- (2) Let $N = n_1 + \cdots + n_m$. Given any $x = \sum_{i=1}^m r_i x_i \in I$, so

$$\begin{aligned}
x^N &= \left(\sum_{i=1}^m r_i x_i \right)^N \\
&= \sum_{k_1 + \cdots + k_m = N} \binom{N}{k_1, \dots, k_m} r_1^{k_1} x_1^{k_1} \cdots r_m^{k_m} x_m^{k_m}.
\end{aligned}$$

- (3) Note that for each term there is some j such that $k_j \geq n_j$. Hence,

$$\begin{aligned}
x_j^{k_j} &= x_j^{k_j - n_j} x_j^{n_j} \in J && (J \text{ is an ideal}) \\
\Rightarrow r_1^{k_1} x_1^{k_1} \cdots r_m^{k_m} x_m^{k_m} &\in J \text{ for each term} && (J \text{ is an ideal}) \\
\Rightarrow x^N &\in J. && (J \text{ is an ideal}) \\
\Rightarrow I^N &\subseteq J.
\end{aligned}$$

□

Supplement. (Exercise 1.13 in the textbook: Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry*.) Suppose that I is an ideal in a commutative ring. Show that if $\text{rad}(I)$ is finitely generated, then for some integer N we have $(\text{rad}(I))^N \subseteq I$. Conclude that in a Noetherian ring the ideals I and J have the same radical iff there is some integer N such that $I^N \subseteq J$ and $J^N \subseteq I$. Use the Nullstellensatz to deduce that if $I, J \subseteq S = k[x_1, \dots, x_n]$ are ideals and k is algebraically closed, then $Z(I) = Z(J)$ iff $I^N \subseteq J$ and $J^N \subseteq I$ for some N .

Proof.

- (1) Show that if $\text{rad}(I)$ is finitely generated, then for some integer N we have $(\text{rad}(I))^N \subseteq I$. Say $x_1, \dots, x_m \in \text{rad}(I)$ generate $\text{rad}(I)$.

(a) For each i , there exists an integer $n_i > 0$ such that $x_i^{n_i} \in I$ (since $\text{rad}(I)$ is radical).

(b) Let $N = n_1 + \dots + n_m$. Given any $x = \sum_{i=1}^m r_i x_i \in \text{rad}(I)$, so

$$\begin{aligned} x^N &= \left(\sum_{i=1}^m r_i x_i \right)^N \\ &= \sum_{k_1 + \dots + k_m = N} \binom{N}{k_1, \dots, k_m} r_1^{k_1} x_1^{k_1} \dots r_m^{k_m} x_m^{k_m}. \end{aligned}$$

(c) Note that for each term there is some j such that $k_j \geq n_j$. Hence,

$$\begin{aligned} x_j^{k_j} &= x_j^{k_j - n_j} x_j^{n_j} \in I && (I \text{ is an ideal}) \\ \implies r_1^{k_1} x_1^{k_1} \dots r_m^{k_m} x_m^{k_m} &\in I \text{ for each term} && (I \text{ is an ideal}) \\ \implies x^N &\in I. && (I \text{ is an ideal}) \\ \implies (\text{rad}(I))^N &\subseteq I. \end{aligned}$$

- (2) Show that in a Noetherian ring the ideals I and J have the same radical iff there is some integer N such that $I^N \subseteq J$ and $J^N \subseteq I$.

(a) (\implies) Since in a Noetherian ring every ideal is finitely generated, $\text{rad}(I)$ and $\text{rad}(J)$ are finitely generated. By (1), there is a common integer N such that

$$(\text{rad}(I))^N \subseteq I \quad \text{and} \quad (\text{rad}(J))^N \subseteq J.$$

Note that $I^N \subseteq (\text{rad}(I))^N$ and $J^N \subseteq (\text{rad}(J))^N$. Since $\text{rad}(I) = \text{rad}(J)$ by assumption,

$$\begin{aligned} I^N &\subseteq (\text{rad}(I))^N = (\text{rad}(J))^N \subseteq J, \\ J^N &\subseteq (\text{rad}(J))^N = (\text{rad}(I))^N \subseteq I. \end{aligned}$$

- (b) (\Longleftarrow) It suffices to show that $\text{rad}(I) \subseteq \text{rad}(J)$. $\text{rad}(J) \subseteq \text{rad}(I)$ is similar. Given any $x \in \text{rad}(I)$, there is an integer $M > 0$ such that $x^M \in I$. Hence $x^{MN} \in I^N \subseteq J$, or $x \in \text{rad}(J)$.
- (3) Show that if $I, J \subseteq S = k[x_1, \dots, x_n]$ are ideals and k is algebraically closed, then $Z(I) = Z(J)$ iff $I^N \subseteq J$ and $J^N \subseteq I$ for some N . Note that S is Noetherian and we can apply part (2). By the Nullstellensatz, $Z(I) = Z(J)$ iff $\text{rad}(I) = \text{rad}(J)$ iff $I^N \subseteq J$ and $J^N \subseteq I$ for some N .

□

2.9. Ideals with a Finite Number of Zeros

2.10. Quotient Modules and Exact Sequences

Problem 2.51.

Let

$$0 \longrightarrow V_1 \longrightarrow \cdots \longrightarrow V_n \longrightarrow 0$$

be an exact sequence of finite-dimensional vector spaces. Show that $\sum (-1)^i \dim(V_i) = 0$.

Proof (Proposition 7 in this section).

- (1) For $i = 0, \dots, n$, by the rank-nullity theorem for a linear transformation $\varphi_i : V_i \rightarrow V_{i+1}$, we have

$$\dim V_i = \dim \text{im}(\varphi_i) + \dim \ker(\varphi_i).$$

(Here $V_0 = V_{n+1} := 0$ by convention.)

- (2) By the exactness of the sequence, we have

- (a) $\text{im}(\varphi_i) = \ker(\varphi_{i+1})$ for $i = 0, \dots, n-1$. In particular, $\ker(\varphi_1) = \text{im}(\varphi_0) = 0$.
- (b) $\ker(\varphi_n) = V_n$.

Hence,

$$\begin{aligned}
\sum_{i=1}^{n-1} (-1)^i \dim(V_i) &= \sum_{i=1}^{n-1} (-1)^i \dim \operatorname{im}(\varphi_i) + \sum_{i=1}^{n-1} (-1)^i \dim \operatorname{ker}(\varphi_i) \\
&= \sum_{i=1}^{n-1} (-1)^i \dim \operatorname{ker}(\varphi_{i+1}) + \sum_{i=1}^{n-1} (-1)^i \dim \operatorname{ker}(\varphi_i) \\
&= (-1)^{n-1} \underbrace{\dim \operatorname{ker}(\varphi_n)}_{=V_n} + (-1)^1 \underbrace{\dim \operatorname{ker}(\varphi_1)}_{=0} \\
&= -(-1)^n \dim V_n,
\end{aligned}$$

or $\sum (-1)^i \dim(V_i) = 0$.

□

2.11. Free Modules

Chapter 3: Local Properties of Plane Curves

3.1. Multiple Points and Tangent Lines

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

3.2. Multiplicities and Local Rings

3.3. Intersection Numbers

Chapter 4: Projective Varieties

4.1. Projective Space

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

4.2. Projective Algebraic Sets

4.3. Affine and Projective Varieties

4.4. Multiprojective Space

Chapter 5: Projective Plane Curves

5.1. Definitions

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

5.2. Linear Systems of Curves

5.3. Bézout's Theorem

5.4. Multiple Points

5.5. Max Noether's Fundamental Theorem

5.6. Applications of Noether's Theorem

Chapter 6: Varieties, Morphisms, and Rational Maps

6.1. The Zariski Topology

6.2. Varieties

6.3. Morphisms of Varieties

6.4. Products and Graphs

6.5. Algebraic Function Fields and Dimension of Varieties

6.6. Rational Maps

Chapter 7: Resolution of Singularities

7.1. Rational Maps of Curves

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

7.2. Blowing up a Point in A^2

7.3. Blowing up a Point in P^2

7.4. Quadratic Transformations

7.5. Nonsingular Models of Curves

Chapter 8: Riemann-Roch Theorem

8.1. Divisors

Problem PLACEHOLDER

PLACEHOLDER

Proof.

(1) PLACEHOLDER

□

8.2. The Vector Spaces $L(D)$

8.3. Riemann's Theorem

8.4. Derivations and Differentials

8.5. Canonical Divisors

8.6. Riemann-Roch Theorem