

Chapter 11: The Lebesgue Theory

Author: Meng-Gen Tsai

Email: plover@gmail.com

Exercise 11.1. If $f \geq 0$ and $\int_E f d\mu = 0$, prove that $f(x) = 0$ almost everywhere on E . (Hint: Let E_n be the subset of E on which $f(x) > \frac{1}{n}$. Write $A = \bigcup E_n$. Then $\mu(A) = 0$ if and only if $\mu(E_n) = 0$ for every n .)

Proof (Hint).

- (1) Define $A = \{x \in E : f(x) > 0\}$. So $f(x) = 0$ almost everywhere on E if and only if $\mu(A) = 0$.

- (2) Define

$$E_n = \left\{ x \in E : f(x) > \frac{1}{n} \right\}$$

for $n = 1, 2, 3, \dots$. Note that $E_1 \subseteq E_2 \subseteq E_3 \subseteq \dots$ and

$$A = \bigcup_{n=1}^{\infty} E_n.$$

Since μ is a measure,

$$\lim_{n \rightarrow \infty} \mu(E_n) = \mu(A)$$

(Theorem 11.3).

- (3) (Reductio ad absurdum) If $\mu(A) > 0$, there is an integer N such that $\mu(E_n) \geq \frac{\mu(A)}{2}$ whenever $n \geq N$ (by (2)). In particular, take $n = N$ to get

$$\begin{aligned} \int_E f d\mu &\geq \int_{E_N} f d\mu && (\mu \text{ is a measure and } E_N \subseteq E) \\ &\geq \frac{1}{N} \cdot \mu(E_N) && (\text{Remarks 11.23(b)}) \\ &\geq \frac{1}{N} \cdot \frac{\mu(A)}{2} \\ &> 0, \end{aligned}$$

contrary to the assumption that $\int_E f d\mu = 0$.

□

Note. Compare to Exercise 6.2.

Exercise 11.2. ...

Proof.

(1)

(2)

□

Exercise 11.3. ...

Proof.

(1)

(2)

□

Exercise 11.4. ...

Proof.

(1)

(2)

□

Exercise 11.5. ...

Proof.

(1)

(2)

□

Exercise 11.6. ...

Proof.

(1)

(2)

□

Exercise 11.7. ...

Proof.

(1)

(2)

□

Exercise 11.8. ...

Proof.

(1)

(2)

□

Exercise 11.9. ...

Proof.

(1)

(2)

□

Exercise 11.10. ...

Proof.

(1)

(2)

□

Exercise 11.11. ...

Proof.

(1)

(2)

□

Exercise 11.12. ...

Proof.

(1)

(2)

□

Exercise 11.13. ...

Proof.

(1)

(2)

□

Exercise 11.14. ...

Proof.

(1)

(2)

□

Exercise 11.15. ...

Proof.

(1)

(2)

□

Exercise 11.16. ...

Proof.

(1)

(2)

□

Exercise 11.17. ...

Proof.

(1)

(2)

□

Exercise 11.18. ...

Proof.

(1)

(2)

□