

# Chapter 1: Curves

*Author: Meng-Gen Tsai*

*Email: plover@gmail.com*

## Section 1-1: Introduction

Classical differential geometry: the study of local properties of curves and surfaces.

Global differential geometry: the study of the influence of the local properties on the behavior of the entire curve and surface.

*No exercises.*

## Section 1-2: Parametrized Curves

**Exercise 1-2.1.** Find a parametrized curve  $\alpha(t)$  whose trace is the circle  $x^2 + y^2 = 1$  such that  $\alpha(t)$  runs clockwise around the circle with  $\alpha(0) = (0, 1)$ .

*Proof.*  $\alpha(t) = (\sin t, \cos t)$ ,  $t \in \mathbb{R}$ .  $\square$

**Exercise 1-2.2.** Let  $\alpha(t)$  be a parametrized curve which does not pass through the origin. If  $\alpha(t_0)$  is the point of the trace of  $\alpha$  closest to the origin and  $\alpha'(t_0) \neq 0$ , show that the position vector  $\alpha(t_0)$  is orthogonal to  $\alpha'(t_0)$ .

*Proof.* Let  $f(t) = |\alpha(t)|^2 = \alpha(t) \cdot \alpha(t)$ .  $f(t)$  is differentiable and  $f(t)$  has a local minimum at a point  $t = t_0 \in I$ . So  $f'(t_0) = 0$ . [Theorem 5.8 in W. Rudin, Principles of Mathematical Analysis, 3rd edition.] Since

$$f'(t) = 2\alpha(t) \cdot \alpha'(t),$$

$f'(t_0) = 2\alpha(t_0) \cdot \alpha'(t_0) = 0$ , or  $\alpha(t_0) \cdot \alpha'(t_0) = 0$ . Since  $\alpha(t_0) \neq 0$  and  $\alpha'(t_0) \neq 0$ ,  $\alpha(t_0)$  is orthogonal to  $\alpha'(t_0)$ .  $\square$

**Exercise 1-2.3.** A parametrized curve  $\alpha(t)$  has a property that its second derivative  $\alpha''(t)$  is identically zero. What can be said about  $\alpha$ ?

$\alpha(t)$  is a straight line.

*Proof.* Since  $\alpha''(t)$  is identically zero,  $\alpha'(t) = a$  is a constant. [Theorem 5.11 in W. Rudin, Principles of Mathematical Analysis, 3rd edition.] Define

$f(t) = \alpha(t) - at$  (on  $I$ ). Since  $f'(t) = \alpha'(t) - a = 0$ ,  $f(t) = \alpha(t) - at = b$  is a constant again. Therefore,  $\alpha(t) = at + b$ , which is a straight line (on  $I$ ).  $\square$

**Exercise 1-2.4.** Let  $\alpha : I \rightarrow \mathbb{R}^3$  be a parametrized curve and let  $v \in \mathbb{R}^3$  be a fixed vector. Assume that  $\alpha'(t)$  is orthogonal to  $v$  for all  $t \in I$  and that  $\alpha(0)$  is orthogonal to  $v$ . Prove that  $\alpha(t)$  is orthogonal to  $v$  for all  $t \in I$ .

Need to assume that  $\alpha(t) \neq 0$  for all  $t \in I$ .

*Proof.* Given any  $t \neq 0 \in I$ . (Nothing to do at  $t = 0$ .) Define  $f : I \rightarrow \mathbb{R}$  by  $f(t) = \alpha(t) \cdot v$ . By the mean value theorem, there exists a point  $\xi$  between 0 and  $t$  such that

$$f(t) - f(0) = f'(\xi)(t - 0),$$

where  $f'(t) = \alpha'(t) \cdot v + \alpha(t) \cdot v' = \alpha'(t) \cdot v$ . Note that  $f(0) = 0$  since  $\alpha(0)$  is orthogonal to  $v$ , and  $f'(\xi) = 0$  since  $\alpha'(\xi)$  is orthogonal to  $v$ . So the identity is reduced to

$$f(t) = 0,$$

or  $\alpha(t) \cdot v = 0$ , or  $\alpha(t)$  is orthogonal to  $v$ .  $\square$

**Exercise 1-2.5.** Let  $\alpha : I \rightarrow \mathbb{R}^3$  be a parametrized curve, with  $\alpha'(t) \neq 0$  for all  $t \in I$ . Show that  $|\alpha(t)|$  is a nonzero constant if and only if  $\alpha(t)$  is orthogonal to  $\alpha'(t)$  for all  $t \in I$ .

The same trick in Exercise 1-2.2.

*Proof.* It is equivalent to show that  $|\alpha(t)|^2$  is a nonzero constant if and only if  $\alpha(t)$  is orthogonal to  $\alpha'(t)$  for all  $t \in I$ . Let

$$f(t) = |\alpha(t)|^2 = \alpha(t) \cdot \alpha(t).$$

Notice that  $\alpha'(t) \neq 0$ , and thus

$$\begin{aligned} & |\alpha(t)| \text{ is a nonzero constant} \\ \iff & f(t) = |\alpha(t)|^2 \text{ is a nonzero constant} \\ \iff & f'(t) = 0 \text{ and } f(t) \text{ is a nonzero constant} \\ \iff & \alpha(t) \cdot \alpha'(t) = 0 \text{ and } \alpha(t) \text{ is a nonzero constant} \\ \iff & \alpha(t) \text{ is orthogonal to } \alpha'(t) \text{ for all } t \in I. \end{aligned}$$

$\square$

## Section 1-3: Regular Curves; Arc Length

**Exercise 1-3.2.** A circular disk of radius 1 in the plane  $xy$  rolls without slipping along the  $x$  axis. The figure described by a point of the circumference of the disk is called a **cycloid** (Figure 1-7 in Mantredo P. do Carmo, *Differential Geometry of Curves and Surfaces*).

- (a) Obtain a parametrized curve  $\alpha : \mathbb{R} \rightarrow \mathbb{R}^2$  the trace of which is the cycloid and determine its singular points.
- (b) Compute the arc length of the cycloid corresponding to a complete rotation of the disk.

*Proof of (a).*

- (1) Since

$$\begin{cases} x = t - \sin t \\ y = 1 - \cos t, \end{cases}$$

we define  $\alpha(t) = (t - \sin t, 1 - \cos t)$ .

- (2)  $\alpha'(t) = (1 - \cos t, \sin t)$ .  $\alpha'(t) = 0$  if and only if  $t = 2n\pi$  where  $n \in \mathbb{Z}$ . That is, all singular points are  $\alpha(2n\pi) = (2n\pi, 0)$  where  $n \in \mathbb{Z}$ .

□

*Proof of (b).* The arc length of the cycloid corresponding to a complete rotation of the disk is

$$\begin{aligned} \int_0^{2\pi} |\alpha'(t)| dt &= \int_0^{2\pi} \sqrt{(1 - \cos t)^2 + (\sin t)^2} dt \\ &= \int_0^{2\pi} \sqrt{2} \sqrt{1 - \cos t} dt \\ &= \int_0^{2\pi} 2 \sin \frac{t}{2} dt \\ &= \left[ -4 \cos \frac{t}{2} \right]_{t=0}^{t=2\pi} \\ &= 8. \end{aligned}$$

□

**Supplement.** The cycloid is not an algebraic curve.

**Exercise 1-3.4.** Let  $\alpha : (0, \pi) \rightarrow \mathbb{R}^2$  be given by

$$\alpha(t) = \left( \sin t, \cos t + \log \tan \frac{t}{2} \right),$$

where  $t$  is the angle that the  $y$  axis makes with the vector  $\alpha(t)$ . The trace of  $\alpha$  is called the **tractrix**. (Figure 1-9 in Mantredo P. do Carmo, *Differential Geometry of Curves and Surfaces*). Show that

- (a)  $\alpha$  is a differentiable parametrized curve, regular except at  $t = \frac{\pi}{2}$ .
- (b) The length of the segment of the tangent of the tractrix between the point of tangency and the  $y$  axis is constantly equal to 1.

*Proof of (a).*

$$\begin{aligned}\alpha'(t) &= \left( \cos t, -\sin t + \frac{1}{\tan \frac{t}{2}} \frac{1}{\cos^2 \frac{t}{2}} \frac{1}{2} \right) \\ &= \left( \cos t, -\sin t + \frac{1}{2 \sin \frac{t}{2} \cos \frac{t}{2}} \right) \\ &= \left( \cos t, \frac{\cos^2 t}{\sin t} \right)\end{aligned}$$

exists. And  $\alpha'(t) = 0$  if and only if  $t = \frac{\pi}{2}$ . That is, there is an unique singular point at  $t = \frac{\pi}{2}$ .  $\square$

*Proof of (b).* The the tangent line of the tractrix through the regular point  $t$  is parametrized by  $\beta : \mathbb{R} \rightarrow \mathbb{R}^2$  which is defined by

$$\begin{aligned}\beta(u) &= u\alpha'(t) + \alpha(t) \\ &= \left( u \cos t + \sin t, u \frac{\cos^2 t}{\sin t} + \cos t + \log \tan \frac{t}{2} \right).\end{aligned}$$

By construction, this tangent line  $\beta(u)$  meets the tractrix at  $u = 0$ , and meets the  $y$ -axis when  $u \cos t + \sin t = 0$  or  $u = -\tan t$ . So the length of the segment is

$$\begin{aligned}|\beta(0) - \beta(-\tan t)| &= \sqrt{(-\tan t \cos t)^2 + \left(-\tan t \frac{\cos^2 t}{\sin t}\right)^2} \\ &= \sqrt{(\sin t)^2 + (\cos t)^2} \\ &= 1.\end{aligned}$$

$\square$

**Exercise 1-3.10.** (*Straight Lines as Shortest.*) Let  $\alpha : I \rightarrow \mathbb{R}^3$  be a parametrized curve. Let  $[a, b] \subseteq I$  and set  $\alpha(a) = p$ ,  $\alpha(b) = q$ .

- (a) Show that, for any constant vector  $v$ ,  $|v| = 1$ ,

$$(q - p) \cdot v = \int_a^b \alpha'(t) \cdot v dt \leq \int_a^b |\alpha'(t)| dt.$$

(b) Set

$$v = \frac{q-p}{|q-p|}$$

and show that

$$|\alpha(b) - \alpha(a)| \leq \int_a^b |\alpha'(t)| dt;$$

that is, the curve of shortest length from  $\alpha(a)$  to  $\alpha(b)$  is the straight line joining these points.

Assume  $p \neq q$  (otherwise  $v = \frac{q-p}{|q-p|}$  is meaningless).

*Proof of (a).* Let  $f(t) = \alpha(t) \cdot v$  defined on  $I$ . By the fundamental theorem of calculus,

$$\int_a^b f'(t) dt = f(b) - f(a).$$

Since  $f'(t) = \alpha'(t) \cdot v$ ,

$$(\alpha(b) - \alpha(a)) \cdot v = \int_a^b \alpha'(t) \cdot v dt.$$

Therefore,

$$\begin{aligned} (q-p) \cdot v &= \int_a^b \alpha'(t) \cdot v dt \\ &\leq \int_a^b |\alpha'(t) \cdot v| dt \\ &\leq \int_a^b |\alpha'(t)| |v| dt \\ &= \int_a^b |\alpha'(t)| dt. \end{aligned}$$

□

*Proof of (b).*  $|v| = \frac{|q-p|}{|q-p|} = 1$ . So,

$$\begin{aligned} (q-p) \cdot \frac{q-p}{|q-p|} &\leq \int_a^b |\alpha'(t)| dt, \\ |q-p| &\leq \int_a^b |\alpha'(t)| dt. \end{aligned}$$

□

## Section 1-4: The Vector Product in $\mathbb{R}^3$

**Exercise 1-4.13.** Let  $u(t) = (u_1(t), u_2(t), u_3(t))$  and  $v(t) = (v_1(t), v_2(t), v_3(t))$  be differentiable maps from the interval  $(a, b)$  into  $\mathbb{R}^3$ . If the derivatives  $u'(t)$  and  $v'(t)$  satisfy the conditions

$$u'(t) = au(t) + bv(t), v'(t) = cu(t) - av(t),$$

where  $a$ ,  $b$ , and  $c$  are constants, show that  $u(t) \wedge v(t)$  is a constant vector.

*Proof.* Since

$$\begin{aligned} \frac{d}{dt}(u(t) \wedge v(t)) &= u'(t) \wedge v(t) + u(t) \wedge v'(t) \\ &= (au(t) + bv(t)) \wedge v(t) + u(t) \wedge (cu(t) - av(t)) \\ &= au(t) \wedge v(t) + u(t) \wedge (-av(t)) \\ &= a(u(t) \wedge v(t)) + (-a)(u(t) \wedge v(t)) \\ &= (0, 0, 0), \end{aligned}$$

$u(t) \wedge v(t)$  is a constant vector.  $\square$

## Section 1-5: The Local Theory of Curves Parametrized by Arc Length

**Exercise 1-5.2.** Show that the torsion  $\tau$  of  $\alpha$  is given by

$$\tau(s) = -\frac{\alpha'(s) \wedge \alpha''(s) \cdot \alpha'''(s)}{|\kappa(s)|^2}.$$

*Proof.*

- (1) Take inner product  $n(s)$  to the definition of torsion  $\tau(s)n(s) = b'(s)$ , we have

$$\tau(s) = b'(s) \cdot n(s).$$

Since  $b'(s) = t(s) \wedge n'(s)$ , we have to compute  $n'(s)$  first.

- (2) Compute  $n'(s)$ .

$$n'(s) = \frac{d}{ds} \left( \frac{\alpha''(s)}{\kappa(s)} \right) = \frac{\alpha'''(s)}{\kappa(s)} - \frac{\alpha''(s)\kappa'(s)}{\kappa(s)^2}.$$

(3) By (1)(2),

$$\begin{aligned}
\tau(s) &= b'(s) \cdot n(s) \\
&= (t(s) \wedge n'(s)) \cdot n(s) \\
&= \left( \alpha'(s) \wedge \left( \frac{\alpha'''(s)}{\kappa(s)} - \frac{\alpha''(s)\kappa'(s)}{\kappa(s)^2} \right) \right) \cdot \frac{\alpha''(s)}{\kappa(s)} \\
&= \left( \alpha'(s) \wedge \frac{\alpha'''(s)}{\kappa(s)} \right) \cdot \frac{\alpha''(s)}{\kappa(s)} \\
&= \frac{\alpha'(s) \wedge \alpha'''(s) \cdot \alpha''(s)}{|\kappa(s)|^2},
\end{aligned}$$

or

$$\tau(s) = \frac{\alpha'(s) \wedge \alpha'''(s) \cdot \alpha''(s)}{\alpha''(s)^2}.$$

□

## Section 1-6: The Local Canonical Form

## Section 1-7: Global Properties of Plane Curves