

Notes on the book:

Jürgen Neukirch, Algebraic Number Theory

Meng-Gen Tsai
plover@gmail.com

July 23, 2021

Contents

Chapter I: Algebraic Integers	3
I.1. The Gaussian Integers	3
Exercise I.1.1.	3
Exercise I.1.4.	3
Exercise I.1.5.	3
I.2. Integrality	4
Exercise I.2.1.	4
Exercise I.2.2.	4
Exercise I.2.3.	5
Exercise I.2.4.	6
Exercise I.2.7. (Stickelberger's discriminant relation)	7
I.3. Ideals	8
Exercise I.3.4.	8
Exercise I.3.5.	9
Exercise I.3.6.	9
I.4. Lattices	10
Exercise I.4.1.	10
Exercise I.4.2.	10
I.5. Minkowski Theory	11
Exercise I.5.2.	11
Exercise I.5.3. (Minkowski bound)	11
I.6. The Class Number	12
Exercise I.6.3.	12
I.11. Localization	13
Exercise I.11.7. (Nakayama's lemma)	13
I.13. One-dimensional Schemes	13
I.14. Function Fields	14

Chapter II: The Theory of Valuations	15
II.1. The p -adic Numbers	15
Exercise II.1.2.	15
II.2. The p -adic Absolute Value	16
Exercise II.2.1.	16
Chapter VII: Zeta Functions and L-series	17
VII.1. The Riemann Zeta Function	17
Exercise VII.1.4.	17

Chapter I: Algebraic Integers

I.1. The Gaussian Integers

Exercise I.1.1.

$\alpha \in \mathbb{Z}[i]$ is a unit if and only if $N(\alpha) = 1$.

Proof.

- (1) (\implies) Since α is a unit, there is $\beta \in \mathbb{Z}[i]$ such that $\alpha\beta = 1$. So $N(\alpha\beta) = N(1)$, or $N(\alpha)N(\beta) = 1$. Since the image of N is nonnegative integers, $N(\alpha) = 1$.
- (2) (\impliedby) $N(\alpha) = \alpha\bar{\alpha}$, or $1 = \alpha\bar{\alpha}$ since $N(\alpha) = 1$. That is, $\bar{\alpha} \in \mathbb{Z}[i]$ is the inverse of $\alpha \in \mathbb{Z}[i]$. (Or we solve the equation $N(\alpha) = a^2 + b^2 = 1$, and show that all four solutions (± 1 and $\pm i$) are units.)
- (3) Conclusion: a unit $\alpha = a + bi$ of $\mathbb{Z}[i]$ is satisfying the equation $N(\alpha) = a^2 + b^2 = 1$ by (1)(2). That is, the only unit of $\mathbb{Z}[i]$ are ± 1 and $\pm i$.

□

Exercise I.1.4.

Show that the ring $\mathbb{Z}[i]$ cannot be ordered.

Proof. Similar to the fact that i cannot be ordered in \mathbb{C} . Thus i cannot be ordered in $\mathbb{Z}[i]$ either. □

Exercise I.1.5.

Show that the only units of the ring $\mathbb{Z}[\sqrt{-d}] = \mathbb{Z} + \mathbb{Z}\sqrt{-d}$, for any rational integer $d > 1$, are ± 1 .

Proof.

- (1) Define the norm N on $\mathbb{Z}[\sqrt{-d}]$ by

$$N(x + y\sqrt{-d}) = (x + y\sqrt{-d})(x - y\sqrt{-d}) = x^2 + y^2d,$$

i.e., by $N(z) = |z|^2$. It is multiplicative.

(2) Similar to Exercise I.1.1,

$$\begin{aligned} x + y\sqrt{-d} \in \mathbb{Z}[\sqrt{-d}] \text{ is a unit} &\iff N(x + y\sqrt{-d}) = x^2 + y^2d = 1 \\ &\iff x^2 = 1 \text{ and } y = 0 \\ &\iff x = \pm 1 \text{ and } y = 0. \end{aligned}$$

Hence the only units of the ring $\mathbb{Z}[\sqrt{-d}]$ are ± 1 ($d > 1$).

□

I.2. Integrality

Exercise I.2.1.

Is $\frac{3+2\sqrt{6}}{1-\sqrt{6}}$ an algebraic integer?

Proof.

- (1) $\alpha := \frac{3+2\sqrt{6}}{1-\sqrt{6}} = -3 - \sqrt{6}$. Since the set of all algebraic integers is a ring, α is an algebraic integer.
- (2) Or show that α satisfies a monic equation $x^2 + 6x + 3 = 0 \in \mathbb{Z}[x]$.

□

Exercise I.2.2.

Show that, if the integral domain A is integrally closed, then so is the polynomial ring $A[t]$.

Proof.

- (1) Suppose A is integrally closed in B . Show that $A[t]$ is integrally closed in $B[t]$. Suppose $f \in B[t]$ is integral over $A[t]$. Write

$$f^n + g_1 f^{n-1} + \cdots + g_{n-1} f + g_n = 0$$

where $n > 0$ and $g_i \in A[t]$. Hence

$$\begin{aligned} f^n + g_1 f^{n-1} + \cdots + g_{n-1} f &= -g_n \in A[t] \\ \implies f \underbrace{(f^{n-1} + g_1 f^{n-2} + \cdots + g_{n-1})}_{:=g} &\in A[t]. \end{aligned}$$

It is possible to show that $fg \in A[t]$ implies that $f \in A[t]$ and $g \in A[t]$ by using the fact that A is integrally closed in B .

- (2) Suppose f, g are monic polynomials in $B[t]$. Show that $fg \in A[t]$ implies that $f \in A[t]$ and $g \in A[t]$. Write

$$f = \prod (t - \xi_i), \quad g = \prod (t - \eta_j)$$

in some splitting field F of f and g containing the quotient field of B . Note that each ξ_i and each η_j is a root of a monic equation fg in $A[t]$. Since A is integrally closed in B , $\xi_i, \eta_j \in A$. Hence $f, g \in A[t]$.

- (3) To apply part (2), we need to remedy leading coefficients of f and g . Take an integer $m > \max\{\deg(f), \deg(g_1), \dots, \deg(g_n)\}$. Let $f_0 = t^m + f$ be a monic polynomial in $B[t]$. Hence

$$\begin{aligned} (f_0 - t^m)^n + g_1(f_0 - t^m)^{n-1} + \dots + g_n &= 0 \\ \implies f_0^n + h_1 f_0^{n-1} + \dots + h_n &= 0 \end{aligned}$$

where

$$h_n = t^{mn} + (-1)^{n-1} g_1 t^{m(n-1)} + \dots + g_n \in A[t]$$

is also monic. So

$$\begin{aligned} f_0^n + h_1 f_0^{n-1} + \dots + h_{n-1} f &= -h_n \text{ is monic in } A[t] \\ \implies f_0 (\underbrace{f_0^{n-1} + h_1 f_0^{n-2} + \dots + h_{n-1}}_{:=h_0}) &\in A[t] \text{ where} \\ f_0 \text{ and } h_0 &\text{ both are monic in } B[t]. \end{aligned}$$

Now we can apply part (2) safely.

- (4) In part (1), we let B be the quotient field of A and thus the quotient field of $A[t]$ is $B(t)$. Hence

$$\begin{aligned} f &\in B(t) \text{ integral over } A[t] \\ \implies f &\in B(t) \text{ integral over } B[t] & (A[t] \subseteq B[t]) \\ \implies f &\in B[t] & (B[t] \text{ is a UFD}) \\ \implies f &\in B[t] \text{ integral over } A[t] \\ \implies f &\in A[t]. & ((1)) \end{aligned}$$

□

Exercise I.2.3.

In the polynomial ring $A = \mathbb{Q}[x, y]$, consider the principal ideal $\mathfrak{p} = (x^2 - y^3)$. Show that \mathfrak{p} is a prime ideal, but A/\mathfrak{p} is not integrally closed.

Proof.

- (1) It is easy to show that $x^2 - y^3$ is irreducible in A . Hence $\mathfrak{p} = (x^2 - y^3)$ is prime since A is a UFD.
- (2) By substituting $x = t^3$, $y = t^2$, $A/\mathfrak{p} \cong \mathbb{Q}[t^3, t^2]$, with quotient field $\mathbb{Q}(t)$ (by noting $t = \frac{x}{y}$). Note that $\mathbb{Q}[t]$ is a UFD, thus is already integrally closed. So the integral closure will be $\mathbb{Q}[t] \supsetneq \mathbb{Q}[t^3, t^2]$. It suggests that A/\mathfrak{p} might not be integrally closed.
- (3) (Reductio ad absurdum) If not, then the element $\frac{x}{y}$ satisfies a monic equation $t^2 - y = 0 \in (A/\mathfrak{p})[t]$. So $\frac{x}{y} \in A/\mathfrak{p}$ or $t \in \mathbb{Q}[t^3, t^2]$, which is absurd.

□

Note.

- (1) Serre's criterion for normality.
- (2) Hence smoothness is the same as normality for affine curves in $\mathbb{Q}[x, y]$. Note that $x^2 - y^3$ is an irreducible cubic with a cusp at the origin $(0, 0)$.
- (3) There is an affine variety $X \in \mathbb{Q}[x, y, z]$ such that X is normal but not smooth. ($X = V(x^2 + y^2 - z^2)$ for example.)

Exercise I.2.4.

Let D be a squarefree rational integer $\neq 0, 1$ and d the discriminant of the quadratic number field $K = \mathbb{Q}(\sqrt{D})$. Show that

$$d = \begin{cases} D & \text{if } D \equiv 1 \pmod{4}, \\ 4D & \text{if } D \equiv 2, 3 \pmod{4}. \end{cases}$$

and that an integral basis of K is given by $\{1, \sqrt{D}\}$ in the second case, by $\left\{1, \frac{1+\sqrt{D}}{2}\right\}$ in the first case, and by $\left\{1, \frac{d+\sqrt{d}}{2}\right\}$ in both case.

Proof.

- (1) The Galois group of $K|\mathbb{Q}$ has two elements, the identity and an automorphism sending \sqrt{D} to $-\sqrt{D}$.
- (2) Note that $\alpha \in \mathcal{O}_K$ iff $\text{Tr}_{K|\mathbb{Q}}(\alpha), N_{K|\mathbb{Q}}(\alpha) \in \mathbb{Z}$ (by noting that the equation $x^2 - \text{Tr}_{K|\mathbb{Q}}(\alpha)x + N_{K|\mathbb{Q}}(\alpha) = 0$ has a root $x = \alpha$). So given $\alpha = x + y\sqrt{D} \in \mathcal{O}_K$, we have

$$\begin{aligned} \text{Tr}_{K|\mathbb{Q}}(\alpha) &= 2x \in \mathbb{Z}, \\ N_{K|\mathbb{Q}}(\alpha) &= x^2 - Dy^2 \in \mathbb{Z}. \end{aligned}$$

(3) So $4(x^2 - Dy^2) = (2x)^2 - D(2y)^2 \in \mathbb{Z}$. So $D(2y)^2 \in \mathbb{Z}$ since $2x \in \mathbb{Z}$. So $2y \in \mathbb{Z}$ since D is squarefree $\neq 0, 1$. Let $r = 2x, s = 2y$. Then $r^2 - Ds^2 \equiv 0 \pmod{4}$. Note that a square $\equiv 0, 1 \pmod{4}$.

(4) If $D \equiv 1 \pmod{4}$, then

$$\begin{aligned} r^2 - Ds^2 &\equiv r^2 - s^2 \pmod{4} \\ \implies r \text{ and } s &\text{ has the same parity} \\ \implies \mathcal{O}_K &= \left\{ \frac{r + s\sqrt{D}}{2} : r \equiv s \pmod{2} \right\} \\ \implies \mathcal{O}_K &= \left\{ \frac{r-s}{2} + s \cdot \frac{1+\sqrt{D}}{2} : r \equiv s \pmod{2} \right\} \\ \implies \mathcal{O}_K &= \mathbb{Z} + \mathbb{Z} \frac{1+\sqrt{D}}{2}. \end{aligned}$$

So $\left\{1, \frac{1+\sqrt{D}}{2}\right\}$ is an integral basis of K . Hence

$$d = \begin{vmatrix} 1 & \frac{1+\sqrt{D}}{2} \\ 1 & \frac{1-\sqrt{D}}{2} \end{vmatrix}^2 = D.$$

(5) If $D \equiv 2, 3 \pmod{4}$, then

$$\begin{aligned} r^2 - Ds^2 &\equiv r^2 + 2s^2 \text{ or } r^2 + s^2 \pmod{4} \\ \implies \text{both } r \text{ and } s &\text{ are even} \\ \implies \text{both } x \text{ and } y &\text{ are rational integers} \\ \implies \mathcal{O}_K &= \mathbb{Z} + \mathbb{Z}\sqrt{D}. \end{aligned}$$

So $\{1, \sqrt{D}\}$ is an integral basis of K . Hence

$$d = \begin{vmatrix} 1 & \sqrt{D} \\ 1 & -\sqrt{D} \end{vmatrix}^2 = 4D.$$

(6) By (4)(5), $\left\{1, \frac{d+\sqrt{d}}{2}\right\}$ is an integral basis of K for any case.

□

Exercise I.2.7. (Stickelberger's discriminant relation)

The discriminant d_K of an algebraic number field K is always $\equiv 0 \pmod{4}$ or $\equiv 1 \pmod{4}$. (Hint: The discriminant $\det(\sigma_i \omega_j)$ of an integral basis ω_j

is a sum of terms, each prefixed by a positive or a negative sign. Writing P (resp. N) for the sum of the positive (resp. negative) terms, one find $d_K = (P - N)^2 = (P + N)^2 - 4PN$.)

Proof (Hint).

- (1) Let S_n be the symmetric group of degree n , and A_n be the alternating group of degree n . So

$$\begin{aligned} \det(\sigma_i \omega_j) &= \sum_{\pi \in S_n} \left(\operatorname{sgn}(\pi) \prod_{i=1}^n \sigma_i \omega_{\pi(i)} \right) \\ &= \underbrace{\sum_{\pi \in A_n} \prod_{i=1}^n \sigma_i \omega_{\pi(i)}}_{:=P} - \underbrace{\sum_{\pi \in S_n - A_n} \prod_{i=1}^n \sigma_i \omega_{\pi(i)}}_{:=N}. \end{aligned}$$

- (2) Note that $\sigma_i(P + N) = P + N$ and $\sigma_i(PN) = PN$ for all σ_i . Hence $P + N, PN \in \mathbb{Q}$. Therefore $P + N, PN \in \mathbb{Q} \cap \mathcal{O}_K = \mathbb{Z}$.

- (3) By (1)(2),

$$\begin{aligned} d_K &= \det(\sigma_i \omega_j)^2 \\ &= (P - N)^2 \\ &= (P + N)^2 - 4PN \\ &\equiv 0, 1 \pmod{4}. \end{aligned}$$

□

I.3. Ideals

Exercise I.3.4.

A Dedekind domain with a finite number of prime ideals is a principal ideal domain. (Hint: If $\mathfrak{a} = \mathfrak{p}_1^{\nu_1} \cdots \mathfrak{p}_r^{\nu_r} \neq 0$ is an ideal, then choose elements $\pi_i \in \mathfrak{p}_i \setminus \mathfrak{p}_i^2$ and apply the Chinese remainder theorem for the cosets $\pi_i^{\nu_i} \pmod{\mathfrak{p}_i^{\nu_i+1}}$.)

Proof.

- (1) The hint gives all.
- (2) The existence of π_i is guaranteed by Theorem I.3.3 (the unique prime factorization). The Chinese remainder theorem shows that there is one element $\pi \in \mathcal{O}$ such that $\pi = \pi_i^{\nu_i} \pmod{\mathfrak{p}_i^{\nu_i+1}}$ for each i .

- (3) Hence $\mathfrak{p} = (\pi)$ since they have the same prime factorization.

□

Exercise I.3.5.

The quotient ring \mathcal{O}/\mathfrak{a} of a Dedekind domain by an ideal $\mathfrak{a} \neq 0$ is a principal ideal domain. (Hint: For $\mathfrak{a} = \mathfrak{p}^n$ the only proper ideals of \mathcal{O}/\mathfrak{a} are given by $\mathfrak{p}/\mathfrak{p}^n, \dots, \mathfrak{p}^{n-1}/\mathfrak{p}^n$. Choose $\pi \in \mathfrak{p} \setminus \mathfrak{p}^2$ and show that $\mathfrak{p}^\nu = \mathcal{O}\pi^\nu + \mathfrak{p}^n$.)

Proof.

- (1) By the Chinese remainder theorem, it suffices to show the case $\mathfrak{a} = \mathfrak{p}^n$ where \mathfrak{p} is prime.
- (2) There is a natural correspondence between

$$\{\text{ideals of } \mathcal{O}/\mathfrak{p}^n\} \longleftrightarrow \{\text{ideals of } \mathcal{O} \text{ containing } \mathfrak{p}^n\}.$$

Hence the proper ideals of $\mathcal{O}/\mathfrak{p}^n$ are given by $\mathfrak{p}/\mathfrak{p}^n, \dots, \mathfrak{p}^{n-1}/\mathfrak{p}^n$.

- (3) Similar to Exercise I.3.4, choose $\pi \in \mathfrak{p} \setminus \mathfrak{p}^2$ and thus $\mathfrak{p}^\nu = \mathcal{O}\pi^\nu + \mathfrak{p}^n$ ($\nu = 1, \dots, n-1$) since they have the same prime factorization. Hence $\mathfrak{p}^\nu/\mathfrak{p}^n = (\pi^\nu + \mathfrak{p}^n)$ is principal.

□

Exercise I.3.6.

Every ideal of a Dedekind domain can be generated by two elements. (Hint: Use Exercise I.3.5.)

Proof.

- (1) Given an ideal $\mathfrak{a} \neq 0$ of a Dedekind domain \mathcal{O} . (Nothing to do if $\mathfrak{a} = 0 = (0)$.) So \mathcal{O}/\mathfrak{a} is a principal ideal domain (Exercise I.3.5).
- (2) Take any $\alpha \in \mathfrak{a} \setminus \{0\}$. So $(\alpha)/\mathfrak{a} = (\beta \pmod{\mathfrak{a}})$ is a principal ideal for some $\beta \in \mathcal{O}$. So $\mathfrak{a} = (\alpha, \beta)$ is generated by two elements.

□

I.4. Lattices

Exercise I.4.1.

Show that a lattice Γ in \mathbb{R}^n is complete if and only if the quotient \mathbb{R}^n/Γ is compact.

Proof.

- (1) (\implies) Define a natural homeomorphism $\varphi : \mathbb{R}^n/\Gamma \rightarrow \mathbb{S}^1 \times \cdots \times \mathbb{S}^1$ by sending (x_1, \dots, x_n) to $(x_1 \pmod{1}, \dots, x_n \pmod{1})$ (where $\mathbb{S}^1 \subseteq \mathbb{R}^2$ is a unit circle). Note that $\mathbb{S}^1 \times \cdots \times \mathbb{S}^1$ is compact.
- (2) (\impliedby) Let V_0 be the linear subspace of V which is spanned by the set Γ . Since the vector space V/V_0 is contained in a compact set V/Γ ,

$$\dim(V/V_0) = 0$$

(otherwise V/V_0 is unbounded). Hence $V_0 = V$ or Γ is complete.

□

Exercise I.4.2.

Show that Minkowski's lattice point theorem cannot be improved, by giving an example of a centrally symmetric convex set $X \subset V$ such that $\text{vol}(X) = 2^n \text{vol}(\Gamma)$ which does not contain any nonzero point of the lattice Γ . If X is compact, however, then the statement $\text{vol}(X) > 2^n \text{vol}(\Gamma)$ does remain true in the case of equality.

Proof.

- (1) Let $V = \mathbb{R}^n$, $\Gamma = \mathbb{Z}^n$ be a complete lattice in V , and $X = (-1, 1)^n \subseteq \mathbb{R}^n$ be a centrally symmetric convex set in V . Hence $\text{vol}(X) = 2^n \text{vol}(\Gamma)$ and X does not contain any nonzero point of Γ .
- (2) Suppose X is compact. Consider $X_\nu = (1 + \frac{1}{\nu})X$ for each $\nu \in \mathbb{Z}_{>0}$. Thus X_ν is again a centrally symmetric convex set in V and

$$\begin{aligned} \text{vol}(X_\nu) &= \left(1 + \frac{1}{\nu}\right) \text{vol}(X) \\ &\geq \left(1 + \frac{1}{\nu}\right) 2^n \text{vol}(\Gamma) \\ &> 2^n \text{vol}(\Gamma). \end{aligned}$$

Minkowski's lattice point theorem shows that there is one nonzero lattice point $\gamma_\nu \in \Gamma$ for $\nu = 1, 2, 3, \dots$

- (3) By the compactness of X_1 , there is a subsequence of $\{\gamma_\nu\}$ converging to $\gamma \in X_1$. Since Γ is discrete (Proposition I.4.2), there are infinitely many ν such that $\gamma = \gamma_\nu \in X_\nu$. (In particular, $\gamma \neq 0$.) Hence $\gamma \in X$ by the compactness of X .

□

I.5. Minkowski Theory

Exercise I.5.2.

Show that the convex, centrally symmetric set

$$X = \left\{ (z_\tau) \in K_{\mathbb{R}} : \sum_{\tau} |z_\tau| < t \right\}$$

has volume $\text{vol}(X) = 2^r \pi^s \frac{t^n}{n!}$.

Proof. It is the same as Lemma III.2.15. □

Exercise I.5.3. (Minkowski bound)

Show that in every ideal $\mathfrak{a} \neq 0$ of \mathcal{O}_K there exists an $a \neq 0$ such that

$$|N_{K|\mathbb{Q}}(a)| \leq M(\mathcal{O}_K : \mathfrak{a}),$$

where $M = \frac{n!}{n^n} \left(\frac{4}{\pi}\right)^s \sqrt{|d_K|}$ (the so-called **Minkowski bound**.)

Proof.

- (1) Let

$$X_t = \left\{ (z_\tau) \in K_{\mathbb{R}} : \sum_{\tau} |z_\tau| \leq t \right\}$$

be a convex, centrally symmetric set for any $t > 0$. Note that $\text{vol}(X_t) = 2^r \pi^s \frac{t^n}{n!}$ (same as Exercise I.5.2).

- (2) In particular, we take $t > 0$ so that

$$\text{vol}(X_t) = 2^r \pi^s \frac{t^n}{n!} = 2^n \text{vol}(\Gamma).$$

Thus the hypothesis of Minkowski's lattice point theorem in Exercise I.4.2 is satisfied. So there does indeed exist a lattice point $ja \in X_t$, $a \neq 0$, $a \in \mathfrak{a}$; in other words, $\sum_{\tau} |\tau a| \leq t$.

(3) Hence

$$\begin{aligned}
|N_{K|\mathbb{Q}}(a)| &= \prod_{\tau} |\tau a| \\
&\leq \left(\frac{1}{n} \sum_{\tau} |\tau a| \right)^n && \text{(AM-GM inequality)} \\
&\leq \frac{t^n}{n^n} && (ja \in X_t) \\
&= \frac{1}{n^n} \frac{n!}{2^r \pi^s} 2^n \text{vol}(\Gamma) && \text{(Definition of } t^n) \\
&= \frac{1}{n^n} \frac{n!}{2^r \pi^s} 2^n \sqrt{|d_K|} (\mathcal{O}_K : \mathfrak{a}) && \text{(Proposition I.5.2)} \\
&= \underbrace{\frac{n!}{n^n} \left(\frac{4}{\pi} \right)^s}_{:=M} \sqrt{|d_K|} (\mathcal{O}_K : \mathfrak{a}). && (n = r + 2s)
\end{aligned}$$

□

I.6. The Class Number

Exercise I.6.3.

Show that in every ideal class of an algebraic number field K of degree n , there exists an integral ideal \mathfrak{a}_1 such that

$$\mathfrak{N}(\mathfrak{a}_1) \leq \frac{n!}{n^n} \left(\frac{4}{\pi} \right)^s \sqrt{|d_K|}$$

(Hint: Use Exercise I.3.5, proceed as in the proof of Theorem I.6.3.)

Proof.

- (1) The hint gives all.
- (2) Take an arbitrary representative \mathfrak{a} of the class in the ideal class group, and a $\gamma \in \mathcal{O}_K$, $\gamma \neq 0$, such that $\mathfrak{b} := \gamma \mathfrak{a}^{-1} \subseteq \mathcal{O}_K$. By Exercise I.3.5, there exists $\alpha \in \mathfrak{b}$, $\alpha \neq 0$, such that

$$|N_{K|\mathbb{Q}}(\alpha)| \cdot \mathfrak{N}(\mathfrak{b})^{-1} = \mathfrak{N}((\alpha)\mathfrak{b}^{-1}) = \mathfrak{N}(\alpha\mathfrak{b}^{-1}) \leq \frac{n!}{n^n} \left(\frac{4}{\pi} \right)^s \sqrt{|d_K|}.$$

The ideal

$$\mathfrak{a}_1 := \alpha \mathfrak{b}^{-1} = \alpha \gamma^{-1} \mathfrak{a} \in [\mathfrak{a}]$$

therefore has the required property.

- (3) This exercise also shows that Cl_K is a finite group.

□

I.11. Localization

Exercise I.11.7. (Nakayama's lemma)

Let A be a local ring with maximal ideal \mathfrak{m} , let M be an A -module and $N \subseteq M$ a submodule such that M/N is finitely generated. Then one has the implication:

$$M = N + \mathfrak{m}M \implies M = N.$$

Proof.

- (1) Note that

$$M = N + \mathfrak{m}M \implies M/N = (N + \mathfrak{m}M)/N = \mathfrak{m}(M/N).$$

So it suffices to show that $M' := M/N = 0$.

- (2) (Reductio ad absurdum) If $M' \neq 0$, then there exists a minimal set of generators $\{x_1, \dots, x_n\}$ for M' . Take $x_n \in M' = \mathfrak{m}(M')$. We have an equation of the form

$$\begin{aligned} x_n &= m_1 x_1 + \dots + m_n x_n \\ \iff (1 - m_n)x_n &= m_1 x_1 + \dots + m_{n-1} x_{n-1}. \end{aligned}$$

where $m_\nu \in \mathfrak{m}$ for all ν . Since \mathfrak{m} is the maximal ideal of a local ring, $1 - m_n$ is a unit. So x_n is in the submodule of M' generated by $\{x_1, \dots, x_{n-1}\}$, contrary to the minimality of n .

□

I.13. One-dimensional Schemes

No exercises.

I.14. Function Fields

No exercises.

Chapter II: The Theory of Valuations

II.1. The p -adic Numbers

Exercise II.1.2.

A p -adic integer $a = a_0 + a_1p + a_2p^2 + \cdots$ is a unit in the ring \mathbb{Z}_p if and only if $a_0 \neq 0$.

Proof.

- (1) (\implies) If $b = b_0 + b_1p + b_2p^2 + \cdots$ is an inverse of a , then $ab = 1$ implies that $a_0b_0 = 1$ so that a_0 is a unit in $\mathbb{Z}/p\mathbb{Z}$ or $a_0 \neq 0$.
- (2) (\impliedby) Our goal is to find

$$b = b_0 + b_1p + b_2p^2 + \cdots \in \mathbb{Z}_p$$

such that the Cauchy product

$$ab = c_0 + c_1p + c_2p^2 + \cdots$$

is equal to $1 \in \mathbb{Z}_p$. Here $c_n = \sum_{\nu=0}^n a_\nu b_{n-\nu}$. By the assumption we have that $c_0 = 1$ and $c_1 = c_2 = \cdots = 0$. Hence

$$\begin{aligned} b_0 &= a_0^{-1} \\ b_1 &= -a_0^{-1}a_1b_0 \\ &\vdots \\ b_n &= a_0^{-1} \sum_{\nu=1}^n a_\nu b_{n-\nu} \\ &\vdots \end{aligned}$$

by induction.

- (3) Also see Exercise 1.5 in the textbook: *Atiyah & Macdonald, Introduction to Commutative Algebra*. Let A be a commutative ring with 1 and $A[[x]]$ be the ring of formal power series $f = \sum_{n=0}^{\infty} a_n x^n$ with coefficients in A . Then f is a unit in $A[[x]]$ if and only if a_0 is a unit in A .

□

II.2. The p -adic Absolute Value

Exercise II.2.1.

$$|x - y|_p \geq ||x|_p - |y|_p|.$$

Proof. Note that $|x + y|_p \leq \max\{|x|_p, |y|_p\} \leq |x|_p + |y|_p$ and $|\pm 1|_p = 1$. \square

Chapter VII: Zeta Functions and L -series

VII.1. The Riemann Zeta Function

Exercise VII.1.4.

For the power sum

$$s_k(n) = 1^k + 2^k + 3^k + \cdots + n^k$$

one has

$$s_k(n) = \frac{1}{k+1}(B_{k+1}(n) - B_{k+1}(0)).$$

Proof. By Exercise VII.1.3,

$$x^k = \frac{1}{k+1}(B_{k+1}(x) - B_{k+1}(x-1)).$$

Hence the telescoping sum is

$$\begin{aligned} s_k(n) &= \sum_{x=1}^n x^k \\ &= \sum_{x=1}^n \frac{1}{k+1}(B_{k+1}(x) - B_{k+1}(x-1)) \\ &= \frac{1}{k+1}(B_{k+1}(n) - B_{k+1}(0)). \end{aligned}$$

□