

Review of Last Lecture

Key concepts and/or techniques:

1. Sample mean: Let X_1, X_2, \dots, X_n be independent and identically distributed with mean μ . Then the sample mean is defined as

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i,$$

and a statistic and also an estimator of mean μ .

2. Mgf technique: Mgf, if exists, uniquely determines the distribution of the RV. Therefore, the distribution of a RV can be equivalently found via its mgf.

Use the mgf technique to derive the distribution of

$$Y = \sum_{i=1}^n a_i X_i$$

Review of Last Lecture

[Theorem 5.4-1]

If X_1, X_2, \dots, X_n are independent RVs with respective mgfs $M_{X_i}(t)$ where $|t| < h_i$ for positive number $h_i, i = 1, 2, \dots, n$. Then the mgf of $Y = \sum_{i=1}^n a_i X_i$ is

$$M_Y(t) = \prod_{i=1}^n M_{X_i}(a_i t),$$

where $|a_i t| < h_i, i = 1, \dots, n$.

Review of Last Lecture

[Theorem 5.4-2]

Let X_1, X_2, \dots, X_n be independent chi-square RVs with r_1, r_2, \dots, r_n degrees of freedom, respectively, i.e., $X_i \sim \chi^2(r_i), i = 1, \dots, n$ Then

$$Y = X_1 + X_2 + \dots + X_n \quad \text{is} \quad \chi^2(r_1 + r_2 + \dots + r_n)$$

[Corollary 5.4-3]

If X_1, X_2, \dots, X_n are independent and have normal distributions $N(\mu_i, \sigma_i^2), i = 1, 2, \dots, n$, respectively, then the distribution of

$$\sum_{i=1}^n \left(\frac{X_i - \mu_i}{\sigma_i} \right)^2 \sim \chi^2(n)$$

STA2001 Probability and Statistics I

Lecture 22

Tianshi Chen

The Chinese University of Hong Kong, Shenzhen

Section 5.5 Random function associated with normal distribution

Theorem 5.5-1, page 200

[Theorem 5.5-1]

If X_1, X_2, \dots, X_n are n independent normal variables with means $\mu_1, \mu_2, \dots, \mu_n$ and variances $\sigma_1^2, \sigma_2^2, \dots, \sigma_n^2$, respectively, then $Y = \sum_{i=1}^n a_i X_i$ has the normal distribution

$$Y \sim N \left(\sum_{i=1}^n a_i \mu_i, \sum_{i=1}^n a_i^2 \sigma_i^2 \right)$$

Proof of Theorem 5.5-1, page 200

By Theorem 5.4-1, we have

$$\begin{aligned} M_Y(t) &= \prod_{i=1}^n M_{X_i}(a_i t) = \prod_{i=1}^n \exp(\mu_i a_i t + \frac{1}{2} \sigma_i^2 a_i^2 t^2) \\ &= \exp \left\{ \left(\sum_{i=1}^n \mu_i a_i \right) t + \frac{1}{2} \left(\sum_{i=1}^n a_i^2 \sigma_i^2 \right) t^2 \right\} \end{aligned}$$

Example 1, page 201

Let X_1 and X_2 be the pounds of butter fat produced by 2 cows, respectively. Assume that

$$X_1 \sim N(693.2, 22820), X_2 \sim N(631.7, 19205)$$

and moreover, X_1 and X_2 are independent. What's the probability $P(X_1 > X_2)$?

Example 1, page 201

Let $Y = X_1 - X_2$. Then

$$Y \sim N(693.2 - 631.7, 22820 + 19205) = N(61.5, 42025)$$

$$P(X_1 > X_2) = P(Y > 0) = P\left(\frac{Y - 61.5}{\sqrt{42025}} > \frac{0 - 61.5}{\sqrt{42025}}\right)$$

$$= 1 - \Phi(-0.3) = 0.6179.$$

Corollary 5.5-1, Page 201

[Corollary 5.5-1]

If X_1, X_2, \dots, X_n is a random sample of size n from the normal distribution $N(\mu, \sigma^2)$, then the sample mean \bar{X} has the following distribution


$$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right) \Leftrightarrow \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

Proof: Let $a_i = \frac{1}{n}, \mu_i = \mu, \sigma_i^2 = \sigma^2, i = 1, \dots, n$. Then by Theorem 5.5-1, we obtain the result.

Example 2, page 201

Let X_1, X_2, \dots, X_n be a random sample of size n from $N(50, 16)$,

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N\left(50, \frac{16}{n}\right)$$



- To illustrate the effect of n :
The larger n , the smaller the variance $\frac{16}{n}$.
- pdf of \bar{X} :
The sharper the peak, the more concentrated in a small interval centered at 50.

Sample Variance

Definition

Let X_1, X_2, \dots, X_n be independent and identically distributed with mean μ and σ^2 . Then the sample variance is defined as

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2,$$

and an estimator of the variance σ^2 , because

$$E(S^2) = \sigma^2.$$

Sample Variance

Note that

$$\begin{aligned}\sum_{i=1}^n (X_i - \mu)^2 &= \sum_{i=1}^n (X_i - \bar{X} + \bar{X} - \mu)^2 \\ &= \sum_{i=1}^n (X_i - \bar{X})^2 + \sum_{i=1}^n (X_i - \bar{X})(\bar{X} - \mu) + \sum_{i=1}^n (\bar{X} - \mu)^2\end{aligned}$$

where

$$\begin{aligned}\sum_{i=1}^n (X_i - \bar{X})(\bar{X} - \mu) &= (\bar{X} - \mu) \sum_{i=1}^n (X_i - \bar{X}) \\ &= (\bar{X} - \mu) \left(\sum_{i=1}^n X_i - n\bar{X} \right) = (\bar{X} - \mu)(n\bar{X} - n\bar{X}) = 0\end{aligned}$$

Therefore,

$$s^2 = \frac{1}{n-1} \left[\sum_{i=1}^n (X_i - \mu)^2 - \sum_{i=1}^n (\bar{X} - \mu)^2 \right]$$

Sample Variance

Then note that

$$E \left(\sum_{i=1}^n (X_i - \mu)^2 \right) = \sum_{i=1}^n E \left[(X_i - \mu)^2 \right] = n \cdot \sigma^2$$
$$E \left(\sum_{i=1}^n (\bar{X} - \mu)^2 \right) = \sum_{i=1}^n E \left[(\bar{X} - \mu)^2 \right] = n \cdot \frac{\sigma^2}{n} = \sigma^2$$

Therefore,

$$E(S^2) = \frac{1}{n-1} (n\sigma^2 - \sigma^2) = \sigma^2$$

Theorem 5.5-2, page 202

[Theorem 5.5-2]

Let X_1, X_2, \dots, X_n be random sample of size n from the normal distribution $N(\mu, \sigma^2)$ with $\sigma^2 > 0$. Then the sample mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$ and the sample variance $S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$ are independent, and

$$\frac{(n-1)S^2}{\sigma^2} = \sum_{i=1}^n \left(\frac{X_i - \bar{X}}{\sigma} \right)^2 \sim \chi^2(n-1)$$

The independence of \bar{X} and S^2 is not proved here but deferred to Section 6.7 on page 294, and we only prove the second part.

Proof of Theorem 5.5-2, page 202

Following the proof of $E(S^2) = \sigma^2$, we have

$$\frac{n-1}{\sigma^2} S^2 = \sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma} \right)^2 - \sum_{i=1}^n \left(\frac{\bar{X} - \mu}{\sigma} \right)^2$$

Now let

$$W = \sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma} \right)^2, \quad Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$$

Then

$$W = \frac{(n-1)S^2}{\sigma^2} + Z^2$$

Further note that $W \sim \chi^2(n)$, $Z^2 \sim \chi^2(1)$, and moreover, S^2 and \bar{X} are independent by assumption.

Proof of Theorem 5.5-2, page 202

Note that $W \sim \chi^2(n)$, $Z^2 \sim \chi^2(1)$, S^2 and \bar{X} are independent

$$E[e^{tw}] = E[e^{t(\frac{(n-1)S^2}{\sigma^2} + Z^2)}] = E[e^{t\frac{(n-1)S^2}{\sigma^2}}]E[e^{tZ^2}]$$

$$(1 - 2t)^{-\frac{n}{2}} = E[e^{t\frac{(n-1)S^2}{\sigma^2}}] \cdot (1 - 2t)^{-\frac{1}{2}}, \quad t < \frac{1}{2}$$

$$\Rightarrow E[e^{t\frac{(n-1)S^2}{\sigma^2}}] = (1 - 2t)^{-\frac{n-1}{2}}, \quad t < \frac{1}{2} \Rightarrow \frac{(n-1)S^2}{\sigma^2} \sim \chi^2(n-1)$$

A remark

Combining Corollary 5.4-3 and Thm 5.5-2 leads to the observation:

If X_1, X_2, \dots, X_n is a random sample of size n from $N(\mu, \sigma^2)$, then

$$\sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma} \right)^2 \sim \chi^2(n), \quad \sum_{i=1}^n \left(\frac{X_i - \bar{X}}{\sigma} \right)^2 \sim \chi^2(n-1)$$

When the mean μ is replaced by the sample mean \bar{X} , one degree of freedom is lost.

This is because there is an additional constraint

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

Example 5.5-3, page 204

Let X_1, X_2, X_3, X_4 be a random sample of size 4 from the normal distribution $N(76.4, 383)$.

$$\sum_{i=1}^4 \frac{(X_i - 76.4)^2}{383} \sim \chi^2(4), \quad \sum_{i=1}^4 \frac{(X_i - \bar{X})^2}{383} \sim \chi^2(3)$$

Student's t Distribution

[Theorem 5.5-3]

Let

$$T = \frac{Z}{\sqrt{U/r}},$$

where $Z \sim N(0, 1)$, $U \sim \chi^2(r)$, and Z and U are independent. Then T has a student's t distribution

$$f(t) = \frac{\Gamma(\frac{r+1}{2})}{\sqrt{\pi r} \Gamma(\frac{r}{2})} \frac{1}{(1 + \frac{t^2}{r})^{\frac{r+1}{2}}}, \quad t \in (-\infty, \infty),$$

where r is called the degrees of freedom, and we simply write $T \sim t(r)$.

Sketch of the Proof of Theorem 5.5-3, page 204

Since Z and U are independent, their joint pdf $g(z, u)$ is

$$g(z, u) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} \cdot \frac{1}{\Gamma(\frac{r}{2}) 2^{\frac{r}{2}}} u^{\frac{r}{2}-1} e^{-\frac{u}{2}}, \quad z \in R, u \in [0, \infty)$$

1. The cdf of T , $F(t)$ is

$$F(t) = P(T \leq t) = P\left(\frac{Z}{\sqrt{\frac{U}{r}}} \leq t\right) = P\left(Z \leq \sqrt{\frac{U}{r}} t\right)$$

2. The pdf of T ,

$$f(t) = F'(t)$$

Student's t Distribution: Heavy-tailed Distribution


Student's t distribution is a heavy tailed distribution

Standard normal distribution :

$$f(x) = \frac{1}{\sqrt{2\pi}} \frac{1}{e^{\frac{1}{2}x^2}}, \quad x \in (-\infty, \infty)$$

Students' t distribution with $r = 1$:

$$f(x) = \frac{1}{\pi(1+x^2)}, \quad x \in (-\infty, \infty)$$



Therefore, Student's t distribution is a better choice than the normal distribution when the data contains outliers.

A Student's t RV based on Random Samples from Normal Distribution

By using the result of Corollary 5.5-1 and Theorems 5.5-2 and 5.5-3, we can construct an important student's t random variable.

Assume that X_1, X_2, \dots, X_n is a random sample of size n from a normal distribution $N(\mu, \sigma^2)$.

A Student's t RV based on Random Samples from Normal Distribution

Let

$$Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}}, \quad U = \frac{(n-1)S^2}{\sigma^2}$$

Then $Z \sim N(0, 1)$ and $U \sim \chi^2(n-1)$. Since Z and U are independent, then

$$T = \frac{Z}{\sqrt{U/(n-1)}} = \frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$

A remark

If X_1, \dots, X_n is a random sample of size n from a normal distribution $N(\mu, \sigma^2)$, then

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1), \quad \frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$