

# STA2001 Probability and Statistics (I)

## Lecture 3

Tianshi Chen

The Chinese University of Hong Kong, Shenzhen

# Review

- For random experiments that satisfy

Assumption 1:  $S$  contains  $m$  possible outcomes

$$e_k, \quad k = 1, 2, \dots, m, \quad \text{i.e.,} \quad S = \{e_1, e_2, \dots, e_m\}.$$

Assumption 2: The  $m$  outcomes are “equally likely”

$$P(\{e_k\}) = \frac{1}{m}, \quad k = 1, \dots, m.$$

$$P(A) = \frac{N(A)}{N(S)},$$

where  $N(X)$  is the number of outcomes in  $X \subseteq S$ .

# Review

- ▶ Revisit the method of enumeration by multiplication principle
  - ▶ permutation
  - ▶ combination
  - ▶ distinguishable permutation

## Section 1.3 Conditional Probability

# Motivation Example

Consider a number of tulip bulbs

	Early(E)	Late(L)	Totals
Red(R)	5	8	13
Yellow(Y)	3	4	7
Totals	8	12	20

**Experiment 1:** Select one bulb randomly.

- ▶ Sample space  $S = \{\text{all bulbs}\}$ .
- ▶ Assumption: all bulbs are “equally likely”.

Consider the event  $R = \{\text{the selected bulb is red}\}$ , what is  $P(R)$ ?

# Motivation Example

Consider a number of tulip bulbs

	Early(E)	Late(L)	Totals
Red(R)	5	8	13
Yellow(Y)	3	4	7
Totals	8	12	20

**Experiment 1:** Select one bulb randomly.

- ▶ Sample space  $S = \{\text{all bulbs}\}$ .
- ▶ Assumption: all bulbs are “equally likely”.

Consider the event  $R = \{\text{the selected bulb is red}\}$ , what is  $P(R)$ ?

$$P(R) = \frac{N(R)}{N(S)} = \frac{13}{20}$$

## Motivation Example

Consider a number of tulip bulbs

	Early(E)	Late(L)	Totals
Red(R)	5	8	13
Yellow(Y)	3	4	7
Totals	8	12	20

**Experiment 2:** Select one bulb from the ones that bloom early.

- ▶ Sample space reduces to  $E = \{\text{all bulbs that bloom early}\}$ .
- ▶ Assumption: all bulbs are “equally likely”.

Consider the event  $R = \{\text{the selected bulb is red}\}$ , what is the probability of the event  $R$ , denoted by  $P(R|E)$ ?

## Motivation Example

Consider a number of tulip bulbs

	Early(E)	Late(L)	Totals
Red(R)	5	8	13
Yellow(Y)	3	4	7
Totals	8	12	20

**Experiment 2:** Select one bulb from the ones that bloom early.

- ▶ Sample space reduces to  $E = \{\text{all bulbs that bloom early}\}$ .
- ▶ Assumption: all bulbs are “equally likely”.

Consider the event  $R = \{\text{the selected bulb is red}\}$ , what is the probability of the event  $R$ , denoted by  $P(R|E)$ ?

$$P(R|E) = \frac{N(R \cap E)}{N(E)} = \frac{5}{8}$$

We have defined a new probability function associated with the reduced sample space  $E$ .



# Motivation Example

We study the problem of how to define a new probability function associated with a reduced sample space  $E \subseteq S$ , where  $S$  is the original sample space.

1. We have defined the probability function associated with the reduced sample space  $E$  directly.
2. We can also define it by linking to the probability function associated with the original sample space  $S$ .

# Motivation Example

Under the assumptions that

1.  $S$  is finite
2. All outcomes are “equally likely”

the above example give us the idea

$$P(R|E) = \frac{N(R \cap E)}{N(E)} = \frac{N(R \cap E)/N(S)}{N(E)/N(S)} = \frac{P(R \cap E)}{P(E)}$$

leading to the next definition

# Conditional Probability

## Definition

The conditional probability of an event  $A$ , given that the event  $B$  has occurred, is defined by

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

provided that  $P(B) > 0$ .

- ▶  $B$  is the sample space for  $P(A|B)$
- ▶ Independent of Assumptions 1 & 2 on the previous slide.

# Conditional Probability

Conditional probability satisfies the probability axioms

1.  $P(A|B) \geq 0$ .
2.  $P(B|B) = 1$ .
3. If  $A_1, A_2, A_3, \dots$  are countable and mutually exclusive events, then

$$P(A_1 \cup A_2 \cup \dots | B) = P(A_1|B) + P(A_2|B) + \dots$$

## Example 2

$$P(A) = 0.4, \quad P(B) = 0.5, \quad P(A \cap B) = 0.3,$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{0.3}{0.5} = 0.6$$

$$P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{0.3}{0.4} = 0.75$$

Can  $P(A|B) > 1$  or  $P(A|B) < 0$ ?

## Example 2

$$P(A) = 0.4, \quad P(B) = 0.5, \quad P(A \cap B) = 0.3,$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{0.3}{0.5} = 0.6$$

$$P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{0.3}{0.4} = 0.75$$

Can  $P(A|B) > 1$  or  $P(A|B) < 0$ ?

No,  $P(A|B)$  is a probability function.

## Example 3 (Shooting Game)

### Question

25 balloons of which, 10 are yellow, 8 red, 7 green.

$$A = \{\text{the first balloon shot is yellow}\}$$

$$B = \{\text{the second balloon shot is yellow}\}$$

What is the probability that the first two balloons shot are all yellow?

## Example 3 (Shooting Game)

### Question

25 balloons of which, 10 are yellow, 8 red, 7 green.

$A = \{\text{the first balloon shot is yellow}\}$

$B = \{\text{the second balloon shot is yellow}\}$

What is the probability that the first two balloons shot are all yellow?

$$P(A) = \frac{10}{25}, \quad P(B|A) = \frac{9}{24}$$

$$\Rightarrow P(A \cap B) = P(A)P(B|A) = \frac{10}{25} \cdot \frac{9}{24}$$



# Multiplication Rule

## Definition

The probability that two events,  $A$  and  $B$  both occur is given by the multiplication rule

$$P(A \cap B) = P(A)P(B|A), \quad \text{provided } P(A) > 0$$

or by


$$P(A \cap B) = P(B)P(A|B), \quad \text{provided } P(B) > 0$$

## Example 4

### Question

A bowl contains 10 chips in total, 7 blue and 3 red. Drawn 2 chips successively at random and without replacement. What is the probability that the 1st draw is red and the 2nd draw is blue?

## Example 4



$$A = \{\text{1st draw is red}\}$$

$$B = \{\text{2nd draw is blue}\}$$

$$P(A) = \frac{3}{10}, \quad P(B|A) = \frac{7}{9}$$

$$P(A \cap B) = P(B|A) \cdot P(A) = \frac{3}{10} \cdot \frac{7}{9} = \frac{7}{30}$$

# Multiplication Rule for Three Events

## Definition

The probability that three events,  $A$ ,  $B$  and  $C$  all occur is given by the multiplication rule

$$P(A \cap B \cap C) = P((A \cap B) \cap C) = P(A \cap B)P(C|A \cap B)$$

$$\text{where } P(A \cap B) = P(A)P(B|A)$$

$$\Rightarrow P(A \cap B \cap C) = P(A)P(B|A)P(C|A \cap B)$$

Induction principle can be used to derive the cases for more than three events.

## Example 5

### Question

Roll a pair of 4-sided dice and observe the sum of the dice

$$A = \{\text{a sum of 3 is rolled}\}$$

$$B = \{\text{a sum of 3 or a sum of 5 is rolled}\}$$

$$C = \{\text{a sum of 3 is rolled before a sum of 5 is rolled}\}$$

What are  $P(A)$ ,  $P(B)$ ,  $P(C)$ ?

## Example 5

Consider  $P(A)$  and  $P(B)$ :

the sample space  $S = \{(1, 1), (1, 2), \dots, (4, 4)\}$

$$P(A) = \frac{N(A)}{N(S)} = \frac{2}{16}, \quad P(B) = \frac{N(B)}{N(S)} = \frac{6}{16}$$

Consider  $P(C)$ :

- ▶ Method 1 [by definition]:
  - A. Figure out the simplified random experiment
  - B. Figure out the corresponding sample space and the event

## Example 5

For A, repeat the experiment of rolling a pair of 4-sided dice and record the sum of dice. For each repetition, we keep rolling the dice till we see either a sum of 3 or a sum of 5. Then we stop because we have an answer to the problem whether a sum of 3 is rolled before a sum of 5 is rolled.

## Example 5

For instance

Repetition 1 : 2, 4, 6, 3.

Repetition 2 : 8, 6, 7, 4, 5

Repetition 3 : 6, 5.

The sums other than 3 and 5 do not matter and we can remove them.

Repetition 1: a sum of 3 first

Repetition 2: a sum of 5 first

Repetition 3: a sum of 5 first

The problem reduces to roll the pair of dice (that gives the sum either 3 or 5) once and compute the probability that the sum is a 3.



## Example 5

For **B**, the reduced sample space

$$S_r = \left\{ \begin{array}{l} (1, 2), (2, 1) \\ (2, 3), (3, 2) \\ (1, 4), (4, 1) \end{array} \right\} \text{ give a sum of 3 or 5}$$

$$P(C) = P(\{\text{roll the pair of dice once and the sum is a 3}\})$$

$$\begin{aligned} &= \frac{N(\{\text{roll the pair of dice once and the sum is 3}\})}{N(S_r)} \\ &= \frac{2}{6} \end{aligned}$$

## Example 5

- ▶ Method 2 [by conditional probability]:

$$P(C) = P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{2/16}{6/16} = \frac{2}{6}$$

Note that the event “ $A|B$ ” is the same as event “ $C$ ”.

This is because

- A. Event  $C$  is concerned with the cases where the sum is either a 3 or a 5. ‘ $B$  happened’ means that the sum is either a 3 or a 5.
- B. If  $B$  happened then  $A|B$  is nothing but the event “roll the pair of dice (that gives the sum either 3 or 5) once, and the sum is 3”.

## Section 1.4 Independent Events

# Motivation

## Motivation

For certain pair of events, the occurrence of one of them does not change the probability of the occurrence of the other.

# Example 1

Experiment: flip a coin twice and observe the sequence of heads and tails.

Sample space:  $S = \{HH, HT, TH, TT\}$

Assumption: the four outcomes are “equally likely”

Events:

$$A = \{\text{heads on the first flip}\} = \{HH, HT\}$$

$$B = \{\text{tails on the second flip}\} = \{HT, TT\}$$

$$C = \{\text{tails on both flips}\} = \{TT\}$$

## Example 1

$$P(A) = \frac{2}{4}, \quad P(B) = \frac{2}{4}, \quad P(C) = \frac{1}{4}$$

## Example 1

$$P(A) = \frac{2}{4}, \quad P(B) = \frac{2}{4}, \quad P(C) = \frac{1}{4}$$

Given that  $C$  has occurred, then

$$P(B|C) = 1 \text{ because } C \subset B \text{ or } \frac{P(B \cap C)}{P(C)} = \frac{P(C)}{P(C)} = 1$$

Given that  $A$  has occurred, then

$$P(B|A) = \frac{P(B \cap A)}{P(A)} = \frac{1/4}{2/4} = \frac{1}{2} = P(B)$$

Given that  $B$  has occurred, then

$$P(A|B) = \frac{1}{2} = P(A)$$

## Example 1

So we have

$$P(B|A) = P(B), \text{ and } P(A|B) = P(A)$$

the occurrence of one of them does not affect the probability of the occurrence of the other. Leading to the definition of independent events.



# Independent Events

## Definition

Events  $A$  and  $B$  are independent if

$$P(A \cap B) = P(A)P(B).$$

Otherwise, events  $A$  and  $B$  are called dependent events

- ▶ When  $P(A) \neq 0$  and  $P(B) \neq 0$ , we have

$$P(A|B) = P(A), \quad P(B|A) = P(B)$$

## Example 2, page 38

### Question

A red die and a white die are rolled.

$$S = \{(1, 1), (1, 2), \dots\}, \quad N(S) = 36$$

$$A = \{4 \text{ on the red die}\}, \quad B = \{\text{sum of dice is odd}\}$$

Assuming the two dice are fair. Are  $A$  and  $B$  independent?

## Example 2, page 38

A red die and a white die are rolled.

$$S = \{(1, 1), (1, 2), \dots\}, \quad N(S) = 36$$

$$A = \{4 \text{ on the red die}\}, \quad B = \{\text{sum of dice is odd}\}$$

Assuming the two dice are fair. Are  $A$  and  $B$  independent?

$$P(A) = \frac{6}{36}, \quad P(B) = \frac{18}{36}, \quad P(A \cap B) = \frac{3}{36}$$

$$P(A \cap B) = \frac{3}{36} = P(A)P(B) = \frac{6}{36} \cdot \frac{18}{36}$$

$\Rightarrow A$  and  $B$  are independent.