

# Review of Last Lecture

## Key concepts and/or techniques:

1. Half-unit correction for continuity

$$P(Y = k) \approx P(k - \frac{1}{2} < Y < k + \frac{1}{2})$$

2. Chebyshev's inequality: If the RV  $X$  has a finite mean  $\mu$  and finite nonzero variance  $\sigma^2$ , then for every  $k \geq 1$ ,

$$P(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2}$$

3. Convergence of sequence of random variables is defined through convergence of sequence of numbers, i.e.,

$$\lim_{n \rightarrow \infty} a_n = a$$

- ▶ convergence in distribution ( $Z_n \xrightarrow{d} Z$ ):  $a_n = P(Z_n \leq z)$  and  $a = P(Z \leq z)$ , where  $P(Z \leq z)$  is continuous
- ▶ convergence in probability ( $Z_n \xrightarrow{p} Z$ ):  $a_n = P(|Z_n - Z| \geq \varepsilon)$  and  $a = 0$ , for any  $\varepsilon > 0$

# Review of Last Lecture

## Half-unit correction for continuity

Now, if  $Y = \sum_{i=1}^n X_i$ , where  $X_1, \dots, X_n$  are i.i.d. random sample drawn from discrete distributions with mean  $\mu$  and variance  $\sigma^2$ .

$$P(Y = k) \approx P(k - \frac{1}{2} < Y < k + \frac{1}{2})$$

discrete RV

approximate by continuous RV

pmf  $f(y)$

by CLT for large  $n$ ,  $Y$  can be approximated by  $N(n\mu, n\sigma^2)$  in the sense that the pdf of the normal distribution is close to the histogram of  $Y$

hard to calculate

easy to calculate

# Review of Last Lecture

## [Chebyshev's inequality]

If the RV  $X$  has a finite mean  $\mu$  and finite nonzero variance  $\sigma^2$ , then for every  $k \geq 1$ ,

$$P(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2}$$

If  $\varepsilon = k\sigma$ , then

$$P(|X - \mu| \geq \varepsilon) \leq \frac{\sigma^2}{\varepsilon^2}$$

# Review of Last Lecture

## Definition

A sequence of RVs  $Z_1, Z_2, \dots$ , is said to converge in probability to a RV  $Z$ , often denoted by,  $Z_n \xrightarrow{P} Z$ , if for any  $\varepsilon > 0$ ,

$$\lim_{n \rightarrow \infty} P(|Z_n - Z| \geq \varepsilon) = 0.$$

# STA2001 Probability and Statistics I

## Lecture 25

Tianshi Chen

The Chinese University of Hong Kong, Shenzhen

# Theorem [Law of Large Number]

## Theorem (Law of Large Numbers)

Let  $X_1, X_2, \dots, X_n$  be a random sample of size  $n$  drawn from a distribution with finite mean  $\mu$  and finite nonzero variance, and let  $\bar{X}$  be the sample mean. Then  $\bar{X}$  converges in probability to  $\mu$ , i.e., for any  $\varepsilon > 0$ ,

$$\lim_{n \rightarrow \infty} P(|\bar{X} - \mu| \geq \varepsilon) = 0$$

# Proof of Law of Large Number

Note that

$$E(\bar{X}) = \mu, \quad \text{Var}(\bar{X}) = \frac{1}{n}\sigma^2$$

By the Chebyshev's inequality, i.e., Corollary 5.8-1, for every  $\varepsilon > 0$ .

$$P(|\bar{X} - \mu| \geq \varepsilon) \leq \frac{1}{\varepsilon^2} \frac{\sigma^2}{n}$$

# Proof of Law of Large Number

Taking limits on both sides yield

$$0 \leq \lim_{n \rightarrow \infty} P(|\bar{X} - \mu| \geq \varepsilon) \leq \lim_{n \rightarrow \infty} \frac{\sigma^2}{\varepsilon^2 n} = 0$$

$$\Rightarrow \lim_{n \rightarrow \infty} P(|\bar{X} - \mu| \geq \varepsilon) = 0$$

or equivalently  $\lim_{n \rightarrow \infty} P(|\bar{X} - \mu| < \varepsilon) = 1$



## Section 5.9 Limiting Moment Generating Functions

# Motivation

Binomial distribution  $b(n, p)$  can be approximated by the Poisson distribution with  $\lambda = np$  when  $n$  is large and  $p$  is fairly small:

- ▶ the approximation is good if  $n \geq 20$  and  $p \leq 0.05$
- ▶ the approximation is very good if  $n \geq 100$  and  $p \leq 0.1$
- ▶ the approximation becomes better with larger  $n$  and smaller  $p$

## Example 1, page 227

### Question

Let  $Y \sim b(50, 1/25)$ . Q:  $P(Y \leq 1)$ ?

## Example 1, page 227

### Question

Let  $Y \sim b(50, 1/25)$ . Q:  $P(Y \leq 1)$ ?

1. By definition,

$$\begin{aligned} P(Y \leq 1) &= P(Y = 0) + P(Y = 1) \\ &= \left(\frac{24}{25}\right)^{50} + 50 \left(\frac{1}{25}\right) \left(\frac{24}{25}\right)^{49} = 0.4 \end{aligned}$$

2. By approximation with Poisson distribution  $\lambda = np = 2$

$$P(Y \leq 1) \approx \frac{2^0 e^{-2}}{0!} + \frac{2^1 e^{-2}}{1!} = 3e^{-2} = 0.406$$

# Why? and an Interesting Observation

First, recall the mgf of  $b(n, p)$  is  $M(t) = (1 - p + pe^t)^n$ .

Then, we will consider the limit of  $M(t) = (1 - p + pe^t)^n$  as  $n \rightarrow \infty$  such that  $np = \lambda$  is a constant.

$$M(t) = \left(1 - \frac{\lambda}{n} + \frac{\lambda}{n}e^t\right)^n = \left[1 + \frac{\lambda(e^t - 1)}{n}\right]^n$$

Since  $\lim_{n \rightarrow \infty} \left(1 + \frac{b}{n}\right)^n = e^b$

$$\lim_{n \rightarrow \infty} M(t) = e^{\lambda(e^t - 1)} \rightarrow \text{mgf for Poisson distribution}$$

## Theorem 5.9-1, page 226

### Limiting mgf technique

Let  $\{M_n(t)\}_{n=1}^{\infty}$  be a sequence of mgfs for  $t$  in an open interval around  $t = 0$ . If  $\lim_{n \rightarrow \infty} M_n(t) = M(t)$ , for  $t$  in the open interval around  $t = 0$ . Then the sequence of RVs

$$Z_n \xrightarrow{d} Z,$$

where  $M_n(t)$  and  $M(t)$  are mgfs of  $Z_n$  and  $Z$ , respectively.

$$\lim_{n \rightarrow \infty} M_n(t) = M(t)$$



$$Z_n \xrightarrow{d} Z$$

# Convergence of $b(n, p)$

$b(n, p)$  converges in distribution to

- ▶ a Poisson distribution according to the limiting mgf technique

# Convergence of $b(n, p)$

$b(n, p)$  converges in distribution to

- ▶ a Poisson distribution according to the limiting mgf technique
- ▶ a standard normal distribution in some sense according to CLT

How to understand?



# Convergence of $b(n, p)$

$b(n, p)$  converges in distribution to

- ▶ a Poisson distribution according to the limiting mgf technique
- ▶ a standard normal distribution in some sense according to CLT

How to understand?

## Convergence of $b(n, p)$

1. as  $n \rightarrow \infty$  with  $\lambda = np$  being a constant, and let  $Z_n \sim b(n, p)$  and  $Z \sim \text{Poisson}(\lambda)$ , then  $Z_n \xrightarrow{d} Z$ .

2. as  $n \rightarrow \infty$  with  $p$  being a constant, and let  $Z_n \sim b(n, p)$  and  $Z \sim N(0, 1)$ , then

$$\frac{Z_n/n - p}{\sqrt{p(1-p)/n}} \xrightarrow{d} Z$$

# Proof of CLT

## CLT

Let  $\bar{X}$  be the sample mean of the random sample of size  $n$ ,  $X_1, X_2, \dots, X_n$  from a distribution with a finite mean  $\mu$  and a finite nonzero variance  $\sigma^2$ , then as  $n \rightarrow \infty$ , the random variable  $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$  converge in distribution to  $N(0, 1)$ .

The idea of the proof:

1. Let

$$W_n = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}},$$

and then show the mgf of  $W_n$ , say  $M_n(t)$ , converges to the mgf of  $N(0, 1)$  for  $t$  in an open interval around  $t = 0$

2. By Theorem 5.9-1,  $W_n \xrightarrow{d} N(0, 1)$

## Proof of CLT, page 208

$$\begin{aligned} E[e^{tW_n}] &= E \left\{ \exp \left[ t \frac{\frac{1}{n} \sum_{i=1}^n (X_i - n\mu)}{\sigma/\sqrt{n}} \right] \right\} \\ &= E \left\{ \exp \left[ \frac{t}{\sqrt{n}} \frac{\sum_{i=1}^n (X_i - n\mu)}{\sigma} \right] \right\} \\ &= E \left\{ \exp \left( \frac{t}{\sqrt{n}} \frac{X_1 - \mu}{\sigma} \right) \cdots \exp \left( \frac{t}{\sqrt{n}} \frac{X_n - \mu}{\sigma} \right) \right\} \\ &= E \left\{ \exp \left( \frac{t}{\sqrt{n}} \frac{X_1 - \mu}{\sigma} \right) \right\} \cdots E \left\{ \exp \left( \frac{t}{\sqrt{n}} \frac{X_n - \mu}{\sigma} \right) \right\} \\ &\quad \text{[independence]} \end{aligned}$$

# Proof of CLT, page 208

Let

$$Z_i = \frac{X_i - \mu}{\sigma}, \quad i = 1, \dots, n$$

Then  $Z_1, \dots, Z_n$  are i.i.d..

Let

$$M(t) = E \{ \exp(tZ_i) \}, \quad |t| < h$$

be the common mgf for  $Z_i, i = 1, \dots, n$ .

Then

$$E[e^{tW_n}] = \left[ M\left(\frac{t}{\sqrt{n}}\right) \right]^n, \quad \left| \frac{t}{\sqrt{n}} \right| < h.$$

## Proof of CLT, page 208

Now consider  $M(t)$ . Actually,  $Z_1, \dots, Z_n$  are i.i.d. with mean 0 and variance 1. Then

$$M(0) = 1, M'(0) = 0, M''(0) = 1.$$

By using Taylor's expansion, there exists  $|t_1| \leq |t|$  such that

$$\begin{aligned} M(t) &= M(0) + M'(0)t + \frac{1}{2}M''(t_1)t^2 \\ &= 1 + \frac{1}{2}M''(t_1)t^2 = 1 + \frac{1}{2}t^2 + \frac{1}{2}t^2[M''(t_1) - 1]. \end{aligned}$$

# Proof of CLT, page 208

Then

$$E(e^{tW_n}) = \left[ M\left(\frac{t}{\sqrt{n}}\right) \right]^n = \left[ 1 + \frac{1}{2} \frac{t^2}{n} + \frac{1}{2} \frac{t^2}{n} [M''(t_1) - 1] \right]^n,$$
$$\left| \frac{t}{\sqrt{n}} \right| < h, \quad |t_1| \leq \frac{|t|}{\sqrt{n}}.$$

Since  $M''(t)$  is continuous at  $t = 0$  and as  $n \rightarrow \infty$ ,  $t_1 \rightarrow 0$ ,

$$\lim_{n \rightarrow \infty} M''(t_1) - 1 = 1 - 1 = 0.$$

Note that

$$\lim_{n \rightarrow \infty} \left( 1 + \frac{b}{n} \right)^n = e^b$$

# Proof of CLT, page 208

we have

$$\begin{aligned}\lim_{n \rightarrow \infty} E(e^{tW_n}) &= \lim_{n \rightarrow \infty} \left[ 1 + \frac{1}{2} \frac{t^2}{n} + \frac{1}{2} \frac{t^2}{n} [M''(t_1) - 1] \right]^n \\ &= e^{\frac{t^2}{2}} \rightarrow \text{mgf of } N(0, 1)\end{aligned}$$

By Theorem 5.9-7,

$$W_n = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \xrightarrow{d} N(0, 1)$$