

## STA2001 Assignment 10

1. (5.4-22). Let  $X_1$  and  $X_2$  be two independent random variables. Let  $X_1$  and  $Y = X_1 + X_2$  be  $\chi^2(r_1)$  and  $\chi^2(r)$ , respectively, where  $r_1 < r$ .

- (a) Find the mgf of  $X_2$ .  
 (b) What is its distribution?

**Solution:**

- (a) We use the mgfs of  $Y$  and  $X_1$  to compute it,

$$\begin{aligned} \frac{1}{(1-2t)^{r/2}} &= E(e^{tY}) \\ &= E(e^{tX_1} e^{tX_2}) \\ &= E(e^{tX_1}) E(e^{tX_2}) \\ &= \frac{1}{(1-2t)^{r_1/2}} M_{X_2}(t) \end{aligned}$$

Solve this equation yields

$$M_{X_2}(t) = \frac{1}{(1-2t)^{(r-r_1)/2}}, \quad \forall t < 1/2$$

- (b) From the answer to (a), we know that  $X_2 \sim \chi^2(r - r_1)$ .

2. (5.4-23). Let  $X$  be  $N(0, 1)$ . Use the mgf technique to show that  $Y = X^2$  is  $\chi^2(1)$ . Hint: Evaluate the integral representing  $E(e^{tX^2})$  by writing  $w = x\sqrt{1-2t}$ .

**Solution:**

$$\begin{aligned} E(e^{tY}) &= E(e^{tX^2}) = \int_{-\infty}^{+\infty} e^{tx^2} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} e^{tx^2 - x^2/2} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} e^{-(x\sqrt{1-2t})^2/2} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} e^{-w^2/2} \frac{1}{\sqrt{1-2t}} dw \\ &= \frac{1}{\sqrt{1-2t}} \int_{-\infty}^{+\infty} \frac{1}{\sqrt{2\pi}} e^{-w^2/2} dw \\ &= \frac{1}{\sqrt{1-2t}}, \quad t < 1/2 \end{aligned}$$

where we have used change of variable at the 4th equality and the final result is obtained by realizing that the last integral is integrating a standard normal pdf.

This shows that  $Y$  has a  $\chi^2(1)$  distribution.

3. (5.5-4). Let  $X$  equal the weight of the soap in a 6-pound box. Assume that the distribution of  $X$  is  $N(6.05, 0.0004)$ .

- (a) Find  $P(X < 6.0171)$ .  
 (b) If nine boxes of soap are selected at random from the production line, find the probability that

at most two boxes weigh less than 6.0171 pounds each. Hint: Let  $Y$  equal the number of boxes that weigh less than 6.0171 pounds.

(c) Let  $\bar{X}$  be the sample mean of the nine boxes. Find  $P(\bar{X} \leq 6.035)$ .

**Solution:**

(a)

$$P(X < 6.071) = P\left(\frac{X - 6.05}{0.02} < \frac{6.0171 - 6.05}{0.02}\right) = P(Z < -1.645) = 0.05$$

(b) Let  $W$  be the number of boxes that weight less than 6.0171 pounds. Then  $W \sim b(9, 0.05)$ , and

$$\begin{aligned} P(W \leq 2) &= \sum_{k=0}^2 \binom{9}{k} (0.05)^k (1 - 0.05)^{9-k} \\ &= 0.9916 \end{aligned}$$

(c) Let  $\bar{X}$  be the mean of a random sample of the boxes with size  $n = 9$ . So  $\bar{X} \sim N(6.05, 0.02^2/n)$ , the probability is given by

$$P(\bar{X} \leq 6.035) = P\left(Z \leq \frac{6.035 - 6.05}{0.02/\sqrt{9}}\right) = 0.0122$$

4. (5.5-14). Let  $T$  have a  $t$  distribution with  $r$  degrees of freedom. Show that  $E(T) = 0$  provided that  $r \geq 2$ , and  $\text{Var}(T) = r/(r-2)$  provided that  $r \geq 3$ , by first finding  $E(Z)$ ,  $E(1/\sqrt{U})$ ,  $E(Z^2)$ , and  $E(1/U)$ .

**Solution:**

Recall Theorem 5.5-3, a random variable  $T$  following  $t$  distribution with degree of freedom  $r$  is constructed by

$$T = \frac{Z}{\sqrt{U/r}}$$

where  $Z \sim N(0, 1)$ ,  $U \sim \chi^2(r)$ , and  $Z$  and  $U$  are independent.

Therefore,  $E(Z) = 0$  and  $E(Z^2) = 1$ . Since  $U \sim \chi^2(r)$ , it follows that

$$\begin{aligned} E\left(1/\sqrt{U}\right) &= \int_0^\infty \frac{1}{\sqrt{u}} \frac{1}{\Gamma(r/2)2^{r/2}} u^{r/2-1} e^{-u/2} du \\ &= \frac{\Gamma[(r-1)/2]}{\sqrt{2}\Gamma(r/2)} \int_0^\infty \frac{1}{\Gamma[(r-1)/2]2^{(r-1)/2}} u^{(r-1)/2-1} e^{-u/2} du \\ &= \frac{\Gamma[(r-1)/2]}{\sqrt{2}\Gamma(r/2)}, \quad r \geq 2 \end{aligned}$$

Note that the last integral is equal to one because the integrand is the pdf of a  $\chi^2(r-1)$  random variable.

To find  $E(1/U)$ , we use a similar approach

$$\begin{aligned} E[1/U] &= \int_0^\infty \frac{1}{u} \frac{1}{\Gamma(r/2)2^{r/2}} u^{r/2-1} e^{-u/2} du \\ &= \frac{\Gamma[(r-2)/2]}{2\Gamma(r/2)} \int_0^\infty \frac{1}{\Gamma[(r-2)/2]2^{(r-2)/2}} u^{(r-2)/2-1} e^{-u/2} du \\ &= \frac{\Gamma(r/2-1)}{2\Gamma(r/2-1)(r/2-1)} \\ &= \frac{1}{r-2}, \quad r \geq 3 \end{aligned}$$

Note that the last integral is equal to one because the integrand is the pdf of a  $\chi^2(r-2)$  random variable.

Hence, we could compute the mean and variance as follows

$$\begin{aligned} E(T) &= E \left[ \frac{Z}{\sqrt{U/r}} \right] \\ &= E[Z] \cdot E[1/\sqrt{U/r}] \\ &= 0 \times \left[ \frac{\sqrt{r}\Gamma[(r-1)/2]}{\sqrt{2}\Gamma(r/2)} \right] \\ &= 0, \quad r \geq 2 \end{aligned}$$

$$\begin{aligned} \text{Var}(T) &= E(T^2) \\ &= E[Z^2] E[r/U] \\ &= 1 \times \left( r \times \frac{1}{r-2} \right) \\ &= \frac{r}{r-2}, \quad r \geq 3 \end{aligned}$$

5. (5.5-16). Let  $n = 9$  in the  $T$  statistic defined in Equation 5.5-2. That is,

$$T = \frac{\frac{\bar{X} - \mu}{\sigma/\sqrt{n}}}{\sqrt{\frac{(n-1)S^2}{\sigma^2}/(n-1)}} = \frac{\bar{X} - \mu}{S/\sqrt{n}}.$$

- (a) Find  $t_{0.025}$  so that  $P(-t_{0.025} \leq T \leq t_{0.025}) = 0.95$ .  
 (b) Solve the inequality  $[-t_{0.025} \leq T \leq t_{0.025}]$  so that  $\mu$  is in the middle.

**Solution:**

(a) From Equation 5.5-2 on the textbook, we know that  $T$  have a student  $t$  distribution with degree of freedom  $r$ , which is given by

$$r = n - 1 = 8$$

where  $n$  is the sample size.

Let's consider a general case by replacing  $t_{0.025}$  with  $t$  (as the hint given by the subscript is too obvious),

$$\begin{aligned} P(-t \leq T \leq t) &= P(T \leq t) - P(T \leq -t) \\ &= P(T \leq t) - (1 - P(T \leq t)) \\ &= 2P(T \leq t) - 1 \\ &= 0.95 \end{aligned}$$

which means we need to find  $t$  such that  $P(T \leq t) = 0.975$ .

In this equation, since  $r = 8$ , the value of  $t$  is exactly the same as  $t_{0.025}(8)$  that can be found in the  $t$  distribution table,

$$t = t_{0.025}(8) = 2.306$$

(b) By Equation 5.5-2 on the textbook,  $T = \frac{\bar{X} - \mu}{S/\sqrt{n}}$ , so we rewrite the inequality as

$$-t_{0.025} \leq \frac{\bar{X} - \mu}{S/\sqrt{n}} \leq t_{0.025}$$

Rearranging the terms in the inequality,

$$\bar{X} - t_{0.025} \frac{S}{\sqrt{n}} \leq \mu \leq \bar{X} + t_{0.025} \frac{S}{\sqrt{n}}$$

Note that  $n = 9$  and  $t_{0.025} = t_{0.025}(8) = 2.306$  in this question, so finally we have

$$\bar{X} - 0.7687 \cdot S \leq \mu \leq \bar{X} + 0.7687 \cdot S$$

6. (5.6-5). Let  $X_1, X_2, \dots, X_{18}$  be a random sample of size 18 from a chi-square distribution with  $r = 1$ . Recall that  $\mu = 1$  and  $\sigma^2 = 2$ .

(a) How is  $Y = \sum_{i=1}^{18} X_i$  distributed?

(b) Using the result of part (a), we see from Table IV in Appendix B that

$$P(Y \leq 9.390) = 0.05$$

and

$$P(Y \leq 34.80) = 0.99$$

Compare these two probabilities with the approximations found with the use of the central limit theorem

**Solution:**

(a)  $Y$  follows a chi-square distribution with 18 degree of freedom.

(b) Each  $X_i$  has  $\mu = 1$  and  $\sigma^2 = 2$  and they are i.i.d (since they forms a random sample), so for  $Y = \sum_{i=1}^{18} X_i$  we have

$$E(Y) = 18 \times 1 = 18, \quad \text{Var}(Y) = 18 \times 2 = 36$$

We use  $N(18, 36)$  to approximate the following

$$P(Y \leq 9.390) \approx P\left(Z \leq \frac{9.390 - 18}{\sqrt{36}}\right) = 0.0756$$

$$P(Y \leq 34.8) \approx P\left(Z \leq \frac{34.8 - 18}{\sqrt{36}}\right) = 0.9974$$

7. (5.6-8). Let  $X$  equal the weight in grams of a miniature candy bar. Assume that  $\mu = E(X) = 24.43$  and  $\sigma^2 = \text{Var}(X) = 2.20$ . Let  $\bar{X}$  be the sample mean of a random sample of  $n = 30$  candy bars. Find

(a)  $E(\bar{X})$ .

(b)  $\text{Var}(\bar{X})$ .

(c)  $P(24.17 \leq \bar{X} \leq 24.82)$ , approximately.

**Solution:**

Note that  $X_1, X_2, \dots, X_{30}$  is a random sample of size 30. So they are i.i.d.

(a)

$$E(\bar{X}) = E\left(\frac{1}{30} \sum_{i=1}^{30} X_i\right) = \frac{1}{30} \sum_{i=1}^{30} E(X_i) = \mu = 24.43$$

(b)

$$\text{Var}(\bar{X}) = \text{Var}\left(\frac{1}{30} \sum_{i=1}^{30} X_i\right) = \frac{1}{30^2} \sum_{i=1}^{30} \text{Var}(X_i) = \frac{1}{30^2} \times 30 \times \sigma^2 = \frac{2.20}{30} = 0.0733$$

(c) Let  $Z \sim N(0, 1)$ . Using central limit theorem, we can approximately compute

$$\begin{aligned} P(24.17 \leq \bar{X} \leq 24.82) &\approx P\left(\frac{24.17 - 24.43}{\sqrt{0.0733}} \leq Z \leq \frac{24.82 - 24.43}{\sqrt{0.0733}}\right) \\ &= P(-0.96 \leq Z < 1.44) \\ &= 0.7566 \end{aligned}$$

8. (5.6-14). Suppose that the sick leave taken by the typical worker per year has  $\mu = 10$ ,  $\sigma = 2$ , measured in days. A firm has  $n = 20$  employees. Assuming independence, how many sick days should the firm budget if the financial officer wants the probability of exceeding the number of days budgeted to be less than 20%?

**Solution:**

Let  $X_i$  be the length of sick leave (measure in days) taken by a typical worker per year. So we have  $E(X_i) = 10$  and  $\text{Var}(X_i) = 4$ .

Let  $Y = \sum_{i=1}^{20} X_i$ , and note that  $X_1, X_2, \dots, X_{20}$  are i.i.d, we have

$$E(Y) = 20 \times 10 = 200$$

$$\text{Var}(Y) = 20 \times 4 = 80$$

Since the distribution of  $Y$  is unknown but it's a sum of i.i.d random variables, we could use the central limit theorem to approximate

$$\begin{aligned} P(Y > y) &= P\left(\frac{Y - 200}{\sqrt{80}} > \frac{y - 200}{\sqrt{80}}\right) \\ &= 1 - P\left(\frac{Y - 200}{\sqrt{80}} \leq \frac{y - 200}{\sqrt{80}}\right) \\ &\approx 1 - P\left(Z \leq \frac{y - 200}{\sqrt{80}}\right) \end{aligned}$$

Setting  $1 - P\left(Z \leq \frac{y - 200}{\sqrt{80}}\right) = 0.2$  and checking the normal distribution table we have

$$\frac{y - 200}{\sqrt{80}} = 0.842 \Rightarrow y = 207.5$$

As we want the probability less than 0.2 and note that  $Y$  is measured in days, we take  $y = 208$ .

9. Let  $\{X_n\}_{n=1}^{\infty}$  be i.i.d. Bernoulli random variables with parameter  $1/2$ . Let

$$Y_n := \frac{\sum_{i=1}^{n^2} X_i - (n^2/2)}{(n/2)}.$$

Let  $\mathbb{Q}$  be the set of rational numbers.

- (i) Show that  $Y_n$  converge in distribution to  $Y \sim \mathcal{N}(0, 1)$ . Hint: consider  $Y_{\sqrt{n}}$ .
- (ii) Find  $\Pr(Y_n \in \mathbb{Q})$  and  $\Pr(Y \in \mathbb{Q})$ . Does  $\Pr(Y_n \in \mathbb{Q})$  converge to  $\Pr(Y \in \mathbb{Q})$ ?

**Solution:**

Observe that  $Y_n = Z_{n^2}$ , where

$$Z_n := \frac{\sum_{i=1}^n X_i - n/2}{\sqrt{n}/2} = \frac{\sum_{i=1}^n (X_i - \mathbb{E}[X_i])}{\sqrt{n} \text{Var}[X_i]}$$

By Central Limit Theorem,  $Z_n \xrightarrow{D} Y$ . Hence, as a subsequence of  $Z_n, Y_n \xrightarrow{D} Y$ .

(ii) Bernoulli random variables are integer valued, hence  $\frac{\sum_{i=1}^{n^2} X_i - (n^2/2)}{(n/2)}$  is always a rational number. We have  $\Pr(Y_n \in \mathbb{Q}) = 1$  for all  $n$ .

$Y \sim \mathcal{N}(0, 1)$  is a continuous random variable. For any rational number  $q$ , we have  $\Pr(Y = q) = 0$ .

The set of rational numbers is countable. Hence by countable additivity of probability measures we have

$$\Pr(Y \in \mathbb{Q}) = \sum_{q \in \mathbb{Q}} \Pr(Y = q) = 0$$

We observe that  $\Pr(Y_n \in \mathbb{Q})$  does not converge to  $\Pr(Y \in \mathbb{Q})$ .

10. Suppose  $X$  is a continuous random variable with probability density function:

$$f(x) = \frac{x^2}{3},$$

where  $-1 \leq x \leq 2$ . Define a random variable  $Y = X^2$ .

- What is the pdf of  $Y$ ? Hint: The function  $Y = X^2$  is not one-to-one mapping for all  $X$ .
- Use the Central Limit Theorem to find the probability  $P(\sum_{i=1}^{175} Y_i \geq 385)$ , where  $Y_1, \dots, Y_{175}$  is a random sample of size 175 from the common distribution of  $Y$ .
- Suppose that two random numbers 0.8 and 0.2 are randomly generated from the uniform distribution  $U(0, 1)$ . Transform the two random numbers 0.8 and 0.2 into two random numbers  $y_1$  and  $y_2$  drawn from the distribution of  $Y$ .

**Solution:**

- The sample space of  $Y$ ,  $\bar{S}_Y$  is  $[0, 4]$ . First consider  $0 \leq y \leq 1$ ,  $P(Y < y) = P(-\sqrt{y} < X < \sqrt{y}) = F_X(\sqrt{y}) - F_X(-\sqrt{y})$ . Then

$$\begin{aligned} f_Y(y) &= f_X(\sqrt{y}) \frac{1}{2\sqrt{y}} - f_X(-\sqrt{y}) \frac{-1}{2\sqrt{y}} \\ &= \frac{y}{3} \times \frac{1}{2\sqrt{y}} - \frac{y}{3} \times \frac{-1}{2\sqrt{y}} \\ &= \frac{\sqrt{y}}{3} \end{aligned}$$

When  $1 < y \leq 4$ , the function is one-to-one mapping, and can easily check that  $f_Y(y) = \frac{\sqrt{y}}{6}$ .

- 

$$\begin{aligned} E(Y) &= \int_0^1 \frac{1}{3} y^{3/2} dy + \int_1^4 \frac{1}{6} y^{3/2} dy = 11/5 \\ E(Y^2) &= \int_0^1 \frac{1}{3} y^{5/2} dy + \int_1^4 \frac{1}{6} y^{5/2} dy = 43/7 \\ Var(Y) &= \frac{1922}{175} \end{aligned}$$

$$\begin{aligned}
P\left(\sum_{i=1}^{175} Y_i \geq 385\right) &= P\left(\frac{\sum_{i=1}^{175} Y_i - 175 \times \frac{11}{5}}{\sqrt{175 \times \frac{1922}{175}}} \geq \frac{385 - 175 \times \frac{11}{5}}{\sqrt{175 \times \frac{1922}{175}}} = 0\right) \\
&= 0.5
\end{aligned}$$

(c) From part(a) we know that the cdf of  $Y$  is

$$F_Y(y) = \begin{cases} \left(\frac{2y}{9}\right)^{3/2}, & 0 \leq y \leq 1 \\ \frac{1}{9} + \frac{1}{9}y^{3/2}, & 1 < y \leq 4 \end{cases} \quad (1)$$

and the inverse of it would be

$$F_Y^{-1}(x) = \begin{cases} \left(\frac{9x}{2}\right)^{2/3}, & 0 \leq x \leq \frac{2}{9} \\ (9x - 1)^{2/3}, & \frac{2}{9} < x \leq 1 \end{cases} \quad (2)$$

Hence, the two random numbers  $y_1 = (9 \times 0.8 - 1)^{2/3} = (31/5)^{2/3} = 3.375$  and  $y_2 = (9 \times 0.2/2)^{2/3} = (0.9)^{2/3} = 0.932$ .