DDA3020 Machine Learning
Lecture 06 Logistic Regression

Haizhou Li
School of Data Science, CUHK-SZ

February 23/28, 2022

Outline

1. Review of last week
2. Classification and representation
3. Logistic regression
4. Regularized logistic regression
5. Probabilistic perspective of logistic regression
6. Summary: linear regression vs. logistic regression

1 Review of last week

2 Classification and representation

3 Logistic regression

4 Regularized logistic regression

5 Probabilistic perspective of logistic regression

6 Summary: linear regression vs. logistic regression

Linear regression: deterministic perspective

- **Linear hypothesis function**: $f_{\mathbf{w},b}(\mathbf{x}) = \mathbf{x}^\top \mathbf{w} + b$, or, simply $f_{\mathbf{w}}(\mathbf{x}) = \mathbf{x}^\top \mathbf{w}$ by concatenating b and $\mathbf{w}$ together and augmenting $\mathbf{x}$ to $[1; \mathbf{x}]$
- **Linear regression** by minimizing residual sum of squares (RSS):

$$\mathbf{w}^* = \arg\min_{\mathbf{w}} J(\mathbf{w}), \text{ where } J(\mathbf{w}) = \frac{1}{2}\sum_{i=1}^{m}(\mathbf{x}_i^\top \mathbf{w} - y_i)^2 = \frac{1}{2}\|\mathbf{X}\mathbf{w} - \mathbf{y}\|^2$$

- **Two solutions**:
  - Closed-form solution: $\mathbf{w}^* = \left(\mathbf{X}^\top \mathbf{X}\right)^{-1} \mathbf{X}^\top \mathbf{y}$
  - Gradient descent: $\mathbf{w} \leftarrow \mathbf{w} - \alpha \mathbf{X}^\top (\mathbf{X}\mathbf{w} - \mathbf{y})$, for multiple iterations until convergence

Linear regression: probabilistic perspective

- We assume that: $y = \mathbf{w}^\top \mathbf{x} + e$, where $e \sim \mathcal{N}(0, \sigma^2)$ is called **observation noise** or **residual error**
- y is also a random variable, and its conditional probability is

$$p(y|\mathbf{x}, \mathbf{w}) = \mathcal{N}(\mathbf{w}^\top \mathbf{x}, \sigma^2)$$

- **Maximum log-likelihood estimation**:

$$\mathbf{w}_{MLE} = \arg\max_{\mathbf{w}} \log \mathcal{L}(\mathbf{w}|D) = \arg\max_{\mathbf{w}} \log \Big(\prod_i^m p(y_i|\mathbf{x}_i, \mathbf{w}) \Big) \tag{1}$$

$$= \arg\max_{\mathbf{w}} \sum_i^m \log p(y_i|\mathbf{x}_i, \mathbf{w}) = \arg\max_{\mathbf{w}} \sum_i^m \log \mathcal{N}(\mathbf{w}^\top \mathbf{x}_i, \sigma^2) \tag{2}$$

$$= \arg\max_{\mathbf{w}} -\log(\sigma^m (2\pi)^{\frac{m}{2}}) - \frac{1}{2\sigma^2} \sum_i^m (y_i - \mathbf{w}^\top \mathbf{x}_i)^2 \tag{3}$$

$$= \arg\min_{\mathbf{w}} \frac{1}{2} \sum_i^m (y_i - \mathbf{w}^\top \mathbf{x}_i)^2, \tag{4}$$

Variants of linear regression

- **Ridge regression** to avoid over-fitting, through MAP estimation:

$$\mathbf{w}_{MAP} = \arg\max_{\mathbf{w}} \sum_{i=1}^{m} \log p(y_i|\mathbf{x}_i, \mathbf{w}) + \log p(\mathbf{w}) \tag{5}$$

$$= \arg\max_{\mathbf{w}} \sum_{i=1}^{m} \log \mathcal{N}(\mathbf{w}^\top \mathbf{x}_i, \sigma^2) + \mathcal{N}(\mathbf{w}|\mathbf{0}, \tau^2\mathbf{I}) \tag{6}$$

$$\equiv \arg\min_{\mathbf{w}} \sum_{i=1}^{m} (\mathbf{w}^\top \mathbf{x}_i - y_i)^2 + \lambda \|\mathbf{w}\|_2^2. \tag{7}$$

- **Polynomial regression**: linear model with basis expansion $\phi(\mathbf{x})$

$$f_{\mathbf{w},b}(\mathbf{x}) = b + \sum_{i=1}^{d} w_i x_i + \sum_{i=1}^{d}\sum_{j=1}^{d} w_{ij} x_i x_j + \sum_{i=1}^{d}\sum_{j=1}^{d}\sum_{k=1}^{d} w_{ijk} x_i x_j x_k + \dots$$

$$= \phi(\mathbf{x})^\top \mathbf{w}, \tag{8}$$

$$\phi(\mathbf{x}) = [1, x_1, \dots, x_d, \dots, x_i x_j, \dots, x_i x_j x_k, \dots]^\top,$$

$$\mathbf{w} = [b, w_1, \dots, w_d, \dots, w_{ij}, \dots, w_{ijk}, \dots]^\top.$$

Variants of linear regression

- **Lasso regression** to obtain sparse model,

$$\mathbf{w}_{MAP} = \arg\max_{\mathbf{w}} \sum_{i}^{m} \log \mathcal{N}(\mathbf{w}^\top \mathbf{x}_i, \sigma^2) + \text{Lap}(\mathbf{w}|\mathbf{0}, b) \tag{9}$$

$$= \arg\min_{\mathbf{w}} \sum_{i=1}^{m} (\mathbf{w}^\top \mathbf{x}_i - y_i)^2 + \lambda \|\mathbf{w}\|_1. \tag{10}$$

ℓ_1


- **Robust regression** for data with outliers:

$$\mathbf{w}_{MLE} = \arg\min_{\mathbf{w}} \sum_{i=1}^{m} |\mathbf{w}^\top \mathbf{x}_i - y_i| \tag{11}$$

Summary of different linear regressions

Note that the uniform distribution will not change the mode of the likelihood. Thus, MAP estimation with a uniform prior corresponds to MLE.

$p(y\|\mathbf{x},\mathbf{w})$	$p(\mathbf{w})$	regression method
Gaussian	Uniform	Least squares
Gaussian	Gaussian	Ridge regression
Gaussian	Laplace	Lasso regression
Laplace	Uniform	Robust regression
~~Student~~	Uniform	~~Robust regression~~



1 Review of last week

2 Classification and representation

3 Logistic regression


4 Regularized logistic regression

5 Probabilistic perspective of logistic regression


6 Summary: linear regression vs. logistic regression

Classification

- Classification: classifying input data into discrete states
  - Email filtering: spam / not spam?
  - Weather forecast: sunny / not sunny?
  - Tumor: malignant / benign?
- The label $y \in \{0, 1\}$:
  - $y = 0$: negative class, *e.g.*, not spam, not sunny, benign
  - $y = 1$: positive class, *e.g.*, spam, sunny, malignant




Threshold classifier with linear regression



- We assume a linear hypothesis function $f_{\mathbf{w},b}(\boldsymbol{x}) = \mathbf{x}^\top \mathbf{w} + b$
- A simple threshold classifier with this hypothesis function is
  - If $f_{\mathbf{w},b}(\boldsymbol{x}) > 0.5$, then $y = 1$, *i.e.*, malignant tumor
  - If $f_{\mathbf{w},b}(\boldsymbol{x}) < 0.5$, then $y = 0$, *i.e.*, benign tumor

Threshold classifier with linear regression



- It seems that the simple threshold classifier with linear regression works well on this classification task
- However, if there is a positive sample with very large tumor size (plot above), what will happen?
- The hypothesis function will be **significantly changed**, causing that some positive samples are mis-classified as negative (not malignant). How to handle it? Adjusting the threshold value, or adopting robust linear regression.

阈值

Threshold classifier with linear regression

- But there is still something wired.
- Our goal is to predict $y \in \{0, 1\}$, but the prediction could be $f_{\mathbf{w},b}(\mathbf{x}) > 1$ or $f_{\mathbf{w},b}(\mathbf{x}) < 0$, which does not serve our purpose.
- A desired hypothesis function for this task should be $f_{\mathbf{w},b}(\mathbf{x}) \in [0, 1]$.

Threshold classifier with linear regression

Exercise: Which statements are true?

F. • If linear regression doesn't work well like the above example, feature scaling may help

特征缩放

both scale good & bad data ⇒ NOT improve seperation

F. • If the training set satisfies that all $y_i \in [0, 1]$ for all points $(\mathbf{x}_i, y_i)$, then the linear hypothesis function $f_{\mathbf{w},b}(\mathbf{x}) \in [0, 1]$ for all values of $\mathbf{x}_i$

construct a model for new data [No constrain]


• None of the above is correct

Hypothesis representation

- A desired hypothesis function for this task should be $f_{\mathbf{w},b}(\mathbf{x}) \in [0, 1]$
- To this end, we introduce a novel function, as follows:

$$f_{\mathbf{w},b}(\mathbf{x}) = g(\mathbf{w}^\top \mathbf{x}) \in [0, 1], \ g(z) = \frac{1}{1 + \exp(-z)},$$


where $g(\cdot)$ is called sigmoid function or logistic function (shown below)




Hypothesis representation


- **Interpretation of sigmoid/logistic function**
  - $f_{\mathbf{w},b}(\mathbf{x}) =$ estimated probability that $y = 1$ of input $\mathbf{x}$.
- For example (plot below), if $f_{\mathbf{w},b}(\mathbf{x}) = 0.8$, then it means that a patient with tumor size $\mathbf{x}$ has 80% chance of tumor being malignant. In this task, larger tumor size has larger chance/probability of being malignant tumor.
- Thus, we can say that

$$f_{\mathbf{w},b}(\mathbf{x}) = P(y = 1|\mathbf{x}; \mathbf{w}).$$





Decision boundary



- In logistic regression, we have

$$f_{\mathbf{w},b}(\mathbf{x}) = g(\mathbf{w}^\top \mathbf{x} + b) = P(y = 1|\mathbf{x}; \mathbf{w}) \in [0, 1], \quad g(z) = \frac{1}{1 + \exp(-z)}.$$


- Suppose that if $f_{\mathbf{w},b}(\mathbf{x}) \geq 0.5$, then we predict $y = 1$; if $f_{\mathbf{w},b}(\mathbf{x}) < 0.5$, then we predict $y = 0$
- Correspondingly, if $\mathbf{w}^\top \mathbf{x} + b \geq 0$, we predict $y = 1$; if $\mathbf{w}^\top \mathbf{x} + b < 0$, then we predict $y = 0$.

↓ still linear here

- It determines the decision boundary, which is the curve/hyper-plane corresponding to $f_{\mathbf{w},b}(\mathbf{x}) = 0.5$, or $\mathbf{w}^\top \mathbf{x} + b = 0$

Decision boundary

↳ still linear



↳ not linear

- $f_{\mathbf{w},b}(\mathbf{x}) = g(b + w_1 x_1 + w_2 x_2) = g(-3 + x_1 + x_2)$
- Predict $y = 1$ if $-3 + x_1 + x_2 \geq 0$ (plot above)

↳ $z \geq 0$ $g(z) = \frac{1}{1+\exp(-z)} > \frac{1}{2} \Rightarrow 1.$

Decision boundary

↳ now not linear




Figure: Non-linear decision boundary

- $f_{\mathbf{w},b}(\mathbf{x}) = g(b + w_1x_1 + w_2x_2 + w_3x_1^2 + w_4x_2^2) = g(-1 + x_1^2 + x_2^2)$
- Predict $y = 1$ if $-1 + x_1^2 + x_2^2 \geq 0$ (plot above)

1 Review of last week

2 Classification and representation

3 Logistic regression

4 Regularized logistic regression

5 Probabilistic perspective of logistic regression

6 Summary: linear regression vs. logistic regression

Cost function

- Training set: m training examples $\{(\mathbf{x}_i, y_i)\}_{i=1}^m$
- Hypothesis function: $f_{\mathbf{w},b}(\mathbf{x}) = g(\mathbf{w}^\top \mathbf{x} + b) = \frac{1}{1+\exp(-\mathbf{w}^\top \mathbf{x} - b)}$
- Cost function:
  - **Linear regression**: $J(\mathbf{w}) = \frac{1}{2m}\sum_{i=1}^m (f_{\mathbf{w},b}(\mathbf{x}_i) - y_i)^2 = \frac{1}{2m}\|\mathbf{X}\mathbf{w} - \mathbf{y}\|^2$, which is called ℓ_2 **loss** or **residual sum of squares**
  - It is convex *w.r.t.* $\mathbf{w}$ for linear regression
  - **Logistic regression**: If we adopt the same cost function for logistic regression, we have

$$J(\mathbf{w}) = \frac{1}{2m}\sum_i^m (g(\mathbf{w}^\top \boldsymbol{x}_i) - y_i)^2.$$

    (y_i: 0 or 1.)

    However, it is non-convex *w.r.t.* $\mathbf{w}$.

Exercise 1: Prove the ℓ_2 loss is convex *w.r.t.* $\mathbf{w}$ for linear regression.
Exercise 2: Prove the ℓ_2 loss is non-convex *w.r.t.* $\mathbf{w}$ for logistic regression.

Cost function

Exercise 1: Prove the ℓ_2 loss is convex *w.r.t.* $\mathbf{w}$ for linear regression.

Exercise 2: Prove the ℓ_2 loss is non-convex *w.r.t.* $\mathbf{w}$ for logistic regression.

for linear: $J = \frac{1}{2m}\|Xw - y\|^2$

$\frac{\partial}{\partial w} = \frac{1}{2m}(Xw - y)^T \cdot X^T$

$\frac{\partial^2}{\partial w^2} = \frac{1}{2m} X^T X > 0 \quad \therefore$ convex

for logistic: $J = \frac{1}{2m}\|g(w^T x) - y\|^2$

$\frac{\partial}{\partial w} = 2[g(w^T x) - y] \cdot \left[\frac{1}{1+\exp(-w^T x - b)}\right]^2 e^{\cdot} \cdot (-x)$

$\frac{\partial^2}{\partial w^2} =$

Cost function

- Cross-entropy: 互熵

$$H(p,q) = -\int_x p(x)\log(q(x))dx \quad \text{or} \quad -\sum_x p(x)\log(q(x)),$$

where $p(x), q(x)$ are **probability density functions** (PDF) of x if x is a continuous random variable, or, **probability mass functions** if x is a discrete random variable

- We set

$$\text{ground-truth posterior probability}: \ y(\mathbf{x}) = P(y=1|\mathbf{x}), \quad \text{— } p(x)$$
$$\text{predicted posterior probability}: \ f_{\mathbf{w},b}(\mathbf{x}) = P(y=1|\mathbf{x};\mathbf{w}). \quad \text{— } q(x)$$

- Cross-entropy loss:

$$\begin{aligned}
&\text{cost}\big(y(\mathbf{x}), f_{\mathbf{w},b}(\mathbf{x})\big) = H\big(y(\mathbf{x}), f_{\mathbf{w},b}(\mathbf{x})\big) \\
&= -P(y=1|\mathbf{x})\cdot\log P(y=1|\mathbf{x};\mathbf{w}) - P(y=0|\mathbf{x})\cdot\log P(y=0|\mathbf{x};\mathbf{w}) \\
&= \begin{cases} -\log(f_{\mathbf{w},b}(\mathbf{x})), & \text{if } y(\mathbf{x})=1 \\ -\log(1-f_{\mathbf{w},b}(\mathbf{x})), & \text{if } y(\mathbf{x})=0 \end{cases}
\end{aligned}$$


Cost function for logistic regression

- Cross-entropy (熵) loss:

$$\text{cost}\big(y(\mathbf{x}), f_{\mathbf{w},b}(\mathbf{x})\big) = \begin{cases} -\log(f_{\mathbf{w},b}(\mathbf{x})), & \text{if } y(\mathbf{x}) = 1 \\ -\log(1 - f_{\mathbf{w},b}(\mathbf{x})), & \text{if } y(\mathbf{x}) = 0 \end{cases}$$

$P(y=1|x;w)$

- For $y = 1$, if $f_{\mathbf{w},b}(\mathbf{x}) = 1$, *i.e.*, $P(y = 1|\mathbf{x}; \mathbf{w}) = 1$, then the prediction equals to the ground-truth label, the cost is 0.
- For $y = 1$, if $f_{\mathbf{w},b}(\mathbf{x}) \to 0$, *i.e.*, $P(y = 1|\mathbf{x}; \mathbf{w}) \to 0$, then it should be penalized with a very large cost. Here we have $\text{cost}(y(\mathbf{x}), f_{\mathbf{w},b}(\mathbf{x})) \to \infty$.




Cost function for logistic regression

- **Cross-entropy loss**:

$$\text{cost}(y(\mathbf{x}), f_{\mathbf{w},b}(\mathbf{x})) = \begin{cases} -\log(f_{\mathbf{w},b}(\mathbf{x})), & \text{if } y(\mathbf{x}) = 1 \\ -\log(1 - f_{\mathbf{w},b}(\mathbf{x})), & \text{if } y(\mathbf{x}) = 0 \end{cases}$$

- For $y = 0$, if $f_{\mathbf{w},b}(\mathbf{x}) = 0$, *i.e.*, $P(y = 1|\mathbf{x}; \mathbf{w}) = 0$, then the prediction equals to the ground-truth label, the cost is 0
- For $y = 0$, if $f_{\mathbf{w},b}(\mathbf{x}) \to 1$, *i.e.*, $P(y = 1|\mathbf{x}; \mathbf{w}) \to 0$, then it should be penalized with a very large cost. Here we have $\text{cost}(y(\mathbf{x}), f_{\mathbf{w},b}(\mathbf{x})) \to \infty$



loss : for 1 data point

cost : for all

Cost function for logistic regression

- **Cross-entropy loss:**

$$\text{cost}(y(\mathbf{x}), f_{\mathbf{w},b}(\mathbf{x})) = \begin{cases} -\log(f_{\mathbf{w},b}(\mathbf{x})), & \text{if } y(\mathbf{x}) = 1 \\ -\log(1 - f_{\mathbf{w},b}(\mathbf{x})), & \text{if } y(\mathbf{x}) = 0 \end{cases}$$

Exercise: Which states are true?

T • If $f_{\mathbf{w},b}(\mathbf{x}) = y$, then $\text{cost}(y(\mathbf{x}), f_{\mathbf{w},b}(\mathbf{x})) = 0$ for both $y = 0$ and $y = 1$

T • If $y = 0$, then $\text{cost}(y(\mathbf{x}), f_{\mathbf{w},b}(\mathbf{x})) \to \infty$ as $f_{\mathbf{w},b}(\mathbf{x}) \to 1$

F • If $y = 0$, then $\text{cost}(y(\mathbf{x}), f_{\mathbf{w},b}(\mathbf{x})) \to \infty$ as $f_{\mathbf{w},b}(\mathbf{x}) \to 0$

T • Regardless whether $y = 0$ or $y = 1$, if $f_{\mathbf{w},b}(\mathbf{x}) = 0.5$, then $\text{cost}(y(\mathbf{x}), f_{\mathbf{w},b}(\mathbf{x})) > 0$

Cost function of logistic regression

- **Cost function of logistic regression**

$$J(\mathbf{w}) = \frac{1}{m}\sum_{i=1}^{m} \text{cost}(y_i, f_{\mathbf{w},b}(\mathbf{x}_i)),$$

$$\text{cost}(y(\mathbf{x}), f_{\mathbf{w},b}(\mathbf{x})) = \begin{cases} -\log(f_{\mathbf{w},b}(\mathbf{x})), & \text{if } y(\mathbf{x}) = 1 \\ -\log(1 - f_{\mathbf{w},b}(\mathbf{x})), & \text{if } y(\mathbf{x}) = 0 \end{cases}$$

- The above cost function can be simplified as follows

$$J(\mathbf{w}) = -\frac{1}{m}\sum_{i=1}^{m} \left[y_i \log(f_{\mathbf{w},b}(\mathbf{x}_i)) + (1 - y_i) \log(1 - f_{\mathbf{w},b}(\mathbf{x}_i)) \right].$$

Exercise: Please prove that $J(\mathbf{w})$ is convex *w.r.t.* $\mathbf{w}$.

(l2 loss $\|\ \|$ is NOT convex wrt w

Gradient descent for logistic regression

- Learning $\mathbf{w}$ by minimize $J(\mathbf{w})$, *i.e.*,

$$\mathbf{w}^* = \arg\min_{\mathbf{w}} J(\mathbf{w}) = -\frac{1}{m}\sum_{i=1}^{m}\left[y_i \log(f_{\mathbf{w},b}(\mathbf{x}_i)) + (1-y_i)\log(1-f_{\mathbf{w},b}(\mathbf{x}_i))\right].$$

- **Gradient descent**: repeat the following update until convergence

$$\mathbf{w} \leftarrow \mathbf{w} - \alpha \nabla_{\mathbf{w}} J(\mathbf{w})$$

$$\nabla_{\mathbf{w}} J(\mathbf{w}) = \frac{1}{m}\sum_{i=1}^{m}[f_{\mathbf{w},b}(\mathbf{x}_i) - y_i]\mathbf{x}_i$$

- How to define convergence? Calculating the changes of $J(\mathbf{w})$ or $\mathbf{w}$ in the last K steps, if the change is lower than a threshold, than it can be seen as convergence. Remember that choosing suitable learning rate α is important to achieve a good converged solution.


$\frac{\partial y \log \hat{y}}{\partial w} = y(1-e^{-z})\left(\frac{1}{1-e^{-z}}\right)^2(-x)$

Gradient descent for logistic regression

`Exercise:` Suppose you are running a logistic regression model, and you should observe the learning procedure to find a suitable learning rate α. Which of the following is reasonable to make sure α is set properly and that the gradient descent is running correctly?

not the correct objective func.

F • Plot $J(\mathbf{w}) = -\frac{1}{m}\sum_i^m (y_i - f_{\mathbf{w},b}(\mathbf{x}_i))^2$ as a function of the number of iterations (*i.e.*, the horizontal axis is the iteration number) and make sure $J(\mathbf{w})$is decreasing on every iteration.




T • Plot $J(\mathbf{w}) = -\frac{1}{m}\sum_i^m \left[y_i \log(f_{\mathbf{w},b}(\mathbf{x}_i)) + (1-y_i)\log(1-f_{\mathbf{w},b}(\mathbf{x}_i))\right]$ as a function of the number of iterations (*i.e.*, the horizontal axis is the iteration number) and make sure $J(\mathbf{w})$ is decreasing on every iteration.

F • Plot $J(\mathbf{w})$ as a function of $\mathbf{w}$ and make sure it is decreasing on every iteration.


F • Plot $J(\mathbf{w})$ as a function of $\mathbf{w}$ and make sure it is convex.

Multi-class classification

- Binary classification: in above examples and derivations, we only consider the binary classification problem, *i.e.*, $y \in \{0, 1\}$.
- Multi-class/multi-category classification: however, many practical problems involve with multi-category outputs, *i.e.*, $y \in \{1, \ldots, C\}$:
  - **Weather forecast**: sunny, cloudy, rain, snow
  - **Email tagging**: work, friends, families, hobby




Multi-class classification



Multi-class classification: one-vs-all


One-vs-all (one-vs-rest):





Class 1: △

Class 2: □


Class 3: ×







Multi-class classification: one-vs-all



One-vs-all logistic regression:

- Train a binary logistic regression $f_{\mathbf{w}_j, b_j}(\cdot)$ for each class j, by setting all samples of other classes as negative class
- For a new testing sample $\mathbf{x}$, predict its class as $\arg\max_j f_{\mathbf{w}_j, b_j}(\mathbf{x})$.

Pros: Easy to implement

Cons: The training cost is too high, and is difficult to scale to tasks with large number of classes. & costly for 很多 classes

Multi-class classification: Softmax regression

- Softmax function:

$f^{(j)}_{\mathbf{W},\mathbf{b}}(\mathbf{x}) \in (0,1)$ [probability]

$\sum f^{(j)}_{\mathbf{W},\mathbf{b}}(\mathbf{x}) = 1$

No longer independent

同时搞出各class的W & b

$$f^{(j)}_{\mathbf{W},\mathbf{b}}(\mathbf{x}) = \frac{\exp(\mathbf{w}_j^\top \mathbf{x} + b_j)}{\sum_{c=1}^{C} \exp(\mathbf{w}_c^\top \mathbf{x} + b_c)} = P(y = j|\mathbf{x}; \mathbf{W}, \mathbf{b}), \tag{12}$$

all the other nodes also influence outcome

where $\mathbf{W} = [\mathbf{w}_1, \ldots, \mathbf{w}_C], \mathbf{b} = [b_1; b_2; \ldots; b_C]$ with C being the number of classes. For simplicity, in the following we write $f^{(j)}_{\mathbf{W},\mathbf{b}}(\cdot)$ as $f_{\mathbf{w}_j,b_j}(\cdot)$

- Cost function:

number of data points

number of classes

$$J(\mathbf{W}) = -\frac{1}{m} \sum_{i}^{m} \sum_{j}^{C} \left[\mathbb{I}(y_i = j) \log(f_{\mathbf{w}_j,b_j}(\mathbf{x}_i)) \right], \tag{13}$$

where $\mathbb{I}(a) = 1$ if a is true, otherwise $\mathbb{I}(a) = 0$.

Multi-class classification: Softmax regression

- It can also be optimized by gradient descent:

$$\mathbf{w}_j \leftarrow \mathbf{w}_j - \alpha \frac{\partial J(\mathbf{W})}{\partial \mathbf{w}_j},$$

$$\text{hint:} \quad \frac{\partial J(\mathbf{W})}{\partial \mathbf{w}_j} = -\frac{1}{m}\sum_{i}^{m}\left[\frac{\mathbb{I}(y_i = j)}{f_{\mathbf{w}_j, b_j}(\mathbf{x}_i)} \cdot \frac{\nabla f_{\mathbf{w}_j, b_j}(\mathbf{x}_i)}{\nabla \mathbf{w}_j} + \sum_{c \neq j}^{C} \frac{\mathbb{I}(y_i = j)}{f_{\mathbf{w}_c, b_c}(\mathbf{x}_i)} \cdot \frac{\nabla f_{\mathbf{w}_c, b_c}(\mathbf{x}_i)}{\nabla \mathbf{w}_j}\right]$$

$$\Longrightarrow \quad \frac{\partial J(\mathbf{W})}{\partial \mathbf{w}_j} = \frac{1}{m}\sum_{i}^{m}\left(f_{\mathbf{w}_j, b_j}(\mathbf{x}_i) - \mathbb{I}(y_i = j)\right)\mathbf{x}_i \tag{14}$$

Note: $\{\mathbf{w}_c\}_{c=1}^{C}$ should be updated in parallel, rather than sequentially.

w_1 变后 $f_{w,b}(x_i)$ 变了, 影响 w_2

1 Review of last week

2 Classification and representation

3 Logistic regression


4 Regularized logistic regression

5 Probabilistic perspective of logistic regression


6 Summary: linear regression vs. logistic regression

Overfitting in linear regression

Example: Linear regression (housing prices)



$b + w_1 x$



$b + w_1 x + w_2 x^2$




$b + w_1 x + w_2 x^2 + w_4 x^3 + w_2 x^4$

Overfitting in linear regression

Addressing overfitting:


$x_1 =$ size of house
$x_2 =$ no. of bedrooms
$x_3 =$ no. of floors
$x_4 =$ age of house
$x_5 =$ average income in neighborhood
$x_6 =$ kitchen size
$\vdots$
x_{100}



Overfitting: If we have too many features, the learned hypothesis may fit the training data very well (low bias), but fail to generalize to new examples.


Overfitting in logistic regression

Example: Logistic regression




$$f_{\mathbf{w},b}(\mathbf{x}) = g(b + w_1 x_1 + w_2 x_2)$$

Under-fitting



$$f_{\mathbf{w},b}(\mathbf{x}) = g(b + w_1 x_1 + w_2 x_2 + w_3 x_1^2 + w_4 x_2^2 + w_5 x_1 x_2)$$

Good-fitting



$$f_{\mathbf{w},b}(\mathbf{x}) = g(b + w_1 x_1 + w_2 x_1^2 + w_3 x_1^2 x_2 + w_4 x_1^2 x_2^2 + w_5 x_1^2 x_2^3 + w_6 x_1^3 x_2 + \dots)$$

Over-fitting

Addressing Overfitting

Generally, there are two approaches to address the overfitting problem, including:

- Reducing the number of features:
  - Feature selection *reduce # elements in input vector x.*
  - Dimensionality reduction (introduced in later lectures)
- Regularization:
  - Keep all features, but reduce magnitude/value of each parameter, such that each feature contributes a bit to predict y

In the following, we will focus on the regularization-based approach.

Regularized logistic regression

- The objective function of the regularized logistic regression is formulated as follows

$$\bar{J}(\mathbf{w}) = J(\mathbf{w}) + \frac{\lambda}{2m}\sum_{j=1}^{d} w_j^2$$

$$= -\frac{1}{m}\sum_{i}^{m}\left[y_i \log(f_{\mathbf{w},b}(\mathbf{x}_i)) + (1-y_i)\log(1-f_{\mathbf{w},b}(\mathbf{x}_i))\right] + \frac{\lambda}{2m}\sum_{j=1}^{d} w_j^2.$$

Note: the bias parameter w_0 (or b) is not regularized/penalized.

- The above objective function can also be solved by gradient descent, as follows

$$w_0 \leftarrow w_0 - \frac{\alpha}{m}\sum_{i=1}^{m}(f_{\mathbf{w},b}(\mathbf{x}_i) - y_i)\cdot \mathbf{x}_i(0), \text{ where } \mathbf{x}_i(0) = 1, \forall i$$

$$w_j \leftarrow w_j - \frac{\alpha}{m}\left[\sum_{i=1}^{m}(f_{\mathbf{w},b}(\mathbf{x}_i) - y_i)\cdot \mathbf{x}_i(j) + \lambda \cdot w_j\right],$$

where $\mathbf{x}_i(j)$ denotes the j-th entry of $\mathbf{x}_i$, and $j = 0, \ldots, d$.

Regularized logistic regression

Exercise: When using regularized logistic regression, which of these is the best way to monitor whether gradient descent is working correctly?

- Plot $J(\mathbf{w})$ as a function of the number of iterations and make sure it's decreasing
- Plot $J(\mathbf{w}) - \frac{\lambda}{2m}\sum_{j=1}^{d} w_j^2$ as a function of the number of iterations and make sure it's decreasing
- ✓ Plot $J(\mathbf{w}) + \frac{\lambda}{2m}\sum_{j=1}^{d} w_j^2$ as a function of the number of iterations and make sure it's decreasing
- Plot $\sum_{j=1}^{d} w_j^2$ as a function of the number of iterations and make sure it's decreasing

1 Review of last week

2 Classification and representation

3 Logistic regression

4 Regularized logistic regression

5 Probabilistic perspective of logistic regression

6 Summary: linear regression vs. logistic regression

Logistic regression: probabilistic modeling

- Behind **logistic regression for binary classification**, we assume that both the feature $\mathbf{x}$ and and the label y are random variables, as follows

$$\mu(\mathbf{x}|\mathbf{w}) = \text{Sigmoid}(\mathbf{w}^\top \mathbf{x}),$$
$$y(\mathbf{x}|\mathbf{w}) \sim \text{Bernoulli}(\mu(\mathbf{x}|\mathbf{w})).$$

→ probability of $y=1$

↳ only 2 possible outcomes

- Then, we have

$$P(y|\mathbf{x};\mathbf{w}) = \begin{cases} \mu & \text{if } y = 1, \\ 1-\mu & \text{if } y = 0. \end{cases}$$

- The log-likelihood function of $P(y|\mathbf{x};\mathbf{w})$ is formulated as

$$\mathcal{L}(\mathbf{w}) = y \log(\mu) + (1-y)\log(1-\mu).$$

- Thus, we obtain

$$\max_{\mathbf{w}} \mathcal{L}(\mathbf{w}) \equiv \min_{\mathbf{w}} J(\mathbf{w}).$$

Logistic regression: probabilistic modeling

- Behind logistic regression, we assume that

$$\mu(\mathbf{x}|\mathbf{w}) = \text{Sigmoid}(\mathbf{w}^\top \mathbf{x}),$$
$$y(\mathbf{x}|\mathbf{w}) \sim \text{Bernoulli}(\mu(\mathbf{x}|\mathbf{w})).$$

- ℓ_2-regularized logistic regression: we further assume $\mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \sigma^2\mathbf{I})$, then we have

$$\max_{\mathbf{w}} \mathcal{L}(\mathbf{w}) + \log \mathcal{N}(\mathbf{w}|\mathbf{0}, \sigma^2\mathbf{I}) \equiv \min_{\mathbf{w}} J(\mathbf{w}) + \frac{\lambda}{2m} \sum_{j=1}^{d} w_j^2.$$

- ℓ_1-regularized logistic regression: if we assume $\mathbf{w} \sim \text{Laplace}(\mathbf{w}|\mathbf{0}, b)$, then we have

$$\max_{\mathbf{w}} \mathcal{L}(\mathbf{w}) + \log \text{Laplace}(\mathbf{w}|\mathbf{0}, b) \equiv \min_{\mathbf{w}} J(\mathbf{w}) + \frac{\lambda}{2m} \sum_{j=1}^{d} |w_j|.$$

1 Review of last week

2 Classification and representation

3 Logistic regression

4 Regularized logistic regression

5 Probabilistic perspective of logistic regression

6 Summary: linear regression vs. logistic regression

Summary: linear regression vs. logistic regression

	Linear regression	Logistic regression
Task	regression	classification
Hypothesis $f_{\mathbf{w},b}(\mathbf{x})$	$\mathbf{w}^\top \mathbf{x} + b \in (-\infty, \infty)$	$g(\mathbf{w}^\top \mathbf{x} + b) \in [0, 1]$
Objective $J(\mathbf{w})$	$\frac{1}{2m} \sum_i^m (y_i - \mathbf{w}^\top \mathbf{x}_i)^2$	$-\frac{1}{m} \sum_{i=1}^m \big[y_i \log(f_{\mathbf{w},b}(\mathbf{x}_i)) + (1 - y_i) \log(1 - f_{\mathbf{w},b}(\mathbf{x}_i))\big]$
Solution	closed-form or gradient descent	gradient descent

Note that: For each variant of linear/logistic regression, you can derive it from both the deterministic and the probabilistic perspectives.

Own reading: Both linear regression and logistic regression are special cases of generalized linear models. If interested, you can find more details from Section 4 of th book "Pattern Recognition and Machine Learning", Bishop, 2006.