Assignment 3

- **Released date: 2024/10/10.**
- **Due: 2024/10/21.**
- Late submission is **NOT** acceptable.
- Please submit your assignment as a PDF file titled "student ID + HW3".

Question 1. Show that $\mathbb{R}^{m\times n}$, together with the usual addition and scalar multiplication of matrices, satisfies the eight axioms of a vector space.

Question 2. Let $\mathbf{x}$, $\mathbf{y}$, and $\mathbf{z}$ be vectors in a vector space V. Prove that if

$$\mathbf{x}+\mathbf{y}=\mathbf{x}+\mathbf{z}$$

then $\mathbf{y}=\mathbf{z}$.

Question 3. Let V be a vector space and let $\mathbf{x}\in V$. Show that

(a) $\beta\mathbf{0}=\mathbf{0}$ for each scaler β.

(b) If $\alpha\mathbf{x}=\mathbf{0}$ then either $\alpha=0$ or $\mathbf{x}=\mathbf{0}$.

Question 4. Let S be the set of all ordered pairs of real numbers. Define scalar multiplication and addition on S by

$$\alpha(x_1,x_2)=(\alpha x_1,\alpha x_2)$$
$$(x_1,x_2)\oplus(y_1,y_2)=(x_1+y_1,0)$$

We use the symbol $\oplus$ to denote the addition operation for this system in order to avoid confusion with the usual addition $\mathbf{x}+\mathbf{y}$ of row vectors. Show that S, together with the ordinary scalar multiplication and the addition operation $\oplus$, is not a vector space. Which of the eight axioms fails to hold?

Question 5. Let V be the set of all ordered pairs of real numbers with addition defined by

$$(x_1, x_2) + (y_1, y_2) = (x_1 + y_1, x_2 + y_2)$$

and scalar multiplication defined by

$$\alpha \circ (x_1, x_2) = (\alpha x_1, x_2)$$

Scalar multiplication for this system is defined in an unusual way, and consequently we use the symbol $\circ$ to avoid confusion with the ordinary scalar multiplication of row vectors. Is V a vector space with these operations? Justify your answer.

Question 6. Let R^+ denote the set of positive real numbers. Define the operation of scalar multiplication, denoted $\circ$, by

$$\alpha \circ x = x^\alpha$$

for each $x \in R^+$ and for any real number α. Define the operation of addition, denoted $\oplus$, by

$$x \oplus y = x \cdot y \quad \text{for all} \quad x, y \in R^+$$

Thus, for this system, the scalar product of -3 times $\frac{1}{2}$ is given by

$$-3 \circ \frac{1}{2} = (\frac{1}{2})^{-3} = 8$$

and the sum of 2 and 5 is given by

$$2 \oplus 5 = 2 \cdot 5 = 10$$

Is R^+ a vector space with these operations? Prove your answer.

Question 7. Let R denote the set of real numbers. Define scalar multiplication by

$$\alpha x = \alpha \cdot x \quad \text{(the usual multiplication of real numbers)}$$

and define addition, denoted $\oplus$, by

$$x \oplus y = \max(x, y) \quad \text{(the maximum of the two numbers)}$$

Is R a vector space with these operations? Prove your answer.

Question 8. Determine whether the following vectors are linearly independent in $\mathbb{R}^2$:

(a)
$$\begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 3 \\ 2 \end{pmatrix}$$

(b)
$$\begin{pmatrix} -2 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 2 \\ 4 \end{pmatrix}$$

Question 9. Determine whether the following vectors are linearly independent in $\mathbb{R}^{2\times 2}$:

(a) $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$.

(b) $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 2 & 3 \\ 0 & 2 \end{pmatrix}$.

Question 10. Let $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_k$ be linearly independent vectors in a vector space V.

(a) If we add a vector $\mathbf{x}_{k+1}$ to the collection, will we still have a linearly independent collection of vectors? Explain.

(b) If we delete a vector, say, $\mathbf{x}_k$, from the collection, will we still have a linearly independent collection of vectors? Explain.

Question 11.

(a) Let $\mathbf{x}_1, \mathbf{x}_2$, and $\mathbf{x}_3$ be linearly independent vectors in $\mathbb{R}^n$ and let

$$\mathbf{y}_1 = \mathbf{x}_1 + \mathbf{x}_2, \quad \mathbf{y}_2 = \mathbf{x}_2 + \mathbf{x}_3, \quad \mathbf{y}_3 = \mathbf{x}_3 + \mathbf{x}_1$$

Are $\mathbf{y}_1, \mathbf{y}_2$, and $\mathbf{y}_3$ linearly independent? Prove your answer.

(b) Let $\mathbf{x}_1, \mathbf{x}_2$, and $\mathbf{x}_3$ be linearly independent vectors in $\mathbb{R}^n$ and let

$$\mathbf{y}_1 = \mathbf{x}_2 - \mathbf{x}_1, \quad \mathbf{y}_2 = \mathbf{x}_3 - \mathbf{x}_2, \quad \mathbf{y}_3 = \mathbf{x}_3 - \mathbf{x}_1$$

Are $\mathbf{y}_1, \mathbf{y}_2$, and $\mathbf{y}_3$ linearly independent? Prove your answer.

Question 12. For each of the following, show that the given vectors are linearly independent in $C[-1, 1]$:

(a) $\cos \pi x, \sin \pi x$

(b) $x^3, |x|^3$

Question 13. Prove that any nonempty subset of a linearly independent set of vectors $\{\mathbf{v}_1, \ldots, \mathbf{v}_n\}$ is also linearly independent.

Question 14. Let $\mathbf{x}_1, \ldots, \mathbf{x}_k$ be linearly independent vectors in $\mathbb{R}^n$, and let A be a non-singular $n \times n$ matrix. Define $\mathbf{y}_i = A\mathbf{x}_i$ for $i = 1, \ldots, k$. Show that $\mathbf{y}_1, \mathbf{y}_2, \ldots, \mathbf{y}_k$ are linearly independent.

Question 15.

(a) Let $\{\mathbf{v}_1, \ldots, \mathbf{v}_n\}$ be a spanning set for the vector space V, and let $\mathbf{v}$ be any other vector in V. Show that $\mathbf{v}, \mathbf{v}_1, \ldots, \mathbf{v}_n$ are linearly dependent.

(b) Let $\mathbf{v}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n$ be linearly independent vectors in a vector space V. Show that $\mathbf{v}_2, \ldots, \mathbf{v}_n$ cannot span V.