

Some of the Big Ideas in Quantum Mechanics

Joseph Cunningham

Contents

I	Quantum Mechanics	3
1	Measurement theory	4
2	Postulates	5
2.1	States, observables and measurement	5
2.1.1	States	5
2.1.2	Effects	5
3	Quantum theory	6
3.1	Quantum statistics	6
3.2	Time evolution	6
3.2.1	The Schrödinger equation	6
3.2.2	Schrödinger and Heisenberg pictures	6
3.2.3	Adiabatic theorem	6
4	Approximations	9
4.1	Approximating eigenvectors	9
4.1.1	Power series expansion of a non-degenerate level	9
4.2	Approximating evolutions	9
5	Investigations of systems	10
5.1	Stepped potentials	10
5.2	Coulomb interaction	10
5.3	Harmonic oscillator	10
II	Quantum Information and computation	11
1	Circuit model	12
1.1	Qubits	12
1.1.1	$\mathfrak{su}(2)$ and the Pauli matrices	12
1.1.2	The Bloch sphere	13
1.2	Gates	13
1.2.1	Controlled gates	13

2	Eigenpath traversal	14
2.1	Quantum Zeno effect	14
2.2	Adiabatic quantum computation	14
2.3	Evolution through measurement	14
2.3.1	Phase randomisation	14
3	Variational quantum algorithms	15
4	Algorithms	16
4.1	Deutsch's problem	16
4.2	Grover's search algorithm	16
4.2.1	Circuit model	16
4.2.1.1	The oracle	16
4.2.1.2	The algorithm	17
4.2.2	Analogue Grover	17
4.2.3	Adiabatic Grover	17
4.2.3.1	Poisson projective measurement	19
A	Bibliography	21

Part I

Quantum Mechanics

Chapter 1

Measurement theory

Chapter 2

Postulates

2.1 States, observables and measurement

2.1.1 States

[A state on a Hilbert space is a positive trace class operator with trace 1.

Proposition I.1. *Let ρ be a state. The following are equivalent:*

1. ρ is extremal;
2. ρ is a projector;
3. $\rho = \rho^2$;
4. $\text{Tr}[\rho^2] = 1$;
5. $\|\rho\| = 1$;
6. $S(\rho) := -\text{Tr}[\rho \ln \rho] = 0$.

The quantity $S(\rho) := -\text{Tr}[\rho \ln \rho]$ is called the von Neumann entropy.

2.1.2 Effects

[An effect on a Hilbert space is a bounded operator in $[0, \text{id}]$.

Chapter 3

Quantum theory

3.1 Quantum statistics

3.2 Time evolution

3.2.1 The Schrödinger equation

$$H = i\hbar \frac{d}{dt}$$

Lemma I.2. $U = e^{Ht/i\hbar} = e^{-iHt/\hbar}$.

3.2.2 Schrödinger and Heisenberg pictures

Lemma I.3. Let \mathcal{H} be a Hilbert space and $\{e_i\}$ an orthonormal basis. Let U be a unitary operator on \mathcal{H} . Then $\{Ue_i\}$ is also an orthonormal basis for \mathcal{H} .

$\{e_i\}$ gives Schrödinger and $\{Ue_i\}$ gives Heisenberg.

3.2.3 Adiabatic theorem

Theorem I.4. Let \mathcal{H} be a Hilbert space and consider a path $H : [0, 1] \rightarrow \mathcal{SA}(\mathcal{H}) : s \mapsto H(s)$ of self-adjoint operators on the Hilbert space.

Take $\epsilon > 0$. If the Hamiltonian of a system is given by $H(\epsilon t)$ for times $t \in [0, \epsilon^{-1}]$. Then the state ρ of the system satisfies

$$i\epsilon \frac{d\rho(s)}{ds} = [H(s), \rho(s)],$$

where $s = \epsilon t \in [0, 1]$.

Assume $\lambda(s) \in \sigma(H(s))$ is an isolated point of the spectrum for all $s \in [0, 1]$. It is an eigenvalue and let $P(s)$ be the orthogonal projector onto the eigenspace. Set $\Delta(s) := \inf d(\lambda, \sigma(H(s)) \setminus \{\lambda(s)\})$.

Assume $\rho(0) \leq P(0)$. Consider the fidelity $F(s) := \text{Tr}(P(s)\rho(s))$. Then

$$F(1) \geq 1 - \epsilon O\left(\frac{\|H'\|^2}{\Delta^3} + \frac{\|H''\|}{\Delta^2}\right).$$

Proof. We first write down a differential equation for the fidelity (where $t = \frac{d}{ds}$)

$$\frac{dF(s)}{ds} = \text{Tr}(P'\rho) + \text{Tr}(P\rho').$$

The second term is zero because P commutes with H :

$$\begin{aligned} \text{Tr}(P\rho') &= -\epsilon^{-1}i \text{Tr}\left(P[H, \rho]\right) \\ &= -\epsilon^{-1}i \text{Tr}\left(P(H\rho - \rho H)\right) \\ &= -\epsilon^{-1}i \text{Tr}\left(PH\rho\right) + \epsilon^{-1}i \text{Tr}\left(\rho HP\right) \\ &= -\epsilon^{-1}i \text{Tr}\left(PH\rho\right) + \epsilon^{-1}i \text{Tr}\left(\rho PH\right) \\ &= -\epsilon^{-1}i \text{Tr}\left([PH, \rho]\right) = 0, \end{aligned}$$

by XIV.305.

Now set $Q(s) = \text{id} - P(s)$ and consider the pseudoinverse $(H - \lambda \text{id})^+$ (TODO by continuous functional calculus?). Then $(H - \lambda \text{id})(H - \lambda \text{id})^+ = Q = (H - \lambda \text{id})^+(H - \lambda \text{id})$ by continuous functional calculus (TODO ref). Now we can expand $P' = PP'Q + QP'P$ (by XII.18.2), so

$$\begin{aligned} \frac{dF(s)}{ds} &= \text{Tr}(P'\rho) \\ &= \text{Tr}(PP'Q\rho + QP'P\rho) \\ &= \text{Tr}\left(PP'(H - \lambda \text{id})^+(H - \lambda \text{id})\rho + (H - \lambda \text{id})(H - \lambda \text{id})^+P'P\rho\right) \\ &= \text{Tr}\left(P'(H - \lambda \text{id})^+(H - \lambda \text{id})\rho P + (H - \lambda \text{id})^+P'P\rho(H - \lambda \text{id})\right) \\ &= \text{Tr}\left(P'(H - \lambda \text{id})^+(H\rho P - \rho\lambda P) + (H - \lambda \text{id})^+P'(P\rho H - \lambda P\rho)\right) \\ &= \text{Tr}\left(P'(H - \lambda \text{id})^+(H\rho - \rho H)P + (H - \lambda \text{id})^+P'P(\rho H - H\rho)\right) \\ &= \text{Tr}\left(P'(H - \lambda \text{id})^+[H, \rho]P - (H - \lambda \text{id})^+P'P[H, \rho]\right) \\ &= \text{Tr}\left(PP'(H - \lambda \text{id})^+[H, \rho] - (H - \lambda \text{id})^+P'P[H, \rho]\right) \\ &= \text{Tr}\left(P'Q(H - \lambda \text{id})^+[H, \rho] - (H - \lambda \text{id})^+QP'[H, \rho]\right) \\ &= \text{Tr}\left(P'(H - \lambda \text{id})^+[H, \rho] - (H - \lambda \text{id})^+P'[H, \rho]\right) \\ &= \text{Tr}\left([P', (H - \lambda \text{id})^+] \cdot [H, \rho]\right) \\ &= i\epsilon \text{Tr}\left([P', (H - \lambda \text{id})^+]\rho'\right) \end{aligned}$$

Integrating this w.r.t. s gives

$$F(1) - F(0) = i\epsilon \int_0^1 \text{Tr}\left([P', (H - \lambda \text{id})^+]\rho'\right) ds.$$

We fill in that $F(0) = 1$ and perform integration by parts to obtain

$$\begin{aligned} F(1) &= 1 + i\epsilon \int_0^1 \text{Tr}\left([P', (H - \lambda \text{id})^+]\rho'\right) ds \\ &= 1 + i\epsilon \text{Tr}\left([(H - \lambda \text{id})^+, P']\rho\right)_0^1 - i\epsilon \int_0^1 \text{Tr}\left([(H - \lambda \text{id})^+, P']'\rho\right) ds \end{aligned}$$



Chapter 4

Approximations

4.1 Approximating eigenvectors

4.1.1 Power series expansion of a non-degenerate level

4.2 Approximating evolutions

Chapter 5

Investigations of systems

5.1 Stepped potentials

(Use density to solve general potentials?)

5.2 Coulomb interaction

5.3 Harmonic oscillator

Part II

Quantum Information and
computation

Chapter 1

Circuit model

1.1 Qubits

[A qubit is an element of $\mathbb{C}P^2$.

1.1.1 $\mathfrak{su}(2)$ and the Pauli matrices

The qubit Hamiltonians are elements of $\mathfrak{su}(2)$. Getting rid of the arbitrary global phase gives us elements of $\mathfrak{su}(2)$.

Lemma II.1. *For all $\sigma \in \mathfrak{su}(2)$, the eigenvalues λ_1, λ_2 are real and $\lambda_1 = -\lambda_2$.*

Proof. For all $\sigma \in \mathfrak{su}(2)$, σ is self-adjoint and $\text{Tr}[\sigma] = \lambda_1 + \lambda_2 = 0$. □

Corollary II.1.1. *For all $\sigma \in \mathfrak{su}(2)$, we have $\|\sigma\|_2 = \sqrt{2}\|\sigma\|$, where $\|\cdot\|_2$ is the Hilbert-Schmidt norm.*

Proof. We have $\|\sigma\|_2 = \sqrt{\lambda_1^2 + \lambda_2^2} = \sqrt{2}|\lambda_1| = \sqrt{2}\|\sigma\|$. □

Corollary II.1.2. *The inner product*

$$\langle \cdot, \cdot \rangle : (\sigma_1, \sigma_2) \mapsto \frac{1}{2} \text{Tr}[\sigma_1 \sigma_2]$$

on $\mathfrak{su}(2)$ yields the operator norm as the norm associated with this inner product.

Corollary II.1.3. *Let σ be a unit vector in $\mathfrak{su}(2)$. Then σ has eigenvalues ± 1 . This means σ is unitary.*

Corollary II.1.4. *There is a bijection between the unit vectors in $\mathfrak{su}(2)$ and the rank-1 projections on \mathbb{C}^2 given by $\mathfrak{su}(2)/\mathbb{R} \rightarrow \sigma \mapsto E_1^\sigma$, where E_1^σ is the eigenspace of σ associated with eigenvalue $+1$.*

Proof. We construct an inverse of the map. Let P_1 be the orthogonal projector on E_1^σ . Then $\sigma = 2P_1 - \text{id}$. □

Proposition II.2. *Let $\sigma \in \mathfrak{su}(2)$. Then $\sigma^2 = \|\sigma\|^2 \mathbf{1}$.*

Proof. We have

$$\sigma^2 = \|\sigma\|^2 \left(\frac{\sigma}{\|\sigma\|} \right)^2 = \|\sigma\|^2 \frac{\sigma}{\|\sigma\|} \left(\frac{\sigma}{\|\sigma\|} \right)^* = \|\sigma\|^2 \mathbf{1},$$

where we have used that $\frac{\sigma}{\|\sigma\|}$ is unitary by II.1.3. \square

Corollary II.2.1. *The real algebra generated by $\mathfrak{su}(2)$ is the Clifford algebra $\text{Cl}_{3,0}$. The unit pseudoscalar is $i\mathbf{1}$.*

Note that $\mathfrak{su}(2)$ is a Lie algebra and thus closed under the Lie bracket, but not an algebra under operator composition. Thus the algebra generated by $\mathfrak{su}(2)$ is larger than $\mathfrak{su}(2)$ (indeed $\sigma^2 = \|\sigma\|^2 \mathbf{1} \notin \mathfrak{su}(2)$).

Proposition II.3 (Pauli matrices). *The Clifford algebra $\mathfrak{su}(2)$ has an orthonormal basis*

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \text{and} \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

1.1.2 The Bloch sphere

Because $\mathfrak{su}(2)$ is the Clifford algebra $\text{Cl}_{3,0}$, we can use the unit sphere in \mathbb{R}^3 to represent the unit vectors in $\mathfrak{su}(2)$. By II.1.4 we can also use it to represent the rank-1 projections on \mathbb{C}^2 . The unit sphere with as x, y, z axes the Pauli matrices $\sigma_x, \sigma_y, \sigma_z$ is called the Bloch sphere.
 TODO image of Bloch sphere.

TODO spherical coordinates: $\cos \theta/2 |0\rangle + e^{i\phi} \sin \theta/2 |1\rangle$

Lemma II.4.

$$e^{\phi/2i\sigma} e^{\theta/2i\sigma^\perp} |1\rangle = \cos \theta/2 |0\rangle + e^{i\phi} \sin \theta/2 |1\rangle$$

1.2 Gates

1.2.1 Controlled gates

Let H_1, H_2 be two Hilbert spaces. Let P be a projector on H_1 and L an operator on H_2 . The operation of L controlled by P is the operation $P \otimes L + (\text{id}_{H_1} - P) \otimes \text{id}_{H_2}$ in $H_1 \otimes H_2 \rightarrow H_1 \otimes H_2$.

Chapter 2

Eigenpath traversal

2.1 Quantum Zeno effect

Proposition II.5. *Consider a path of states $|\psi(s)\rangle$ where $s \in [0, 1]$. Assume that, for fixed d and all δ ,*

$$|\langle\psi(s)|\psi(s+\delta)\rangle|^2 \geq 1 - d^2\delta^2.$$

Then

2.2 Adiabatic quantum computation

2.3 Evolution through measurement

2.3.1 Phase randomisation

$$|\psi_0\rangle = |E_0(0)\rangle = \sum_i \alpha_i |E_i(s_1)\rangle$$

$$\begin{aligned} \rho(s_1) &= \frac{1}{T} \int_0^T e^{-iH(s_1)t} |\psi_0\rangle \langle\psi_0| e^{iH(s_1)t} dt \\ &= \sum_{i,j} \alpha_i \alpha_j^* \frac{1}{T} \left(\int_0^T e^{-i(E_i - E_j)t} dt \right) |E_i\rangle \langle E_j| \\ &= \sum_{i,j} \alpha_i \alpha_j^* \left(\frac{i(e^{-i(E_i - E_j)T} - 1)}{T(E_i - E_j)} \right) |E_i\rangle \langle E_j| \end{aligned}$$

Theorem II.6 (Randomised dephasing). *Let $|\psi(s)\rangle$ be a nondegenerate eigenstate of $H(s)$ and $\{\omega_j\}$ the energy differences to the other eigenstates $|\psi_j(s)\rangle$. Let T be a random variable. Then, for all states ρ , we have*

$$\left\| (M_l - e^{-iH(s)T} \rho e^{-iH(s)T}) \right\|_{tr} \leq \epsilon = \sup_{\omega_j} |\Phi(\omega_j)|$$

Chapter 3

Variational quantum algorithms

Chapter 4

Algorithms

4.1 Deutsch's problem

4.2 Grover's search algorithm

Suppose we have a set \mathcal{N} of N items, some of which are marked. WLOG we can take \mathcal{N} to be the set of bit strings of length ν . The marked items form a subset \mathcal{M} of size M . Suppose we have an oracle that tells us whether an object is marked or not:

$$f : \mathcal{N} \rightarrow \{0, 1\} : x \mapsto \begin{cases} 0 & (x \in \mathcal{M}) \\ 1 & (x \notin \mathcal{M}) \end{cases}$$

Now we want to find a marked item. Classically we need to check $\Theta(N/M)$ items on average.

4.2.1 Circuit model

Let $\{|n\rangle \mid n \in \mathcal{N}\}$ be a basis of a Hilbert space \mathcal{H}_N and let \mathcal{H}_2 be a qubit space.

4.2.1.1 The oracle

We would like to use the operator

$$U_f : \mathcal{H}_N \rightarrow \mathcal{H}_N : |n\rangle \mapsto (-1)^{f(n)} |n\rangle.$$

This can be constructed from the oracle

$$O_f : \mathcal{H}_N \otimes \mathcal{H}_2 \rightarrow \mathcal{H}_N \otimes \mathcal{H}_2 : |n\rangle \otimes |q\rangle \mapsto |n\rangle \otimes |f(n) \oplus q\rangle,$$

where \oplus is addition modulo 2, by preparing the qubit in the superposition $\frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$. Then

$$\begin{aligned} O_f \left(|n\rangle \otimes \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle) \right) &= |n\rangle \otimes \frac{1}{\sqrt{2}}(|f(n)\rangle - |f(n) \oplus 1\rangle) \\ &= (-1)^{f(n)} |n\rangle \otimes \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle). \end{aligned}$$

The operator U_f can also be written

$$U_f = \text{id}_{\mathcal{H}_N} - 2 \sum_{m \in \mathcal{M}} |m\rangle \langle m|.$$

4.2.1.2 The algorithm

We construct the an initial state which is a superposition of all possible solutions

$$|\mathcal{N}\rangle = \frac{1}{\sqrt{N}} \sum_{n \in \mathcal{N}} |n\rangle.$$

This can be prepared by applying a Hadamard gate W to each qubit in the register. We also define the state

$$|\mathcal{M}\rangle = \frac{1}{\sqrt{M}} \sum_{m \in \mathcal{M}} |m\rangle.$$

We define the operation

$$U_0 = W^{\otimes \nu} (\text{id} - 2|0\rangle\langle 0|) W^{\otimes \nu} = \text{id} - 2|\mathcal{N}\rangle\langle \mathcal{N}|.$$

Now both U_0 and U_f leave $\text{span}\{\}$

4.2.2 Analogue Grover

Why does analogue Grover work?

4.2.3 Adiabatic Grover

$$H_0 = \mathbf{1} - |\mathcal{N}\rangle\langle \mathcal{N}| = \mathbf{1} - \mathbb{J}/N$$

$$H_f = \mathbf{1} - \sum_{m \in \mathcal{M}} |m\rangle\langle m| = \begin{pmatrix} 0 & 0 \\ 0 & \mathbf{1}_{N-M} \end{pmatrix}.$$

Then

$$H(s) = (1-s)H_0 + sH_f = \begin{pmatrix} (1-s)\mathbf{1} + \frac{s-1}{N}\mathbb{J} & \frac{s-1}{N}\mathbb{J} \\ \frac{s-1}{N}\mathbb{J} & \mathbf{1} + \frac{s-1}{N}\mathbb{J} \end{pmatrix} = \begin{pmatrix} (1-s)\mathbf{1}_M & 0 \\ 0 & \mathbf{1}_{N-M} \end{pmatrix} + \frac{s-1}{N}\mathbb{J}.$$

Then, setting $A = \begin{pmatrix} (1-s-\lambda)\mathbf{1}_M & 0 \\ 0 & (1-\lambda)\mathbf{1}_{N-M} \end{pmatrix}$

$$\begin{aligned} \det(H(s) - \lambda) &= \det\left(A + \frac{s-1}{N}\mathbb{J}^{n \times 1} \mathbb{J}^{1 \times n}\right) \\ &= \det(A) \det\left(\mathbf{1}_N + \frac{s-1}{N}A^{-1}\mathbb{J}^{n \times 1} \mathbb{J}^{1 \times n}\right) \\ &= \det(A) \det\left(1 + \frac{s-1}{N}\mathbb{J}^{1 \times n} A^{-1} \mathbb{J}^{n \times 1}\right) \\ &= (1-s-\lambda)^M (1-\lambda)^{N-M} \left(1 + \frac{M(s-1)}{(1-s-\lambda)N} + \frac{(N-M)(s-1)}{(1-\lambda)N}\right) \\ &= (1-s-\lambda)^{M-1} (1-\lambda)^{N-M-1} \left(\lambda^2 - \lambda + s(1-s)\frac{N-M}{N}\right), \end{aligned}$$

where we have used the matrix determinant lemma to simplify the calculation.

We see that there are four distinct eigenvalues:

$$\begin{aligned}\lambda_{0,1} &= \frac{1}{2} \left(1 \pm \frac{\sqrt{N + 4(Ns^2 - Ns - Ms^2 + Ms)}}{\sqrt{N}} \right) && \text{with multiplicity 1} \\ \lambda_2 &= 1 - s && \text{with multiplicity } M - 1 \\ \lambda_3 &= 1 && \text{with multiplicity } N - M - 1.\end{aligned}$$

Next we are interested in the eigen vectors associated to $\lambda_{0,1}$. Let Q_1 be a unitary transfor-

mation that maps $\mathbb{J}^{M \times 1}$ to $\begin{pmatrix} \sqrt{M} \\ 0 \\ 0 \\ \vdots \end{pmatrix} \in \mathbb{C}^M$. Similarly let Q_2 be a unitary transformation that maps $\mathbb{J}^{(N-M) \times 1}$ to $\begin{pmatrix} \sqrt{N-M} \\ 0 \\ 0 \\ \vdots \end{pmatrix} \in \mathbb{C}^{(N-M)}$ and define $Q = \begin{pmatrix} Q_1 & 0 \\ 0 & Q_2 \end{pmatrix}$, which is also unitary.

Then we have

$$\begin{aligned}QH(s)Q^{-1} &= \begin{pmatrix} (1-s)\mathbb{1}_M & 0 \\ 0 & \mathbb{1}_{N-M} \end{pmatrix} + \frac{s-1}{N} Q \mathbb{J}^{N \times 1} (\mathbb{J}^{N \times 1})^* Q^* \\ &= \begin{pmatrix} (1-s)\mathbb{1}_M & 0 \\ 0 & \mathbb{1}_{N-M} \end{pmatrix} + \frac{s-1}{N} \begin{pmatrix} \sqrt{M} \\ 0 \\ \vdots \\ 0 \\ \sqrt{N-M} \\ 0 \\ \vdots \end{pmatrix} \begin{pmatrix} \sqrt{M} \\ 0 \\ \vdots \\ 0 \\ \sqrt{N-M} \\ 0 \\ \vdots \end{pmatrix}^*.\end{aligned}$$

All but two of the eigenvectors are just elements of \mathcal{N} . To study the other two we can simplify by “removing the zeros”. Now the eigenvalue problem becomes

$$\left(\begin{pmatrix} 1-s-\lambda_{0,1} & 0 \\ 0 & 1-\lambda_{0,1} \end{pmatrix} + \frac{s-1}{N} \begin{pmatrix} \sqrt{M} \\ \sqrt{N-M} \end{pmatrix} \begin{pmatrix} \sqrt{M} \\ \sqrt{N-M} \end{pmatrix}^* \right) \mathbf{v} = 0.$$

Setting $\mathbf{v} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$ gives us the equations

$$\begin{aligned}0 &= (1-s-\lambda_{0,1})x_1 + \frac{s-1}{N}(Mx_1 + \sqrt{M}\sqrt{N-M}x_2) \\ 0 &= (1-\lambda_{0,1})x_2 + \frac{s-1}{N}(\sqrt{M}\sqrt{N-M}x_1 + (N-M)x_2)\end{aligned}$$

reshuffling and dividing these equations gives

$$\frac{(1-s-\lambda_{0,1})x_1}{(1-s-\lambda_{0,1})x_2} = \frac{-\frac{s-1}{N}(Mx_1 + \sqrt{M}\sqrt{N-M}x_2)}{-\frac{s-1}{N}(\sqrt{M}\sqrt{N-M}x_1 + (N-M)x_2)} = \frac{\sqrt{M}(\sqrt{M}x_1 + \sqrt{N-M}x_2)}{\sqrt{N-M}(\sqrt{M}x_1 + \sqrt{N-M}x_2)} = \frac{\sqrt{M}}{\sqrt{N-M}}.$$

So we have eigenvectors $\mathbf{v}_{0,1} = (\sqrt{M}(1 - \lambda_{0,1}), \sqrt{N - M}(1 - s - \lambda_{0,1}))^T$ of $QH(s)Q^{-1}$. The corresponding eigenvectors of $H(s)$ are then given by

$$Q^* \mathbf{v}_{0,1} = \begin{pmatrix} (1 - \lambda_{0,1}) \mathbb{J}^{M \times 1} \\ (1 - s - \lambda_{0,1}) \mathbb{J}^{(N-M) \times 1} \end{pmatrix}.$$

For computational ease we will keep on working with the eigenvectors $\mathbf{v}_{0,1}$ of $QH(s)Q^{-1}$. We denote by $|0\rangle$ and $|1\rangle$ the normalisation of $\mathbf{v}_{0,1}$.

4.2.3.1 Poisson projective measurement

The dynamics of the system is governed by the differential equation

$$\frac{d\rho}{ds} = \Lambda(|0\rangle\langle 0|\rho|0\rangle\langle 0| + |1\rangle\langle 1|\rho|1\rangle\langle 1| - \rho).$$

Now we can rewrite $\frac{d\rho}{ds}$ in the basis $|0\rangle, |1\rangle$: let $i, j, k, l = 0, 1 \pmod 2$ with implicit summation

$$\begin{aligned} \frac{d\rho}{ds} &= \frac{d}{ds} (|i\rangle\langle i|\rho|j\rangle\langle j|) \\ &= \frac{d}{ds} (\langle i|\rho|j\rangle) |i\rangle\langle j| + \langle i|\rho|j\rangle \frac{d}{ds} (|i\rangle\langle j|) \\ &= \frac{d}{ds} (\langle i|\rho|j\rangle) |i\rangle\langle j| + \langle i|\rho|j\rangle |k\rangle\langle k| \frac{d}{ds} |i\rangle\langle j| - \langle i|\rho|j\rangle |i\rangle\langle j| \frac{d}{ds} |l\rangle\langle l| \\ &= \frac{d}{ds} (\langle i|\rho|j\rangle) |i\rangle\langle j| + \langle i|\rho|j\rangle |i+1\rangle\langle i+1| \frac{d}{ds} |i\rangle\langle j| - \langle i|\rho|j\rangle |i\rangle\langle j| \frac{d}{ds} |j+1\rangle\langle j+1| \\ &= \left(\frac{d}{ds} (\langle i|\rho|j\rangle) + \langle i+1|\rho|j\rangle \langle i| \frac{d}{ds} |i+1\rangle - \langle i|\rho|j+1\rangle \langle j+1| \frac{d}{ds} |j\rangle \right) |i\rangle\langle j|. \end{aligned}$$

For each $|i\rangle\langle j|$ we get an equation, four in total. One of these is redundant by the zero trace requirement. Writing $y_{i,j} := \langle i|\rho|j\rangle$ and $\omega_{ij} := \langle i| \frac{d}{ds} |j\rangle$ the three remaining equations are

$$\begin{aligned} \frac{d\rho_{00}}{ds} &= -\rho_{10}\omega_{01} + \rho_{01}\omega_{10} \\ \frac{d\rho_{01}}{ds} &= -\Lambda\rho_{01} - (1 - \rho_{00})\omega_{01} + \rho_{00}\omega_{01} \\ \frac{d\rho_{10}}{ds} &= -\Lambda\rho_{10} - \rho_{00}\omega_{10} + (1 - \rho_{00})\omega_{10} \end{aligned}$$

Setting $\omega = \omega_{10} = -\omega_{01}$ and $y = \rho_{00} - 1/2$ we obtain the equation

$$\frac{d^2 y}{ds^2} = \left(\frac{\frac{d\omega}{ds}}{\omega} - \Lambda \right) \frac{dy}{ds} - 4\omega^2 y.$$

Setting $\mathbf{y} = \begin{pmatrix} y \\ y' \end{pmatrix}$ this second order differential equation is equivalent to the first order system

$$\begin{pmatrix} y \\ y' \end{pmatrix}' = \begin{pmatrix} 0 & 1 \\ -4 & -\Lambda/\omega \end{pmatrix} \mathbf{y},$$

which is in the form $\mathbf{y}' = A\mathbf{y}$. Now A is similar to $\begin{pmatrix} \omega/\Lambda & 0 \\ -4 & -\Lambda/\omega \end{pmatrix}$, so it is bounded if both Λ and Λ^{-1} are bounded functions.

Then by the Picard-Lindelöf theorem the initial value problem has a unique solution on $[0, 1]$. In addition let y_1 be the solution obtained using Λ_1 and y_2 using Λ_2 . Then

$$\Lambda_1 \leq \Lambda_2 \implies y_1 \leq y_2.$$

To see this, we may first remark that

$$\{t \in [0, 1] \mid y_1(t) \leq y_2(t) \wedge y'_1(t) \leq y'_2(t)\} = (y_2 - y_1)^{-1}[[0, +\infty[] \cap (y'_2 - y'_1)^{-1}[[0, +\infty[]$$

is a closed and bounded set that contains 0. Let t_1 be its supremum. This means that $y_1(t_1) \leq y_2(t_1)$ and $y'_1(t_1) \leq y'_2(t_1)$, but $y_1(t) > y_2(t)$ or $y'_1(t) > y'_2(t)$ on some open set $]t_1, t_1 + \delta[$.

Appendix A

Bibliography

- Quantum Theory - Peres
- QM - Schwabl