

Lecture notes in Derived Algebraic Geometry

Session 3

Course by F. Binda, notes by E. Hecky

17 February 2025

1 A word about simplicial rings and \mathbb{E}_∞ -rings

Recall from the previous session that if C is a category with coproducts, then $\mathrm{Ani}(C) = \mathcal{P}_\Sigma(C^{1-\omega\mathrm{proj}})$ is its animation. In particular, $\mathrm{CAlg}^{\mathrm{ani}} = \mathrm{Fun}^\Pi(\mathrm{Poly}_{\mathbf{Z}}^{\mathrm{op}}, \mathcal{S}\mathrm{pc})$ is the ∞ -category of animated rings, and it contains fully faithfully the usual category CAlg^\heartsuit as the subcategory of 0-truncated objects.

Remark 1.1. Denoting by SCR the simplicial category of simplicial commutative rings, there is an equivalence

$$\infty(\mathrm{N}_\Delta(\mathrm{SCR})) \simeq \mathrm{CAlg}^{\mathrm{ani}}$$

between animated rings and the ∞ -category associated to the simplicial nerve of SCR . That is the main reason why the simplicial approach of Toën-Vezzosi gives the same theory in the end as the more modern approach using animated rings.

Another approach that is linked to animated rings is the theory of \mathbb{E}_∞ -rings. Recall that the ∞ -category $\mathcal{S}\mathrm{p}$ of spectra admits a structure of a symmetric monoidal stable ∞ -category. In particular, it is possible to consider commutative algebra objects, and we get the ∞ -category $\mathrm{CAlg}(\mathcal{S}\mathrm{p})$ of \mathbb{E}_∞ -rings spectra.

One can associate to any polynomial algebra a connective ring spectrum, yielding a functor

$$\mathrm{Poly}_{\mathbf{Z}} \rightarrow \mathrm{CAlg}(\mathcal{S}\mathrm{p}_{\geq 0})$$

which then extends to

$$\mathrm{CAlg}^{\mathrm{ani}} \rightarrow \mathbb{E}_\infty\text{-rings}.$$

However, this functor is not essentially surjective. Indeed, if A is an animated ring, then $\pi_*(A)$ is a graded commutative ring, each $\pi_n(A)$ is a module over the classical ring $\pi_0(A)$. More importantly, the even homotopy groups $\pi_{2*}(A)$ have a structure of a divided power algebra¹.

This is not the case for every \mathbb{E}_∞ -ring spectrum. For example, the algebra $\mathbb{F}_p[X]$ with X of degree 2 does not admit a PD-structure, so it doesn't come from any animated ring.

That being said, this phenomenon purely comes from torsion, and if we consider rational algebras, we get an equivalence :

$$\mathrm{CAlg}_{\mathbf{Q}}^{\mathrm{ani}} \simeq \mathrm{CAlg}(\mathcal{S}\mathrm{p}_{\geq 0})_{\mathbf{Q}}/.$$

¹if A is an algebra over the rational numbers \mathbf{Q} , then this structure is just the usual division by integers.

2 Base change

Lemma 2.1. *Let $\varphi : R \rightarrow R'$ be a morphism in \mathbf{CAlg}^\heartsuit . It induces a functor $\mathbf{Poly}_R \rightarrow \mathbf{Poly}_{R'}$ sending P to $P \otimes_R R'$, inducing an adjunction :*

$$\mathbf{CAlg}_R^{\text{ani}} \rightleftarrows \mathbf{CAlg}_{R'}^{\text{ani}}$$

between extension and restriction of scalars.

Proof. In the following diagram, the diagonal composition descends to a unique extension of scalars functor :

$$\begin{array}{ccc} \mathbf{Poly}_R & \xrightarrow{- \otimes_R R'} & \mathbf{Poly}_{R'} \\ j \downarrow & \searrow & \downarrow j \\ \mathbf{CAlg}_R^{\text{ani}} & \xrightarrow[\exists!]{-} & \mathbf{CAlg}_{R'}^{\text{ani}} \end{array}$$

as the image of $j \circ (- \otimes_R R')$ by the equivalence

$$\mathbf{Fun}_\Sigma(\mathbf{Poly}_R, \mathbf{CAlg}_{R'}^{\text{ani}}) \simeq \mathbf{Fun}'(\mathbf{CAlg}_R^{\text{ani}}, \mathbf{CAlg}_{R'}^{\text{ani}}).$$

The existence of a right adjoint comes from presentability and the adjoint functor theorem. \square

This construction yields the same result as the “naive” idea of taking $R \in \mathbf{CAlg}^\heartsuit$ and $S, T \in \mathbf{CAlg}_R^{\text{ani}}$, and defining $S \otimes_R^{\mathbb{L}} T$ as the corresponding pushout of animated algebras. To see this, just notice that they have the exact same behavior on polynomial algebras, and both constructions preserve enough colimits to extend to the same functor.

Remark 2.2. There is an equivalence $\mathbf{Ani}(\mathbf{CAlg}_R^\heartsuit) \simeq \mathbf{CAlg}_{R/}^{\text{ani}}$ between the animation of the category of (classical) R -algebras and the slice category of animated algebras under R .

This remark being true for static rings R motivates the following definition for all animated rings.

Definition 2.3. Let $A \in \mathbf{CAlg}^{\text{ani}}$ be an animated ring. Then define the ∞ -category of animated A -algebras as :

$$\mathbf{CAlg}_A^{\text{ani}} = \mathbf{Ani}(\mathbf{CAlg}^\heartsuit)_{A/}.$$

In fact this is even functorial in A , in the sense that a functor

$$\mathbf{CAlg}_{(-)}^{\text{ani}} : \mathbf{Ani}(\mathbf{CAlg}^\heartsuit) \rightarrow \mathcal{C}at_\infty$$

can be obtained by straightening the extension/restriction of scalars adjunction constructed above.

3 Animated modules

Let $R \in \mathbf{CAlg}^\heartsuit$ be a classical ring. Then the compact projective R -modules $(\mathbf{Mod}_R^\heartsuit)^{\omega\text{proj}}$ are the retracts of finite free modules. One then gets the ∞ -category of animated R -modules by taking Σ -presheaves of spaces on those modules.

It is possible to promote this construction in two different ways :

- allow R to be an animated ring ;
- make this construction functorial in R .

This section's goal is to do both at once.

Let $\mathbf{CRMod}^\heartsuit$ be the following 1-category. Its objects are pairs (A, M) with $A \in \mathbf{CAlg}^\heartsuit$ a ring and $M \in \mathbf{Mod}_A^\heartsuit$ an A -module. A morphism $(A, M) \rightarrow (B, N)$ consists of a ring map $f : A \rightarrow B$, and of a B -linear map $\varphi : M \otimes_A B \rightarrow N$ (or equivalently by adjunction, an A -linear map from M to the restriction of N).

Proposition 3.1. *The compact projective objects of $\mathbf{CRMod}^\heartsuit$ are the (retracts of) finite free modules over (retracts of) finite polynomial algebras over \mathbf{Z} . In other words, up to retracts, every compact projective object in this category looks like $(\mathbf{Z}[\underline{X}], \mathbf{Z}[\underline{X}]^{\oplus n})$ with \underline{X} a finite set of unknowns and $n \geq 0$.*

Ideas of the proof. Let C be the category which is claimed to be $(\mathbf{CRMod}^\heartsuit)^{\omega\text{proj}}$ in the statement. Notice that every object in $\mathbf{CRMod}^\heartsuit$ is a 1-sifted colimit of objects of C . Also, C is the closure under finite coproducts of the category spanned by the two generating objects $(\mathbf{Z}[T], 0)$ and (\mathbf{Z}, \mathbf{Z}) . Finally, both these objects are themselves compact projective. \square

Keeping this notation, the animation $\mathbf{Ani}(\mathbf{CRMod}^\heartsuit)$ is the ∞ -category of presheaves of spaces on this C . We then get two projection functors :

$$\mathbf{CAlg}^{\text{ani}} \xleftarrow{\text{pr}_1} \mathbf{Ani}(\mathbf{CRMod}^\heartsuit) \xrightarrow{\text{pr}_2} \mathbf{Mod}_{\mathbf{Z}}^{\text{ani}}.$$

Notice that both functors have sections.

Remark 3.2. For every static ring $R \in \mathbf{CAlg}^\heartsuit$, the forgetful functor

$$\mathbf{Ani}(\mathbf{CRMod}_R^\heartsuit) \rightarrow \mathbf{CAlg}_R^{\text{ani}} \times \mathbf{Mod}_R^{\text{ani}}$$

is conservative.

This ∞ -category together with the inner fibration pr_1 gives the desired definition and functoriality of the ∞ -categories of animated modules.

Definition 3.3. For every animated ring $A \in \mathbf{CAlg}^{\text{ani}}$, let \mathbf{Mod}_A be the fiber of pr_1 over A . In other words, it is the following pullback

$$\begin{array}{ccc} \mathbf{Mod}_A & \longrightarrow & \mathbf{Ani}(\mathbf{CRMod}^\heartsuit) \\ \downarrow & \lrcorner & \downarrow \text{pr}_1 \\ \Delta^0 & \xrightarrow{A} & \mathbf{CAlg}^{\text{ani}} \end{array}$$

and it is indeed an ∞ -category because pr_1 is an inner fibration.

Actually, pr_1 is fortunately more than an inner fibration. It is both a Cartesian and a coCartesian fibration, whence a functoriality of $\mathbf{Mod}_(-)$ both covariantly and contravariantly.

Theorem 3.4. pr_1 is a biCartesian fibration.

Proof. Let us prove that it is coCartesian. Concretely, for every animated ring map $f : A \rightarrow B$, we need to find $\mathcal{A}, \mathcal{B} \in \text{Ani}(\text{CRMod}^\heartsuit)$ whose pr_1 are respectively A and B , and a map $\phi : \mathcal{A} \rightarrow \mathcal{B}$ such that for every $X \in \text{Ani}(\text{CRMod}^\heartsuit)$, the following diagram is homotopy-cartesian in Spc :

$$\begin{array}{ccc} \text{Map}(\mathcal{B}, X) & \xrightarrow{\phi} & \text{Map}(\mathcal{A}, X) \\ \text{pr}_1 \downarrow & & \downarrow \text{pr}_1 \\ \text{Map}(B, \text{pr}_1(X)) & \xrightarrow{f} & \text{Map}(A, \text{pr}_1(X)) \end{array}$$

The homotopy-cartesianity condition is equivalent to the fibers of the vertical arrows being homotopy equivalent.

Since every possible choice of \mathcal{A} and \mathcal{B} are sifted colimits of compact projective objects, and since pullbacks commute with cosifted limits, it will suffice to treat the case of A and B compact projective, and to look for \mathcal{A} and \mathcal{B} compact projective as well. By definition of compact projectivity, it also suffices to treat the case where X itself is compact projective.

Now the mapping spaces in the diagram have all become discrete sets ! Write $\mathcal{A} = (A, M)$, $\mathcal{B} = (B, N)$ and $X = (\text{pr}_1(X), M_X)$. Take $\psi : B \rightarrow \text{pr}_1(X)$ and compare $\text{pr}_1^{-1}(\psi) \in \text{Map}(\mathcal{B}, X)$ which we can describe as

$$\text{pr}_1^{-1}(\psi) = \{\tilde{\psi} : (B, N) \rightarrow (\text{pr}_1(X), M_X) \mid \text{pr}_1(\tilde{\psi}) = \psi\} \simeq \text{Hom}_{\text{Mod}_{\text{pr}_1(X)}}(N \otimes_B \text{pr}_1(X), M_X),$$

with $\text{pr}_1^{-1}(\psi \circ f) \in \text{Map}(\mathcal{A}, X)$ which can similarly be described as

$$\text{pr}_1^{-1}(\psi \circ f) \simeq \text{Hom}_{\text{Mod}_{\text{pr}_1(X)}}((M \otimes_A B) \otimes_B \text{pr}_1(X), M_X).$$

Of course, only $f : A \rightarrow B$ is given, and we are free to choose M and N . We just need a morphism $\text{pr}_2(\phi) : M \otimes_A B \rightarrow N$ such that $\text{pr}_2(\phi) \otimes \text{pr}_1(X)$ is an equivalence for every choice of X .

One can take $M = A$ and $N = B$ (and the fibers are both M_X), or even more stupidly, one can take $M = 0$ and $N = 0$ (and the fibers are both singletons). \square

This is the exact theorem that was needed to show the functoriality of the ∞ -category of animated modules. We get two functors :

- $\text{Mod} \in \text{Fun}(\text{CAlg}^{\text{ani, op}}, \text{Pr}^{\text{R}})$ sending $\phi : A \rightarrow A'$ to the ∞ -functor $\text{Mod}(\phi) : \text{Mod}_{A'} \rightarrow \text{Mod}_A$ given by (generalized) restriction of scalars ;
- $\text{Mod}^* \in \text{Fun}(\text{CAlg}^{\text{ani}}, \text{Pr}^{\text{L}, \otimes})$ sending $A \rightarrow A'$ to the symmetric monoidal ∞ -functor of (generalized) extension of scalars $-\otimes_A^{\mathbb{L}} A' : \text{Mod}_A \rightarrow \text{Mod}_{A'}$.

Remark 3.5. There is another possible construction for animated modules, which is to define finite free animated A -modules for $A \in \text{CAlg}^{\text{ani}}$ as $\text{FFree}_A = A \otimes_{\mathbb{Z}} \text{FFree}_{\mathbb{Z}}$ and then animate this to get Mod_A .