

# Group Theory

Arifa Alam

November 25, 2024

## 1 Basics

Laws of compositions:

- **Associative Law:**  $(ab)c = a(bc)$
- **Commutative Law:**  $ab = ba$

These two laws shown using the product operation. However, a group may have other operation such as addition.

Let  $g \circ f$  denote as the composition of two functions  $g$  and  $f$ . Then for functions  $g$ ,  $f$ , and  $h$  the associativity holds but comutativity necessearly doesn't (take matrix multiplication, for example). In other words–

$$(g \circ f) \circ h = g \circ (f \circ h)$$

Identity of law of compostions is defined as–

$$ae = ea = a$$

(and yes,  $a = e$ )

### Definition 1.1: Groups

A group is a set  $G$  which should satisfy these following properties–

- Associative for every element in  $G$ .
- Exist identity element, unique.
- Exist inverse for all element in  $G$ .

A group is *abelian* if its law of composton is commutative. Such as  $\mathbb{Z}^+$ ,  $\mathbb{R}^+$ ,  $\mathbb{R}^\times$ ,  $\mathbb{C}^+$ ,  $\mathbb{C}^\times$   
The  $\times$  and  $+$  signs are defining the operations, not the set.

*Order* of a group is defined as its cardinality,  $|G|$ .

### Theorem 1.1: Cancellation Law

Let  $a$ ,  $b$ , and  $c$  be elements of  $G$  whose law of composition is written multiplicatively. If  $ab = ac$  or  $ba = ca$  then  $b = c$ . And if  $ab = a$  and  $ba = a$  then  $b = 1$ .

*Proof.* Just multiply with  $a^{-1}$  in both left and right side of the term.  $\square$

### Definition 1.2: Subgroup

A subset  $H$  of a group  $G$  must have the following properties:

- *Closure*: If  $a \in H \implies ab \in H$ .
- *Identity*:  $1 \in H$
- *Inverse*:  $a \in H \implies a^{-1} \in H$

A subgroup is proper subgroup if it's not trivial subgroup. And they are–

- The group itself
- The subgroup only with identity

Some examples of subgroup are–

- *circle group*: Let group  $G = \mathbb{C}$  for which absolute value of every element in  $\mathbb{C}$  is equals to 1. The unit circle of the plane is the subgroup of  $G$ , which is the circle group.
- *special linear group*:  $G$  is the set of  $n \times n$  matrices. Matrices with determinant 1 are subgroup of  $G$ , which is a special linear group also known as  $SL_n$ . And  $G$  is known as *general linear group*  $GL_n$

### Theorem 1.2

Let  $S$  be a subgraoup of the additive group of  $\mathbb{Z}^+$ . Either  $S$  is the trivial subgroup 0, or else it has to be in the form  $\mathbb{Z}a$ , where  $a$  is the smallest positive integer in  $S$ .