

Participants with Drug-Related Adverse Events
(Incidence > 0 % in One or More Treatment Groups)
Weeks 0 to 12
All Participants as Treated

Low Dose		Placebo		Difference in % Low Dose vs. Placebo		
n	(%)	n	(%)	Estimate	(95% CI)	p-value
Participants in population ⁸⁴		86			(-4.4, 4.3)	
with one or more drug-related adverse events ⁷³	(86.9)	44	(51.2)	35.7	(22.4, 48.0)	<0.001
with no drug-related adverse events ¹¹	(13.1)	42	(48.8)	-35.7	(-48.0, -22.4)	1.000
Cardiac disorders⁷	(8.3)	6	(7.0)	1.4	(-7.3, 10.2)	0.370
Atrial fibrillation ⁰	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Atrial flutter ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Atrioventricular block first degree ⁰	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Atrioventricular block second degree ⁰	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Bradycardia ⁰	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Bundle branch block right ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Cardiac failure congestive ⁰	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Myocardial infarction ¹	(1.2)	2	(2.3)	-1.1	(-7.1, 4.4)	0.712
Palpitations ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Sinus arrhythmia ⁰	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Sinus bradycardia ²	(2.4)	2	(2.3)	0.1	(-6.0, 6.2)	0.491

Participants with Drug-Related Adverse Events
(Incidence > 0 % in One or More Treatment Groups)
Weeks 0 to 12

All Participants as Treated

Low Dose		Placebo		Difference in % Low Dose vs. Placebo		
n	(%)	n	(%)	Estimate	(95% CI)	p-value
Supraventricular extrasystoles1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Ventricular extrasystoles1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Wolff-parkinson-white syndrome1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Congenital, familial and genetic disorders1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Ventricular septal defect1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Ear and labyrinth disorders2	(2.4)	0	(0.0)	2.4	(-2.0, 8.3)	0.076
Tinnitus1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Vertigo1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Eye disorders1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Vision blurred1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Gastrointestinal disorders8	(9.5)	4	(4.7)	4.9	(-3.2, 13.7)	0.108
Abdominal pain1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Diarrhoea3	(3.6)	3	(3.5)	0.1	(-6.7, 7.0)	0.488
Dyspepsia1	(1.2)	1	(1.2)	0.0	(-5.2, 5.4)	0.493
Gastrooesophageal reflux disease0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Nausea3	(3.6)	0	(0.0)	3.6	(-0.8, 10.0)	0.039

Participants with Drug-Related Adverse Events (Incidence > 0 % in One or More Treatment Groups) Weeks 0 to 12 All Participants as Treated

Low Dose		Placebo		Difference in % Low Dose vs. Placebo		
n	(%)	n	(%)	Estimate	(95% CI)	p-value
Vomiting2	(2.4)	0	(0.0)	2.4	(-2.0, 8.3)	0.076

General disorders and administration site conditions ⁴³	(51.2)	18	(20.9)	30.3	(16.1, 43.4)	<0.001
Application site bleeding ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Application site dermatitis ⁹	(10.7)	5	(5.8)	4.9	(-3.7, 14.1)	0.123
Application site desquamation ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Application site discolouration ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Application site erythema ¹²	(14.3)	3	(3.5)	10.8	(2.5, 20.4)	0.007
Application site induration ⁰	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Application site irritation ⁹	(10.7)	3	(3.5)	7.2	(-0.6, 16.1)	0.033
Application site pruritus ²²	(26.2)	6	(7.0)	19.2	(8.4, 30.5)	<0.001
Application site reaction ⁰	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Application site swelling ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Application site urticaria ²	(2.4)	0	(0.0)	2.4	(-2.0, 8.3)	0.076
Application site vesicles ⁴	(4.8)	1	(1.2)	3.6	(-2.1, 10.6)	0.083
Application site warmth ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Asthenia ⁰	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838

Participants with Drug-Related Adverse Events
(Incidence > 0 % in One or More Treatment Groups)
Weeks 0 to 12
All Participants as Treated

Low Dose		Placebo		Difference in % Low Dose vs. Placebo		
n	(%)	n	(%)	Estimate	(95% CI)	p-value
Chills ¹	(1.2)	1	(1.2)	0.0	(-5.2, 5.4)	0.493
Fatigue ²	(2.4)	1	(1.2)	1.2	(-4.2, 7.3)	0.274

Malaise1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Oedema2	(2.4)	0	(0.0)	2.4	(-2.0, 8.3)	0.076
Pain1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Injury, poisoning and procedural complications 2	(2.4)	0	(0.0)	2.4	(-2.0, 8.3)	0.076
Fall1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Skin laceration1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Wound1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Investigations2	(2.4)	4	(4.7)	-2.3	(-9.3, 4.2)	0.788
Blood creatine phosphokinase increased0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Body temperature increased1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Electrocardiogram st segment depression1	(1.2)	1	(1.2)	0.0	(-5.2, 5.4)	0.493
Heart rate increased0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Heart rate irregular0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Metabolism and nutrition disorders0	(0.0)	3	(3.5)	-3.5	(-9.8, 1.0)	0.957

Participants with Drug-Related Adverse Events
 (Incidence > 0 % in One or More Treatment Groups)
 Weeks 0 to 12
 All Participants as Treated

Low Dose		Placebo		Difference in % Low Dose vs. Placebo		
n	(%)	n	(%)	Estimate	(95% CI)	p-value
Decreased appetite0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Food craving0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Increased appetite0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838

Musculoskeletal and connective tissue disorders ⁰	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Shoulder pain ⁰	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Nervous system disorders ¹²	(14.3)	5	(5.8)	8.5	(-0.6, 18.3)	0.033
Balance disorder ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Complex partial seizures ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Coordination abnormal ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Dizziness ⁶	(7.1)	2	(2.3)	4.8	(-1.9, 12.7)	0.070
Headache ¹	(1.2)	2	(2.3)	-1.1	(-7.1, 4.4)	0.712
Lethargy ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Paraesthesia oral ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Somnolence ⁰	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Stupor ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Syncope ⁴	(4.8)	0	(0.0)	4.8	(0.4, 11.6)	0.021

Participants with Drug-Related Adverse Events
 (Incidence > 0 % in One or More Treatment Groups)
 Weeks 0 to 12
 All Participants as Treated

Low Dose		Placebo		Difference in % Low Dose vs. Placebo		
n	(%)	n	(%)	Estimate	(95% CI)	p-value
Transient ischaemic attack ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Psychiatric disorders ⁹	(10.7)	2	(2.3)	8.4	(1.1, 17.1)	0.013
Agitation ²	(2.4)	0	(0.0)	2.4	(-2.0, 8.3)	0.076
Anxiety ³	(3.6)	0	(0.0)	3.6	(-0.8, 10.0)	0.039
Confusional state ²	(2.4)	1	(1.2)	1.2	(-4.2, 7.3)	0.274
Depressed mood ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Irritability ¹	(1.2)	1	(1.2)	0.0	(-5.2, 5.4)	0.493
Restlessness ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Renal and urinary disorders ¹	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156

Enuresis1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Reproductive system and breast disorders0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Pelvic pain0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Respiratory, thoracic and mediastinal disorders0	(0.0)	2	(2.3)	-2.3	(-8.1, 2.1)	0.920
Dyspnoea0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Emphysema0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Skin and subcutaneous tissue disorders39	(46.4)	17	(19.8)	26.7	(12.7, 39.8)	<0.001

Participants with Drug-Related Adverse Events
(Incidence > 0 % in One or More Treatment Groups)
Weeks 0 to 12
All Participants as Treated

Low Dose		Placebo		Difference in % Low Dose vs. Placebo		
n	(%)	n	(%)	Estimate	(95% CI)	p-value
Blister5	(6.0)	0	(0.0)	6.0	(1.5, 13.2)	0.011
Cold sweat0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Dermatitis contact1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Erythema13	(15.5)	9	(10.5)	5.0	(-5.3, 15.6)	0.166
Hyperhidrosis 4	(4.8)	1	(1.2)	3.6	(-2.1, 10.6)	0.083
Pruritus21	(25.0)	7	(8.1)	16.9	(5.9, 28.2)	0.002
Pruritus generalised1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Rash11	(13.1)	3	(3.5)	9.6	(1.5, 19.0)	0.012
Rash erythematous2	(2.4)	0	(0.0)	2.4	(-2.0, 8.3)	0.076
Rash pruritic1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Skin exfoliation1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Skin irritation6	(7.1)	2	(2.3)	4.8	(-1.9, 12.7)	0.070
Skin ulcer0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838
Urticaria1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156
Vascular disorders2	(2.4)	1	(1.2)	1.2	(-4.2, 7.3)	0.274
Hypertension1	(1.2)	0	(0.0)	1.2	(-3.1, 6.5)	0.156

Participants with Drug-Related Adverse Events
(Incidence > 0 % in One or More Treatment Groups)
Weeks 0 to 12
All Participants as Treated

Low Dose		Placebo		Difference in % Low Dose vs. Placebo		
n	(%)	n	(%)	Estimate	(95% CI)	p-value
Hypotension1	(1.2)	1	(1.2)	0.0	(-5.2, 5.4)	0.493
Orthostatic hypotension0	(0.0)	1	(1.2)	-1.2	(-6.3, 3.3)	0.838

Every participant is counted a single time for each applicable row and column.

A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding.

Adverse event terms are from MedDRA Version 24.0.Source: [CDISCpilot: adam-adsl; adae]