

Participants with Drug-Related Adverse Events
(Incidence > 0 % in One or More Treatment Groups)
Weeks 0 to 12
All Participants as Treated

Low Dose			Placebo		
n	(%)	Mean Duration (SE)	n	(%)	Mean Duration (SE)
Participants in population84			86		
with one or more drug-related adverse events73	(86.9)	27.2 (3.2)	44	(51.2)	29.0 (3.5)
with no drug- related adverse events11	(13.1)		42	(48.8)	
Cardiac disorders7	(8.3)	16.1 (3.5)	6	(7.0)	27.1 (5.9)
Atrial fibrillation0	(0.0)		1	(1.2)	6.0
Atrial flutter1	(1.2)		0	(0.0)	
Atrioventricular block first degree0	(0.0)		1	(1.2)	
Atrioventricular block second degree0	(0.0)		1	(1.2)	
Bradycardia0	(0.0)		1	(1.2)	44.5 (8.4)
Bundle branch block right1	(1.2)		0	(0.0)	
Cardiac failure congestive0	(0.0)		1	(1.2)	
Myocardial infarction1	(1.2)	14.0 (0.0)	2	(2.3)	1.0 (0.0)
Palpitations1	(1.2)	1.0	0	(0.0)	
Sinus arrhythmia0	(0.0)		1	(1.2)	29.0 (0.0)
Sinus bradycardia2	(2.4)		2	(2.3)	
Supraventricular extrasystoles1	(1.2)	17.0 (0.0)	0	(0.0)	
Ventricular extrasystoles1	(1.2)	5.0 (0.0)	0	(0.0)	
Wolff-parkinson- white syndrome1	(1.2)	36.0 (0.0)	0	(0.0)	
Congenital, familial and genetic disorders1	(1.2)		0	(0.0)	

Ventricular septal defect1	(1.2)		0	(0.0)	
Ear and labyrinth disorders2	(2.4)	3.0	0	(0.0)	

Participants with Drug-Related Adverse Events
(Incidence > 0 % in One or More Treatment Groups)
Weeks 0 to 12
All Participants as Treated

Low Dose			Placebo		
n	(%)	Mean Duration (SE)	n	(%)	Mean Duration (SE)
Tinnitus1	(1.2)		0	(0.0)	
Vertigo1	(1.2)	3.0	0	(0.0)	
Eye disorders1	(1.2)	1.0	0	(0.0)	
Vision blurred1	(1.2)	1.0	0	(0.0)	
Gastrointestinal disorders8	(9.5)	15.1 (5.1)	4	(4.7)	3.6 (0.9)
Abdominal pain1	(1.2)	9.0	0	(0.0)	
Diarrhoea3	(3.6)	17.0 (7.1)	3	(3.5)	4.2 (1.0)
Dyspepsia1	(1.2)	13.0 (0.0)	1	(1.2)	1.0
Gastrooesophageal reflux disease0	(0.0)		1	(1.2)	
Nausea3	(3.6)	24.0 (13.9)	0	(0.0)	
Vomiting2	(2.4)	1.0 (0.0)	0	(0.0)	
General disorders and administration site conditions43	(51.2)	34.8 (3.4)	18	(20.9)	19.1 (3.6)
Application site bleeding1	(1.2)	11.0	0	(0.0)	
Application site dermatitis9	(10.7)	43.8 (9.8)	5	(5.8)	1.3 (0.2)
Application site desquamation1	(1.2)		0	(0.0)	
Application site discolouration1	(1.2)		0	(0.0)	
Application site erythema12	(14.3)	39.0 (7.0)	3	(3.5)	
Application site induration0	(0.0)		1	(1.2)	1.0
Application site irritation9	(10.7)	52.2 (11.6)	3	(3.5)	23.2 (8.3)
Application site pruritus22	(26.2)	34.2 (3.7)	6	(7.0)	32.4 (9.8)
Application site reaction0	(0.0)		1	(1.2)	52.0 (0.0)

Application site swelling1	(1.2)		0	(0.0)	
----------------------------	-------	--	---	-------	--

Participants with Drug-Related Adverse Events
 (Incidence > 0 % in One or More Treatment Groups)
 Weeks 0 to 12
 All Participants as Treated

Low Dose			Placebo		
n	(%)	Mean Duration (SE)	n	(%)	Mean Duration (SE)
Application site urticaria2	(2.4)		0	(0.0)	
Application site vesicles4	(4.8)	12.6 (4.6)	1	(1.2)	27.0 (0.0)
Application site warmth1	(1.2)	12.0 (0.0)	0	(0.0)	
Asthenia0	(0.0)		1	(1.2)	
Chills1	(1.2)	7.0 (0.0)	1	(1.2)	1.0 (0.0)
Fatigue2	(2.4)		1	(1.2)	18.0 (0.0)
Malaise1	(1.2)	23.0 (0.0)	0	(0.0)	
Oedema2	(2.4)		0	(0.0)	
Pain1	(1.2)	2.0	0	(0.0)	
Injury, poisoning and procedural complications2	(2.4)	4.5 (2.0)	0	(0.0)	
Fall1	(1.2)	1.0	0	(0.0)	
Skin laceration1	(1.2)	1.0	0	(0.0)	
Wound1	(1.2)	8.0 (0.0)	0	(0.0)	
Investigations2	(2.4)	10.7 (4.3)	4	(4.7)	5.0 (1.9)
Blood creatine phosphokinase increased0	(0.0)		1	(1.2)	14.0 (0.0)
Body temperature increased1	(1.2)	2.0	0	(0.0)	
Electrocardiogram st segment depression1	(1.2)	15.0 (0.0)	1	(1.2)	
Heart rate increased0	(0.0)		1	(1.2)	4.0 (0.0)
Heart rate irregular0	(0.0)		1	(1.2)	1.0 (0.0)
Metabolism and nutrition disorders0	(0.0)		3	(3.5)	25.0 (0.0)
Decreased appetite0	(0.0)		1	(1.2)	

Food craving0	(0.0)		1	(1.2)	
---------------	-------	--	---	-------	--

Participants with Drug-Related Adverse Events
(Incidence > 0 % in One or More Treatment Groups)
Weeks 0 to 12
All Participants as Treated

Low Dose			Placebo		
n	(%)	Mean Duration (SE)	n	(%)	Mean Duration (SE)
Increased appetite0	(0.0)		1	(1.2)	25.0 (0.0)
Musculoskeletal and connective tissue disorders0	(0.0)		1	(1.2)	
Shoulder pain0	(0.0)		1	(1.2)	
Nervous system disorders12	(14.3)	7.2 (2.5)	5	(5.8)	19.0 (8.1)
Balance disorder1	(1.2)	38.0 (0.0)	0	(0.0)	
Complex partial seizures1	(1.2)	2.0	0	(0.0)	
Coordination abnormal1	(1.2)	1.0	0	(0.0)	
Dizziness6	(7.1)	3.6 (1.4)	2	(2.3)	2.3 (1.3)
Headache1	(1.2)	1.0	2	(2.3)	
Lethargy1	(1.2)		0	(0.0)	
Paraesthesia oral1	(1.2)		0	(0.0)	
Somnolence0	(0.0)		1	(1.2)	44.0 (0.0)
Stupor1	(1.2)	1.0	0	(0.0)	
Syncope4	(4.8)	2.0 (0.7)	0	(0.0)	
Transient ischaemic attack1	(1.2)	1.0	0	(0.0)	
Psychiatric disorders9	(10.7)	2.7 (0.3)	2	(2.3)	196.0 (0.0)
Agitation2	(2.4)		0	(0.0)	
Anxiety3	(3.6)	2.3 (0.3)	0	(0.0)	
Confusional state2	(2.4)	1.0	1	(1.2)	
Depressed mood1	(1.2)		0	(0.0)	
Irritability1	(1.2)		1	(1.2)	196.0 (0.0)
Restlessness1	(1.2)	4.0 (0.0)	0	(0.0)	
Renal and urinary disorders1	(1.2)		0	(0.0)	

Enuresis1	(1.2)		0	(0.0)	
Reproductive system and breast disorders0	(0.0)		1	(1.2)	101.0 (0.0)

Participants with Drug-Related Adverse Events
(Incidence > 0 % in One or More Treatment Groups)
Weeks 0 to 12
All Participants as Treated

Low Dose			Placebo		
n	(%)	Mean Duration (SE)	n	(%)	Mean Duration (SE)
Pelvic pain0	(0.0)		1	(1.2)	101.0 (0.0)
Respiratory, thoracic and mediastinal disorders0	(0.0)		2	(2.3)	1.0
Dyspnoea0	(0.0)		1	(1.2)	1.0
Emphysema0	(0.0)		1	(1.2)	
Skin and subcutaneous tissue disorders39	(46.4)	41.6 (8.3)	17	(19.8)	40.2 (7.3)
Blister5	(6.0)	16.4 (4.1)	0	(0.0)	
Cold sweat0	(0.0)		1	(1.2)	1.0 (0.0)
Dermatitis contact1	(1.2)	15.0 (0.0)	0	(0.0)	
Erythema13	(15.5)	48.3 (11.5)	9	(10.5)	17.8 (2.7)
Hyperhidrosis4	(4.8)	34.2 (16.9)	1	(1.2)	
Pruritus21	(25.0)	96.2 (28.5)	7	(8.1)	101.5 (17.0)
Pruritus generalised1	(1.2)	1.2 (0.2)	0	(0.0)	
Rash11	(13.1)	21.0 (1.9)	3	(3.5)	47.0 (10.1)
Rash erythematous2	(2.4)		0	(0.0)	
Rash pruritic1	(1.2)	5.0 (0.0)	0	(0.0)	
Skin exfoliation1	(1.2)	5.0 (0.0)	0	(0.0)	
Skin irritation6	(7.1)	22.4 (4.4)	2	(2.3)	
Skin ulcer0	(0.0)		1	(1.2)	15.0 (0.0)
Urticaria1	(1.2)		0	(0.0)	
Vascular disorders2	(2.4)	8.0 (6.0)	1	(1.2)	37.3 (17.7)

Participants with Drug-Related Adverse Events
(Incidence > 0 % in One or More Treatment Groups)

Weeks 0 to 12
All Participants as Treated

Low Dose			Placebo		
n	(%)	Mean Duration (SE)	n	(%)	Mean Duration (SE)
Hypertension1	(1.2)	2.0	0	(0.0)	
Hypotension1	(1.2)	14.0	1	(1.2)	2.0
Orthostatic hypotension0	(0.0)		1	(1.2)	55.0 (0.0)

Every participant is counted a single time for each applicable row and column.

A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding.

Adverse event terms are from MedDRA Version 24.0.Source: [CDISCpilot: adam-adsl; adae]