

Participants with Drug-Related Adverse Events
(Incidence ≥ 6 % in One or More Treatment Groups)
Weeks 0 to 12
All Participants as Treated

Low Dose		Placebo		Total	
n	(%)	n	(%)	n	(%)
Participants in population ⁸⁴		86		170	
with one or more drug-related adverse events ⁷³	(86.9)	44	(51.2)	117	(68.8)
with no drug-related adverse events ¹¹	(13.1)	42	(48.8)	53	(31.2)
Cardiac disorders⁷	(8.3)	6	(7.0)	13	(7.6)
Gastrointestinal disorders⁸	(9.5)	4	(4.7)	12	(7.1)
General disorders and administration site conditions⁴³	(51.2)	18	(20.9)	61	(35.9)
Application site dermatitis ⁹	(10.7)	5	(5.8)	14	(8.2)
Application site erythema ¹²	(14.3)	3	(3.5)	15	(8.8)
Application site irritation ⁹	(10.7)	3	(3.5)	12	(7.1)
Application site pruritus ²²	(26.2)	6	(7.0)	28	(16.5)
Nervous system disorders¹²	(14.3)	5	(5.8)	17	(10.0)
Dizziness ⁶	(7.1)	2	(2.3)	8	(4.7)
Psychiatric disorders⁹	(10.7)	2	(2.3)	11	(6.5)
Skin and subcutaneous tissue disorders³⁹	(46.4)	17	(19.8)	56	(32.9)
Blister ⁵	(6.0)	0	(0.0)	5	(2.9)
Erythema ¹³	(15.5)	9	(10.5)	22	(12.9)
Pruritus ²¹	(25.0)	7	(8.1)	28	(16.5)
Rash ¹¹	(13.1)	3	(3.5)	14	(8.2)
Skin irritation ⁶	(7.1)	2	(2.3)	8	(4.7)

Every participant is counted a single time for each applicable row and column.

A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding.

Adverse event terms are from MedDRA Version 24.0.Source: [CDISCpilot: adam-adsl; adae]